

Table 7.69C Types of Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Standard Errors of Numbers in Thousands, Annual Averages Based on 2004-2005 and 2006-2007

| Drug | Total ¹ | | PREGNANCY STATUS | | | |
|--|--------------------|-----------|------------------|-----------|--------------|-----------|
| | | | Pregnant | | Not Pregnant | |
| | 2004-2005 | 2006-2007 | 2004-2005 | 2006-2007 | 2004-2005 | 2006-2007 |
| ILLCIT DRUGS² | 132 | 127 | 13 | 18 | 131 | 125 |
| Marijuana and Hashish | 106 | 100 | 11 | 15 | 106 | 97 |
| Cocaine | 43 | 42 | 2 | 6 | 43 | 41 |
| Crack | 25 | 18 | 1 | 6 | 25 | 17 |
| Heroin | 8 | 10 | 2 | 1 | 8 | 9 |
| Hallucinogens | 19 | 29 | 2 | 2 | 18 | 28 |
| LSD | 5 | 6 | 2 | 0 | 5 | 6 |
| PCP | 3 | 3 | * | 0 | 3 | 3 |
| Ecstasy | 15 | 25 | 1 | 1 | 15 | 25 |
| Inhalants | 13 | 18 | 1 | 4 | 13 | 17 |
| Nonmedical Use of Psychotherapeutics ^{3,4} | 84 | 79 | 7 | 8 | 83 | 78 |
| Pain Relievers | 68 | 65 | 7 | 4 | 68 | 64 |
| OxyContin [®] | 17 | 12 | 1 | 1 | 17 | 12 |
| Tranquilizers | 44 | 41 | 2 | 3 | 44 | 41 |
| Stimulants ⁴ | 39 | 34 | 2 | 7 | 39 | 34 |
| Methamphetamine ⁴ | 29 | 26 | 1 | 0 | 29 | 26 |
| Sedatives | 17 | 20 | 1 | 1 | 17 | 20 |
| ILLCIT DRUGS OTHER THAN MARIJUANA² | 94 | 89 | 8 | 11 | 93 | 88 |

*Low precision; no estimate reported.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004, 2005, 2006, and 2007.

Table 7.69D Types of Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Standard Errors of Percentages, Annual Averages Based on 2004-2005 and 2006-2007

| Drug | Total ¹ | | PREGNANCY STATUS | | | |
|--|--------------------|-----------|------------------|-----------|--------------|-----------|
| | | | Pregnant | | Not Pregnant | |
| | 2004-2005 | 2006-2007 | 2004-2005 | 2006-2007 | 2004-2005 | 2006-2007 |
| ILLCIT DRUGS² | 0.20 | 0.19 | 0.51 | 0.67 | 0.21 | 0.20 |
| Marijuana and Hashish | 0.17 | 0.16 | 0.44 | 0.56 | 0.18 | 0.16 |
| Cocaine | 0.07 | 0.07 | 0.09 | 0.24 | 0.07 | 0.07 |
| Crack | 0.04 | 0.03 | 0.05 | 0.23 | 0.04 | 0.03 |
| Heroin | 0.01 | 0.02 | 0.07 | 0.05 | 0.01 | 0.02 |
| Hallucinogens | 0.03 | 0.05 | 0.09 | 0.06 | 0.03 | 0.05 |
| LSD | 0.01 | 0.01 | 0.07 | 0.02 | 0.01 | 0.01 |
| PCP | 0.00 | 0.01 | * | 0.02 | 0.01 | 0.01 |
| Ecstasy | 0.02 | 0.04 | 0.06 | 0.04 | 0.02 | 0.04 |
| Inhalants | 0.02 | 0.03 | 0.06 | 0.14 | 0.02 | 0.03 |
| Nonmedical Use of Psychotherapeutics ^{3,4} | 0.13 | 0.12 | 0.29 | 0.31 | 0.14 | 0.13 |
| Pain Relievers | 0.11 | 0.10 | 0.28 | 0.17 | 0.11 | 0.11 |
| OxyContin [®] | 0.03 | 0.02 | 0.04 | 0.02 | 0.03 | 0.02 |
| Tranquilizers | 0.07 | 0.07 | 0.10 | 0.11 | 0.07 | 0.07 |
| Stimulants ⁴ | 0.06 | 0.06 | 0.08 | 0.26 | 0.07 | 0.06 |
| Methamphetamine ⁴ | 0.05 | 0.04 | 0.06 | 0.02 | 0.05 | 0.04 |
| Sedatives | 0.03 | 0.03 | 0.04 | 0.03 | 0.03 | 0.03 |
| ILLCIT DRUGS OTHER THAN MARIJUANA² | 0.15 | 0.14 | 0.31 | 0.41 | 0.15 | 0.14 |

*Low precision; no estimate reported.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004, 2005, 2006, and 2007.

Table 7.69P Types of Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: P Values from Tests of Differences of Percentages, 2006-2007 Versus 2004-2005

| Drug | Total ¹ | | PREGNANCY STATUS | | | |
|--|--------------------|-----------|------------------|-----------|--------------|-----------|
| | | | Pregnant | | Not Pregnant | |
| | 2004-2005 | 2006-2007 | 2004-2005 | 2006-2007 | 2004-2005 | 2006-2007 |
| ILLCIT DRUGS² | 0.7234 | | 0.1237 | | 0.5156 | |
| Marijuana and Hashish | 0.4811 | | 0.1982 | | 0.3099 | |
| Cocaine | 0.2930 | | 0.6008 | | 0.2903 | |
| Crack | 0.0738 | | 0.3522 | | 0.0476 | |
| Heroin | 0.4898 | | 0.6528 | | 0.4387 | |
| Hallucinogens | 0.0418 | | 0.3522 | | 0.0300 | |
| LSD | 0.6189 | | 0.1720 | | 0.4346 | |
| PCP | 0.6771 | | * | | 0.7095 | |
| Ecstasy | 0.0324 | | 0.6419 | | 0.0235 | |
| Inhalants | 0.0934 | | 0.2232 | | 0.1213 | |
| Nonmedical Use of Psychotherapeutics ^{3,4} | 0.3012 | | 0.7013 | | 0.3112 | |
| Pain Relievers | 0.4561 | | 0.2087 | | 0.5065 | |
| OxyContin [®] | 0.1732 | | 0.5451 | | 0.2019 | |
| Tranquilizers | 0.9582 | | 0.7383 | | 0.9241 | |
| Stimulants ⁴ | 0.0961 | | 0.5338 | | 0.0825 | |
| Methamphetamine ⁴ | 0.1305 | | 0.2935 | | 0.1449 | |
| Sedatives | 0.5837 | | 0.7696 | | 0.5866 | |
| ILLCIT DRUGS OTHER THAN MARIJUANA² | 0.7004 | | 0.6016 | | 0.6633 | |

*Low precision; no estimate reported.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004, 2005, 2006, and 2007.

Table 7.70C Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Numbers in Thousands, Annual Averages Based on 2004-2005 and 2006-2007

| Demographic/Pregnancy Characteristic | Total ¹ | | PREGNANCY STATUS | | | |
|---|--------------------|-----------|------------------|-----------|--------------|-----------|
| | | | Pregnant | | Not Pregnant | |
| | 2004-2005 | 2006-2007 | 2004-2005 | 2006-2007 | 2004-2005 | 2006-2007 |
| TOTAL | 132 | 127 | 13 | 18 | 131 | 125 |
| AGE | | | | | | |
| 15-17 | 30 | 28 | 3 | 4 | 29 | 28 |
| 18-25 | 53 | 54 | 8 | 8 | 58 | 63 |
| 26-44 | 109 | 104 | 9 | 16 | 109 | 103 |
| HISPANIC ORIGIN AND RACE | | | | | | |
| Not Hispanic or Latino | 122 | 121 | 10 | 17 | 122 | 119 |
| White | 109 | 104 | 8 | 17 | 108 | 102 |
| Black or African American | 46 | 51 | 6 | 5 | 46 | 51 |
| American Indian or Alaska Native | 9 | 8 | * | * | 9 | 7 |
| Native Hawaiian or Other Pacific Islander | * | * | * | * | * | * |
| Asian | 12 | 23 | * | * | 12 | 23 |
| Two or More Races | 18 | 15 | * | * | 18 | 15 |
| Hispanic or Latino | 50 | 44 | 8 | 5 | 49 | 44 |
| TRIMESTER² | | | | | | |
| First | N/A | N/A | 8 | 7 | N/A | N/A |
| Second | N/A | N/A | 8 | 13 | N/A | N/A |
| Third | N/A | N/A | 6 | 10 | N/A | N/A |

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, based on data from original questions not including methamphetamine items added in 2005 and 2006.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004, 2005, 2006, and 2007.

Table 7.70D Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Percentages, Annual Averages Based on 2004-2005 and 2006-2007

| Demographic/Pregnancy Characteristic | Total ¹ | | PREGNANCY STATUS | | | |
|---|--------------------|-----------|------------------|-----------|--------------|-----------|
| | | | Pregnant | | Not Pregnant | |
| | 2004-2005 | 2006-2007 | 2004-2005 | 2006-2007 | 2004-2005 | 2006-2007 |
| TOTAL | 0.20 | 0.19 | 0.51 | 0.67 | 0.21 | 0.20 |
| AGE | | | | | | |
| 15-17 | 0.45 | 0.41 | 3.63 | 4.33 | 0.45 | 0.41 |
| 18-25 | 0.33 | 0.33 | 0.87 | 0.82 | 0.34 | 0.35 |
| 26-44 | 0.26 | 0.25 | 0.61 | 0.99 | 0.28 | 0.26 |
| HISPANIC ORIGIN AND RACE | | | | | | |
| Not Hispanic or Latino | 0.22 | 0.21 | 0.52 | 0.82 | 0.23 | 0.22 |
| White | 0.26 | 0.25 | 0.56 | 1.08 | 0.27 | 0.25 |
| Black or African American | 0.50 | 0.54 | 1.75 | 1.26 | 0.52 | 0.56 |
| American Indian or Alaska Native | 2.27 | 2.47 | * | * | 2.32 | 2.48 |
| Native Hawaiian or Other Pacific Islander | * | * | * | * | * | * |
| Asian | 0.38 | 0.71 | * | * | 0.40 | 0.75 |
| Two or More Races | 2.36 | 1.85 | * | * | 2.42 | 1.90 |
| Hispanic or Latino | 0.50 | 0.41 | 1.50 | 0.85 | 0.52 | 0.43 |
| TRIMESTER² | | | | | | |
| First | N/A | N/A | 1.09 | 0.83 | N/A | N/A |
| Second | N/A | N/A | 0.97 | 1.28 | N/A | N/A |
| Third | N/A | N/A | 0.64 | 1.27 | N/A | N/A |

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, based on data from original questions not including methamphetamine items added in 2005 and 2006.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004, 2005, 2006, and 2007.

Table 7.70P Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2006-2007 Versus 2004-2005

| Demographic/Pregnancy Characteristic | Total ¹ | | PREGNANCY STATUS | | | |
|---|--------------------|-----------|------------------|-----------|--------------|-----------|
| | | | Pregnant | | Not Pregnant | |
| | 2004-2005 | 2006-2007 | 2004-2005 | 2006-2007 | 2004-2005 | 2006-2007 |
| TOTAL | 0.7234 | | 0.1237 | | 0.5156 | |
| AGE | | | | | | |
| 15-17 | 0.0410 | | 0.0609 | | 0.0211 | |
| 18-25 | 0.9672 | | 0.8580 | | 0.8662 | |
| 26-44 | 0.9014 | | 0.2400 | | 0.7638 | |
| HISPANIC ORIGIN AND RACE | | | | | | |
| Not Hispanic or Latino | 0.5064 | | 0.0909 | | 0.7059 | |
| White | 0.8325 | | 0.0691 | | 0.9254 | |
| Black or African American | 0.1182 | | 0.6064 | | 0.1157 | |
| American Indian or Alaska Native | 0.6862 | | * | | 0.8042 | |
| Native Hawaiian or Other Pacific Islander | * | | * | | * | |
| Asian | 0.1871 | | * | | 0.2219 | |
| Two or More Races | 0.0411 | | * | | 0.0303 | |
| Hispanic or Latino | 0.0221 | | 0.9692 | | 0.0178 | |
| TRIMESTER² | | | | | | |
| First | N/A | | 0.2399 | | N/A | |
| Second | N/A | | 0.1111 | | N/A | |
| Third | N/A | | 0.4056 | | N/A | |

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, based on data from original questions not including methamphetamine items added in 2005 and 2006.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004, 2005, 2006, and 2007.

Table 7.71C Marijuana Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Numbers in Thousands, Annual Averages Based on 2004-2005 and 2006-2007

| Demographic/Pregnancy Characteristic | Total ¹ | | PREGNANCY STATUS | | | |
|---|--------------------|-----------|------------------|-----------|--------------|-----------|
| | | | Pregnant | | Not Pregnant | |
| | 2004-2005 | 2006-2007 | 2004-2005 | 2006-2007 | 2004-2005 | 2006-2007 |
| TOTAL | 106 | 100 | 11 | 15 | 106 | 97 |
| AGE | | | | | | |
| 15-17 | 25 | 25 | 3 | 4 | 25 | 24 |
| 18-25 | 47 | 50 | 7 | 7 | 50 | 55 |
| 26-44 | 87 | 77 | 8 | 13 | 87 | 76 |
| HISPANIC ORIGIN AND RACE | | | | | | |
| Not Hispanic or Latino | 99 | 95 | 8 | 15 | 99 | 92 |
| White | 90 | 82 | 8 | 14 | 90 | 80 |
| Black or African American | 36 | 43 | 3 | 5 | 36 | 43 |
| American Indian or Alaska Native | 8 | 6 | * | * | 8 | 6 |
| Native Hawaiian or Other Pacific Islander | 3 | 3 | * | * | 3 | 3 |
| Asian | 10 | 15 | * | * | 10 | 15 |
| Two or More Races | 12 | 14 | * | * | 12 | 14 |
| Hispanic or Latino | 36 | 33 | 7 | 3 | 35 | 33 |
| TRIMESTER² | | | | | | |
| First | N/A | N/A | 7 | 6 | N/A | N/A |
| Second | N/A | N/A | 8 | 11 | N/A | N/A |
| Third | N/A | N/A | 4 | 7 | N/A | N/A |

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004, 2005, 2006, and 2007.

Table 7.71D Marijuana Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Percentages, Annual Averages Based on 2004-2005 and 2006-2007

| Demographic/Pregnancy Characteristic | Total ¹ | | PREGNANCY STATUS | | | |
|---|--------------------|-----------|------------------|-----------|--------------|-----------|
| | | | Pregnant | | Not Pregnant | |
| | 2004-2005 | 2006-2007 | 2004-2005 | 2006-2007 | 2004-2005 | 2006-2007 |
| TOTAL | 0.17 | 0.16 | 0.44 | 0.56 | 0.18 | 0.16 |
| AGE | | | | | | |
| 15-17 | 0.38 | 0.36 | 3.50 | 4.00 | 0.39 | 0.36 |
| 18-25 | 0.29 | 0.31 | 0.73 | 0.67 | 0.31 | 0.32 |
| 26-44 | 0.22 | 0.19 | 0.55 | 0.81 | 0.23 | 0.20 |
| HISPANIC ORIGIN AND RACE | | | | | | |
| Not Hispanic or Latino | 0.19 | 0.17 | 0.42 | 0.69 | 0.20 | 0.18 |
| White | 0.22 | 0.20 | 0.50 | 0.91 | 0.23 | 0.21 |
| Black or African American | 0.39 | 0.46 | 0.99 | 1.16 | 0.41 | 0.48 |
| American Indian or Alaska Native | 1.98 | 2.01 | * | * | 2.02 | 2.09 |
| Native Hawaiian or Other Pacific Islander | 1.56 | 1.05 | * | * | 1.62 | 1.10 |
| Asian | 0.32 | 0.47 | * | * | 0.33 | 0.51 |
| Two or More Races | 1.68 | 1.79 | * | * | 1.73 | 1.83 |
| Hispanic or Latino | 0.37 | 0.31 | 1.43 | 0.52 | 0.38 | 0.33 |
| TRIMESTER² | | | | | | |
| First | N/A | N/A | 0.93 | 0.78 | N/A | N/A |
| Second | N/A | N/A | 0.94 | 1.11 | N/A | N/A |
| Third | N/A | N/A | 0.39 | 0.88 | N/A | N/A |

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004, 2005, 2006, and 2007.

Table 7.71P Marijuana Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2006-2007 Versus 2004-2005

| Demographic/Pregnancy Characteristic | Total ¹ | | PREGNANCY STATUS | | | |
|---|--------------------|-----------|------------------|-----------|--------------|-----------|
| | | | Pregnant | | Not Pregnant | |
| | 2004-2005 | 2006-2007 | 2004-2005 | 2006-2007 | 2004-2005 | 2006-2007 |
| TOTAL | 0.4811 | | 0.1982 | | 0.3099 | |
| AGE | | | | | | |
| 15-17 | 0.0731 | | 0.1791 | | 0.0360 | |
| 18-25 | 0.5674 | | 0.6789 | | 0.7180 | |
| 26-44 | 0.3458 | | 0.4080 | | 0.2649 | |
| HISPANIC ORIGIN AND RACE | | | | | | |
| Not Hispanic or Latino | 0.9752 | | 0.1000 | | 0.7249 | |
| White | 0.5793 | | 0.1247 | | 0.3862 | |
| Black or African American | 0.1376 | | 0.5670 | | 0.1765 | |
| American Indian or Alaska Native | 0.9626 | | * | | 0.9974 | |
| Native Hawaiian or Other Pacific Islander | 0.4116 | | * | | 0.4349 | |
| Asian | 0.5732 | | * | | 0.5497 | |
| Two or More Races | 0.3420 | | * | | 0.2681 | |
| Hispanic or Latino | 0.1075 | | 0.6167 | | 0.1010 | |
| TRIMESTER² | | | | | | |
| First | N/A | | 0.7921 | | N/A | |
| Second | N/A | | 0.2729 | | N/A | |
| Third | N/A | | 0.7911 | | N/A | |

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004, 2005, 2006, and 2007.

Table 7.72C Tobacco Product and Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Standard Errors of Numbers in Thousands, Annual Averages Based on 2004-2005 and 2006-2007

| Substance | Total ¹ | | PREGNANCY STATUS | | | |
|-------------------------------------|--------------------|-----------|------------------|-----------|--------------|-----------|
| | | | Pregnant | | Not Pregnant | |
| | 2004-2005 | 2006-2007 | 2004-2005 | 2006-2007 | 2004-2005 | 2006-2007 |
| TOBACCO PRODUCTS² | 255 | 256 | 30 | 34 | 250 | 251 |
| Cigarettes | 251 | 250 | 29 | 33 | 246 | 245 |
| Smokeless Tobacco | 20 | 20 | 2 | 2 | 20 | 19 |
| Cigars | 68 | 70 | 6 | 8 | 67 | 69 |
| Pipe Tobacco | 23 | 19 | 1 | 1 | 23 | 19 |
| ALCOHOL | 364 | 394 | 32 | 31 | 360 | 390 |
| Binge Alcohol Use ³ | 213 | 236 | 15 | 14 | 211 | 234 |
| Heavy Alcohol Use ³ | 90 | 98 | 6 | 6 | 90 | 97 |

*Low precision; no estimate reported.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004, 2005, 2006, and 2007.

Table 7.72D Tobacco Product and Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Standard Errors of Percentages, Annual Averages Based on 2004-2005 and 2006-2007

| Substance | Total ¹ | | PREGNANCY STATUS | | | |
|-------------------------------------|--------------------|-----------|------------------|-----------|--------------|-----------|
| | | | Pregnant | | Not Pregnant | |
| | 2004-2005 | 2006-2007 | 2004-2005 | 2006-2007 | 2004-2005 | 2006-2007 |
| TOBACCO PRODUCTS² | 0.36 | 0.35 | 1.15 | 1.20 | 0.37 | 0.35 |
| Cigarettes | 0.35 | 0.34 | 1.14 | 1.20 | 0.36 | 0.35 |
| Smokeless Tobacco | 0.03 | 0.03 | 0.08 | 0.07 | 0.03 | 0.03 |
| Cigars | 0.11 | 0.11 | 0.22 | 0.30 | 0.11 | 0.11 |
| Pipe Tobacco | 0.04 | 0.03 | 0.05 | 0.05 | 0.04 | 0.03 |
| ALCOHOL | 0.38 | 0.42 | 1.19 | 1.12 | 0.39 | 0.43 |
| Binge Alcohol Use ³ | 0.30 | 0.33 | 0.58 | 0.53 | 0.31 | 0.34 |
| Heavy Alcohol Use ³ | 0.14 | 0.16 | 0.25 | 0.21 | 0.15 | 0.16 |

*Low precision; no estimate reported.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004, 2005, 2006, and 2007.

Table 7.72P Tobacco Product and Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: *P* Values from Tests of Differences of Percentages, 2006-2007 Versus 2004-2005

| Substance | Total ¹ | | PREGNANCY STATUS | | | |
|-------------------------------------|--------------------|-----------|------------------|-----------|--------------|-----------|
| | | | Pregnant | | Not Pregnant | |
| | 2004-2005 | 2006-2007 | 2004-2005 | 2006-2007 | 2004-2005 | 2006-2007 |
| TOBACCO PRODUCTS² | 0.0525 | | 0.7910 | | 0.0484 | |
| Cigarettes | 0.0283 | | 0.8722 | | 0.0257 | |
| Smokeless Tobacco | 0.3404 | | 0.5555 | | 0.4551 | |
| Cigars | 0.3200 | | 0.5771 | | 0.3519 | |
| Pipe Tobacco | 0.4098 | | 0.7802 | | 0.4002 | |
| ALCOHOL | 0.9638 | | 0.7810 | | 0.9133 | |
| Binge Alcohol Use ³ | 0.0930 | | 0.8354 | | 0.0840 | |
| Heavy Alcohol Use ³ | 0.2931 | | 0.9579 | | 0.2999 | |

*Low precision; no estimate reported.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004, 2005, 2006, and 2007.

Table 7.73C Cigarette Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Numbers in Thousands, Annual Averages Based on 2004-2005 and 2006-2007

| Demographic/Pregnancy Characteristic | Total ¹ | | PREGNANCY STATUS | | | |
|---|--------------------|-----------|------------------|-----------|--------------|-----------|
| | | | Pregnant | | Not Pregnant | |
| | 2004-2005 | 2006-2007 | 2004-2005 | 2006-2007 | 2004-2005 | 2006-2007 |
| TOTAL | 251 | 250 | 29 | 33 | 246 | 245 |
| AGE | | | | | | |
| 15-17 | 34 | 31 | 4 | 4 | 34 | 31 |
| 18-25 | 72 | 70 | 16 | 16 | 87 | 86 |
| 26-44 | 228 | 220 | 24 | 29 | 226 | 216 |
| HISPANIC ORIGIN AND RACE | | | | | | |
| Not Hispanic or Latino | 238 | 239 | 28 | 32 | 232 | 233 |
| White | 210 | 215 | 26 | 24 | 205 | 212 |
| Black or African American | 95 | 91 | 9 | 11 | 94 | 85 |
| American Indian or Alaska Native | 20 | 13 | * | * | 19 | 13 |
| Native Hawaiian or Other Pacific Islander | * | 14 | * | * | * | 13 |
| Asian | 36 | 47 | * | * | 36 | 43 |
| Two or More Races | 25 | 30 | * | * | 25 | 30 |
| Hispanic or Latino | 86 | 80 | 9 | 13 | 86 | 80 |
| TRIMESTER² | | | | | | |
| First | N/A | N/A | 18 | 24 | N/A | N/A |
| Second | N/A | N/A | 11 | 20 | N/A | N/A |
| Third | N/A | N/A | 16 | 12 | N/A | N/A |

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004, 2005, 2006, and 2007.

Table 7.73D Cigarette Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Percentages, Annual Averages Based on 2004-2005 and 2006-2007

| Demographic/Pregnancy Characteristic | Total ¹ | | PREGNANCY STATUS | | | |
|---|--------------------|-----------|------------------|-----------|--------------|-----------|
| | | | Pregnant | | Not Pregnant | |
| | 2004-2005 | 2006-2007 | 2004-2005 | 2006-2007 | 2004-2005 | 2006-2007 |
| TOTAL | 0.35 | 0.34 | 1.14 | 1.20 | 0.36 | 0.35 |
| AGE | | | | | | |
| 15-17 | 0.50 | 0.44 | 4.73 | 4.11 | 0.51 | 0.44 |
| 18-25 | 0.44 | 0.43 | 1.54 | 1.52 | 0.46 | 0.44 |
| 26-44 | 0.50 | 0.48 | 1.55 | 1.76 | 0.52 | 0.49 |
| HISPANIC ORIGIN AND RACE | | | | | | |
| Not Hispanic or Latino | 0.39 | 0.38 | 1.34 | 1.39 | 0.40 | 0.39 |
| White | 0.44 | 0.44 | 1.64 | 1.52 | 0.45 | 0.45 |
| Black or African American | 0.94 | 0.87 | 2.52 | 2.70 | 0.98 | 0.87 |
| American Indian or Alaska Native | 4.27 | 4.11 | * | * | 4.35 | 4.29 |
| Native Hawaiian or Other Pacific Islander | * | 4.62 | * | * | * | 4.77 |
| Asian | 1.08 | 1.36 | * | * | 1.13 | 1.36 |
| Two or More Races | 2.74 | 3.14 | * | * | 2.81 | 3.23 |
| Hispanic or Latino | 0.80 | 0.69 | 1.84 | 2.39 | 0.84 | 0.72 |
| TRIMESTER² | | | | | | |
| First | N/A | N/A | 2.38 | 2.55 | N/A | N/A |
| Second | N/A | N/A | 1.41 | 1.91 | N/A | N/A |
| Third | N/A | N/A | 1.76 | 1.43 | N/A | N/A |

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004, 2005, 2006, and 2007.

Table 7.73P Cigarette Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2006-2007 Versus 2004-2005

| Demographic/Pregnancy Characteristic | Total ¹ | | PREGNANCY STATUS | | | |
|---|--------------------|-----------|------------------|-----------|--------------|-----------|
| | | | Pregnant | | Not Pregnant | |
| | 2004-2005 | 2006-2007 | 2004-2005 | 2006-2007 | 2004-2005 | 2006-2007 |
| TOTAL | 0.0283 | | 0.8722 | | 0.0257 | |
| AGE | | | | | | |
| 15-17 | 0.0003 | | 0.7371 | | 0.0002 | |
| 18-25 | 0.0018 | | 0.1677 | | 0.0034 | |
| 26-44 | 0.5134 | | 0.6109 | | 0.4530 | |
| HISPANIC ORIGIN AND RACE | | | | | | |
| Not Hispanic or Latino | 0.2118 | | 0.8083 | | 0.2166 | |
| White | 0.4407 | | 0.4619 | | 0.5139 | |
| Black or African American | 0.1459 | | 0.9850 | | 0.1124 | |
| American Indian or Alaska Native | 0.8473 | | * | | 0.8876 | |
| Native Hawaiian or Other Pacific Islander | * | | * | | * | |
| Asian | 0.0915 | | * | | 0.1863 | |
| Two or More Races | 0.2758 | | * | | 0.2544 | |
| Hispanic or Latino | 0.0243 | | 0.8841 | | 0.0187 | |
| TRIMESTER² | | | | | | |
| First | N/A | | 0.4790 | | N/A | |
| Second | N/A | | 0.2620 | | N/A | |
| Third | N/A | | 0.2095 | | N/A | |

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004, 2005, 2006, and 2007.

Table 7.74C Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Numbers in Thousands, Annual Averages Based on 2004-2005 and 2006-2007

| Demographic/Pregnancy Characteristic | Total ¹ | | PREGNANCY STATUS | | | |
|---|--------------------|-----------|------------------|-----------|--------------|-----------|
| | | | Pregnant | | Not Pregnant | |
| | 2004-2005 | 2006-2007 | 2004-2005 | 2006-2007 | 2004-2005 | 2006-2007 |
| TOTAL | 364 | 394 | 32 | 31 | 360 | 390 |
| AGE | | | | | | |
| 15-17 | 43 | 41 | 3 | 3 | 42 | 41 |
| 18-25 | 76 | 83 | 10 | 11 | 122 | 160 |
| 26-44 | 331 | 335 | 30 | 29 | 328 | 332 |
| HISPANIC ORIGIN AND RACE | | | | | | |
| Not Hispanic or Latino | 346 | 371 | 28 | 30 | 341 | 367 |
| White | 312 | 326 | 24 | 27 | 309 | 323 |
| Black or African American | 124 | 136 | 10 | 12 | 122 | 134 |
| American Indian or Alaska Native | 19 | 19 | * | * | 19 | 19 |
| Native Hawaiian or Other Pacific Islander | * | 16 | * | * | * | 16 |
| Asian | 72 | 72 | * | * | 72 | 71 |
| Two or More Races | 29 | 38 | * | * | 29 | 38 |
| Hispanic or Latino | 120 | 132 | 15 | 6 | 118 | 131 |
| TRIMESTER² | | | | | | |
| First | N/A | N/A | 22 | 21 | N/A | N/A |
| Second | N/A | N/A | 15 | 17 | N/A | N/A |
| Third | N/A | N/A | 16 | 16 | N/A | N/A |

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004, 2005, 2006, and 2007.

Table 7.74D Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Percentages, Annual Averages Based on 2004-2005 and 2006-2007

| Demographic/Pregnancy Characteristic | Total ¹ | | PREGNANCY STATUS | | | |
|---|--------------------|-----------|------------------|-----------|--------------|-----------|
| | | | Pregnant | | Not Pregnant | |
| | 2004-2005 | 2006-2007 | 2004-2005 | 2006-2007 | 2004-2005 | 2006-2007 |
| TOTAL | 0.38 | 0.42 | 1.19 | 1.12 | 0.39 | 0.43 |
| AGE | | | | | | |
| 15-17 | 0.58 | 0.55 | 3.76 | 3.65 | 0.58 | 0.56 |
| 18-25 | 0.47 | 0.51 | 1.02 | 1.05 | 0.48 | 0.52 |
| 26-44 | 0.55 | 0.58 | 1.85 | 1.76 | 0.56 | 0.59 |
| HISPANIC ORIGIN AND RACE | | | | | | |
| Not Hispanic or Latino | 0.41 | 0.44 | 1.31 | 1.33 | 0.42 | 0.45 |
| White | 0.45 | 0.47 | 1.49 | 1.66 | 0.45 | 0.48 |
| Black or African American | 0.99 | 1.04 | 2.78 | 2.88 | 1.01 | 1.09 |
| American Indian or Alaska Native | 4.38 | 4.97 | * | * | 4.48 | 5.14 |
| Native Hawaiian or Other Pacific Islander | * | 5.37 | * | * | * | 5.70 |
| Asian | 1.90 | 2.01 | * | * | 1.93 | 2.11 |
| Two or More Races | 2.66 | 3.26 | * | * | 2.71 | 3.34 |
| Hispanic or Latino | 0.96 | 0.98 | 2.79 | 1.09 | 0.99 | 1.01 |
| TRIMESTER² | | | | | | |
| First | N/A | N/A | 2.67 | 2.27 | N/A | N/A |
| Second | N/A | N/A | 1.79 | 1.64 | N/A | N/A |
| Third | N/A | N/A | 1.66 | 1.91 | N/A | N/A |

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004, 2005, 2006, and 2007.

Table 7.74P Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2006-2007 Versus 2004-2005

| Demographic/Pregnancy Characteristic | Total ¹ | | PREGNANCY STATUS | | | |
|---|--------------------|-----------|------------------|-----------|--------------|-----------|
| | | | Pregnant | | Not Pregnant | |
| | 2004-2005 | 2006-2007 | 2004-2005 | 2006-2007 | 2004-2005 | 2006-2007 |
| TOTAL | 0.9638 | | 0.7810 | | 0.9133 | |
| AGE | | | | | | |
| 15-17 | 0.0034 | | 0.7354 | | 0.0034 | |
| 18-25 | 0.0096 | | 0.9327 | | 0.0051 | |
| 26-44 | 0.7553 | | 0.7134 | | 0.8687 | |
| HISPANIC ORIGIN AND RACE | | | | | | |
| Not Hispanic or Latino | 0.9886 | | 0.9561 | | 0.8974 | |
| White | 0.7996 | | 0.7381 | | 0.7973 | |
| Black or African American | 0.1688 | | 0.5587 | | 0.1443 | |
| American Indian or Alaska Native | 0.6446 | | * | | 0.6815 | |
| Native Hawaiian or Other Pacific Islander | * | | * | | * | |
| Asian | 0.9825 | | * | | 0.9617 | |
| Two or More Races | 0.1977 | | * | | 0.2065 | |
| Hispanic or Latino | 0.5060 | | 0.3570 | | 0.4563 | |
| TRIMESTER² | | | | | | |
| First | N/A | | 0.3937 | | N/A | |
| Second | N/A | | 0.4257 | | N/A | |
| Third | N/A | | 0.5916 | | N/A | |

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004, 2005, 2006, and 2007.

Table 7.75C Binge Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Numbers in Thousands, Annual Averages Based on 2004-2005 and 2006-2007

| Demographic/Pregnancy Characteristic | Total ¹ | | PREGNANCY STATUS | | | |
|---|--------------------|-----------|------------------|-----------|--------------|-----------|
| | | | Pregnant | | Not Pregnant | |
| | 2004-2005 | 2006-2007 | 2004-2005 | 2006-2007 | 2004-2005 | 2006-2007 |
| TOTAL | 213 | 236 | 15 | 14 | 211 | 234 |
| AGE | | | | | | |
| 15-17 | 32 | 31 | 2 | 2 | 31 | 31 |
| 18-25 | 71 | 80 | 7 | 8 | 91 | 116 |
| 26-44 | 184 | 189 | 12 | 11 | 183 | 187 |
| HISPANIC ORIGIN AND RACE | | | | | | |
| Not Hispanic or Latino | 200 | 219 | 12 | 13 | 199 | 218 |
| White | 174 | 199 | 9 | 10 | 173 | 198 |
| Black or African American | 76 | 71 | 9 | 9 | 75 | 69 |
| American Indian or Alaska Native | 16 | 18 | * | * | 16 | 18 |
| Native Hawaiian or Other Pacific Islander | * | 14 | * | * | * | 14 |
| Asian | 41 | 36 | * | * | 41 | 35 |
| Two or More Races | 20 | 20 | * | * | 20 | 20 |
| Hispanic or Latino | 75 | 83 | 8 | 4 | 74 | 82 |
| TRIMESTER² | | | | | | |
| First | N/A | N/A | 10 | 10 | N/A | N/A |
| Second | N/A | N/A | 7 | 7 | N/A | N/A |
| Third | N/A | N/A | 7 | 3 | N/A | N/A |

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004, 2005, 2006, and 2007.

Table 7.75D Binge Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Percentages, Annual Averages Based on 2004-2005 and 2006-2007

| Demographic/Pregnancy Characteristic | Total ¹ | | PREGNANCY STATUS | | | |
|---|--------------------|-----------|------------------|-----------|--------------|-----------|
| | | | Pregnant | | Not Pregnant | |
| | 2004-2005 | 2006-2007 | 2004-2005 | 2006-2007 | 2004-2005 | 2006-2007 |
| TOTAL | 0.30 | 0.33 | 0.58 | 0.53 | 0.31 | 0.34 |
| AGE | | | | | | |
| 15-17 | 0.47 | 0.45 | 2.78 | 2.75 | 0.48 | 0.46 |
| 18-25 | 0.44 | 0.49 | 0.76 | 0.80 | 0.46 | 0.51 |
| 26-44 | 0.41 | 0.43 | 0.82 | 0.73 | 0.42 | 0.45 |
| HISPANIC ORIGIN AND RACE | | | | | | |
| Not Hispanic or Latino | 0.33 | 0.35 | 0.61 | 0.63 | 0.34 | 0.36 |
| White | 0.37 | 0.41 | 0.57 | 0.67 | 0.38 | 0.42 |
| Black or African American | 0.76 | 0.71 | 2.53 | 2.16 | 0.79 | 0.74 |
| American Indian or Alaska Native | 3.92 | 4.74 | * | * | 4.01 | 4.89 |
| Native Hawaiian or Other Pacific Islander | * | 4.72 | * | * | * | 5.00 |
| Asian | 1.24 | 1.09 | * | * | 1.27 | 1.15 |
| Two or More Races | 2.56 | 2.58 | * | * | 2.63 | 2.67 |
| Hispanic or Latino | 0.71 | 0.73 | 1.55 | 0.87 | 0.75 | 0.76 |
| TRIMESTER² | | | | | | |
| First | N/A | N/A | 1.37 | 1.24 | N/A | N/A |
| Second | N/A | N/A | 0.79 | 0.68 | N/A | N/A |
| Third | N/A | N/A | 0.76 | 0.35 | N/A | N/A |

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004, 2005, 2006, and 2007.

Table 7.75P Binge Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2006-2007 Versus 2004-2005

| Demographic/Pregnancy Characteristic | Total ¹ | | PREGNANCY STATUS | | | |
|---|--------------------|-----------|------------------|-----------|--------------|-----------|
| | | | Pregnant | | Not Pregnant | |
| | 2004-2005 | 2006-2007 | 2004-2005 | 2006-2007 | 2004-2005 | 2006-2007 |
| TOTAL | 0.0930 | | 0.8354 | | 0.0840 | |
| AGE | | | | | | |
| 15-17 | 0.0196 | | 0.6148 | | 0.0207 | |
| 18-25 | 0.0078 | | 0.8991 | | 0.0048 | |
| 26-44 | 0.2622 | | 0.6388 | | 0.2592 | |
| HISPANIC ORIGIN AND RACE | | | | | | |
| Not Hispanic or Latino | 0.0492 | | 0.9140 | | 0.0438 | |
| White | 0.0036 | | 0.8152 | | 0.0037 | |
| Black or African American | 0.6371 | | 0.7603 | | 0.7129 | |
| American Indian or Alaska Native | 0.9812 | | * | | 0.9791 | |
| Native Hawaiian or Other Pacific Islander | * | | * | | * | |
| Asian | 0.5881 | | * | | 0.5099 | |
| Two or More Races | 0.4017 | | * | | 0.3869 | |
| Hispanic or Latino | 0.9374 | | 0.5008 | | 0.9167 | |
| TRIMESTER² | | | | | | |
| First | N/A | | 0.6431 | | N/A | |
| Second | N/A | | 0.7597 | | N/A | |
| Third | N/A | | 0.4169 | | N/A | |

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004, 2005, 2006, and 2007.

Table 7.76C Heavy Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Numbers in Thousands, Annual Averages Based on 2004-2005 and 2006-2007

| Demographic/Pregnancy Characteristic | Total ¹ | | PREGNANCY STATUS | | | |
|---|--------------------|-----------|------------------|-----------|--------------|-----------|
| | | | Pregnant | | Not Pregnant | |
| | 2004-2005 | 2006-2007 | 2004-2005 | 2006-2007 | 2004-2005 | 2006-2007 |
| TOTAL | 90 | 98 | 6 | 6 | 90 | 97 |
| AGE | | | | | | |
| 15-17 | 14 | 15 | 1 | 1 | 14 | 15 |
| 18-25 | 41 | 45 | 2 | 3 | 45 | 51 |
| 26-44 | 77 | 79 | 6 | 5 | 76 | 78 |
| HISPANIC ORIGIN AND RACE | | | | | | |
| Not Hispanic or Latino | 87 | 92 | 5 | 5 | 87 | 91 |
| White | 80 | 86 | 5 | 5 | 79 | 86 |
| Black or African American | 29 | 27 | 1 | 2 | 29 | 26 |
| American Indian or Alaska Native | 6 | 9 | * | * | 6 | 9 |
| Native Hawaiian or Other Pacific Islander | 5 | 3 | * | * | 5 | 3 |
| Asian | 13 | 15 | * | * | 13 | 15 |
| Two or More Races | 8 | 11 | * | * | 8 | 11 |
| Hispanic or Latino | 25 | 31 | * | 2 | 25 | 31 |
| TRIMESTER² | | | | | | |
| First | N/A | N/A | 5 | 5 | N/A | N/A |
| Second | N/A | N/A | 4 | 1 | N/A | N/A |
| Third | N/A | N/A | 1 | 1 | N/A | N/A |

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on each of 5 or more days in the past 30 days.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004, 2005, 2006, and 2007.

Table 7.76D Heavy Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Percentages, Annual Averages Based on 2004-2005 and 2006-2007

| Demographic/Pregnancy Characteristic | Total ¹ | | PREGNANCY STATUS | | | |
|---|--------------------|-----------|------------------|-----------|--------------|-----------|
| | | | Pregnant | | Not Pregnant | |
| | 2004-2005 | 2006-2007 | 2004-2005 | 2006-2007 | 2004-2005 | 2006-2007 |
| TOTAL | 0.14 | 0.16 | 0.25 | 0.21 | 0.15 | 0.16 |
| AGE | | | | | | |
| 15-17 | 0.23 | 0.23 | 1.51 | 1.63 | 0.23 | 0.23 |
| 18-25 | 0.25 | 0.28 | 0.26 | 0.26 | 0.27 | 0.29 |
| 26-44 | 0.19 | 0.20 | 0.38 | 0.31 | 0.20 | 0.21 |
| HISPANIC ORIGIN AND RACE | | | | | | |
| Not Hispanic or Latino | 0.16 | 0.17 | 0.24 | 0.26 | 0.17 | 0.18 |
| White | 0.20 | 0.21 | 0.31 | 0.35 | 0.21 | 0.22 |
| Black or African American | 0.33 | 0.31 | 0.22 | 0.43 | 0.35 | 0.32 |
| American Indian or Alaska Native | 1.46 | 2.61 | * | * | 1.49 | 2.72 |
| Native Hawaiian or Other Pacific Islander | 2.11 | 0.97 | * | * | 2.18 | 1.00 |
| Asian | 0.41 | 0.46 | * | * | 0.43 | 0.49 |
| Two or More Races | 1.16 | 1.46 | * | * | 1.20 | 1.51 |
| Hispanic or Latino | 0.25 | 0.30 | * | 0.33 | 0.27 | 0.31 |
| TRIMESTER² | | | | | | |
| First | N/A | N/A | 0.63 | 0.63 | N/A | N/A |
| Second | N/A | N/A | 0.53 | 0.14 | N/A | N/A |
| Third | N/A | N/A | 0.07 | 0.18 | N/A | N/A |

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on each of 5 or more days in the past 30 days.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004, 2005, 2006, and 2007.

Table 7.76P Heavy Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2006-2007 Versus 2004-2005

| Demographic/Pregnancy Characteristic | Total ¹ | | PREGNANCY STATUS | | | |
|---|--------------------|-----------|------------------|-----------|--------------|-----------|
| | | | Pregnant | | Not Pregnant | |
| | 2004-2005 | 2006-2007 | 2004-2005 | 2006-2007 | 2004-2005 | 2006-2007 |
| TOTAL | 0.2931 | | 0.9579 | | 0.2999 | |
| AGE | | | | | | |
| 15-17 | 0.5085 | | 0.9495 | | 0.4394 | |
| 18-25 | 0.0090 | | 0.6493 | | 0.0082 | |
| 26-44 | 0.9308 | | 0.7696 | | 0.9261 | |
| HISPANIC ORIGIN AND RACE | | | | | | |
| Not Hispanic or Latino | 0.4404 | | 0.8110 | | 0.4568 | |
| White | 0.4276 | | 0.9373 | | 0.4361 | |
| Black or African American | 0.8759 | | 0.3948 | | 0.9614 | |
| American Indian or Alaska Native | 0.1865 | | * | | 0.1816 | |
| Native Hawaiian or Other Pacific Islander | 0.4455 | | * | | 0.4765 | |
| Asian | 0.6333 | | * | | 0.6116 | |
| Two or More Races | 0.7160 | | * | | 0.7561 | |
| Hispanic or Latino | 0.1324 | | * | | 0.1190 | |
| TRIMESTER² | | | | | | |
| First | N/A | | 0.9538 | | N/A | |
| Second | N/A | | 0.3420 | | N/A | |
| Third | N/A | | 0.2253 | | N/A | |

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on each of 5 or more days in the past 30 days.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004, 2005, 2006, and 2007.