

Table 8.35C Serious Psychological Distress in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2004-2007

Demographic Characteristic	2004	2005	2006	2007
TOTAL	750	505	492	485
AGE				
18-25	159	105	116	114
26-49	516	334	336	322
50 or Older	481	332	340	350
GENDER				
Male	454	301	314	307
Female	606	390	367	386
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	677	451	457	451
White	642	423	436	430
Black or African American	253	188	168	186
American Indian or Alaska Native	31	48	72	36
Native Hawaiian or Other Pacific Islander	*	*	28	23
Asian	152	88	103	85
Two or More Races	79	53	98	42
Hispanic or Latino	307	201	194	195

*Low precision; no estimate reported.

NOTE: Estimates for 2004 in this table are based on a subsample of respondents aged 18 or older. Due to the use of alternative 2004 subsample data, these 2004 estimates may differ from 2004 estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD) is defined as having a score of 13 or higher on the K6 scale. Due to questionnaire changes, these estimates are not comparable with SPD estimates published in 2004 and prior years. See Section B.4.4 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004, 2005, 2006, and 2007.

Table 8.35D Serious Psychological Distress in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2004-2007

Demographic Characteristic	2004	2005	2006	2007
TOTAL	0.35	0.23	0.22	0.22
AGE				
18-25	0.50	0.32	0.36	0.35
26-49	0.52	0.33	0.33	0.32
50 or Older	0.58	0.39	0.39	0.39
GENDER				
Male	0.44	0.29	0.29	0.29
Female	0.54	0.35	0.32	0.34
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.36	0.24	0.24	0.23
White	0.40	0.26	0.26	0.26
Black or African American	1.02	0.69	0.59	0.68
American Indian or Alaska Native	3.07	3.78	4.67	3.07
Native Hawaiian or Other Pacific Islander	*	*	3.29	3.59
Asian	1.75	0.95	1.09	0.88
Two or More Races	3.96	2.19	3.72	1.81
Hispanic or Latino	1.15	0.72	0.67	0.66

*Low precision; no estimate reported.

NOTE: Estimates for 2004 in this table are based on a subsample of respondents aged 18 or older. Due to the use of alternative 2004 subsample data, these 2004 estimates may differ from 2004 estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD) is defined as having a score of 13 or higher on the K6 scale. Due to questionnaire changes, these estimates are not comparable with SPD estimates published in 2004 and prior years. See Section B.4.4 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004, 2005, 2006, and 2007.

Table 8.35P Serious Psychological Distress in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2007 Versus 2004, 2005, and 2006

Demographic Characteristic	2004	2005	2006	2007
TOTAL	0.0022	0.2501	0.2265	
AGE				
18-25	0.0001	0.1002	0.7068	
26-49	0.0038	0.5951	0.0810	
50 or Older	0.9482	0.7396	0.9631	
GENDER				
Male	0.0263	0.7360	0.2901	
Female	0.0357	0.2220	0.5120	
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.0072	0.5470	0.3071	
White	0.0644	0.8616	0.8475	
Black or African American	0.2479	0.8322	0.9624	
American Indian or Alaska Native	0.5031	0.1433	0.0369	
Native Hawaiian or Other Pacific Islander	*	*	0.8025	
Asian	0.1543	0.5343	0.3263	
Two or More Races	0.0516	0.3117	0.0061	
Hispanic or Latino	0.1374	0.1281	0.5509	

*Low precision; no estimate reported.

NOTE: Estimates for 2004 in this table are based on a subsample of respondents aged 18 or older. Due to the use of alternative 2004 subsample data, these 2004 estimates may differ from 2004 estimates published in prior NSDUH reports. See Section B.4.4 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD) is defined as having a score of 13 or higher on the K6 scale. Due to questionnaire changes, these estimates are not comparable with SPD estimates published in 2004 and prior years. See Section B.4.4 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004, 2005, 2006, and 2007.