

**Table 8.22C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Gender: Standard Errors of Numbers in Thousands, 2002-2007**

Gender/Substance	2002	2003	2004	2005	2006	2007
<b>TOTAL</b>						
<b>TOBACCO PRODUCTS<sup>1</sup></b>	832	803	794	845	848	844
Cigarettes	795	789	768	777	804	808
Smokeless Tobacco	279	284	245	267	284	281
Cigars	350	331	323	354	352	337
Pipe Tobacco	166	131	147	146	166	170
<b>ALCOHOL</b>	989	919	957	975	969	1,018
Binge Alcohol Use <sup>2</sup>	732	682	698	701	746	767
Heavy Alcohol Use <sup>2</sup>	392	380	382	379	427	429
<b>MALE</b>						
<b>TOBACCO PRODUCTS<sup>1</sup></b>	572	573	570	576	584	595
Cigarettes	527	545	537	515	543	574
Smokeless Tobacco	264	256	234	241	273	251
Cigars	327	305	290	324	319	311
Pipe Tobacco	139	127	139	141	160	168
<b>ALCOHOL</b>	656	599	626	621	611	667
Binge Alcohol Use <sup>2</sup>	540	517	513	529	559	585
Heavy Alcohol Use <sup>2</sup>	346	321	319	321	355	375
<b>FEMALE</b>						
<b>TOBACCO PRODUCTS<sup>1</sup></b>	513	507	497	543	511	511
Cigarettes	501	488	491	538	502	493
Smokeless Tobacco	100	126	65	87	76	113
Cigars	114	123	122	112	131	116
Pipe Tobacco	90	35	43	42	47	30
<b>ALCOHOL</b>	644	600	664	637	644	674
Binge Alcohol Use <sup>2</sup>	415	382	390	402	399	394
Heavy Alcohol Use <sup>2</sup>	177	174	174	163	168	171

\*Low precision; no estimate reported.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, 2005, 2006, and 2007.

**Table 8.22D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Gender: Standard Errors of Percentages, 2002-2007**

<b>Gender/Substance</b>	<b>2002</b>	<b>2003</b>	<b>2004</b>	<b>2005</b>	<b>2006</b>	<b>2007</b>
<b>TOTAL</b>						
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.35	0.34	0.33	0.35	0.34	0.34
Cigarettes	0.34	0.33	0.32	0.32	0.33	0.33
Smokeless Tobacco	0.12	0.12	0.10	0.11	0.12	0.11
Cigars	0.15	0.14	0.13	0.15	0.14	0.14
Pipe Tobacco	0.07	0.06	0.06	0.06	0.07	0.07
<b>ALCOHOL</b>	0.42	0.39	0.40	0.40	0.39	0.41
Binge Alcohol Use <sup>2</sup>	0.31	0.29	0.29	0.29	0.30	0.31
Heavy Alcohol Use <sup>2</sup>	0.17	0.16	0.16	0.16	0.17	0.17
<b>MALE</b>						
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.50	0.50	0.49	0.49	0.49	0.50
Cigarettes	0.46	0.47	0.46	0.44	0.45	0.48
Smokeless Tobacco	0.23	0.22	0.20	0.20	0.23	0.21
Cigars	0.29	0.27	0.25	0.27	0.27	0.26
Pipe Tobacco	0.12	0.11	0.12	0.12	0.13	0.14
<b>ALCOHOL</b>	0.58	0.52	0.54	0.53	0.51	0.55
Binge Alcohol Use <sup>2</sup>	0.48	0.45	0.44	0.45	0.47	0.49
Heavy Alcohol Use <sup>2</sup>	0.30	0.28	0.27	0.27	0.30	0.31
<b>FEMALE</b>						
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.42	0.41	0.40	0.43	0.40	0.40
Cigarettes	0.41	0.40	0.40	0.43	0.40	0.39
Smokeless Tobacco	0.08	0.10	0.05	0.07	0.06	0.09
Cigars	0.09	0.10	0.10	0.09	0.10	0.09
Pipe Tobacco	0.07	0.03	0.03	0.03	0.04	0.02
<b>ALCOHOL</b>	0.53	0.49	0.54	0.51	0.51	0.53
Binge Alcohol Use <sup>2</sup>	0.34	0.31	0.31	0.32	0.32	0.31
Heavy Alcohol Use <sup>2</sup>	0.15	0.14	0.14	0.13	0.13	0.13

\*Low precision; no estimate reported.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, 2005, 2006, and 2007.

**Table 8.22P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Gender: *P* Values from Tests of Differences of Percentages, 2007 Versus 2002, 2003, 2004, 2005, and 2006**

Gender/Substance	2002	2003	2004	2005	2006	2007
<b>TOTAL</b>						
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.0003	0.0171	0.2151	0.1027	0.0306	
Cigarettes	0.0002	0.0107	0.1444	0.1525	0.0749	
Smokeless Tobacco	0.7005	0.9921	0.0727	0.5750	0.5340	
Cigars	0.7229	0.7984	0.0626	0.1998	0.2695	
Pipe Tobacco	0.5905	0.1016	0.4980	0.3992	0.2149	
<b>ALCOHOL</b>	0.7484	0.0530	0.1314	0.2379	0.6941	
Binge Alcohol Use <sup>2</sup>	0.3187	0.1027	0.1881	0.1124	0.4697	
Heavy Alcohol Use <sup>2</sup>	0.6223	0.7631	0.7474	0.2575	0.9193	
<b>MALE</b>						
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.0143	0.3354	0.5120	0.4311	0.0928	
Cigarettes	0.0153	0.1597	0.3674	0.6538	0.2536	
Smokeless Tobacco	0.8367	0.6484	0.0661	0.4472	0.3864	
Cigars	0.4461	0.8358	0.0627	0.1609	0.6016	
Pipe Tobacco	0.3179	0.1214	0.4516	0.5866	0.2984	
<b>ALCOHOL</b>	0.3260	0.3442	0.6806	0.0515	0.6120	
Binge Alcohol Use <sup>2</sup>	0.4694	0.2438	0.3373	0.0751	0.5079	
Heavy Alcohol Use <sup>2</sup>	0.7799	0.5796	0.9917	0.4603	0.8720	
<b>FEMALE</b>						
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.0012	0.0049	0.1943	0.0806	0.1053	
Cigarettes	0.0007	0.0098	0.1804	0.0831	0.1099	
Smokeless Tobacco	0.4735	0.2664	0.8483	0.6927	0.5855	
Cigars	0.4109	0.1629	0.5701	0.9796	0.0798	
Pipe Tobacco	0.3279	0.6589	0.7786	0.1855	0.3523	
<b>ALCOHOL</b>	0.1585	0.0001	0.0096	0.9214	0.2724	
Binge Alcohol Use <sup>2</sup>	0.4870	0.1980	0.2941	0.6817	0.6754	
Heavy Alcohol Use <sup>2</sup>	0.1147	0.6066	0.4217	0.2261	0.9213	

\*Low precision; no estimate reported.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, 2005, 2006, and 2007.

**Table 8.23C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Gender: Standard Errors of Numbers in Thousands, 2002-2007**

Gender/Substance	2002	2003	2004	2005	2006	2007
<b>TOTAL</b>						
<b>TOBACCO PRODUCTS<sup>1</sup></b>	81	80	80	78	72	76
Cigarettes	74	73	76	71	65	67
Smokeless Tobacco	27	30	32	29	31	35
Cigars	47	44	44	46	41	44
Pipe Tobacco	14	18	19	17	18	19
<b>ALCOHOL</b>	80	82	81	81	82	87
Binge Alcohol Use <sup>2</sup>	67	67	73	65	69	68
Heavy Alcohol Use <sup>2</sup>	31	33	36	32	33	37
<b>MALE</b>						
<b>TOBACCO PRODUCTS<sup>1</sup></b>	59	56	55	59	54	57
Cigarettes	53	47	52	53	46	48
Smokeless Tobacco	26	29	30	28	29	34
Cigars	37	36	37	38	34	37
Pipe Tobacco	13	17	15	14	15	16
<b>ALCOHOL</b>	55	57	59	57	57	61
Binge Alcohol Use <sup>2</sup>	46	46	52	48	47	51
Heavy Alcohol Use <sup>2</sup>	24	23	27	27	26	28
<b>FEMALE</b>						
<b>TOBACCO PRODUCTS<sup>1</sup></b>	51	52	53	50	47	48
Cigarettes	49	50	52	48	45	45
Smokeless Tobacco	9	6	11	8	10	9
Cigars	26	23	25	24	23	22
Pipe Tobacco	9	7	13	11	10	11
<b>ALCOHOL</b>	58	60	59	58	59	61
Binge Alcohol Use <sup>2</sup>	46	46	50	43	49	45
Heavy Alcohol Use <sup>2</sup>	19	23	23	20	21	23

\*Low precision; no estimate reported.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, 2005, 2006, and 2007.

**Table 8.23D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Gender: Standard Errors of Percentages, 2002-2007**

<b>Gender/Substance</b>	<b>2002</b>	<b>2003</b>	<b>2004</b>	<b>2005</b>	<b>2006</b>	<b>2007</b>
<b>TOTAL</b>						
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.33	0.32	0.32	0.31	0.29	0.30
Cigarettes	0.30	0.29	0.30	0.28	0.26	0.26
Smokeless Tobacco	0.11	0.12	0.13	0.12	0.12	0.14
Cigars	0.19	0.17	0.18	0.18	0.16	0.18
Pipe Tobacco	0.06	0.07	0.08	0.07	0.07	0.08
<b>ALCOHOL</b>	0.32	0.33	0.32	0.32	0.32	0.34
Binge Alcohol Use <sup>2</sup>	0.27	0.27	0.29	0.26	0.27	0.27
Heavy Alcohol Use <sup>2</sup>	0.12	0.13	0.14	0.13	0.13	0.15
<b>MALE</b>						
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.47	0.44	0.43	0.46	0.41	0.44
Cigarettes	0.42	0.37	0.40	0.41	0.36	0.37
Smokeless Tobacco	0.21	0.23	0.23	0.22	0.22	0.26
Cigars	0.30	0.28	0.28	0.29	0.26	0.29
Pipe Tobacco	0.10	0.13	0.11	0.11	0.12	0.12
<b>ALCOHOL</b>	0.43	0.45	0.45	0.44	0.44	0.47
Binge Alcohol Use <sup>2</sup>	0.37	0.36	0.40	0.37	0.36	0.39
Heavy Alcohol Use <sup>2</sup>	0.19	0.18	0.21	0.21	0.20	0.22
<b>FEMALE</b>						
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.42	0.42	0.43	0.40	0.38	0.39
Cigarettes	0.40	0.41	0.42	0.39	0.37	0.37
Smokeless Tobacco	0.08	0.05	0.09	0.07	0.08	0.07
Cigars	0.21	0.19	0.20	0.19	0.19	0.18
Pipe Tobacco	0.07	0.06	0.10	0.09	0.08	0.09
<b>ALCOHOL</b>	0.48	0.49	0.48	0.47	0.47	0.49
Binge Alcohol Use <sup>2</sup>	0.38	0.38	0.41	0.35	0.39	0.37
Heavy Alcohol Use <sup>2</sup>	0.16	0.19	0.19	0.16	0.17	0.18

\*Low precision; no estimate reported.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, 2005, 2006, and 2007.

**Table 8.23P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Gender: *P* Values from Tests of Differences of Percentages, 2007 Versus 2002, 2003, 2004, 2005, and 2006**

Gender/Substance	2002	2003	2004	2005	2006	2007
<b>TOTAL</b>						
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.0000	0.0000	0.0000	0.0955	0.2113	
Cigarettes	0.0000	0.0000	0.0000	0.0148	0.1372	
Smokeless Tobacco	0.0078	0.0287	0.3406	0.0337	0.6704	
Cigars	0.2914	0.3734	0.0350	0.8653	0.6137	
Pipe Tobacco	0.0834	0.4559	0.9030	0.3566	0.5667	
<b>ALCOHOL</b>	0.0003	0.0002	0.0004	0.2044	0.1349	
Binge Alcohol Use <sup>2</sup>	0.0110	0.0138	0.0005	0.5638	0.1172	
Heavy Alcohol Use <sup>2</sup>	0.2860	0.1745	0.1149	0.7000	0.8640	
<b>MALE</b>						
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.0036	0.0182	0.0473	0.8224	0.8274	
Cigarettes	0.0000	0.0003	0.0161	0.1750	0.8885	
Smokeless Tobacco	0.0049	0.0619	0.3207	0.0304	0.5972	
Cigars	0.5985	0.7227	0.1479	0.5382	0.1944	
Pipe Tobacco	0.1890	0.8346	0.7578	0.6813	0.8347	
<b>ALCOHOL</b>	0.0167	0.0559	0.0407	0.9803	0.5282	
Binge Alcohol Use <sup>2</sup>	0.1356	0.3141	0.0702	0.6673	0.8382	
Heavy Alcohol Use <sup>2</sup>	0.3237	0.7418	0.2746	0.6926	0.9938	
<b>FEMALE</b>						
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.0000	0.0000	0.0000	0.0214	0.0265	
Cigarettes	0.0000	0.0000	0.0000	0.0350	0.0413	
Smokeless Tobacco	0.9149	0.0717	0.9593	0.7596	0.7932	
Cigars	0.2301	0.2472	0.0886	0.4910	0.2410	
Pipe Tobacco	0.2879	0.0551	0.8429	0.3117	0.4686	
<b>ALCOHOL</b>	0.0063	0.0008	0.0037	0.0818	0.1366	
Binge Alcohol Use <sup>2</sup>	0.0294	0.0095	0.0014	0.1823	0.0377	
Heavy Alcohol Use <sup>2</sup>	0.6213	0.0839	0.2313	0.9340	0.7812	

\*Low precision; no estimate reported.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, 2005, 2006, and 2007.

**Table 8.24C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Gender: Standard Errors of Numbers in Thousands, 2002-2007**

Gender/Substance	2002	2003	2004	2005	2006	2007
<b>TOTAL</b>						
<b>TOBACCO PRODUCTS<sup>1</sup></b>	150	151	161	155	159	162
Cigarettes	150	151	159	152	155	159
Smokeless Tobacco	59	56	60	64	65	63
Cigars	83	84	97	92	94	92
Pipe Tobacco	24	25	30	37	33	32
<b>ALCOHOL</b>	164	160	164	166	168	170
Binge Alcohol Use <sup>2</sup>	161	155	166	171	169	170
Heavy Alcohol Use <sup>2</sup>	112	114	120	120	125	115
<b>MALE</b>						
<b>TOBACCO PRODUCTS<sup>1</sup></b>	100	106	106	110	108	116
Cigarettes	99	109	105	107	108	116
Smokeless Tobacco	56	55	58	60	62	61
Cigars	73	72	83	79	79	81
Pipe Tobacco	22	21	27	33	30	30
<b>ALCOHOL</b>	111	100	110	101	108	110
Binge Alcohol Use <sup>2</sup>	110	105	113	111	115	115
Heavy Alcohol Use <sup>2</sup>	87	91	88	96	95	94
<b>FEMALE</b>						
<b>TOBACCO PRODUCTS<sup>1</sup></b>	101	93	102	101	102	96
Cigarettes	102	92	101	101	98	94
Smokeless Tobacco	8	11	10	12	13	13
Cigars	38	43	44	41	44	43
Pipe Tobacco	12	12	12	16	11	14
<b>ALCOHOL</b>	102	114	104	112	111	114
Binge Alcohol Use <sup>2</sup>	99	100	96	104	103	109
Heavy Alcohol Use <sup>2</sup>	55	59	60	56	63	59

\*Low precision; no estimate reported.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, 2005, 2006, and 2007.

**Table 8.24D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Gender: Standard Errors of Percentages, 2002-2007**

<b>Gender/Substance</b>	<b>2002</b>	<b>2003</b>	<b>2004</b>	<b>2005</b>	<b>2006</b>	<b>2007</b>
<b>TOTAL</b>						
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.48	0.48	0.50	0.48	0.49	0.49
Cigarettes	0.48	0.47	0.49	0.47	0.47	0.48
Smokeless Tobacco	0.19	0.18	0.19	0.20	0.20	0.19
Cigars	0.27	0.26	0.30	0.28	0.29	0.28
Pipe Tobacco	0.08	0.08	0.09	0.11	0.10	0.10
<b>ALCOHOL</b>	0.53	0.50	0.51	0.51	0.51	0.52
Binge Alcohol Use <sup>2</sup>	0.52	0.49	0.52	0.53	0.52	0.52
Heavy Alcohol Use <sup>2</sup>	0.36	0.36	0.37	0.37	0.38	0.35
<b>MALE</b>						
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.64	0.67	0.66	0.67	0.66	0.71
Cigarettes	0.64	0.69	0.65	0.66	0.66	0.70
Smokeless Tobacco	0.36	0.34	0.36	0.37	0.37	0.37
Cigars	0.47	0.45	0.51	0.48	0.48	0.49
Pipe Tobacco	0.14	0.13	0.17	0.20	0.18	0.18
<b>ALCOHOL</b>	0.71	0.63	0.68	0.62	0.65	0.67
Binge Alcohol Use <sup>2</sup>	0.71	0.66	0.70	0.68	0.70	0.70
Heavy Alcohol Use <sup>2</sup>	0.56	0.57	0.54	0.59	0.58	0.57
<b>FEMALE</b>						
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.65	0.59	0.64	0.63	0.63	0.59
Cigarettes	0.66	0.58	0.63	0.63	0.60	0.58
Smokeless Tobacco	0.05	0.07	0.06	0.08	0.08	0.08
Cigars	0.25	0.27	0.27	0.26	0.27	0.26
Pipe Tobacco	0.08	0.08	0.08	0.10	0.07	0.08
<b>ALCOHOL</b>	0.66	0.72	0.65	0.69	0.69	0.70
Binge Alcohol Use <sup>2</sup>	0.64	0.63	0.60	0.65	0.63	0.67
Heavy Alcohol Use <sup>2</sup>	0.35	0.37	0.37	0.35	0.39	0.36

\*Low precision; no estimate reported.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, 2005, 2006, and 2007.



**Table 8.24P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Gender: *P* Values from Tests of Differences of Percentages, 2007 Versus 2002, 2003, 2004, 2005, and 2006**

Gender/Substance	2002	2003	2004	2005	2006	2007
<b>TOTAL</b>						
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.0000	0.0000	0.0001	0.0007	0.0020	
Cigarettes	0.0000	0.0000	0.0000	0.0001	0.0005	
Smokeless Tobacco	0.1228	0.0274	0.2274	0.6206	0.8680	
Cigars	0.0358	0.2821	0.0272	0.6927	0.4589	
Pipe Tobacco	0.2772	0.0249	0.8393	0.0990	0.4299	
<b>ALCOHOL</b>	0.3175	0.8429	0.3242	0.6759	0.2807	
Binge Alcohol Use <sup>2</sup>	0.2485	0.7644	0.4452	0.9214	0.5430	
Heavy Alcohol Use <sup>2</sup>	0.7152	0.4636	0.5135	0.2599	0.0937	
<b>MALE</b>						
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.0238	0.0822	0.0674	0.1096	0.2922	
Cigarettes	0.0000	0.0002	0.0015	0.0148	0.1291	
Smokeless Tobacco	0.2855	0.0441	0.3842	0.7006	0.9866	
Cigars	0.0140	0.0917	0.0897	0.8840	0.7586	
Pipe Tobacco	0.3609	0.0379	0.4762	0.1609	0.3303	
<b>ALCOHOL</b>	0.9748	0.0731	0.7247	0.2633	0.4904	
Binge Alcohol Use <sup>2</sup>	0.7059	0.1194	0.7488	0.0583	0.6437	
Heavy Alcohol Use <sup>2</sup>	0.1174	0.1099	0.0880	0.0284	0.1701	
<b>FEMALE</b>						
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.0000	0.0000	0.0000	0.0002	0.0002	
Cigarettes	0.0000	0.0000	0.0000	0.0002	0.0001	
Smokeless Tobacco	0.0200	0.3647	0.0895	0.5988	0.4612	
Cigars	0.7336	0.3536	0.0886	0.2146	0.3139	
Pipe Tobacco	0.6457	0.4318	0.2970	0.3742	0.7686	
<b>ALCOHOL</b>	0.1449	0.1841	0.2558	0.0980	0.3647	
Binge Alcohol Use <sup>2</sup>	0.0335	0.0425	0.1201	0.0706	0.6643	
Heavy Alcohol Use <sup>2</sup>	0.1051	0.3138	0.2079	0.1691	0.2755	

\*Low precision; no estimate reported.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, 2005, 2006, and 2007.

**Table 8.25C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Gender: Standard Errors of Numbers in Thousands, 2002-2007**

Gender/Substance	2002	2003	2004	2005	2006	2007
<b>TOTAL</b>						
<b>TOBACCO PRODUCTS<sup>1</sup></b>	797	748	752	798	798	801
Cigarettes	754	738	718	726	757	762
Smokeless Tobacco	270	274	236	256	268	266
Cigars	331	332	311	335	332	311
Pipe Tobacco	160	126	142	141	161	166
<b>ALCOHOL</b>	953	887	933	936	915	985
Binge Alcohol Use <sup>2</sup>	702	639	654	650	697	712
Heavy Alcohol Use <sup>2</sup>	355	349	356	347	393	403
<b>MALE</b>						
<b>TOBACCO PRODUCTS<sup>1</sup></b>	553	546	552	554	566	572
Cigarettes	506	521	512	493	525	547
Smokeless Tobacco	252	247	226	232	259	237
Cigars	311	302	280	309	305	289
Pipe Tobacco	132	124	135	136	156	164
<b>ALCOHOL</b>	630	577	610	598	584	641
Binge Alcohol Use <sup>2</sup>	527	492	492	507	541	562
Heavy Alcohol Use <sup>2</sup>	323	306	307	302	333	356
<b>FEMALE</b>						
<b>TOBACCO PRODUCTS<sup>1</sup></b>	488	477	469	515	477	486
Cigarettes	475	460	466	508	471	469
Smokeless Tobacco	99	125	63	85	74	111
Cigars	99	117	107	98	117	104
Pipe Tobacco	89	32	39	37	44	26
<b>ALCOHOL</b>	628	582	654	615	614	657
Binge Alcohol Use <sup>2</sup>	393	347	361	370	356	364
Heavy Alcohol Use <sup>2</sup>	158	155	160	146	147	157

\*Low precision; no estimate reported.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, 2005, 2006, and 2007.

**Table 8.25D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Gender: Standard Errors of Percentages, 2002-2007**

<b>Gender/Substance</b>	<b>2002</b>	<b>2003</b>	<b>2004</b>	<b>2005</b>	<b>2006</b>	<b>2007</b>
<b>TOTAL</b>						
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.44	0.41	0.41	0.43	0.42	0.42
Cigarettes	0.42	0.41	0.39	0.39	0.40	0.40
Smokeless Tobacco	0.15	0.15	0.13	0.14	0.14	0.14
Cigars	0.18	0.18	0.17	0.18	0.18	0.16
Pipe Tobacco	0.09	0.07	0.08	0.08	0.09	0.09
<b>ALCOHOL</b>	0.53	0.49	0.51	0.51	0.49	0.52
Binge Alcohol Use <sup>2</sup>	0.39	0.35	0.36	0.35	0.37	0.37
Heavy Alcohol Use <sup>2</sup>	0.20	0.19	0.19	0.19	0.21	0.21
<b>MALE</b>						
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.65	0.63	0.63	0.62	0.63	0.63
Cigarettes	0.59	0.60	0.59	0.56	0.58	0.60
Smokeless Tobacco	0.29	0.29	0.26	0.26	0.29	0.26
Cigars	0.36	0.35	0.32	0.35	0.34	0.32
Pipe Tobacco	0.15	0.14	0.15	0.15	0.17	0.18
<b>ALCOHOL</b>	0.74	0.67	0.70	0.67	0.65	0.70
Binge Alcohol Use <sup>2</sup>	0.62	0.57	0.56	0.57	0.60	0.62
Heavy Alcohol Use <sup>2</sup>	0.38	0.35	0.35	0.34	0.37	0.39
<b>FEMALE</b>						
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.52	0.50	0.49	0.53	0.49	0.49
Cigarettes	0.51	0.49	0.49	0.53	0.48	0.47
Smokeless Tobacco	0.11	0.13	0.07	0.09	0.08	0.11
Cigars	0.11	0.12	0.11	0.10	0.12	0.11
Pipe Tobacco	0.10	0.03	0.04	0.04	0.05	0.03
<b>ALCOHOL</b>	0.67	0.61	0.68	0.64	0.63	0.66
Binge Alcohol Use <sup>2</sup>	0.42	0.37	0.38	0.38	0.36	0.37
Heavy Alcohol Use <sup>2</sup>	0.17	0.16	0.17	0.15	0.15	0.16

\*Low precision; no estimate reported.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, 2005, 2006, and 2007.

**Table 8.25P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Gender: *P* Values from Tests of Differences of Percentages, 2007 Versus 2002, 2003, 2004, 2005, and 2006**

Gender/Substance	2002	2003	2004	2005	2006	2007
<b>TOTAL</b>						
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.0200	0.2004	0.9453	0.3751	0.1247	
Cigarettes	0.0488	0.3238	0.9655	0.6541	0.2993	
Smokeless Tobacco	0.2781	0.4480	0.1403	0.8243	0.4545	
Cigars	0.4202	0.7024	0.3551	0.2123	0.3232	
Pipe Tobacco	0.8535	0.2452	0.4633	0.5491	0.2348	
<b>ALCOHOL</b>	0.8069	0.0298	0.1293	0.1787	0.5151	
Binge Alcohol Use <sup>2</sup>	0.3528	0.0570	0.1114	0.0699	0.2857	
Heavy Alcohol Use <sup>2</sup>	0.4983	0.4976	0.9748	0.1036	0.6845	
<b>MALE</b>						
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.0519	0.6100	0.8544	0.6099	0.1183	
Cigarettes	0.1588	0.6762	0.9197	0.8682	0.3884	
Smokeless Tobacco	0.3938	0.8003	0.1159	0.6685	0.3237	
Cigars	0.1602	0.8769	0.2122	0.1098	0.5486	
Pipe Tobacco	0.4617	0.2157	0.3813	0.7586	0.3626	
<b>ALCOHOL</b>	0.2925	0.4736	0.6294	0.0472	0.6670	
Binge Alcohol Use <sup>2</sup>	0.4104	0.1190	0.2323	0.0289	0.4273	
Heavy Alcohol Use <sup>2</sup>	0.9084	0.3055	0.5921	0.1474	0.8455	
<b>FEMALE</b>						
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.0966	0.1258	0.9526	0.3920	0.4838	
Cigarettes	0.1020	0.2536	0.9793	0.3879	0.4879	
Smokeless Tobacco	0.3412	0.1857	0.9791	0.6187	0.6385	
Cigars	0.1655	0.4017	0.5933	0.4917	0.2002	
Pipe Tobacco	0.2190	0.6917	0.5451	0.1589	0.1981	
<b>ALCOHOL</b>	0.1801	0.0001	0.0083	0.9113	0.1637	
Binge Alcohol Use <sup>2</sup>	0.6875	0.2349	0.2362	0.9035	0.3799	
Heavy Alcohol Use <sup>2</sup>	0.2202	0.5154	0.2639	0.3696	0.5377	

\*Low precision; no estimate reported.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, 2005, 2006, and 2007.

**Table 8.26C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 or Older, by Gender: Standard Errors of Numbers in Thousands, 2002-2007**

Gender/Substance	2002	2003	2004	2005	2006	2007
<b>TOTAL</b>						
<b>TOBACCO PRODUCTS<sup>1</sup></b>	829	784	793	839	837	838
Cigarettes	790	774	765	772	796	802
Smokeless Tobacco	276	283	243	264	281	278
Cigars	346	333	323	351	347	331
Pipe Tobacco	163	130	145	145	164	169
<b>ALCOHOL</b>	972	909	950	969	955	1,010
Binge Alcohol Use <sup>2</sup>	730	675	699	701	741	763
Heavy Alcohol Use <sup>2</sup>	389	376	383	378	425	426
<b>MALE</b>						
<b>TOBACCO PRODUCTS<sup>1</sup></b>	565	561	568	576	580	592
Cigarettes	521	537	532	514	540	570
Smokeless Tobacco	260	255	233	239	271	248
Cigars	322	306	291	324	316	306
Pipe Tobacco	136	127	138	141	159	168
<b>ALCOHOL</b>	643	590	616	616	600	658
Binge Alcohol Use <sup>2</sup>	540	517	511	529	561	585
Heavy Alcohol Use <sup>2</sup>	346	320	318	322	353	374
<b>FEMALE</b>						
<b>TOBACCO PRODUCTS<sup>1</sup></b>	509	498	494	541	505	505
Cigarettes	497	481	488	537	497	487
Smokeless Tobacco	99	126	64	87	75	112
Cigars	110	121	119	109	129	112
Pipe Tobacco	90	35	41	40	46	28
<b>ALCOHOL</b>	640	593	662	635	640	671
Binge Alcohol Use <sup>2</sup>	411	375	389	401	395	391
Heavy Alcohol Use <sup>2</sup>	175	170	173	162	168	169

\*Low precision; no estimate reported.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, 2005, 2006, and 2007.

**Table 8.26D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 or Older, by Gender: Standard Errors of Percentages, 2002-2007**

Gender/Substance	2002	2003	2004	2005	2006	2007
<b>TOTAL</b>						
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.39	0.37	0.37	0.39	0.38	0.38
Cigarettes	0.38	0.36	0.36	0.35	0.36	0.36
Smokeless Tobacco	0.13	0.13	0.11	0.12	0.13	0.13
Cigars	0.16	0.16	0.15	0.16	0.16	0.15
Pipe Tobacco	0.08	0.06	0.07	0.07	0.07	0.08
<b>ALCOHOL</b>	0.46	0.43	0.44	0.44	0.43	0.45
Binge Alcohol Use <sup>2</sup>	0.35	0.32	0.32	0.32	0.34	0.34
Heavy Alcohol Use <sup>2</sup>	0.19	0.18	0.18	0.17	0.19	0.19
<b>MALE</b>						
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.56	0.55	0.55	0.55	0.55	0.55
Cigarettes	0.52	0.53	0.51	0.49	0.51	0.53
Smokeless Tobacco	0.26	0.25	0.22	0.23	0.25	0.23
Cigars	0.32	0.30	0.28	0.31	0.30	0.28
Pipe Tobacco	0.13	0.12	0.13	0.13	0.15	0.16
<b>ALCOHOL</b>	0.64	0.58	0.59	0.59	0.56	0.61
Binge Alcohol Use <sup>2</sup>	0.53	0.51	0.49	0.50	0.53	0.54
Heavy Alcohol Use <sup>2</sup>	0.34	0.31	0.31	0.31	0.33	0.35
<b>FEMALE</b>						
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.47	0.45	0.44	0.48	0.44	0.44
Cigarettes	0.45	0.44	0.44	0.48	0.43	0.42
Smokeless Tobacco	0.09	0.11	0.06	0.08	0.07	0.10
Cigars	0.10	0.11	0.11	0.10	0.11	0.10
Pipe Tobacco	0.08	0.03	0.04	0.04	0.04	0.02
<b>ALCOHOL</b>	0.59	0.54	0.59	0.56	0.56	0.58
Binge Alcohol Use <sup>2</sup>	0.38	0.34	0.35	0.36	0.35	0.34
Heavy Alcohol Use <sup>2</sup>	0.16	0.15	0.16	0.14	0.15	0.15

\*Low precision; no estimate reported.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, 2005, 2006, and 2007.

**Table 8.26P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 or Older, by Gender: *P* Values from Tests of Differences of Percentages, 2007 Versus 2002, 2003, 2004, 2005, and 2006**

<b>Gender/Substance</b>	<b>2002</b>	<b>2003</b>	<b>2004</b>	<b>2005</b>	<b>2006</b>	<b>2007</b>
<b>TOTAL</b>						
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.0015	0.0357	0.3617	0.1145	0.0342	
Cigarettes	0.0014	0.0298	0.2710	0.1905	0.0861	
Smokeless Tobacco	0.4733	0.7743	0.0944	0.7615	0.4896	
Cigars	0.8134	0.8744	0.1069	0.1796	0.2331	
Pipe Tobacco	0.7178	0.1199	0.5055	0.3357	0.1883	
<b>ALCOHOL</b>	0.6869	0.0396	0.1045	0.2124	0.6667	
Binge Alcohol Use <sup>2</sup>	0.2631	0.0779	0.1194	0.1173	0.4100	
Heavy Alcohol Use <sup>2</sup>	0.6073	0.7238	0.8075	0.2623	0.9076	
<b>MALE</b>						
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.0205	0.3823	0.5622	0.3911	0.0785	
Cigarettes	0.0293	0.2222	0.4309	0.6884	0.2377	
Smokeless Tobacco	0.5619	0.8452	0.0894	0.6293	0.3403	
Cigars	0.4539	0.8289	0.0843	0.1287	0.4889	
Pipe Tobacco	0.3839	0.1164	0.4743	0.5504	0.2846	
<b>ALCOHOL</b>	0.3056	0.3108	0.6705	0.0334	0.5835	
Binge Alcohol Use <sup>2</sup>	0.4755	0.2625	0.3102	0.1010	0.5269	
Heavy Alcohol Use <sup>2</sup>	0.7708	0.6171	0.9953	0.4748	0.8477	
<b>FEMALE</b>						
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.0075	0.0138	0.3864	0.1127	0.1488	
Cigarettes	0.0052	0.0297	0.3672	0.1107	0.1480	
Smokeless Tobacco	0.4776	0.2214	0.8503	0.6734	0.5648	
Cigars	0.2606	0.2331	0.8248	0.9003	0.1169	
Pipe Tobacco	0.2581	0.9486	0.8376	0.0929	0.2545	
<b>ALCOHOL</b>	0.1218	0.0001	0.0059	0.8694	0.2385	
Binge Alcohol Use <sup>2</sup>	0.3639	0.1164	0.1585	0.5967	0.5154	
Heavy Alcohol Use <sup>2</sup>	0.1015	0.7538	0.5033	0.2230	0.9005	

\*Low precision; no estimate reported.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, 2005, 2006, and 2007.