

Table 1.11A Illicit Drug Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2007 and 2008

Age Category	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	114,275 ^a	117,325	35,692	35,525	19,857	20,077
12	398	432	216 ^a	278	107	118
13	675	599	419	389	163	136
14	885	883	608	609	278	274
15	1,273	1,239	939	888	483	458
16	1,673	1,552	1,272	1,191	657	584
17	1,717	1,815	1,272	1,372	720	735
18	2,236	2,327	1,658	1,702	987	996
19	2,337	2,285	1,606	1,635	978	966
20	2,332	2,210	1,495	1,465	898	883
21	2,463	2,348	1,539	1,417	944	830
22	2,261	2,424	1,242	1,404	757	833
23	2,482	2,412	1,250	1,176	692	704
24	2,446	2,306	1,130	1,137	636	669
25	2,239	2,331	944	1,084	572	583
26-29	9,484 ^a	10,437	3,772	3,989	2,092	2,216
30-34	10,493	10,291	3,201	3,052	1,772	1,781
35-39	11,811	11,174	2,931	2,950	1,543	1,738
40-44	12,437	12,891	2,773	2,765	1,480	1,342
45-49	13,778	14,114	2,689	2,582	1,615	1,569
50-54	12,794	12,756	2,297	1,789	1,242	940
55-59	9,505	9,233	1,472	1,361	751	885
60-64	4,658 ^b	6,248	588	782	251	454
65 or Older	3,899 ^a	5,016	378	508	238	384

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, based on data from original questions not including methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 1.11B Illicit Drug Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2007 and 2008

Age Category	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	46.1	47.0	14.4	14.2	8.0	8.0
12	9.9	11.2	5.4 ^a	7.2	2.7	3.1
13	16.4	15.2	10.2	9.9	4.0	3.4
14	21.4	21.5	14.7	14.9	6.7	6.7
15	29.0	28.4	21.4	20.4	11.0	10.5
16	37.6	35.8	28.6	27.5	14.8	13.5
17	41.5	42.0	30.8	31.8	17.4	17.0
18	46.9	47.5	34.8	34.7	20.7	20.3
19	53.3	52.5	36.6	37.6	22.3	22.2
20	57.1	55.7	36.6	36.9	22.0	22.3
21	60.3	59.0	37.6	35.6	23.1	20.9
22	59.6	60.1	32.8	34.8	20.0	20.7
23	62.7 ^a	59.1	31.6	28.8	17.5	17.3
24	62.1	59.9	28.7	29.6	16.1	17.4
25	60.1	61.7	25.4 ^a	28.7	15.4	15.4
26-29	57.8 ^a	61.1	23.0	23.4	12.8	13.0
30-34	55.5	55.4	16.9	16.4	9.4	9.6
35-39	56.1	55.2	13.9	14.6	7.3	8.6
40-44	58.6	60.2	13.1	12.9	7.0	6.3
45-49	61.0	62.7	11.9	11.5	7.2	7.0
50-54	58.9	58.0	10.6	8.1	5.7	4.3
55-59	51.6	51.9	8.0	7.7	4.1	5.0
60-64	35.0 ^a	41.3	4.4	5.2	1.9	3.0
65 or Older	10.7 ^a	13.5	1.0	1.4	0.7	1.0

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, based on data from original questions not including methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 1.12A Marijuana Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2007 and 2008

Age Category	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	100,518	102,404	25,085	25,768	14,448	15,203
12	51	56	41	44	23	20
13	167	182	120	151	52	54
14	410	400	321	333	154	147
15	787	837	630	645	334	336
16	1,224	1,149	970	916	506	473
17	1,444	1,484	1,065	1,149	617	626
18	1,883	1,988	1,348	1,436	814	872
19	2,050	2,001	1,376	1,403	829	844
20	2,075	1,973	1,287	1,250	786	750
21	2,201	2,099	1,267	1,211	773	709
22	2,014	2,162	1,037	1,150	631	697
23	2,256	2,178	992	915	578	572
24	2,174	2,084	921	897	499	511
25	1,966	2,107	756	825	467	471
26-29	8,528	9,329	2,851	3,107	1,600	1,789
30-34	9,220	8,962	2,102	2,191	1,199	1,340
35-39	10,494	9,817	1,874	2,032	1,145	1,226
40-44	11,340	11,796	1,704	1,701	949	869
45-49	12,849	12,969	1,775	1,705	1,110	1,069
50-54	11,910	11,692	1,465	1,166	822	668
55-59	8,507	8,444	818	910	395	692
60-64	3,936 ^b	5,236	255	479	85 ^a	349
65 or Older	3,031	3,456	108	152	78	119

*Low precision; no estimate reported.

^a Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 1.12B Marijuana Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2007 and 2008

Age Category	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	40.6	41.0	10.1	10.3	5.8	6.1
12	1.3	1.5	1.0	1.1	0.6	0.5
13	4.1	4.6	2.9	3.8	1.3	1.4
14	9.9	9.8	7.8	8.1	3.7	3.6
15	17.9	19.2	14.4	14.8	7.6	7.7
16	27.5	26.5	21.8	21.2	11.4	10.9
17	34.9	34.4	25.8	26.6	14.9	14.5
18	39.5	40.6	28.3	29.3	17.1	17.8
19	46.8	46.0	31.4	32.2	18.9	19.4
20	50.8	49.8	31.5	31.5	19.2	18.9
21	53.8	52.7	31.0	30.4	18.9	17.8
22	53.1	53.6	27.4	28.5	16.7	17.3
23	57.0	53.4	25.1	22.4	14.6	14.0
24	55.2	54.2	23.4	23.3	12.7	13.3
25	52.8	55.8	20.3	21.9	12.5	12.5
26-29	52.0	54.6	17.4	18.2	9.8	10.5
30-34	48.8	48.3	11.1	11.8	6.3	7.2
35-39	49.8	48.5	8.9	10.0	5.4	6.1
40-44	53.5	55.1	8.0	7.9	4.5	4.1
45-49	56.9	57.6	7.9	7.6	4.9	4.7
50-54	54.8	53.2	6.7	5.3	3.8	3.0
55-59	46.2	47.5	4.4	5.1	2.1	3.9
60-64	29.6	34.6	1.9	3.2	0.6 ^a	2.3
65 or Older	8.4	9.3	0.3	0.4	0.2	0.3

*Low precision; no estimate reported.

^a Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 1.13A Cocaine Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2007 and 2008

Age Category	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	35,882	36,773	5,738	5,255	2,075	1,855
12	6	2	3	1	2	*
13	14	15	14	8	5	1
14	26	20	23	14	6	5
15	78	81	63	67	12	27
16	169 ^a	122	108	78	32	22
17	250	232	173	138	47	43
18	384	367	242	210	61	49
19	490	538	280	296	85	82
20	569	536	299	257	83	80
21	593	572	286	236	84	93
22	657	666	306	267	65	46
23	784	717	266	214	69	53
24	753	680	201	190	65	54
25	688	653	203	149	60	49
26-29	2,947	3,287	718	900	246	302
30-34	2,910	2,675	543	521	252	247
35-39	4,011	3,677	431	450	166	158
40-44	5,272	5,181	403	422	188	155
45-49	6,131	6,486	547	420	311	181
50-54	5,291	5,112	397	210	114	98
55-59	2,589 ^a	3,409	159	96	68	58
60-64	854 ^a	1,341	2	95	*	36
65 or Older	418	402	71	16	53	16

*Low precision; no estimate reported.

^a Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 1.13B Cocaine Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2007 and 2008

Age Category	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	14.5	14.7	2.3	2.1	0.8	0.7
12	0.2	0.0	0.1	0.0	0.0	*
13	0.3	0.4	0.3	0.2	0.1	0.0
14	0.6	0.5	0.5	0.3	0.1	0.1
15	1.8	1.8	1.4	1.5	0.3	0.6
16	3.8	2.8	2.4	1.8	0.7	0.5
17	6.0	5.4	4.2	3.2	1.1	1.0
18	8.1	7.5	5.1	4.3	1.3	1.0
19	11.2	12.3	6.4	6.8	2.0	1.9
20	13.9	13.5	7.3	6.5	2.0	2.0
21	14.5	14.4	7.0	5.9	2.1	2.3
22	17.3	16.5	8.1	6.6	1.7	1.1
23	19.8	17.6	6.7	5.2	1.7	1.3
24	19.1	17.7	5.1	4.9	1.6	1.4
25	18.5	17.3	5.5	3.9	1.6	1.3
26-29	18.0	19.3	4.4	5.3	1.5	1.8
30-34	15.4	14.4	2.9	2.8	1.3	1.3
35-39	19.0	18.2	2.0	2.2	0.8	0.8
40-44	24.9	24.2	1.9	2.0	0.9	0.7
45-49	27.2	28.8	2.4	1.9	1.4	0.8
50-54	24.4	23.2	1.8	1.0	0.5	0.4
55-59	14.1 ^b	19.2	0.9	0.5	0.4	0.3
60-64	6.4	8.9	0.0	0.6	*	0.2
65 or Older	1.2	1.1	0.2	0.0	0.1	0.0

*Low precision; no estimate reported.

^a Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 1.14A Crack Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2007 and 2008

Age Category	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	8,581	8,445	1,451 ^a	1,109	610 ^b	359
12	*	*	*	*	*	*
13	5	*	3	*	0	*
14	9	6	7	1	2	*
15	18	16	16	11	3	3
16	31	20	21	8	6	2
17	43	36	17	13	4	1
18	57	59	22	29	7	8
19	85	75	30	28	9	6
20	102	109	40	33	7	8
21	128	106	24	24	3	12
22	149	155	36	43	9	2
23	144	120	30	16	7	6
24	156	148	37	22	6	4
25	160	131	28	20	10	7
26-29	669	823	115	133	59	20
30-34	885	873	89	126	37	71
35-39	1,076	1,028	172	135	77	24
40-44	1,407	1,417	186	171	103	57
45-49	1,554	1,621	297	167	174	83
50-54	1,232	948	170	71	42	33
55-59	470	639	76	11	33	*
60-64	77	94	*	32	*	11
65 or Older	125	19	32	16	13	*

*Low precision; no estimate reported.

^a Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 1.14B Crack Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2007 and 2008

Age Category	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	3.5	3.4	0.6 ^a	0.4	0.2 ^b	0.1
12	*	*	*	*	*	*
13	0.1	*	0.1	*	0.0	*
14	0.2	0.2	0.2	0.0	0.1	*
15	0.4	0.4	0.4	0.3	0.1	0.1
16	0.7	0.5	0.5	0.2	0.1	0.0
17	1.0	0.8	0.4	0.3	0.1	0.0
18	1.2	1.2	0.5	0.6	0.1	0.2
19	1.9	1.7	0.7	0.6	0.2	0.1
20	2.5	2.8	1.0	0.8	0.2	0.2
21	3.1	2.7	0.6	0.6	0.1	0.3
22	3.9	3.9	1.0	1.1	0.2	0.1
23	3.6	2.9	0.8	0.4	0.2	0.1
24	4.0	3.8	0.9	0.6	0.1	0.1
25	4.3	3.5	0.8	0.5	0.3	0.2
26-29	4.1	4.8	0.7	0.8	0.4	0.1
30-34	4.7	4.7	0.5	0.7	0.2	0.4
35-39	5.1	5.1	0.8	0.7	0.4	0.1
40-44	6.6	6.6	0.9	0.8	0.5	0.3
45-49	6.9	7.2	1.3	0.7	0.8	0.4
50-54	5.7	4.3	0.8	0.3	0.2	0.2
55-59	2.6	3.6	0.4	0.1	0.2	*
60-64	0.6	0.6	*	0.2	*	0.1
65 or Older	0.3	0.1	0.1	0.0	0.0	*

*Low precision; no estimate reported.

^a Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 1.15A Hallucinogen Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2007 and 2008

Age Category	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	34,215 ^a	35,963	3,762	3,678	996	1,060
12	20	32	14	21	1	8
13	45	40	31	34	18	14
14	85	84	59	59	15	30
15	196	179	143	143	37	56
16	239	269	163	196	45	69
17	311	368	219	237	60	65
18	499	547	278	352	87	101
19	641	669	362	378	81	91
20	716	658	341	354	73	110
21	844 ^a	717	311	289	64	74
22	777	807	244	291	47	75
23	958 ^b	758	172	155	41	29
24	1,001 ^b	818	193	155	40	46
25	852	862	147	134	40	23
26-29	3,942	4,292	463	343	84	128
30-34	3,812	3,604	359	253	147	87
35-39	3,788	3,943	124	145	46	40
40-44	3,648	3,741	52	50	24	*
45-49	4,152	4,683	51	39	34	5
50-54	4,472	3,830	30	11	10	2
55-59	2,417 ^a	3,331	*	33	*	8
60-64	590 ^b	1,460	*	*	*	*
65 or Older	211	271	9	4	*	*

*Low precision; no estimate reported.

^a Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 1.15B Hallucinogen Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2007 and 2008

Age Category	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	13.8	14.4	1.5	1.5	0.4	0.4
12	0.5	0.8	0.3	0.5	0.0	0.2
13	1.1	1.0	0.8	0.9	0.4	0.3
14	2.0	2.0	1.4	1.4	0.4	0.7
15	4.5	4.1	3.2	3.3	0.8	1.3
16	5.4	6.2	3.7	4.5	1.0 ^a	1.6
17	7.5	8.5	5.3	5.5	1.5	1.5
18	10.5	11.2	5.8	7.2	1.8	2.1
19	14.6	15.4	8.3	8.7	1.9	2.1
20	17.5	16.6	8.3	8.9	1.8	2.8
21	20.7	18.0	7.6	7.3	1.6	1.9
22	20.5	20.0	6.4	7.2	1.2	1.9
23	24.2 ^b	18.6	4.3	3.8	1.0	0.7
24	25.4 ^b	21.3	4.9	4.0	1.0	1.2
25	22.9	22.8	3.9	3.6	1.1	0.6
26-29	24.0	25.1	2.8	2.0	0.5	0.7
30-34	20.2	19.4	1.9	1.4	0.8	0.5
35-39	18.0	19.5	0.6	0.7	0.2	0.2
40-44	17.2	17.5	0.2	0.2	0.1	*
45-49	18.4 ^a	20.8	0.2	0.2	0.1	0.0
50-54	20.6	17.4	0.1	0.1	0.0	0.0
55-59	13.1 ^b	18.7	*	0.2	*	0.0
60-64	4.4 ^b	9.7	*	*	*	*
65 or Older	0.6	0.7	0.0	0.0	*	*

*Low precision; no estimate reported.

^a Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 1.16A Inhalant Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2007 and 2008

Age Category	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	22,477	22,274	2,080	2,047	616	640
12	245	264	106	143	41	52
13	418	365	198	187	50	42
14	431	451	203	209	55	62
15	486	426	186	167	60	49
16	424	413	166	156	52	41
17	418	388	135	117	36	21
18	369	421	113	118	23	29
19	422	411	116	99	23	13
20	400	409	77	103	21	25
21	430	383	62	66	14	10
22	477	423	60	40	24 ^a	5
23	514	487	38	37	14	7
24	523 ^a	426	28	50	3	10
25	556 ^a	451	31	19	8	5
26-29	2,283	2,400	132	99	43	21
30-34	2,644 ^a	2,170	99	76	26	22
35-39	2,675	2,572	123	76	5	35
40-44	2,502	2,609	71	64	28	50
45-49	2,550	2,860	40	91	30	78
50-54	2,124	1,593	69	7	33	5
55-59	1,010	1,358	25	48	25	48
60-64	209 ^b	592	*	*	*	2
65 or Older	368	399	*	8	*	8

*Low precision; no estimate reported.

^a Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 1.16B Inhalant Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2007 and 2008

Age Category	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	9.1	8.9	0.8	0.8	0.2	0.3
12	6.1	6.9	2.6 ^a	3.7	1.0	1.4
13	10.1	9.3	4.8	4.7	1.2	1.1
14	10.4	11.0	4.9	5.1	1.3	1.5
15	11.1	9.8	4.2	3.8	1.4	1.1
16	9.5	9.5	3.7	3.6	1.2	1.0
17	10.1	9.0	3.3	2.7	0.9	0.5
18	7.7	8.6	2.4	2.4	0.5	0.6
19	9.6	9.4	2.6	2.3	0.5	0.3
20	9.8	10.3	1.9	2.6	0.5	0.6
21	10.5	9.6	1.5	1.6	0.3	0.3
22	12.6	10.5	1.6	1.0	0.6 ^a	0.1
23	13.0	11.9	1.0	0.9	0.4	0.2
24	13.3	11.1	0.7	1.3	0.1	0.3
25	14.9 ^a	12.0	0.8	0.5	0.2	0.1
26-29	13.9	14.1	0.8	0.6	0.3	0.1
30-34	14.0 ^a	11.7	0.5	0.4	0.1	0.1
35-39	12.7	12.7	0.6	0.4	0.0	0.2
40-44	11.8	12.2	0.3	0.3	0.1	0.2
45-49	11.3	12.7	0.2	0.4	0.1	0.3
50-54	9.8	7.2	0.3	0.0	0.2	0.0
55-59	5.5	7.6	0.1	0.3	0.1	0.3
60-64	1.6 ^a	3.9	*	*	*	0.0
65 or Older	1.0	1.1	*	0.0	*	0.0

*Low precision; no estimate reported.

^a Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 1.17A Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2007 and 2008

Age Category	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	50,415	51,970	16,280 ^a	15,166	6,895 ^a	6,224
12	171	201	97	124	53	55
13	262	216	160	148	65	59
14	357	348	244	228	108	92
15	610 ^a	512	463 ^a	367	183	160
16	761	704	535	513	188	170
17	788	791	535	533	232 ^a	175
18	1,141	1,000	762	672	305	237
19	1,156	1,133	700	689	285	271
20	1,247	1,190	666	651	260	281
21	1,344 ^a	1,158	711 ^a	569	277	244
22	1,214	1,329	556	662	239	271
23	1,267	1,289	563	566	197	225
24	1,274	1,288	507	569	197	234
25	1,146	1,239	449	489	190	180
26-29	4,822	5,273	1,669	1,646	644	587
30-34	4,769	4,713	1,534	1,331	594	541
35-39	5,145	4,938	1,469	1,250	564	584
40-44	5,111	5,289	1,281	1,268	625	541
45-49	5,790	6,080	1,077	1,106	520	507
50-54	5,157	5,325	883	706	468	273
55-59	4,011	3,802	727	471	331	184
60-64	1,681	2,159	387	274	193	113
65 or Older	1,191 ^b	1,994	307	333	178	242

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 1.17B Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2007 and 2008

Age Category	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	20.3	20.8	6.6 ^a	6.1	2.8 ^a	2.5
12	4.3	5.2	2.4	3.2	1.3	1.4
13	6.4	5.5	3.9	3.8	1.6	1.5
14	8.6	8.5	5.9	5.6	2.6	2.2
15	13.9 ^a	11.7	10.5 ^a	8.4	4.2	3.7
16	17.1	16.3	12.0	11.9	4.2	3.9
17	19.1	18.3	12.9	12.3	5.6 ^a	4.0
18	23.9 ^b	20.4	16.0	13.7	6.4 ^a	4.8
19	26.4	26.0	16.0	15.8	6.5	6.2
20	30.5	30.0	16.3	16.4	6.4	7.1
21	32.9 ^a	29.1	17.4 ^a	14.3	6.8	6.1
22	32.0	33.0	14.7	16.4	6.3	6.7
23	32.0	31.6	14.2	13.9	5.0	5.5
24	32.3	33.5	12.9	14.8	5.0	6.1
25	30.8	32.8	12.1	13.0	5.1	4.8
26-29	29.4	30.9	10.2	9.6	3.9	3.4
30-34	25.2	25.4	8.1	7.2	3.1	2.9
35-39	24.4	24.4	7.0	6.2	2.7	2.9
40-44	24.1	24.7	6.0	5.9	2.9	2.5
45-49	25.7	27.0	4.8	4.9	2.3	2.3
50-54	23.7	24.2	4.1	3.2	2.2	1.2
55-59	21.8	21.4	3.9	2.6	1.8	1.0
60-64	12.6	14.3	2.9	1.8	1.4	0.7
65 or Older	3.3 ^b	5.4	0.8	0.9	0.5	0.6

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 1.18A Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2007 and 2008

Age Category	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	33,060 ^a	34,861	12,466	11,885	5,174	4,747
12	139	175	87	114	45	48
13	195	175	130	119	52	49
14	311	289	200	196	82	72
15	498	449	370	321	135	128
16	653	594	448	425	154	130
17	656	665	446	433	200 ^a	140
18	926	881	610	568	237	200
19	995	978	590	554	239	224
20	1,041	1,003	519	512	203	215
21	1,146	994	582 ^a	466	217	187
22	1,032	1,088	442	534	188	201
23	1,055	1,074	436	460	153	172
24	1,003	1,041	423	445	143	173
25	930	982	368	403	137	129
26-29	3,713	4,141	1,279	1,318	522	464
30-34	3,479	3,458	1,173	1,040	464	406
35-39	3,505	3,237	1,120	922	406	426
40-44	3,086	3,386	959	963	432	398
45-49	3,227	3,377	808	829	393	366
50-54	2,380	2,756	643	476	350	213
55-59	1,857	1,865	427	283	226	103
60-64	762	1,170	201	209	93	77
65 or Older	471 ^b	1,083	205	296	102	226

*Low precision; no estimate reported.

^a Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 1.18B Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2007 and 2008

Age Category	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	13.3	14.0	5.0	4.8	2.1	1.9
12	3.4	4.5	2.2	3.0	1.1	1.2
13	4.7	4.4	3.1	3.0	1.3	1.2
14	7.5	7.0	4.8	4.8	2.0	1.8
15	11.4	10.3	8.4	7.4	3.1	2.9
16	14.7	13.7	10.1	9.8	3.5	3.0
17	15.9	15.4	10.8	10.0	4.8 ^b	3.2
18	19.4	18.0	12.8	11.6	5.0	4.1
19	22.7	22.5	13.5	12.7	5.5	5.1
20	25.5	25.3	12.7	12.9	5.0	5.4
21	28.0	25.0	14.2 ^a	11.7	5.3	4.7
22	27.2	27.0	11.6	13.2	5.0	5.0
23	26.7	26.3	11.0	11.3	3.9	4.2
24	25.5	27.1	10.7	11.6	3.6	4.5
25	25.0	26.0	9.9	10.7	3.7	3.4
26-29	22.6	24.3	7.8	7.7	3.2	2.7
30-34	18.4	18.6	6.2	5.6	2.5	2.2
35-39	16.6	16.0	5.3	4.6	1.9	2.1
40-44	14.5	15.8	4.5	4.5	2.0	1.9
45-49	14.3	15.0	3.6	3.7	1.7	1.6
50-54	11.0	12.5	3.0	2.2	1.6	1.0
55-59	10.1	10.5	2.3	1.6	1.2	0.6
60-64	5.7	7.7	1.5	1.4	0.7	0.5
65 or Older	1.3 ^b	2.9	0.6	0.8	0.3	0.6

*Low precision; no estimate reported.

^a Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.