

Table 1.59A (REVISED) Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics:
Numbers in Thousands, 2007 and 2008

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	114,509 ^a	117,815	35,789	35,643	19,892	20,123
GEOGRAPHIC DIVISION						
Northeast	21,018	21,840	6,498	6,310	3,595	3,812
New England	5,975	6,308	1,827	1,859	1,054	1,200
Middle Atlantic	15,044	15,532	4,670	4,451	2,541	2,612
Midwest	26,012	26,107	7,608	7,505	4,303	4,172
East North Central	18,344	18,348	5,574	5,511	3,144	3,136
West North Central	7,668	7,759	2,034	1,994	1,160	1,036
South	39,781	40,988	12,265	12,049	6,649	6,450
South Atlantic	21,451	22,591	6,516	6,676	3,507	3,588
East South Central	6,499	6,117	2,100	1,882	1,190	1,067
West South Central	11,831	12,280	3,649	3,491	1,952	1,796
West	27,698	28,880	9,419	9,779	5,345	5,688
Mountain	8,705	8,742	2,791	2,777	1,617	1,625
Pacific	18,994	20,137	6,628	7,002	3,727	4,063
COUNTY TYPE						
Large Metro	62,848	63,996	19,887	19,965	11,046	11,361
Small Metro	34,396	35,691	10,784	10,708	6,080	6,158
250K – 1 Mil. Pop.	22,910	23,821	7,238	7,102	4,108	4,255
< 250K Pop.	11,486	11,870	3,546	3,606	1,972	1,903
Nonmetro	17,265	18,127	5,118	4,969	2,766	2,605
Urbanized	7,732	8,123	2,427	2,267	1,255	1,237
Less Urbanized	7,882	8,416	2,387	2,325	1,324	1,122
Completely Rural	1,652	1,588	304	377	187	245

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, based on data from original questions not including methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 1.59B (REVISED) Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2007 and 2008

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	46.2	47.2	14.4	14.3	8.0	8.1
GEOGRAPHIC DIVISION						
Northeast	45.8	47.4	14.2	13.7	7.8	8.3
New England	49.7	52.3	15.2	15.4	8.8	10.0
Middle Atlantic	44.4	45.6	13.8	13.1	7.5	7.7
Midwest	47.5	47.5	13.9	13.7	7.9	7.6
East North Central	47.9	47.8	14.6	14.4	8.2	8.2
West North Central	46.5	46.8	12.3	12.0	7.0	6.3
South	44.2	45.1	13.6	13.2	7.4	7.1
South Atlantic	45.1	47.1	13.7	13.9	7.4	7.5
East South Central	44.1	41.2	14.3	12.7	8.1	7.2
West South Central	42.8	43.7	13.2	12.4	7.1	6.4
West	48.4	50.0	16.5	16.9	9.3	9.8
Mountain	50.4	49.7	16.2	15.8	9.4	9.2
Pacific	47.5	50.1	16.6	17.4	9.3	10.1
COUNTY TYPE						
Large Metro	47.4	48.1	15.0	15.0	8.3	8.5
Small Metro	46.4	47.3	14.6	14.2	8.2	8.2
250K – 1 Mil. Pop.	46.2	47.6	14.6	14.2	8.3	8.5
< 250K Pop.	46.8	46.6	14.4	14.2	8.0	7.5
Nonmetro	42.0	43.9	12.5	12.0	6.7	6.3
Urbanized	46.0	47.3	14.4	13.2	7.5	7.2
Less Urbanized	40.1	42.0	12.1	11.6	6.7	5.6
Completely Rural	35.9	39.4	6.6	9.4	4.1	6.1

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, based on data from original questions not including methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 1.60A (REVISED) Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics:
Numbers in Thousands, 2007 and 2008

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	6,638	6,565	4,737	4,751	2,415	2,312
GEOGRAPHIC DIVISION						
Northeast	1,058	1,080	786	822	410	459
New England	324	291	248	220	135	122
Middle Atlantic	734	789	537	602	275 ^a	337
Midwest	1,502 ^a	1,374	1,070	1,015	537	501
East North Central	1,105 ^a	993	783	734	392	375
West North Central	397	381	287	281	145	126
South	2,428	2,365	1,696	1,618	851 ^a	719
South Atlantic	1,234	1,325	844	906	416	405
East South Central	392 ^a	331	267 ^a	215	120	94
West South Central	802	709	585 ^a	497	315 ^b	220
West	1,650	1,746	1,185	1,297	617	633
Mountain	526	554	391	409	203	211
Pacific	1,124	1,193	795	888	414	422
COUNTY TYPE						
Large Metro	3,547	3,425	2,497	2,534	1,300	1,241
Small Metro	1,985	2,064	1,476	1,455	757	746
250K – 1 Mil. Pop.	1,318	1,405	981	1,000	506	495
< 250K Pop.	667	659	495	455	251	252
Nonmetro	1,106	1,075	764	762	358	325
Urbanized	467	491	328	365	173	159
Less Urbanized	544	499	369	344	149	139
Completely Rural	94	85	67	53	37	26

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, based on data from original questions not including methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 1.60B (REVISED) Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Percentages, 2007 and 2008

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	26.3	26.4	18.8	19.1	9.6	9.3
GEOGRAPHIC DIVISION						
Northeast	23.7	24.7	17.6	18.8	9.2	10.5
New England	28.0	25.7	21.5	19.4	11.7	10.8
Middle Atlantic	22.2	24.3	16.3	18.6	8.3 ^a	10.4
Midwest	26.7	24.9	19.1	18.4	9.6	9.1
East North Central	28.0 ^a	25.7	19.9	19.0	9.9	9.7
West North Central	23.7	23.2	17.2	17.1	8.7	7.7
South	26.6	26.1	18.6	17.9	9.3 ^a	7.9
South Atlantic	26.6	29.0	18.2	19.8	9.0	8.9
East South Central	26.4 ^a	22.5	18.0 ^a	14.6	8.1	6.4
West South Central	26.7	23.6	19.5 ^a	16.5	10.5 ^b	7.3
West	27.3	29.3	19.6	21.8	10.2	10.6
Mountain	29.1	30.6	21.6	22.6	11.2	11.7
Pacific	26.6	28.7	18.8	21.4	9.8	10.2
COUNTY TYPE						
Large Metro	25.9	25.8	18.2	19.0	9.5	9.3
Small Metro	26.9	27.9	20.0	19.6	10.3	10.1
250K – 1 Mil. Pop.	26.4	28.7	19.7	20.4	10.2	10.1
< 250K Pop.	27.9	26.3	20.7	18.1	10.5	10.0
Nonmetro	26.5	25.7	18.3	18.2	8.6	7.8
Urbanized	27.2	27.5	19.1	20.4	10.1	8.9
Less Urbanized	26.9	24.8	18.2	17.1	7.4	6.9
Completely Rural	22.0	22.6	15.8	14.1	8.6	6.8

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, based on data from original questions not including methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 1.61A (REVISED) Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics:
Numbers in Thousands, 2007 and 2008

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	18,831	18,772	10,889	11,090	6,475	6,490
GEOGRAPHIC DIVISION						
Northeast	3,399 ^a	3,600	2,090	2,243	1,262 ^a	1,421
New England	1,005	982	663	631	399	407
Middle Atlantic	2,394 ^b	2,618	1,427 ^a	1,611	864 ^a	1,015
Midwest	4,351	4,213	2,437	2,414	1,448	1,370
East North Central	3,082 ^b	2,901	1,767	1,707	1,051	984
West North Central	1,269	1,312	670	707	397	385
South	6,527	6,377	3,650	3,609	2,104	2,064
South Atlantic	3,419	3,276	2,005	1,898	1,173	1,075
East South Central	1,023	970	566	562	346	350
West South Central	2,085	2,131	1,079	1,150	585	638
West	4,554	4,581	2,712	2,824	1,661	1,635
Mountain	1,376	1,452	781	814	492	484
Pacific	3,178	3,129	1,931	2,010	1,169	1,151
COUNTY TYPE						
Large Metro	9,773	10,081	5,775	6,086	3,515	3,584
Small Metro	6,131	5,904	3,520	3,485	2,021	2,045
250K – 1 Mil. Pop.	4,035	3,788	2,345	2,207	1,365	1,294
< 250K Pop.	2,096	2,116	1,175	1,278	656	750
Nonmetro	2,927	2,787	1,594	1,519	939	861
Urbanized	1,333	1,350	758	741	460	404
Less Urbanized	1,405	1,263	726	688	422	399
Completely Rural	190	174	109	90	57	58

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, based on data from original questions not including methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 1.61B (REVISED) Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Percentages, 2007 and 2008

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	57.5	57.0	33.3	33.7	19.8	19.7
GEOGRAPHIC DIVISION						
Northeast	57.6	60.1	35.4	37.5	21.4	23.7
New England	65.8	63.1	43.4	40.6	26.1	26.1
Middle Atlantic	54.7 ^a	59.1	32.6 ^a	36.4	19.7 ^a	22.9
Midwest	59.7	57.9	33.4	33.2	19.9	18.8
East North Central	61.3 ^b	57.6	35.2	33.9	20.9	19.6
West North Central	56.1	58.6	29.6	31.6	17.5	17.2
South	55.8	54.2	31.2	30.7	18.0	17.5
South Atlantic	57.9	54.9	33.9	31.8	19.9	18.0
East South Central	54.7	51.8	30.3	30.0	18.5	18.7
West South Central	53.3	54.3	27.6	29.3	15.0	16.3
West	58.0	57.9	34.5	35.7	21.2	20.7
Mountain	58.4	61.3	33.2	34.3	20.9	20.4
Pacific	57.8	56.5	35.1	36.3	21.3	20.8
COUNTY TYPE						
Large Metro	56.5	56.5	33.4	34.1	20.3	20.1
Small Metro	59.4	58.5	34.1	34.5	19.6	20.2
250K – 1 Mil. Pop.	59.6	57.9	34.6	33.7	20.2	19.8
< 250K Pop.	59.0	59.5	33.1	35.9	18.5	21.1
Nonmetro	57.4	55.8	31.3	30.4	18.4	17.2
Urbanized	58.5	59.1	33.2	32.4	20.2	17.7
Less Urbanized	57.4	53.7	29.7	29.2	17.3	16.9
Completely Rural	51.0	48.9	29.5	25.4	15.3	16.2

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, based on data from original questions not including methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 1.62A (REVISED) Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics:
Numbers in Thousands, 2007 and 2008

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	89,040 ^b	92,478	20,163	19,801	11,001	11,321
GEOGRAPHIC DIVISION						
Northeast	16,561	17,160	3,622	3,245	1,922	1,932
New England	4,645	5,035	916	1,008	520	672
Middle Atlantic	11,916	12,125	2,707	2,237	1,402	1,260
Midwest	20,159	20,520	4,100	4,076	2,318	2,302
East North Central	14,157	14,454	3,024	3,069	1,700	1,777
West North Central	6,002	6,066	1,076	1,006	618	525
South	30,826	32,245	6,919	6,822	3,694	3,668
South Atlantic	16,797	17,991	3,667	3,872	1,918	2,108
East South Central	5,084	4,815	1,267	1,105	724	622
West South Central	8,944	9,440	1,985	1,845	1,052	938
West	21,494	22,552	5,522	5,659	3,067	3,420
Mountain	6,803	6,737	1,619	1,555	923	930
Pacific	14,691	15,815	3,903	4,103	2,144	2,490
COUNTY TYPE						
Large Metro	49,528	50,490	11,614	11,346	6,232	6,535
Small Metro	26,280	27,723	5,789	5,768	3,301	3,367
250K – 1 Mil. Pop.	17,558	18,628	3,912	3,895	2,237	2,466
< 250K Pop.	8,722	9,095	1,876	1,873	1,064	901
Nonmetro	13,232	14,265	2,760	2,688	1,468	1,420
Urbanized	5,932	6,283	1,341	1,160	622	674
Less Urbanized	5,932	6,653	1,292	1,293	753	584
Completely Rural	1,368	1,329	127	234	93	162

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, based on data from original questions not including methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 1.62B (REVISED) Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Percentages, 2007 and 2008

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	46.9	48.2	10.6	10.3	5.8	5.9
GEOGRAPHIC DIVISION						
Northeast	46.6	48.0	10.2	9.1	5.4	5.4
New England	49.7	53.7	9.8	10.8	5.6	7.2
Middle Atlantic	45.5	46.0	10.3	8.5	5.4	4.8
Midwest	48.1	48.7	9.8	9.7	5.5	5.5
East North Central	48.3	49.0	10.3	10.4	5.8	6.0
West North Central	47.8	47.8	8.6	7.9	4.9	4.1
South	44.6	46.0	10.0	9.7	5.3	5.2
South Atlantic	45.4	48.0	9.9	10.3	5.2	5.6
East South Central	44.7	41.8	11.1	9.6	6.4	5.4
West South Central	43.2	44.6	9.6	8.7	5.1	4.4
West	49.6	51.3	12.7	12.9	7.1	7.8
Mountain	51.9	50.2	12.3	11.6	7.0	6.9
Pacific	48.6	51.8	12.9	13.4	7.1	8.2
COUNTY TYPE						
Large Metro	48.7	49.5	11.4	11.1	6.1	6.4
Small Metro	46.6	47.8	10.3	10.0	5.9	5.8
250K – 1 Mil. Pop.	46.4	48.3	10.3	10.1	5.9	6.4
< 250K Pop.	46.9	46.9	10.1	9.6	5.7	4.6
Nonmetro	41.6	44.5	8.7	8.4	4.6	4.4
Urbanized	46.3	48.0	10.5	8.9	4.9	5.1
Less Urbanized	39.0	42.4	8.5	8.2	5.0	3.7
Completely Rural	36.0	40.3	3.3	7.1	2.4	4.9

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, based on data from original questions not including methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 1.63A (REVISED) Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics:
Numbers in Thousands, 2007 and 2008

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	107,871 ^a	111,250	31,052	30,892	17,477	17,811
GEOGRAPHIC DIVISION						
Northeast	19,961	20,760	5,712	5,488	3,184	3,353
New England	5,651	6,017	1,579	1,639	919	1,078
Middle Atlantic	14,310	14,743	4,133	3,848	2,266	2,275
Midwest	24,510	24,733	6,537	6,490	3,767	3,672
East North Central	17,239	17,355	4,791	4,777	2,752	2,761
West North Central	7,271	7,378	1,746	1,714	1,015	911
South	37,353	38,623	10,569	10,431	5,798	5,731
South Atlantic	20,216	21,267	5,671	5,770	3,091	3,183
East South Central	6,107	5,785	1,833	1,667	1,070	973
West South Central	11,029	11,571	3,064	2,994	1,637	1,576
West	26,048	27,133	8,234	8,482	4,728	5,055
Mountain	8,178	8,189	2,400	2,369	1,414	1,414
Pacific	17,870	18,945	5,833	6,114	3,313	3,641
COUNTY TYPE						
Large Metro	59,301	60,571	17,389	17,432	9,746	10,119
Small Metro	32,411	33,627	9,308	9,253	5,323	5,412
250K – 1 Mil. Pop.	21,593	22,416	6,257	6,102	3,602	3,761
< 250K Pop.	10,819	11,211	3,051	3,151	1,721	1,651
Nonmetro	16,159	17,052	4,354	4,207	2,407	2,280
Urbanized	7,264	7,632	2,099	1,901	1,082	1,078
Less Urbanized	7,337	7,917	2,018	1,981	1,176	983
Completely Rural	1,558	1,503	236	324	150	220

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, based on data from original questions not including methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 1.63B (REVISED) Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Percentages, 2007 and 2008

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	48.5	49.5	13.9	13.7	7.9	7.9
GEOGRAPHIC DIVISION						
Northeast	48.2	49.8	13.8	13.2	7.7	8.0
New England	52.0	55.1	14.5	15.0	8.5	9.9
Middle Atlantic	46.8	47.9	13.5	12.5	7.4	7.4
Midwest	49.8	50.0	13.3	13.1	7.7	7.4
East North Central	50.2	50.3	13.9	13.8	8.0	8.0
West North Central	49.0	49.4	11.8	11.5	6.8	6.1
South	46.2	47.2	13.1	12.7	7.2	7.0
South Atlantic	47.1	49.0	13.2	13.3	7.2	7.3
East South Central	46.1	43.2	13.8	12.4	8.1	7.3
West South Central	44.8	46.1	12.4	11.9	6.6	6.3
West	50.9	52.3	16.1	16.4	9.2	9.8
Mountain	52.9	51.9	15.5	15.0	9.1	9.0
Pacific	50.0	52.5	16.3	17.0	9.3	10.1
COUNTY TYPE						
Large Metro	49.8	50.6	14.6	14.6	8.2	8.4
Small Metro	48.6	49.4	14.0	13.6	8.0	8.0
250K – 1 Mil. Pop.	48.4	49.7	14.0	13.5	8.1	8.3
< 250K Pop.	48.8	48.8	13.8	13.7	7.8	7.2
Nonmetro	43.8	46.0	11.8	11.3	6.5	6.2
Urbanized	48.1	49.6	13.9	12.4	7.2	7.0
Less Urbanized	41.6	43.9	11.4	11.0	6.7	5.4
Completely Rural	37.3	41.2	5.7	8.9	3.6	6.0

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, based on data from original questions not including methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.