

**Table 1.76A (REVISED) Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics:**  
**Numbers in Thousands, 2007 and 2008**

<b>Geographic Characteristic</b>	<b>Lifetime (2007)</b>	<b>Lifetime (2008)</b>	<b>Past Year (2007)</b>	<b>Past Year (2008)</b>	<b>Past Month (2007)</b>	<b>Past Month (2008)</b>
<b>TOTAL</b>	22,558	22,430	2,085	2,065	617	644
<b>GEOGRAPHIC DIVISION</b>						
Northeast	4,322	4,238	390	352	139	78
New England	1,081	1,146	92	83	45	29
Middle Atlantic	3,241	3,092	298	269	95	49
Midwest	5,007	5,155	427	377	99	103
East North Central	3,474	3,626	314	264	66	78
West North Central	1,533	1,529	112	113	34	25
South	7,242	7,238	647	612	168	180
South Atlantic	3,885	4,302	342	290	89	81
East South Central	1,138	994	58	79	13	25
West South Central	2,220	1,942	248	242	67	73
West	5,987	5,799	621	724	210	284
Mountain	1,866	1,783	200	153	67	56
Pacific	4,121	4,016	421	571	143	228
<b>COUNTY TYPE</b>						
Large Metro	12,550	12,382	1,197	1,183	362	348
Small Metro	6,751	6,703	627	562	184	201
250K – 1 Mil. Pop.	4,493	4,659	392	380	101	158
< 250K Pop.	2,259	2,044	235	182	83	43
Nonmetro	3,256	3,345	261	319	71	96
Urbanized	1,536	1,561	114	164	31	47
Less Urbanized	1,512	1,517	121	134	33	43
Completely Rural	208	267	26	21	7	6

\*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

<sup>a</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 1.76B (REVISED) Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2007 and 2008**

<b>Geographic Characteristic</b>	<b>Lifetime (2007)</b>	<b>Lifetime (2008)</b>	<b>Past Year (2007)</b>	<b>Past Year (2008)</b>	<b>Past Month (2007)</b>	<b>Past Month (2008)</b>
<b>TOTAL</b>	9.1	9.0	0.8	0.8	0.2	0.3
<b>GEOGRAPHIC DIVISION</b>						
Northeast	9.4	9.2	0.8	0.8	0.3	0.2
New England	9.0	9.5	0.8	0.7	0.4	0.2
Middle Atlantic	9.6	9.1	0.9	0.8	0.3	0.1
Midwest	9.1	9.4	0.8	0.7	0.2	0.2
East North Central	9.1	9.4	0.8	0.7	0.2	0.2
West North Central	9.3	9.2	0.7	0.7	0.2	0.2
South	8.1	8.0	0.7	0.7	0.2	0.2
South Atlantic	8.2	9.0	0.7	0.6	0.2	0.2
East South Central	7.7	6.7	0.4	0.5	0.1	0.2
West South Central	8.0	6.9	0.9	0.9	0.2	0.3
West	10.5	10.0	1.1	1.3	0.4	0.5
Mountain	10.8	10.1	1.2	0.9	0.4	0.3
Pacific	10.3	10.0	1.1	1.4	0.4	0.6
<b>COUNTY TYPE</b>						
Large Metro	9.5	9.3	0.9	0.9	0.3	0.3
Small Metro	9.1	8.9	0.8	0.7	0.2	0.3
250K – 1 Mil. Pop.	9.1	9.3	0.8	0.8	0.2	0.3
< 250K Pop.	9.2	8.0	1.0	0.7	0.3	0.2
Nonmetro	7.9	8.1	0.6	0.8	0.2	0.2
Urbanized	9.1	9.1	0.7	1.0	0.2	0.3
Less Urbanized	7.7	7.6	0.6	0.7	0.2	0.2
Completely Rural	4.5	6.6	0.6	0.5	0.1	0.1

\*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

<sup>a</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 1.77A (REVISED) Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics:**  
**Numbers in Thousands, 2007 and 2008**

<b>Geographic Characteristic</b>	<b>Lifetime (2007)</b>	<b>Lifetime (2008)</b>	<b>Past Year (2007)</b>	<b>Past Year (2008)</b>	<b>Past Month (2007)</b>	<b>Past Month (2008)</b>
<b>TOTAL</b>	2,428	2,326	995	986	294	270
<b>GEOGRAPHIC DIVISION</b>						
Northeast	377	355	150	151	42	37
New England	113	98	44	44	17	14
Middle Atlantic	264	257	106	107	25	22
Midwest	520	484	226	208	54	56
East North Central	371	337	164	141	38	38
West North Central	148	147	62	67	16	18
South	883	813	350	308	97	74
South Atlantic	453	464	182	149	52	31
East South Central	137	121	45	47	9	12
West South Central	293 <sup>a</sup>	227	123	111	36	30
West	648	674	268	319	101	104
Mountain	210	236	90	91	31	30
Pacific	438	438	178	228	70	74
<b>COUNTY TYPE</b>						
Large Metro	1,290	1,177	511	519	148	146
Small Metro	716	706	323	274	99	85
250K – 1 Mil. Pop.	477	460	198	185	59	67
< 250K Pop.	239	247	125	90	40	19
Nonmetro	422	442	160	192	48	39
Urbanized	171	195	64	91	24	16
Less Urbanized	205	204	75	85	17	17
Completely Rural	46	43	21	16	6	6

\*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

<sup>a</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 1.77B (REVISED) Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics:**  
**Percentages, 2007 and 2008**

<b>Geographic Characteristic</b>	<b>Lifetime (2007)</b>	<b>Lifetime (2008)</b>	<b>Past Year (2007)</b>	<b>Past Year (2008)</b>	<b>Past Month (2007)</b>	<b>Past Month (2008)</b>
<b>TOTAL</b>	9.6	9.3	3.9	4.0	1.2	1.1
<b>GEOGRAPHIC DIVISION</b>						
Northeast	8.5	8.1	3.4	3.5	0.9	0.8
New England	9.7	8.7	3.8	3.9	1.5	1.3
Middle Atlantic	8.0	7.9	3.2	3.3	0.8	0.7
Midwest	9.3	8.8	4.0	3.8	1.0	1.0
East North Central	9.4	8.7	4.2	3.6	1.0	1.0
West North Central	8.9	8.9	3.7	4.1	0.9	1.1
South	9.7	9.0	3.8	3.4	1.1	0.8
South Atlantic	9.8	10.1	3.9	3.3	1.1	0.7
East South Central	9.3	8.3	3.1	3.2	0.6	0.8
West South Central	9.7 <sup>a</sup>	7.6	4.1	3.7	1.2	1.0
West	10.7	11.3	4.4	5.4	1.7	1.7
Mountain	11.6	13.1	5.0	5.1	1.7	1.7
Pacific	10.4	10.6	4.2	5.5	1.7	1.8
<b>COUNTY TYPE</b>						
Large Metro	9.4	8.8	3.7	3.9	1.1	1.1
Small Metro	9.7	9.5	4.4	3.7	1.3	1.2
250K – 1 Mil. Pop.	9.6	9.4	4.0	3.8	1.2	1.4
< 250K Pop.	10.0	9.8	5.2 <sup>a</sup>	3.6	1.7	0.7
Nonmetro	10.1	10.6	3.8	4.6	1.1	0.9
Urbanized	10.0	10.9	3.8	5.1	1.4	0.9
Less Urbanized	10.1	10.1	3.7	4.2	0.9	0.9
Completely Rural	10.7	11.5	5.0	4.3	1.5	1.5

\*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

<sup>a</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.