

Table 1.78A (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Numbers in Thousands, 2007 and 2008

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	50,541	52,154	16,334 ^a	15,190	6,907 ^a	6,210
GEOGRAPHIC DIVISION						
Northeast	8,038	8,806	2,583	2,427	987	1,037
New England	2,168	2,418	665	723	287	382
Middle Atlantic	5,871	6,387	1,919	1,705	700	655
Midwest	10,869	11,179	3,395	3,353	1,468	1,402
East North Central	7,637	7,866	2,499	2,462	1,051	1,057
West North Central	3,233	3,313	896	891	417	345
South	18,417	18,123	6,155 ^a	5,361	2,676 ^a	2,203
South Atlantic	9,350	9,753	2,880	2,692	1,174	1,104
East South Central	3,317	2,890	1,271 ^a	967	633	463
West South Central	5,751	5,480	2,005	1,703	870 ^a	635
West	13,216	14,047	4,200	4,048	1,777	1,568
Mountain	4,306	4,288	1,374	1,248	563	437
Pacific	8,910	9,759	2,826	2,801	1,214	1,131
COUNTY TYPE						
Large Metro	27,288	27,857	8,750	7,999	3,653	3,233
Small Metro	15,302	16,242	4,900	4,804	2,100	1,998
250K – 1 Mil. Pop.	10,290	10,943	3,190	3,148	1,297	1,399
< 250K Pop.	5,012	5,299	1,710	1,657	803	599
Nonmetro	7,951	8,055	2,684	2,386	1,154	978
Urbanized	3,522	3,494	1,187	1,019	440	437
Less Urbanized	3,723	3,845	1,346	1,223	635	484
Completely Rural	707	716	152	144	78	57

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 1.78B (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2007 and 2008

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	20.4	20.9	6.6 ^a	6.1	2.8 ^a	2.5
GEOGRAPHIC DIVISION						
Northeast	17.5	19.1	5.6	5.3	2.2	2.2
New England	18.0	20.1	5.5	6.0	2.4	3.2
Middle Atlantic	17.3	18.8	5.7	5.0	2.1	1.9
Midwest	19.8	20.3	6.2	6.1	2.7	2.6
East North Central	19.9	20.5	6.5	6.4	2.7	2.8
West North Central	19.6	20.0	5.4	5.4	2.5	2.1
South	20.5	19.9	6.8 ^b	5.9	3.0 ^a	2.4
South Atlantic	19.7	20.3	6.1	5.6	2.5	2.3
East South Central	22.5	19.4	8.6 ^a	6.5	4.3	3.1
West South Central	20.8	19.5	7.3	6.1	3.1 ^a	2.3
West	23.1	24.3	7.3	7.0	3.1	2.7
Mountain	24.9	24.4	8.0	7.1	3.3	2.5
Pacific	22.3	24.3	7.1	7.0	3.0	2.8
COUNTY TYPE						
Large Metro	20.6	20.9	6.6	6.0	2.8	2.4
Small Metro	20.7	21.5	6.6	6.4	2.8	2.6
250K – 1 Mil. Pop.	20.8	21.9	6.4	6.3	2.6	2.8
< 250K Pop.	20.4	20.8	7.0	6.5	3.3 ^a	2.4
Nonmetro	19.4	19.5	6.5	5.8	2.8	2.4
Urbanized	21.0	20.4	7.1	5.9	2.6	2.5
Less Urbanized	18.9	19.2	6.8	6.1	3.2	2.4
Completely Rural	15.4	17.8	3.3	3.6	1.7	1.4

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 1.79A (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Numbers in Thousands, 2007 and 2008

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	2,954	2,783	2,037	1,919	830 ^a	711
GEOGRAPHIC DIVISION						
Northeast	406	355	268	259	90	112
New England	118	90	80	63	23	19
Middle Atlantic	287	265	188	196	68	93
Midwest	672	635	469	445	190	175
East North Central	479	450	339	317	131	134
West North Central	194	185	130	128	59	41
South	1,149	1,066	812 ^a	677	351 ^b	253
South Atlantic	531	553	354	340	135	133
East South Central	202	160	140 ^a	100	57	40
West South Central	417	353	318 ^a	237	158 ^b	80
West	726	727	487	538	198	170
Mountain	233	234	173	179	68	72
Pacific	494	493	314	358	130	98
COUNTY TYPE						
Large Metro	1,538 ^a	1,362	1,067	966	409	358
Small Metro	898	888	631	601	278 ^a	215
250K – 1 Mil. Pop.	584	575	399	394	166	131
< 250K Pop.	314	313	232	208	112	84
Nonmetro	518	533	339	351	143	138
Urbanized	219	224	142	159	57	60
Less Urbanized	248	266	160	168	64	68
Completely Rural	50	44	37	24	22 ^a	10

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 1.79B (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Percentages, 2007 and 2008

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	11.7	11.2	8.1	7.7	3.3 ^a	2.9
GEOGRAPHIC DIVISION						
Northeast	9.1	8.1	6.0	5.9	2.0	2.6
New England	10.2	8.0	6.9	5.6	2.0	1.7
Middle Atlantic	8.7	8.2	5.7	6.0	2.1	2.9
Midwest	12.0	11.5	8.4	8.1	3.4	3.2
East North Central	12.1	11.6	8.6	8.2	3.3	3.5
West North Central	11.6	11.3	7.8	7.8	3.5	2.5
South	12.6	11.8	8.9 ^a	7.5	3.8 ^b	2.8
South Atlantic	11.4	12.1	7.6	7.4	2.9	2.9
East South Central	13.6	10.9	9.5 ^a	6.8	3.9	2.7
West South Central	13.9	11.7	10.6 ^a	7.9	5.3 ^b	2.6
West	12.0	12.2	8.1	9.0	3.3	2.8
Mountain	12.9	13.0	9.6	9.9	3.8	4.0
Pacific	11.7	11.9	7.4	8.6	3.1	2.4
COUNTY TYPE						
Large Metro	11.2	10.2	7.8	7.3	3.0	2.7
Small Metro	12.2	12.0	8.5	8.1	3.8 ^a	2.9
250K – 1 Mil. Pop.	11.7	11.7	8.0	8.0	3.3	2.7
< 250K Pop.	13.1	12.5	9.7	8.3	4.7	3.3
Nonmetro	12.4	12.8	8.1	8.4	3.4	3.3
Urbanized	12.8	12.5	8.3	8.9	3.3	3.4
Less Urbanized	12.3	13.2	7.9	8.4	3.2	3.3
Completely Rural	11.8	11.7	8.6	6.4	5.2	2.7

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 1.80A (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Numbers in Thousands, 2007 and 2008

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	9,807	9,694	4,920	4,891	1,947	1,936
GEOGRAPHIC DIVISION						
Northeast	1,586	1,737	807	870	294	356
New England	507	472	266	241	94	92
Middle Atlantic	1,079 ^b	1,265	541	628	200	264
Midwest	2,168	2,155	1,055	1,067	438	416
East North Central	1,546	1,473	791	753	319	312
West North Central	623	682	265	314	119	105
South	3,503	3,341	1,792	1,741	745	731
South Atlantic	1,733 ^a	1,547	912	801	377	324
East South Central	630	558	330	308	149	145
West South Central	1,140	1,235	550	633	220	263
West	2,550	2,461	1,266	1,214	470	432
Mountain	808	861	397	370	158	142
Pacific	1,742	1,600	870	843	312	290
COUNTY TYPE						
Large Metro	4,967	5,008	2,485	2,499	998	960
Small Metro	3,242	3,177	1,597	1,641	593	653
250K – 1 Mil. Pop.	2,166	2,049	1,084	1,027	389	421
< 250K Pop.	1,076	1,128	512	614	204	232
Nonmetro	1,598	1,509	838	751	356	323
Urbanized	735	692	381	319	160	141
Less Urbanized	757	731	394	385	168	161
Completely Rural	106	85	63	47	29	21

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 1.80B (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Percentages, 2007 and 2008

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	30.0	29.4	15.0	14.9	5.9	5.9
GEOGRAPHIC DIVISION						
Northeast	26.9	29.0	13.7	14.5	5.0	5.9
New England	33.2	30.3	17.4	15.5	6.2	5.9
Middle Atlantic	24.7 ^a	28.5	12.4	14.2	4.6	6.0
Midwest	29.8	29.6	14.5	14.7	6.0	5.7
East North Central	30.8	29.2	15.7	15.0	6.3	6.2
West North Central	27.5	30.4	11.7	14.0	5.3	4.7
South	30.0	28.4	15.3	14.8	6.4	6.2
South Atlantic	29.3 ^a	25.9	15.4	13.4	6.4	5.4
East South Central	33.7	29.8	17.6	16.4	8.0	7.7
West South Central	29.1	31.5	14.1	16.1	5.6	6.7
West	32.5	31.1	16.1	15.3	6.0	5.5
Mountain	34.3	36.3	16.8	15.6	6.7	6.0
Pacific	31.7	28.9	15.8	15.2	5.7	5.2
COUNTY TYPE						
Large Metro	28.7	28.1	14.4	14.0	5.8	5.4
Small Metro	31.4	31.5	15.5	16.2	5.7	6.5
250K – 1 Mil. Pop.	32.0	31.3	16.0	15.7	5.7	6.4
< 250K Pop.	30.3	31.7	14.4 ^a	17.3	5.7	6.5
Nonmetro	31.3	30.2	16.4	15.0	7.0	6.5
Urbanized	32.2	30.3	16.7	14.0	7.0	6.2
Less Urbanized	30.9	31.1	16.1	16.4	6.9	6.8
Completely Rural	28.4	24.0	17.0	13.1	7.7	6.0

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 1.81A (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Numbers in Thousands, 2007 and 2008

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	37,781	39,677	9,377 ^a	8,380	4,130	3,563
GEOGRAPHIC DIVISION						
Northeast	6,046	6,713	1,508	1,299	602	569
New England	1,542	1,856	319	418	170	270
Middle Atlantic	4,505	4,857	1,189	881	433	299
Midwest	8,029	8,389	1,871	1,840	840	810
East North Central	5,612	5,943	1,370	1,392	602	612
West North Central	2,416	2,446	501	449	238	198
South	13,766	13,715	3,551 ^a	2,943	1,579	1,218
South Atlantic	7,086	7,652	1,614	1,551	661	647
East South Central	2,485	2,171	801 ^a	559	427	278
West South Central	4,194	3,892	1,137	833	492 ^a	293
West	9,940	10,858	2,447	2,297	1,109	966
Mountain	3,265	3,192	804	698	337	223
Pacific	6,675	7,666	1,642	1,599	771	743
COUNTY TYPE						
Large Metro	20,782	21,486	5,198	4,534	2,246	1,916
Small Metro	11,163	12,178	2,672	2,562	1,230	1,130
250K – 1 Mil. Pop.	7,540	8,319	1,706	1,727	743	847
< 250K Pop.	3,623	3,858	966	835	487 ^a	283
Nonmetro	5,836	6,013	1,507	1,284	655	518
Urbanized	2,568	2,578	664	541	224	236
Less Urbanized	2,717	2,848	791	670	403	256
Completely Rural	551	587	52	73	27	25

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 1.81B (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Percentages, 2007 and 2008

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	19.9	20.7	4.9 ^a	4.4	2.2	1.9
GEOGRAPHIC DIVISION						
Northeast	17.0	18.8	4.2	3.6	1.7	1.6
New England	16.5	19.8	3.4	4.5	1.8	2.9
Middle Atlantic	17.2	18.4	4.5	3.3	1.7	1.1
Midwest	19.2	19.9	4.5	4.4	2.0	1.9
East North Central	19.1	20.2	4.7	4.7	2.1	2.1
West North Central	19.2	19.3	4.0	3.5	1.9	1.6
South	19.9	19.6	5.1 ^a	4.2	2.3 ^a	1.7
South Atlantic	19.1	20.4	4.4	4.1	1.8	1.7
East South Central	21.8	18.8	7.0 ^a	4.9	3.7	2.4
West South Central	20.2	18.4	5.5 ^a	3.9	2.4 ^a	1.4
West	22.9	24.7	5.6	5.2	2.6	2.2
Mountain	24.9	23.8	6.1	5.2	2.6	1.7
Pacific	22.1	25.1	5.4	5.2	2.6	2.4
COUNTY TYPE						
Large Metro	20.4	21.1	5.1	4.4	2.2	1.9
Small Metro	19.8	21.0	4.7	4.4	2.2	1.9
250K – 1 Mil. Pop.	19.9	21.6	4.5	4.5	2.0	2.2
< 250K Pop.	19.5	19.9	5.2	4.3	2.6 ^a	1.5
Nonmetro	18.3	18.7	4.7	4.0	2.1	1.6
Urbanized	20.0	19.7	5.2	4.1	1.7	1.8
Less Urbanized	17.9	18.2	5.2	4.3	2.7	1.6
Completely Rural	14.5	17.8	1.4	2.2	0.7	0.8

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 1.82A (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Numbers in Thousands, 2007 and 2008

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	47,588	49,371	14,297 ^a	13,271	6,077	5,499
GEOGRAPHIC DIVISION						
Northeast	7,633	8,450	2,315	2,169	897	925
New England	2,049	2,328	585	660	264	362
Middle Atlantic	5,583	6,122	1,731	1,509	633	562
Midwest	10,197	10,544	2,926	2,907	1,278	1,227
East North Central	7,158	7,416	2,160	2,145	920	923
West North Central	3,039	3,128	766	762	358	303
South	17,268	17,056	5,343 ^a	4,684	2,325	1,950
South Atlantic	8,819	9,200	2,526	2,351	1,038	971
East South Central	3,115	2,729	1,130 ^a	867	575	423
West South Central	5,334	5,127	1,687	1,466	711	556
West	12,490	13,320	3,713	3,511	1,578	1,398
Mountain	4,073	4,053	1,201	1,068	495	365
Pacific	8,416	9,267	2,512	2,442	1,083	1,033
COUNTY TYPE						
Large Metro	25,749	26,495	7,683	7,033	3,244	2,875
Small Metro	14,405	15,355	4,269	4,203	1,823	1,783
250K – 1 Mil. Pop.	9,706	10,369	2,791	2,754	1,132	1,268
< 250K Pop.	4,698	4,986	1,478	1,449	691	515
Nonmetro	7,434	7,522	2,345	2,035	1,011	840
Urbanized	3,303	3,270	1,045	860	383	377
Less Urbanized	3,474	3,579	1,185	1,055	571	417
Completely Rural	656	672	115	120	56	47

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 1.82B (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Percentages, 2007 and 2008

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	21.4	22.0	6.4 ^a	5.9	2.7	2.4
GEOGRAPHIC DIVISION						
Northeast	18.4	20.3	5.6	5.2	2.2	2.2
New England	18.9	21.3	5.4	6.0	2.4	3.3
Middle Atlantic	18.3	19.9	5.7	4.9	2.1	1.8
Midwest	20.7	21.3	5.9	5.9	2.6	2.5
East North Central	20.8	21.5	6.3	6.2	2.7	2.7
West North Central	20.5	20.9	5.2	5.1	2.4	2.0
South	21.4	20.8	6.6 ^a	5.7	2.9 ^a	2.4
South Atlantic	20.5	21.2	5.9	5.4	2.4	2.2
East South Central	23.5	20.4	8.5 ^a	6.5	4.3	3.2
West South Central	21.7	20.4	6.8	5.8	2.9	2.2
West	24.4	25.7	7.3	6.8	3.1	2.7
Mountain	26.3	25.7	7.8	6.8	3.2	2.3
Pacific	23.6	25.7	7.0	6.8	3.0	2.9
COUNTY TYPE						
Large Metro	21.6	22.1	6.5	5.9	2.7	2.4
Small Metro	21.6	22.6	6.4	6.2	2.7	2.6
250K – 1 Mil. Pop.	21.8	23.0	6.3	6.1	2.5	2.8
< 250K Pop.	21.2	21.7	6.7	6.3	3.1	2.2
Nonmetro	20.1	20.3	6.4	5.5	2.7	2.3
Urbanized	21.9	21.3	6.9	5.6	2.5	2.5
Less Urbanized	19.7	19.8	6.7	5.8	3.2	2.3
Completely Rural	15.7	18.4	2.8	3.3	1.3	1.3

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.