

Table 1.75C (REVISED) Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics:
Standard Errors of Numbers in Thousands, 2007 and 2008

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	640	647	140	130	75	68
GEOGRAPHIC DIVISION						
Northeast	243	294	55	47	37	29
New England	103	141	19	23	11	10
Middle Atlantic	219	258	51	41	35	27
Midwest	235	252	46	58	21	24
East North Central	201	208	39	44	17	17
West North Central	126	137	22	39	11	18
South	384	375	90	72	45	39
South Atlantic	290	320	54	55	21	32
East South Central	138	107	34	26	18	16
West South Central	198	173	61	37	35	15
West	386	399	80	75	44	42
Mountain	180	182	46	35	28	17
Pacific	342	355	65	64	33	39
COUNTY TYPE						
Large Metro	581	613	107	105	58	59
Small Metro	397	425	85	63	44	32
250K – 1 Mil. Pop.	350	374	73	54	37	28
< 250K Pop.	218	229	41	35	23	18
Nonmetro	219	262	39	43	17	16
Urbanized	157	168	25	35	12	11
Less Urbanized	156	174	29	25	12	11
Completely Rural	49	100	6	6	2	5

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 1.75D (REVISED) Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics:
Standard Errors of Percentages, 2007 and 2008**

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	0.26	0.26	0.06	0.05	0.03	0.03
GEOGRAPHIC DIVISION						
Northeast	0.53	0.64	0.12	0.10	0.08	0.06
New England	0.86	1.17	0.16	0.19	0.09	0.08
Middle Atlantic	0.65	0.76	0.15	0.12	0.10	0.08
Midwest	0.43	0.46	0.08	0.11	0.04	0.04
East North Central	0.52	0.54	0.10	0.11	0.04	0.04
West North Central	0.76	0.82	0.14	0.24	0.07	0.11
South	0.43	0.41	0.10	0.08	0.05	0.04
South Atlantic	0.61	0.67	0.11	0.11	0.04	0.07
East South Central	0.93	0.72	0.23	0.18	0.12	0.11
West South Central	0.72	0.62	0.22	0.13	0.13	0.05
West	0.68	0.69	0.14	0.13	0.08	0.07
Mountain	1.04	1.04	0.26	0.20	0.16	0.09
Pacific	0.86	0.88	0.16	0.16	0.08	0.10
COUNTY TYPE						
Large Metro	0.38	0.40	0.08	0.08	0.04	0.04
Small Metro	0.45	0.46	0.11	0.08	0.06	0.04
250K – 1 Mil. Pop.	0.57	0.58	0.14	0.11	0.07	0.06
< 250K Pop.	0.78	0.79	0.17	0.14	0.10	0.07
Nonmetro	0.48	0.53	0.10	0.10	0.04	0.04
Urbanized	0.74	0.76	0.15	0.20	0.07	0.06
Less Urbanized	0.72	0.75	0.15	0.12	0.06	0.05
Completely Rural	0.94	2.12	0.13	0.15	0.04	0.11

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 1.75P (REVISED) Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics:
***P* Values from Tests of Differences of Percentages, 2008 Versus 2007**

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	0.0829		0.5237		0.5985	
GEOGRAPHIC DIVISION						
Northeast	0.1790		0.8765		0.8833	
New England	0.1431		0.6550		0.7613	
Middle Atlantic	0.4289		0.7064		0.9572	
Midwest	0.1216		0.7953		0.3974	
East North Central	0.1501		0.5863		0.7369	
West North Central	0.5369		0.2426		0.3741	
South	0.6284		0.0948		0.4038	
South Atlantic	0.0949		0.8710		0.7280	
East South Central	0.1467		0.3659		0.9778	
West South Central	0.5110		0.0501		0.1009	
West	0.6416		0.5344		0.1627	
Mountain	0.8345		0.9659		0.9108	
Pacific	0.6790		0.4738		0.1074	
COUNTY TYPE						
Large Metro	0.3368		0.8151		0.7114	
Small Metro	0.2856		0.5230		0.8573	
250K – 1 Mil. Pop.	0.9390		0.1542		0.7633	
< 250K Pop.	0.0671		0.2265		0.8963	
Nonmetro	0.2088		0.6950		0.5893	
Urbanized	0.6256		0.9772		0.5630	
Less Urbanized	0.8323		0.4200		0.2966	
Completely Rural	0.0291		0.6370		0.4338	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.