

Table 1.78C (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2007 and 2008

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	698	757	381	352	251	225
GEOGRAPHIC DIVISION						
Northeast	277	330	152	148	84	95
New England	144	136	62	76	37	64
Middle Atlantic	237	301	139	126	76	70
Midwest	285	291	143	150	97	88
East North Central	234	244	124	126	83	71
West North Central	163	154	69	79	48	52
South	439	468	251	211	161	130
South Atlantic	317	366	177	163	111	97
East South Central	168	168	112	82	80	61
West South Central	237	225	131	113	83	67
West	400	417	204	199	153	133
Mountain	179	170	100	84	71	48
Pacific	359	383	177	181	136	124
COUNTY TYPE						
Large Metro	657	695	310	286	209	177
Small Metro	481	541	216	198	138	131
250K – 1 Mil. Pop.	415	482	171	171	105	119
< 250K Pop.	274	282	134	114	92	57
Nonmetro	308	359	154	149	89	76
Urbanized	225	239	109	81	55	47
Less Urbanized	213	244	114	122	73	57
Completely Rural	110	135	23	35	15	18

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 1.78D (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2007 and 2008

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	0.28	0.30	0.15	0.14	0.10	0.09
GEOGRAPHIC DIVISION						
Northeast	0.60	0.72	0.33	0.32	0.18	0.21
New England	1.20	1.13	0.52	0.63	0.31	0.53
Middle Atlantic	0.70	0.88	0.41	0.37	0.22	0.21
Midwest	0.52	0.53	0.26	0.27	0.18	0.16
East North Central	0.61	0.64	0.32	0.33	0.22	0.19
West North Central	0.99	0.93	0.42	0.48	0.29	0.31
South	0.49	0.51	0.28	0.23	0.18	0.14
South Atlantic	0.67	0.76	0.37	0.34	0.23	0.20
East South Central	1.14	1.13	0.76	0.55	0.54	0.41
West South Central	0.86	0.80	0.48	0.40	0.30	0.24
West	0.70	0.72	0.36	0.34	0.27	0.23
Mountain	1.04	0.97	0.58	0.48	0.41	0.27
Pacific	0.90	0.95	0.44	0.45	0.34	0.31
COUNTY TYPE						
Large Metro	0.42	0.42	0.22	0.21	0.15	0.13
Small Metro	0.50	0.58	0.27	0.25	0.17	0.17
250K – 1 Mil. Pop.	0.62	0.73	0.31	0.32	0.19	0.24
< 250K Pop.	0.88	0.90	0.52	0.40	0.37	0.22
Nonmetro	0.60	0.66	0.35	0.32	0.21	0.18
Urbanized	0.97	0.97	0.57	0.40	0.30	0.25
Less Urbanized	0.85	0.96	0.50	0.54	0.35	0.28
Completely Rural	1.73	2.60	0.45	0.79	0.31	0.45

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 1.78P (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2008 Versus 2007

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	0.2483		0.0151		0.0242	
GEOGRAPHIC DIVISION						
Northeast	0.1070		0.4275		0.7223	
New England	0.2269		0.5574		0.1877	
Middle Atlantic	0.2314		0.2351		0.6398	
Midwest	0.4536		0.7952		0.5915	
East North Central	0.4890		0.8054		0.9704	
West North Central	0.7459		0.9247		0.3018	
South	0.4454		0.0083		0.0127	
South Atlantic	0.5195		0.3651		0.5863	
East South Central	0.0586		0.0167		0.0861	
West South Central	0.2947		0.0565		0.0121	
West	0.2303		0.4961		0.2704	
Mountain	0.6969		0.2297		0.1031	
Pacific	0.1341		0.8628		0.6288	
COUNTY TYPE						
Large Metro	0.5530		0.0502		0.1061	
Small Metro	0.2553		0.4828		0.4458	
250K – 1 Mil. Pop.	0.2277		0.7394		0.5471	
< 250K Pop.	0.7655		0.4784		0.0288	
Nonmetro	0.8476		0.1198		0.1103	
Urbanized	0.6569		0.1192		0.8542	
Less Urbanized	0.8439		0.3270		0.0714	
Completely Rural	0.4466		0.7553		0.5897	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 1.79C (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2007 and 2008

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	74	69	63	58	40	33
GEOGRAPHIC DIVISION						
Northeast	23	23	21	20	12	13
New England	13	10	11	8	5	4
Middle Atlantic	20	20	17	18	11	12
Midwest	31	27	25	24	15	14
East North Central	24	23	21	20	12	13
West North Central	18	15	13	13	9	7
South	47	45	40	35	26	20
South Atlantic	35	32	29	24	18	15
East South Central	17	14	15	12	9	8
West South Central	27	28	24	22	18	11
West	42	38	34	35	22	19
Mountain	18	17	16	16	11	11
Pacific	37	33	30	31	20	15
COUNTY TYPE						
Large Metro	64	58	53	47	32	27
Small Metro	44	42	38	35	23	18
250K – 1 Mil. Pop.	36	36	30	29	17	14
< 250K Pop.	26	24	22	20	15	12
Nonmetro	29	34	23	25	14	14
Urbanized	20	22	15	18	9	10
Less Urbanized	21	24	16	17	9	10
Completely Rural	8	11	7	6	5	3

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 1.79D (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Percentages, 2007 and 2008

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	0.29	0.28	0.25	0.23	0.16	0.13
GEOGRAPHIC DIVISION						
Northeast	0.52	0.52	0.47	0.45	0.26	0.29
New England	1.09	0.87	0.99	0.71	0.43	0.38
Middle Atlantic	0.59	0.63	0.53	0.56	0.32	0.37
Midwest	0.55	0.49	0.44	0.43	0.28	0.26
East North Central	0.62	0.59	0.53	0.52	0.31	0.33
West North Central	1.08	0.91	0.75	0.77	0.57	0.40
South	0.51	0.50	0.44	0.39	0.29	0.22
South Atlantic	0.75	0.69	0.63	0.53	0.38	0.33
East South Central	1.17	0.96	1.00	0.84	0.62	0.53
West South Central	0.90	0.94	0.81	0.75	0.60	0.37
West	0.69	0.63	0.56	0.59	0.37	0.32
Mountain	1.00	0.95	0.90	0.90	0.58	0.62
Pacific	0.89	0.80	0.70	0.75	0.46	0.36
COUNTY TYPE						
Large Metro	0.42	0.38	0.35	0.33	0.22	0.19
Small Metro	0.51	0.49	0.44	0.43	0.29	0.23
250K – 1 Mil. Pop.	0.61	0.62	0.52	0.53	0.32	0.27
< 250K Pop.	0.93	0.85	0.81	0.72	0.61	0.46
Nonmetro	0.62	0.65	0.51	0.54	0.33	0.33
Urbanized	0.90	0.97	0.75	0.86	0.50	0.53
Less Urbanized	0.88	0.99	0.71	0.76	0.45	0.47
Completely Rural	1.95	2.08	1.61	1.44	1.25	0.86

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 1.79P (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2008 Versus 2007

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	0.2026		0.2862		0.0324	
GEOGRAPHIC DIVISION						
Northeast	0.1740		0.8637		0.1611	
New England	0.0997		0.2661		0.6932	
Middle Atlantic	0.5326		0.6446		0.0914	
Midwest	0.5272		0.6588		0.5872	
East North Central	0.5337		0.5917		0.7513	
West North Central	0.8208		0.9818		0.1357	
South	0.2742		0.0154		0.0029	
South Atlantic	0.5455		0.8280		0.9965	
East South Central	0.0623		0.0233		0.1633	
West South Central	0.1096		0.0146		0.0002	
West	0.8513		0.2221		0.3675	
Mountain	0.9463		0.7795		0.7911	
Pacific	0.8639		0.2096		0.2045	
COUNTY TYPE						
Large Metro	0.0835		0.2723		0.3098	
Small Metro	0.7844		0.4628		0.0216	
250K – 1 Mil. Pop.	0.9901		0.9674		0.1288	
< 250K Pop.	0.6078		0.1810		0.0760	
Nonmetro	0.6926		0.6970		0.7560	
Urbanized	0.8419		0.5764		0.9246	
Less Urbanized	0.4778		0.6659		0.7727	
Completely Rural	0.9807		0.2640		0.0832	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 1.80C (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2007 and 2008

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	145	142	110	112	69	71
GEOGRAPHIC DIVISION						
Northeast	60	57	43	42	24	28
New England	26	28	23	20	14	11
Middle Atlantic	53	50	35	38	20	25
Midwest	59	52	42	40	26	26
East North Central	45	41	34	32	21	22
West North Central	36	31	23	24	16	14
South	89	88	68	69	43	48
South Atlantic	64	59	48	48	31	35
East South Central	36	31	25	26	16	18
West South Central	51	53	41	40	27	26
West	85	80	66	63	40	36
Mountain	37	39	29	29	19	17
Pacific	77	68	59	56	36	32
COUNTY TYPE						
Large Metro	158	147	105	100	60	56
Small Metro	115	106	72	72	40	41
250K – 1 Mil. Pop.	92	88	59	58	31	33
< 250K Pop.	72	66	43	45	26	26
Nonmetro	77	82	48	49	27	31
Urbanized	52	61	32	31	18	19
Less Urbanized	55	54	36	37	20	23
Completely Rural	22	14	15	9	8	6

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 1.80D (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Percentages, 2007 and 2008

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	0.44	0.43	0.34	0.34	0.21	0.22
GEOGRAPHIC DIVISION						
Northeast	1.02	0.96	0.72	0.71	0.41	0.46
New England	1.73	1.79	1.51	1.27	0.93	0.70
Middle Atlantic	1.22	1.13	0.80	0.85	0.45	0.57
Midwest	0.81	0.72	0.57	0.56	0.36	0.36
East North Central	0.89	0.82	0.68	0.64	0.42	0.44
West North Central	1.60	1.39	1.02	1.07	0.72	0.61
South	0.76	0.75	0.58	0.59	0.37	0.41
South Atlantic	1.08	1.00	0.81	0.81	0.52	0.59
East South Central	1.94	1.64	1.35	1.40	0.84	0.96
West South Central	1.31	1.36	1.04	1.01	0.68	0.66
West	1.09	1.01	0.84	0.80	0.51	0.46
Mountain	1.56	1.66	1.25	1.22	0.82	0.70
Pacific	1.40	1.23	1.08	1.01	0.65	0.58
COUNTY TYPE						
Large Metro	0.63	0.61	0.50	0.48	0.31	0.30
Small Metro	0.81	0.73	0.57	0.60	0.36	0.39
250K – 1 Mil. Pop.	0.95	0.93	0.69	0.76	0.42	0.48
< 250K Pop.	1.49	1.17	0.99	0.95	0.68	0.69
Nonmetro	1.08	0.91	0.77	0.73	0.50	0.53
Urbanized	1.72	1.34	1.20	0.97	0.76	0.71
Less Urbanized	1.51	1.37	1.12	1.15	0.74	0.85
Completely Rural	3.92	2.57	3.21	1.90	1.91	1.49

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 1.80P (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2008 Versus 2007

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	0.3802		0.7013		0.8090	
GEOGRAPHIC DIVISION						
Northeast	0.1193		0.4129		0.1321	
New England	0.2582		0.3718		0.8215	
Middle Atlantic	0.0163		0.1247		0.0633	
Midwest	0.8991		0.8170		0.5767	
East North Central	0.1953		0.3924		0.8085	
West North Central	0.1721		0.1366		0.5029	
South	0.1223		0.5077		0.7682	
South Atlantic	0.0162		0.0684		0.2348	
East South Central	0.0977		0.5045		0.8458	
West South Central	0.2044		0.1541		0.2328	
West	0.3435		0.4876		0.4502	
Mountain	0.3512		0.4470		0.5178	
Pacific	0.1257		0.6818		0.6131	
COUNTY TYPE						
Large Metro	0.4617		0.6036		0.3751	
Small Metro	0.9679		0.3504		0.1731	
250K – 1 Mil. Pop.	0.5828		0.7500		0.2902	
< 250K Pop.	0.4779		0.0459		0.3959	
Nonmetro	0.4362		0.1998		0.4717	
Urbanized	0.4078		0.0928		0.4171	
Less Urbanized	0.9541		0.8698		0.9765	
Completely Rural	0.3781		0.3086		0.4318	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 1.81C (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2007 and 2008

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	662	730	346	310	237	204
GEOGRAPHIC DIVISION						
Northeast	267	313	141	131	81	87
New England	134	124	50	67	35	61
Middle Atlantic	232	287	132	112	74	62
Midwest	272	280	131	134	93	82
East North Central	224	236	115	114	80	67
West North Central	159	146	62	69	47	49
South	420	455	229	192	150	117
South Atlantic	304	357	157	153	99	88
East South Central	160	163	109	72	78	57
West South Central	222	214	118	94	77	56
West	381	408	182	175	142	122
Mountain	174	169	86	77	65	42
Pacific	338	372	158	160	126	115
COUNTY TYPE						
Large Metro	603	651	281	248	193	159
Small Metro	435	504	183	178	126	121
250K – 1 Mil. Pop.	382	443	142	154	96	111
< 250K Pop.	237	262	118	96	82	49
Nonmetro	276	323	133	130	80	66
Urbanized	192	211	92	67	47	40
Less Urbanized	190	216	97	109	67	49
Completely Rural	104	125	14	28	10	16

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 1.81D (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2007 and 2008

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	0.35	0.38	0.18	0.16	0.12	0.11
GEOGRAPHIC DIVISION						
Northeast	0.75	0.87	0.40	0.37	0.23	0.24
New England	1.43	1.32	0.53	0.71	0.38	0.65
Middle Atlantic	0.89	1.09	0.51	0.42	0.28	0.23
Midwest	0.65	0.66	0.31	0.32	0.22	0.19
East North Central	0.76	0.80	0.39	0.39	0.27	0.23
West North Central	1.27	1.15	0.49	0.54	0.38	0.38
South	0.61	0.65	0.33	0.27	0.22	0.17
South Atlantic	0.82	0.95	0.42	0.41	0.27	0.23
East South Central	1.41	1.41	0.96	0.63	0.68	0.49
West South Central	1.07	1.01	0.57	0.45	0.37	0.26
West	0.88	0.93	0.42	0.40	0.33	0.28
Mountain	1.33	1.26	0.66	0.57	0.49	0.31
Pacific	1.12	1.22	0.52	0.52	0.42	0.38
COUNTY TYPE						
Large Metro	0.52	0.54	0.27	0.24	0.19	0.15
Small Metro	0.62	0.72	0.31	0.30	0.21	0.21
250K – 1 Mil. Pop.	0.78	0.90	0.35	0.39	0.24	0.29
< 250K Pop.	1.05	1.14	0.62	0.46	0.43	0.25
Nonmetro	0.74	0.82	0.40	0.39	0.24	0.20
Urbanized	1.18	1.21	0.67	0.48	0.35	0.29
Less Urbanized	1.05	1.17	0.59	0.65	0.42	0.31
Completely Rural	2.19	3.15	0.38	0.83	0.28	0.48

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 1.81P (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2008 Versus 2007

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	0.1455		0.0190		0.0509	
GEOGRAPHIC DIVISION						
Northeast	0.1439		0.2656		0.7581	
New England	0.0974		0.2347		0.1481	
Middle Atlantic	0.4100		0.0749		0.1647	
Midwest	0.3952		0.8147		0.7790	
East North Central	0.3158		0.9252		0.9447	
West North Central	0.9804		0.5122		0.5358	
South	0.6935		0.0274		0.0414	
South Atlantic	0.3169		0.7012		0.8667	
East South Central	0.1463		0.0409		0.1155	
West South Central	0.2342		0.0393		0.0298	
West	0.1715		0.4671		0.4057	
Mountain	0.5457		0.2807		0.1131	
Pacific	0.0724		0.7910		0.8358	
COUNTY TYPE						
Large Metro	0.4108		0.0612		0.1755	
Small Metro	0.2070		0.4402		0.4410	
250K – 1 Mil. Pop.	0.1661		0.9440		0.5257	
< 250K Pop.	0.7975		0.2570		0.0237	
Nonmetro	0.7187		0.1926		0.1641	
Urbanized	0.8297		0.2282		0.9023	
Less Urbanized	0.8563		0.2907		0.0529	
Completely Rural	0.3978		0.3468		0.9362	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 1.82C (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2007 and 2008

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	692	753	377	345	248	221
GEOGRAPHIC DIVISION						
Northeast	276	331	152	146	84	94
New England	142	136	61	76	37	63
Middle Atlantic	238	302	139	125	75	69
Midwest	284	288	141	146	96	87
East North Central	233	241	121	123	83	72
West North Central	165	151	69	76	48	50
South	434	465	248	208	159	128
South Atlantic	314	364	174	162	108	96
East South Central	167	168	111	82	80	62
West South Central	231	224	128	108	81	65
West	398	414	201	193	150	130
Mountain	177	170	96	84	69	48
Pacific	358	380	175	175	133	122
COUNTY TYPE						
Large Metro	645	683	307	278	206	173
Small Metro	470	532	207	191	135	131
250K – 1 Mil. Pop.	405	472	164	166	102	118
< 250K Pop.	267	278	128	109	90	56
Nonmetro	302	348	150	144	87	75
Urbanized	218	232	106	78	54	46
Less Urbanized	209	235	110	119	72	57
Completely Rural	107	131	21	34	13	17

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 1.82D (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2007 and 2008

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	0.31	0.33	0.17	0.15	0.11	0.10
GEOGRAPHIC DIVISION						
Northeast	0.67	0.79	0.37	0.35	0.20	0.23
New England	1.30	1.25	0.56	0.69	0.34	0.58
Middle Atlantic	0.78	0.98	0.46	0.40	0.25	0.22
Midwest	0.58	0.58	0.29	0.30	0.20	0.18
East North Central	0.68	0.70	0.35	0.36	0.24	0.21
West North Central	1.11	1.01	0.47	0.51	0.33	0.33
South	0.54	0.57	0.31	0.25	0.20	0.16
South Atlantic	0.73	0.84	0.41	0.37	0.25	0.22
East South Central	1.26	1.25	0.84	0.61	0.61	0.46
West South Central	0.94	0.89	0.52	0.43	0.33	0.26
West	0.78	0.80	0.39	0.37	0.29	0.25
Mountain	1.14	1.08	0.62	0.53	0.45	0.31
Pacific	1.00	1.05	0.49	0.49	0.37	0.34
COUNTY TYPE						
Large Metro	0.46	0.47	0.25	0.23	0.17	0.14
Small Metro	0.55	0.64	0.29	0.27	0.19	0.19
250K – 1 Mil. Pop.	0.68	0.80	0.33	0.35	0.21	0.26
< 250K Pop.	0.97	0.99	0.56	0.43	0.40	0.24
Nonmetro	0.67	0.73	0.38	0.36	0.23	0.19
Urbanized	1.06	1.06	0.63	0.44	0.34	0.27
Less Urbanized	0.94	1.05	0.55	0.60	0.38	0.30
Completely Rural	1.94	2.90	0.47	0.86	0.31	0.47

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 1.82P (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2008 Versus 2007

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	0.2168		0.0221		0.0520	
GEOGRAPHIC DIVISION						
Northeast	0.0925		0.4381		0.8659	
New England	0.1867		0.4513		0.1767	
Middle Atlantic	0.2229		0.2102		0.4696	
Midwest	0.4284		0.8600		0.6533	
East North Central	0.4602		0.8796		0.9940	
West North Central	0.7415		0.9282		0.4214	
South	0.4962		0.0229		0.0414	
South Atlantic	0.5725		0.3849		0.5830	
East South Central	0.0795		0.0341		0.1216	
West South Central	0.3757		0.1363		0.0770	
West	0.2452		0.3624		0.3215	
Mountain	0.6756		0.2000		0.0891	
Pacific	0.1406		0.7014		0.7442	
COUNTY TYPE						
Large Metro	0.4822		0.0728		0.1387	
Small Metro	0.2491		0.5553		0.6775	
250K – 1 Mil. Pop.	0.2334		0.7344		0.4085	
< 250K Pop.	0.7207		0.6043		0.0557	
Nonmetro	0.8790		0.1000		0.1135	
Urbanized	0.6763		0.0982		0.8377	
Less Urbanized	0.9132		0.2891		0.0616	
Completely Rural	0.4559		0.5830		0.8989	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.