

Table 1.83C (REVISED) Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2007 and 2008

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	563	597	316	294	214	196
GEOGRAPHIC DIVISION						
Northeast	218	271	115	128	73	84
New England	106	127	51	71	32	62
Middle Atlantic	190	239	102	106	65	56
Midwest	233	227	127	132	86	81
East North Central	188	195	112	110	76	63
West North Central	136	110	58	70	39	49
South	331	364	203	171	131	108
South Atlantic	242	286	141	130	90	77
East South Central	140	122	102	75	75	59
West South Central	171	184	103	92	57	50
West	319	322	176	165	128	116
Mountain	138	143	92	71	68	43
Pacific	290	289	150	150	109	108
COUNTY TYPE						
Large Metro	482	520	250	250	169	159
Small Metro	344	378	181	164	119	104
250K – 1 Mil. Pop.	286	333	132	130	82	89
< 250K Pop.	205	209	124	107	86	56
Nonmetro	229	267	124	105	77	64
Urbanized	164	190	85	67	46	39
Less Urbanized	163	174	94	77	65	47
Completely Rural	74	92	20	31	12	17

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 1.83D (REVISED) Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2007 and 2008

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	0.23	0.24	0.13	0.12	0.09	0.08
GEOGRAPHIC DIVISION						
Northeast	0.48	0.59	0.25	0.28	0.16	0.18
New England	0.88	1.06	0.42	0.59	0.27	0.51
Middle Atlantic	0.56	0.70	0.30	0.31	0.19	0.17
Midwest	0.42	0.41	0.23	0.24	0.16	0.15
East North Central	0.49	0.51	0.29	0.29	0.20	0.16
West North Central	0.82	0.67	0.35	0.42	0.24	0.29
South	0.37	0.40	0.23	0.19	0.15	0.12
South Atlantic	0.51	0.59	0.30	0.27	0.19	0.16
East South Central	0.95	0.82	0.69	0.50	0.51	0.40
West South Central	0.62	0.66	0.37	0.33	0.21	0.18
West	0.56	0.56	0.31	0.29	0.22	0.20
Mountain	0.80	0.81	0.53	0.40	0.39	0.25
Pacific	0.73	0.72	0.37	0.37	0.27	0.27
COUNTY TYPE						
Large Metro	0.33	0.34	0.18	0.18	0.13	0.12
Small Metro	0.39	0.43	0.23	0.21	0.15	0.14
250K – 1 Mil. Pop.	0.48	0.53	0.25	0.25	0.16	0.18
< 250K Pop.	0.72	0.72	0.48	0.38	0.34	0.21
Nonmetro	0.49	0.53	0.29	0.23	0.18	0.15
Urbanized	0.77	0.85	0.46	0.35	0.26	0.21
Less Urbanized	0.68	0.72	0.43	0.35	0.31	0.23
Completely Rural	1.47	1.95	0.41	0.71	0.26	0.42

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 1.83P (REVISED) Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2008 Versus 2007

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	0.0520		0.1091		0.0828	
GEOGRAPHIC DIVISION						
Northeast	0.0715		0.9156		0.6879	
New England	0.1058		0.4334		0.1418	
Middle Atlantic	0.2425		0.7602		0.5204	
Midwest	0.9974		0.9941		0.7420	
East North Central	0.5083		0.7907		0.8587	
West North Central	0.3202		0.6498		0.7374	
South	0.6540		0.0139		0.0129	
South Atlantic	0.3016		0.3140		0.2720	
East South Central	0.1101		0.0733		0.1692	
West South Central	0.6911		0.0905		0.0345	
West	0.0984		0.7922		0.6288	
Mountain	0.6418		0.2729		0.1140	
Pacific	0.1097		0.7744		0.7871	
COUNTY TYPE						
Large Metro	0.1939		0.4137		0.3680	
Small Metro	0.1990		0.6666		0.4888	
250K – 1 Mil. Pop.	0.2163		0.8863		0.7875	
< 250K Pop.	0.6338		0.6087		0.1504	
Nonmetro	0.5008		0.0332		0.0553	
Urbanized	0.7244		0.0656		0.4285	
Less Urbanized	0.6210		0.1300		0.0737	
Completely Rural	0.9210		0.7795		0.6924	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 1.84C (REVISED) Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2007 and 2008

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	69	66	59	54	36	29
GEOGRAPHIC DIVISION						
Northeast	22	20	20	18	11	12
New England	11	8	10	6	3	3
Middle Atlantic	19	19	17	17	10	12
Midwest	28	26	22	22	14	12
East North Central	23	22	19	18	11	11
West North Central	15	14	11	12	8	6
South	43	41	37	31	24	17
South Atlantic	31	29	27	23	17	12
East South Central	17	13	14	11	9	7
West South Central	25	25	22	17	16	10
West	39	36	31	32	19	16
Mountain	16	17	15	16	9	8
Pacific	35	31	27	28	16	14
COUNTY TYPE						
Large Metro	57	53	47	43	28	22
Small Metro	40	38	33	31	21	16
250K – 1 Mil. Pop.	33	32	26	26	15	12
< 250K Pop.	25	22	21	18	15	10
Nonmetro	26	32	21	23	13	12
Urbanized	17	21	13	16	8	8
Less Urbanized	18	22	14	15	8	9
Completely Rural	8	11	7	5	5	3

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 1.84D (REVISED) Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Percentages, 2007 and 2008

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	0.27	0.26	0.23	0.22	0.14	0.12
GEOGRAPHIC DIVISION						
Northeast	0.49	0.47	0.44	0.41	0.24	0.28
New England	0.99	0.71	0.86	0.55	0.26	0.31
Middle Atlantic	0.56	0.58	0.51	0.51	0.31	0.36
Midwest	0.49	0.46	0.39	0.39	0.25	0.22
East North Central	0.58	0.56	0.48	0.47	0.29	0.28
West North Central	0.89	0.84	0.68	0.70	0.51	0.37
South	0.48	0.46	0.41	0.34	0.27	0.19
South Atlantic	0.67	0.64	0.59	0.50	0.36	0.27
East South Central	1.12	0.91	0.94	0.75	0.60	0.49
West South Central	0.82	0.82	0.72	0.58	0.52	0.34
West	0.65	0.60	0.51	0.54	0.31	0.28
Mountain	0.91	0.92	0.83	0.88	0.51	0.43
Pacific	0.83	0.75	0.64	0.67	0.38	0.34
COUNTY TYPE						
Large Metro	0.38	0.36	0.32	0.30	0.20	0.16
Small Metro	0.48	0.45	0.41	0.39	0.27	0.21
250K – 1 Mil. Pop.	0.57	0.56	0.47	0.47	0.28	0.25
< 250K Pop.	0.91	0.81	0.79	0.69	0.60	0.40
Nonmetro	0.56	0.63	0.46	0.50	0.29	0.29
Urbanized	0.84	0.93	0.67	0.80	0.44	0.44
Less Urbanized	0.82	0.96	0.66	0.70	0.38	0.45
Completely Rural	1.87	2.09	1.52	1.35	1.23	0.73

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 1.84P (REVISED) Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2008 Versus 2007

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	0.4612		0.5302		0.0347	
GEOGRAPHIC DIVISION						
Northeast	0.2175		0.6182		0.1984	
New England	0.1600		0.4222		0.7223	
Middle Atlantic	0.5015		0.8808		0.2168	
Midwest	0.7992		0.9610		0.6090	
East North Central	0.9131		0.8348		0.9854	
West North Central	0.7625		0.6738		0.3692	
South	0.3697		0.0110		0.0006	
South Atlantic	0.4834		0.6888		0.1736	
East South Central	0.1849		0.1017		0.2190	
West South Central	0.0899		0.0028		0.0018	
West	0.5778		0.0469		0.7185	
Mountain	0.5263		0.2833		0.9841	
Pacific	0.7654		0.0933		0.6608	
COUNTY TYPE						
Large Metro	0.1233		0.2704		0.2471	
Small Metro	0.9706		0.6844		0.0524	
250K – 1 Mil. Pop.	0.7545		0.8640		0.2848	
< 250K Pop.	0.6061		0.3659		0.0832	
Nonmetro	0.2527		0.2654		0.6230	
Urbanized	0.9241		0.2731		0.5155	
Less Urbanized	0.1160		0.2588		0.3272	
Completely Rural	0.9685		0.1560		0.0340	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 1.85C (REVISED) Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2007 and 2008

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	136	138	97	101	60	61
GEOGRAPHIC DIVISION						
Northeast	54	53	37	38	21	23
New England	24	25	21	18	11	9
Middle Atlantic	47	47	31	33	18	22
Midwest	53	52	37	39	22	24
East North Central	42	41	31	31	19	21
West North Central	31	31	21	23	13	12
South	81	83	60	62	38	42
South Atlantic	60	55	43	44	28	31
East South Central	32	27	23	22	14	15
West South Central	46	51	34	36	20	21
West	81	77	58	57	38	31
Mountain	34	38	26	27	17	16
Pacific	73	67	52	50	34	27
COUNTY TYPE						
Large Metro	140	131	91	88	52	48
Small Metro	97	97	60	65	33	35
250K – 1 Mil. Pop.	81	80	52	52	28	26
< 250K Pop.	57	61	31	41	19	25
Nonmetro	70	74	43	45	25	26
Urbanized	46	57	29	31	16	15
Less Urbanized	51	45	30	31	17	21
Completely Rural	21	13	14	8	8	6

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 1.85D (REVISED) Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Percentages, 2007 and 2008

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	0.42	0.42	0.30	0.31	0.18	0.19
GEOGRAPHIC DIVISION						
Northeast	0.91	0.89	0.63	0.64	0.36	0.39
New England	1.59	1.61	1.35	1.19	0.70	0.59
Middle Atlantic	1.08	1.05	0.70	0.75	0.41	0.49
Midwest	0.73	0.72	0.51	0.54	0.31	0.33
East North Central	0.83	0.82	0.61	0.62	0.37	0.41
West North Central	1.36	1.39	0.91	1.03	0.56	0.52
South	0.70	0.70	0.51	0.52	0.32	0.35
South Atlantic	1.01	0.93	0.73	0.73	0.48	0.52
East South Central	1.69	1.45	1.25	1.19	0.75	0.82
West South Central	1.17	1.30	0.87	0.91	0.52	0.54
West	1.03	0.97	0.74	0.72	0.48	0.39
Mountain	1.43	1.59	1.09	1.14	0.72	0.66
Pacific	1.33	1.20	0.95	0.90	0.62	0.48
COUNTY TYPE						
Large Metro	0.59	0.59	0.44	0.44	0.27	0.26
Small Metro	0.71	0.69	0.50	0.54	0.31	0.33
250K – 1 Mil. Pop.	0.86	0.87	0.64	0.69	0.39	0.39
< 250K Pop.	1.23	1.14	0.77	0.91	0.52	0.64
Nonmetro	1.04	0.84	0.71	0.66	0.45	0.48
Urbanized	1.53	1.31	1.14	0.94	0.68	0.62
Less Urbanized	1.54	1.19	0.97	1.02	0.64	0.79
Completely Rural	3.93	2.40	3.11	1.74	1.87	1.50

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 1.85P (REVISED) Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2008 Versus 2007

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	0.6071		0.7176		0.6465	
GEOGRAPHIC DIVISION						
Northeast	0.2220		0.4820		0.1757	
New England	0.4152		0.4370		0.8758	
Middle Atlantic	0.0677		0.1896		0.1475	
Midwest	0.7848		0.5836		0.9776	
East North Central	0.3669		0.5683		0.8073	
West North Central	0.1153		0.0886		0.7065	
South	0.4117		0.6848		0.4874	
South Atlantic	0.1661		0.1847		0.1043	
East South Central	0.0890		0.7560		0.8255	
West South Central	0.2247		0.2749		0.2331	
West	0.2815		0.3060		0.3677	
Mountain	0.5345		0.9773		0.9105	
Pacific	0.1357		0.2601		0.3367	
COUNTY TYPE						
Large Metro	0.5673		0.4744		0.3775	
Small Metro	0.6832		0.2784		0.2520	
250K – 1 Mil. Pop.	0.5286		0.5811		0.6752	
< 250K Pop.	0.1374		0.0091		0.1911	
Nonmetro	0.4904		0.3416		0.3830	
Urbanized	0.7268		0.1822		0.2085	
Less Urbanized	0.7582		0.7961		0.9837	
Completely Rural	0.3586		0.4623		0.7867	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 1.86C (REVISED) Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2007 and 2008

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	526	567	290	259	200	177
GEOGRAPHIC DIVISION						
Northeast	209	256	102	113	70	78
New England	100	119	40	65	31	60
Middle Atlantic	182	226	93	92	63	49
Midwest	221	214	115	117	82	76
East North Central	178	186	102	99	74	59
West North Central	132	102	53	60	36	46
South	309	353	188	153	121	95
South Atlantic	222	273	128	115	81	67
East South Central	134	118	99	69	73	56
West South Central	159	175	92	80	50	41
West	296	314	158	151	118	107
Mountain	128	141	83	65	63	36
Pacific	269	282	134	138	100	101
COUNTY TYPE						
Large Metro	437	483	225	220	155	145
Small Metro	305	345	157	146	108	95
250K – 1 Mil. Pop.	251	299	110	118	75	83
< 250K Pop.	180	191	112	92	77	48
Nonmetro	204	236	106	87	70	55
Urbanized	140	164	70	53	39	33
Less Urbanized	142	151	82	64	61	40
Completely Rural	70	88	13	28	8	15

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 1.86D (REVISED) Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2007 and 2008

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	0.28	0.30	0.15	0.13	0.11	0.09
GEOGRAPHIC DIVISION						
Northeast	0.59	0.72	0.29	0.32	0.20	0.22
New England	1.07	1.27	0.43	0.69	0.34	0.64
Middle Atlantic	0.70	0.86	0.36	0.35	0.24	0.18
Midwest	0.53	0.51	0.27	0.28	0.20	0.18
East North Central	0.61	0.63	0.35	0.34	0.25	0.20
West North Central	1.05	0.80	0.42	0.48	0.29	0.36
South	0.45	0.50	0.27	0.22	0.18	0.14
South Atlantic	0.60	0.73	0.35	0.31	0.22	0.18
East South Central	1.18	1.03	0.87	0.60	0.64	0.49
West South Central	0.77	0.83	0.45	0.38	0.24	0.19
West	0.68	0.72	0.36	0.34	0.27	0.24
Mountain	0.97	1.05	0.63	0.48	0.48	0.27
Pacific	0.89	0.92	0.44	0.45	0.33	0.33
COUNTY TYPE						
Large Metro	0.40	0.43	0.22	0.21	0.15	0.14
Small Metro	0.48	0.53	0.27	0.25	0.19	0.16
250K – 1 Mil. Pop.	0.58	0.65	0.28	0.30	0.19	0.22
< 250K Pop.	0.88	0.90	0.59	0.45	0.41	0.24
Nonmetro	0.59	0.65	0.33	0.27	0.21	0.17
Urbanized	0.95	1.05	0.53	0.40	0.29	0.25
Less Urbanized	0.82	0.87	0.50	0.39	0.39	0.25
Completely Rural	1.80	2.40	0.34	0.81	0.23	0.46

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 1.86P (REVISED) Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2008 Versus 2007

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	0.0240		0.1396		0.1823	
GEOGRAPHIC DIVISION						
Northeast	0.0854		0.9530		0.8520	
New England	0.0486		0.2103		0.1503	
Middle Atlantic	0.3812		0.4506		0.1558	
Midwest	0.9744		0.8710		0.8089	
East North Central	0.3736		0.9574		0.8076	
West North Central	0.1576		0.8210		0.9486	
South	0.4336		0.0441		0.0715	
South Atlantic	0.1959		0.5809		0.7450	
East South Central	0.2633		0.1153		0.2230	
West South Central	0.7263		0.1054		0.0460	
West	0.0544		0.7810		0.8510	
Mountain	0.7966		0.1752		0.0882	
Pacific	0.0440		0.6526		0.5111	
COUNTY TYPE						
Large Metro	0.1154		0.5977		0.5749	
Small Metro	0.1658		0.5626		0.5109	
250K – 1 Mil. Pop.	0.1239		0.8189		0.6645	
< 250K Pop.	0.8048		0.2583		0.1117	
Nonmetro	0.4203		0.0296		0.1046	
Urbanized	0.6244		0.0891		0.8476	
Less Urbanized	0.6085		0.0647		0.0480	
Completely Rural	0.8318		0.3971		0.8165	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 1.87C (REVISED) Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2007 and 2008

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	558	593	312	288	211	192
GEOGRAPHIC DIVISION						
Northeast	217	271	114	127	72	83
New England	105	127	49	71	32	61
Middle Atlantic	189	239	102	104	64	55
Midwest	232	224	125	128	86	80
East North Central	186	193	109	108	76	63
West North Central	137	108	58	66	38	47
South	327	362	201	169	129	107
South Atlantic	240	283	140	128	89	76
East South Central	139	122	101	75	75	59
West South Central	168	184	100	89	54	48
West	317	319	173	162	126	113
Mountain	135	143	88	70	66	43
Pacific	288	286	148	147	107	105
COUNTY TYPE						
Large Metro	474	510	246	243	167	156
Small Metro	334	371	173	160	114	104
250K – 1 Mil. Pop.	276	327	126	128	79	88
< 250K Pop.	199	203	119	103	82	55
Nonmetro	225	257	122	100	77	63
Urbanized	158	183	83	63	45	38
Less Urbanized	159	167	92	75	66	47
Completely Rural	72	90	19	30	12	16

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 1.87D (REVISED) Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2007 and 2008

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	0.25	0.26	0.14	0.13	0.09	0.09
GEOGRAPHIC DIVISION						
Northeast	0.52	0.65	0.27	0.30	0.17	0.20
New England	0.97	1.16	0.45	0.65	0.30	0.56
Middle Atlantic	0.62	0.77	0.33	0.34	0.21	0.18
Midwest	0.47	0.45	0.25	0.26	0.17	0.16
East North Central	0.54	0.56	0.32	0.31	0.22	0.18
West North Central	0.92	0.72	0.39	0.44	0.26	0.32
South	0.40	0.44	0.25	0.21	0.16	0.13
South Atlantic	0.56	0.65	0.33	0.30	0.21	0.18
East South Central	1.05	0.91	0.76	0.56	0.57	0.44
West South Central	0.68	0.73	0.41	0.35	0.22	0.19
West	0.62	0.62	0.34	0.31	0.25	0.22
Mountain	0.87	0.90	0.57	0.45	0.43	0.27
Pacific	0.81	0.79	0.42	0.41	0.30	0.29
COUNTY TYPE						
Large Metro	0.36	0.38	0.20	0.20	0.14	0.13
Small Metro	0.43	0.47	0.25	0.23	0.17	0.15
250K – 1 Mil. Pop.	0.52	0.59	0.27	0.28	0.17	0.20
< 250K Pop.	0.79	0.78	0.52	0.41	0.36	0.24
Nonmetro	0.54	0.58	0.31	0.25	0.20	0.16
Urbanized	0.85	0.93	0.51	0.38	0.28	0.23
Less Urbanized	0.75	0.79	0.47	0.38	0.35	0.25
Completely Rural	1.62	2.16	0.44	0.77	0.27	0.45

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 1.87P (REVISED) Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2008 Versus 2007

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	0.0435		0.1337		0.1606	
GEOGRAPHIC DIVISION						
Northeast	0.0590		0.8499		0.8262	
New England	0.0886		0.3653		0.1475	
Middle Atlantic	0.2240		0.7805		0.3882	
Midwest	0.9915		1.0000		0.8101	
East North Central	0.5046		0.8187		0.8653	
West North Central	0.3313		0.6909		0.8550	
South	0.5812		0.0436		0.0557	
South Atlantic	0.3451		0.3527		0.3823	
East South Central	0.1434		0.1107		0.2174	
West South Central	0.5377		0.2536		0.1663	
West	0.1119		0.5324		0.6597	
Mountain	0.7025		0.1831		0.1080	
Pacific	0.1153		0.9970		0.7355	
COUNTY TYPE						
Large Metro	0.1477		0.5364		0.4651	
Small Metro	0.1960		0.7185		0.7086	
250K – 1 Mil. Pop.	0.2292		0.8708		0.6489	
< 250K Pop.	0.5814		0.7050		0.2427	
Nonmetro	0.5868		0.0175		0.0652	
Urbanized	0.7218		0.0382		0.4950	
Less Urbanized	0.7707		0.0843		0.0557	
Completely Rural	0.9185		0.5493		0.8568	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.