

**Table 2.62A (REVISED) Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics:**  
**Numbers in Thousands, 2007 and 2008**

<b>Geographic Characteristic</b>	<b>Lifetime (2007)</b>	<b>Lifetime (2008)</b>	<b>Past Year (2007)</b>	<b>Past Year (2008)</b>	<b>Past Month (2007)</b>	<b>Past Month (2008)</b>
<b>TOTAL</b>	88,522	89,765	26,649	26,282	13,300	13,225
<b>GEOGRAPHIC DIVISION</b>						
Northeast	15,373	15,765	4,541	4,384	2,227	2,103
New England	4,442	4,386	1,310	1,310	593	536
Middle Atlantic	10,931	11,379	3,231	3,074	1,634	1,567
Midwest	21,508	22,155	6,453	6,793	3,166	3,331
East North Central	14,874	15,107	4,563	4,835	2,290	2,443
West North Central	6,634	7,048	1,890	1,958	875	887
South	30,851	31,002	9,717	8,979	5,138	5,128
South Atlantic	16,308	16,553	5,142	4,711	2,823	2,750
East South Central	5,129	5,234	1,564	1,540	865	853
West South Central	9,414	9,215	3,011	2,727	1,451	1,524
West	20,790	20,843	5,938	6,127	2,769	2,665
Mountain	6,927	6,645	2,088	1,865	1,030	796
Pacific	13,862	14,198	3,850	4,262	1,739	1,869
<b>COUNTY TYPE</b>						
Large Metro	45,674	46,993	14,344	14,389	6,853	7,332
Small Metro	27,583	27,867	8,287	7,944	4,290	3,910
250K – 1 Mil. Pop.	18,152	18,283	5,622	5,184	2,998 <sup>a</sup>	2,554
< 250K Pop.	9,431	9,584	2,665	2,760	1,292	1,356
Nonmetro	15,265	14,905	4,018	3,950	2,157	1,983
Urbanized	6,652	6,443	1,767	1,794	955	888
Less Urbanized	7,086	7,053	1,920	1,879	1,052	941
Completely Rural	1,527	1,409	331	277	150	155

\*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

<sup>a</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 2.62B (REVISED) Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics:**  
**Percentages, 2007 and 2008**

<b>Geographic Characteristic</b>	<b>Lifetime (2007)</b>	<b>Lifetime (2008)</b>	<b>Past Year (2007)</b>	<b>Past Year (2008)</b>	<b>Past Month (2007)</b>	<b>Past Month (2008)</b>
<b>TOTAL</b>	35.7	35.9	10.8	10.5	5.4	5.3
<b>GEOGRAPHIC DIVISION</b>						
Northeast	33.5	34.2	9.9	9.5	4.9	4.6
New England	36.9	36.4	10.9	10.9	4.9	4.4
Middle Atlantic	32.3	33.4	9.5	9.0	4.8	4.6
Midwest	39.2	40.3	11.8	12.4	5.8	6.1
East North Central	38.8	39.4	11.9	12.6	6.0	6.4
West North Central	40.2	42.5	11.5	11.8	5.3	5.4
South	34.3	34.1	10.8 <sup>a</sup>	9.9	5.7	5.6
South Atlantic	34.3	34.5	10.8	9.8	5.9	5.7
East South Central	34.8	35.2	10.6	10.4	5.9	5.7
West South Central	34.1	32.8	10.9	9.7	5.3	5.4
West	36.3	36.1	10.4	10.6	4.8	4.6
Mountain	40.1	37.8	12.1	10.6	6.0 <sup>a</sup>	4.5
Pacific	34.7	35.3	9.6	10.6	4.4	4.6
<b>COUNTY TYPE</b>						
Large Metro	34.4	35.3	10.8	10.8	5.2	5.5
Small Metro	37.2	36.9	11.2	10.5	5.8	5.2
250K – 1 Mil. Pop.	36.6	36.6	11.3	10.4	6.0 <sup>a</sup>	5.1
< 250K Pop.	38.4	37.6	10.9	10.8	5.3	5.3
Nonmetro	37.2	36.1	9.8	9.6	5.3	4.8
Urbanized	39.6	37.5	10.5	10.4	5.7	5.2
Less Urbanized	36.0	35.2	9.8	9.4	5.3	4.7
Completely Rural	33.2	35.0	7.2	6.9	3.3	3.8

\*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

<sup>a</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 2.63A (REVISED) Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Numbers in Thousands, 2007 and 2008**

<b>Geographic Characteristic</b>	<b>Lifetime (2007)</b>	<b>Lifetime (2008)</b>	<b>Past Year (2007)</b>	<b>Past Year (2008)</b>	<b>Past Month (2007)</b>	<b>Past Month (2008)</b>
<b>TOTAL</b>	3,290	3,119	2,373 <sup>b</sup>	2,127	1,073 <sup>a</sup>	951
<b>GEOGRAPHIC DIVISION</b>						
Northeast	542	523	394	379	185	166
New England	173	155	130	120	53	44
Middle Atlantic	369	368	264	259	131	122
Midwest	796	773	599 <sup>a</sup>	523	271	251
East North Central	580	560	434 <sup>a</sup>	364	211	180
West North Central	217	213	165	160	60	71
South	1,232 <sup>a</sup>	1,071	870 <sup>b</sup>	691	403 <sup>a</sup>	329
South Atlantic	574	570	419	371	196	168
East South Central	253 <sup>b</sup>	188	168 <sup>b</sup>	101	77	53
West South Central	405 <sup>a</sup>	312	283 <sup>a</sup>	219	130	108
West	720	752	510	534	214	205
Mountain	257	272	173	191	82	66
Pacific	463	480	337	343	132	139
<b>COUNTY TYPE</b>						
Large Metro	1,606	1,529	1,146	1,104	516	491
Small Metro	1,023	1,024	740	655	323	285
250K – 1 Mil. Pop.	659	655	475	421	206	177
< 250K Pop.	364	369	265	234	117	107
Nonmetro	661 <sup>a</sup>	566	486 <sup>b</sup>	369	234 <sup>a</sup>	176
Urbanized	273	256	204	177	101	82
Less Urbanized	334 <sup>a</sup>	259	252 <sup>b</sup>	161	123 <sup>a</sup>	80
Completely Rural	53	51	30	30	11	14

\*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

<sup>a</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 2.63B (REVISED) Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics:**  
**Percentages, 2007 and 2008**

<b>Geographic Characteristic</b>	<b>Lifetime (2007)</b>	<b>Lifetime (2008)</b>	<b>Past Year (2007)</b>	<b>Past Year (2008)</b>	<b>Past Month (2007)</b>	<b>Past Month (2008)</b>
<b>TOTAL</b>	13.0	12.5	9.4 <sup>a</sup>	8.5	4.3	3.8
<b>GEOGRAPHIC DIVISION</b>						
Northeast	12.2	12.0	8.8	8.7	4.1	3.8
New England	15.0	13.7	11.3	10.6	4.6	3.9
Middle Atlantic	11.2	11.4	8.0	8.0	4.0	3.7
Midwest	14.2	14.0	10.7	9.5	4.8	4.6
East North Central	14.7	14.5	11.0 <sup>a</sup>	9.4	5.4	4.6
West North Central	12.9	13.0	9.8	9.7	3.6	4.3
South	13.5 <sup>a</sup>	11.8	9.5 <sup>b</sup>	7.6	4.4 <sup>a</sup>	3.6
South Atlantic	12.4	12.5	9.0	8.1	4.2	3.7
East South Central	17.1 <sup>b</sup>	12.8	11.3 <sup>b</sup>	6.9	5.2	3.6
West South Central	13.5 <sup>a</sup>	10.4	9.4 <sup>a</sup>	7.3	4.3	3.6
West	11.9	12.6	8.4	9.0	3.5	3.4
Mountain	14.2	15.0	9.6	10.6	4.5	3.7
Pacific	10.9	11.6	8.0	8.3	3.1	3.4
<b>COUNTY TYPE</b>						
Large Metro	11.7	11.5	8.4	8.3	3.8	3.7
Small Metro	13.9	13.8	10.0	8.8	4.4	3.8
250K – 1 Mil. Pop.	13.2	13.4	9.5	8.6	4.1	3.6
< 250K Pop.	15.2	14.7	11.1	9.3	4.9	4.3
Nonmetro	15.9 <sup>a</sup>	13.5	11.7 <sup>b</sup>	8.8	5.6 <sup>a</sup>	4.2
Urbanized	15.9	14.3	11.9	9.9	5.9	4.6
Less Urbanized	16.5 <sup>b</sup>	12.9	12.4 <sup>b</sup>	8.0	6.1 <sup>b</sup>	3.9
Completely Rural	12.5	13.4	7.1	8.1	2.5	3.8

\*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

<sup>a</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 2.64A (REVISED) Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Numbers in Thousands, 2007 and 2008**

<b>Geographic Characteristic</b>	<b>Lifetime (2007)</b>	<b>Lifetime (2008)</b>	<b>Past Year (2007)</b>	<b>Past Year (2008)</b>	<b>Past Month (2007)</b>	<b>Past Month (2008)</b>
<b>TOTAL</b>	13,887	13,792	7,756	7,601	3,887	3,770
<b>GEOGRAPHIC DIVISION</b>						
Northeast	2,380	2,484	1,348	1,344	626	670
New England	710	723	428	385	196	185
Middle Atlantic	1,670	1,761	920	959	429	485
Midwest	3,511	3,483	1,980	1,996	1,002	986
East North Central	2,401	2,343	1,369	1,338	706	669
West North Central	1,109	1,139	611	657	297	317
South	4,704	4,512	2,593	2,588	1,400	1,374
South Atlantic	2,425 <sup>a</sup>	2,219	1,349	1,240	697	653
East South Central	764	780	445	470	260	259
West South Central	1,515	1,513	799	877	443	462
West	3,293	3,313	1,836	1,674	859	740
Mountain	1,069	1,079	611	573	284	248
Pacific	2,224	2,234	1,225	1,101	575	492
<b>COUNTY TYPE</b>						
Large Metro	6,975	7,050	3,867	3,877	1,971	1,939
Small Metro	4,576	4,524	2,539	2,493	1,221	1,230
250K – 1 Mil. Pop.	2,955	2,866	1,624	1,544	799	776
< 250K Pop.	1,621	1,658	915	949	422	454
Nonmetro	2,337	2,219	1,350	1,231	695	600
Urbanized	1,111	1,070	676	586	351	289
Less Urbanized	1,068	1,004	594	576	304	289
Completely Rural	157	145	81	69	40	22

\*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

<sup>a</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 2.64B (REVISED) Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics:**  
**Percentages, 2007 and 2008**

<b>Geographic Characteristic</b>	<b>Lifetime (2007)</b>	<b>Lifetime (2008)</b>	<b>Past Year (2007)</b>	<b>Past Year (2008)</b>	<b>Past Month (2007)</b>	<b>Past Month (2008)</b>
<b>TOTAL</b>	42.4	41.9	23.7	23.1	11.9	11.4
<b>GEOGRAPHIC DIVISION</b>						
Northeast	40.3	41.5	22.8	22.4	10.6	11.2
New England	46.5	46.4	28.0	24.7	12.9	11.9
Middle Atlantic	38.2	39.8	21.0	21.6	9.8	10.9
Midwest	48.2	47.9	27.2	27.4	13.8	13.6
East North Central	47.8	46.5	27.2	26.6	14.0	13.3
West North Central	49.0	50.9	27.0	29.3	13.1	14.2
South	40.2	38.4	22.2	22.0	12.0	11.7
South Atlantic	41.1 <sup>b</sup>	37.2	22.8	20.8	11.8	10.9
East South Central	40.9	41.6	23.8	25.1	13.9	13.8
West South Central	38.7	38.6	20.4	22.3	11.3	11.8
West	41.9	41.9	23.4	21.2	10.9	9.4
Mountain	45.4	45.6	25.9	24.2	12.1	10.5
Pacific	40.5	40.3	22.3	19.9	10.5	8.9
<b>COUNTY TYPE</b>						
Large Metro	40.3	39.5	22.3	21.7	11.4	10.9
Small Metro	44.3	44.8	24.6	24.7	11.8	12.2
250K – 1 Mil. Pop.	43.6	43.8	24.0	23.6	11.8	11.9
< 250K Pop.	45.6	46.6	25.7	26.7	11.9	12.8
Nonmetro	45.8	44.4	26.5	24.7	13.6	12.0
Urbanized	48.7	46.8	29.7 <sup>a</sup>	25.7	15.4 <sup>a</sup>	12.7
Less Urbanized	43.7	42.7	24.3	24.5	12.4	12.3
Completely Rural	42.3	40.9	21.7	19.3	10.8	6.3

\*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

<sup>a</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 2.65A (REVISED) Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics:**  
**Numbers in Thousands, 2007 and 2008**

<b>Geographic Characteristic</b>	<b>Lifetime (2007)</b>	<b>Lifetime (2008)</b>	<b>Past Year (2007)</b>	<b>Past Year (2008)</b>	<b>Past Month (2007)</b>	<b>Past Month (2008)</b>
<b>TOTAL</b>	71,345	72,853	16,519	16,554	8,340	8,504
<b>GEOGRAPHIC DIVISION</b>						
Northeast	12,450	12,757	2,799	2,661	1,417	1,267
New England	3,558	3,509	752	805	343	306
Middle Atlantic	8,892	9,249	2,048	1,856	1,074	960
Midwest	17,201	17,899	3,874	4,274	1,892	2,094
East North Central	11,892	12,203	2,760	3,133	1,374	1,595
West North Central	5,309	5,695	1,114	1,141	518	499
South	24,916	25,419	6,254	5,700	3,335	3,424
South Atlantic	13,310	13,764	3,374	3,100	1,930	1,929
East South Central	4,111	4,266	951	969	527	541
West South Central	7,495	7,389	1,929	1,631	877	954
West	16,777	16,778	3,592	3,919	1,696	1,719
Mountain	5,602	5,294	1,303	1,101	664	482
Pacific	11,175	11,484	2,289	2,817	1,032	1,237
<b>COUNTY TYPE</b>						
Large Metro	37,093	38,414	9,331	9,408	4,367	4,902
Small Metro	21,984	22,319	5,007	4,796	2,746	2,395
250K – 1 Mil. Pop.	14,538	14,762	3,523	3,219	1,993	1,601
< 250K Pop.	7,446	7,557	1,484	1,577	753	795
Nonmetro	12,268	12,119	2,181	2,350	1,227	1,207
Urbanized	5,267	5,117	887	1,030	503	517
Less Urbanized	5,684	5,789	1,074	1,143	625	573
Completely Rural	1,316	1,213	220	178	99	118

\*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

<sup>a</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 2.65B (REVISED) Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Percentages, 2007 and 2008**

<b>Geographic Characteristic</b>	<b>Lifetime (2007)</b>	<b>Lifetime (2008)</b>	<b>Past Year (2007)</b>	<b>Past Year (2008)</b>	<b>Past Month (2007)</b>	<b>Past Month (2008)</b>
<b>TOTAL</b>	37.6	37.9	8.7	8.6	4.4	4.4
<b>GEOGRAPHIC DIVISION</b>						
Northeast	35.1	35.7	7.9	7.4	4.0	3.5
New England	38.1	37.4	8.0	8.6	3.7	3.3
Middle Atlantic	34.0	35.1	7.8	7.0	4.1	3.6
Midwest	41.1	42.4	9.2	10.1	4.5	5.0
East North Central	40.5	41.4	9.4	10.6	4.7	5.4
West North Central	42.3	44.9	8.9	9.0	4.1	3.9
South	36.0	36.2	9.0	8.1	4.8	4.9
South Atlantic	35.9	36.7	9.1	8.3	5.2	5.1
East South Central	36.1	37.0	8.4	8.4	4.6	4.7
West South Central	36.2	34.9	9.3	7.7	4.2	4.5
West	38.7	38.2	8.3	8.9	3.9	3.9
Mountain	42.7	39.5	9.9	8.2	5.1	3.6
Pacific	37.0	37.6	7.6	9.2	3.4	4.1
<b>COUNTY TYPE</b>						
Large Metro	36.5	37.7	9.2	9.2	4.3	4.8
Small Metro	39.0	38.5	8.9	8.3	4.9	4.1
250K – 1 Mil. Pop.	38.5	38.3	9.3	8.3	5.3 <sup>a</sup>	4.2
< 250K Pop.	40.0	38.9	8.0	8.1	4.1	4.1
Nonmetro	38.6	37.8	6.9	7.3	3.9	3.8
Urbanized	41.1	39.1	6.9	7.9	3.9	3.9
Less Urbanized	37.4	36.9	7.1	7.3	4.1	3.7
Completely Rural	34.6	36.8	5.8	5.4	2.6	3.6

\*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

<sup>a</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.



**Table 2.66A (REVISED) Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics:**  
**Numbers in Thousands, 2007 and 2008**

<b>Geographic Characteristic</b>	<b>Lifetime (2007)</b>	<b>Lifetime (2008)</b>	<b>Past Year (2007)</b>	<b>Past Year (2008)</b>	<b>Past Month (2007)</b>	<b>Past Month (2008)</b>
<b>TOTAL</b>	85,231	86,645	24,276	24,155	12,227	12,274
<b>GEOGRAPHIC DIVISION</b>						
Northeast	14,830	15,241	4,147	4,005	2,043	1,937
New England	4,269	4,231	1,180	1,190	540	492
Middle Atlantic	10,562	11,010	2,967	2,815	1,503	1,445
Midwest	20,711	21,382	5,854	6,270	2,895	3,080
East North Central	14,294	14,547	4,128	4,471	2,079	2,264
West North Central	6,418	6,835	1,725	1,798	815	816
South	29,620	29,931	8,847	8,288	4,735	4,798
South Atlantic	15,735	15,983	4,723	4,340	2,627	2,582
East South Central	4,875	5,046	1,396	1,439	787	800
West South Central	9,010	8,902	2,728	2,508	1,321	1,417
West	20,070	20,091	5,428	5,593	2,555	2,459
Mountain	6,671	6,373	1,915	1,674	948	730
Pacific	13,399	13,718	3,514	3,919	1,607	1,729
<b>COUNTY TYPE</b>						
Large Metro	44,067	45,464	13,198	13,285	6,338	6,841
Small Metro	26,560	26,843	7,547	7,289	3,967	3,625
250K – 1 Mil. Pop.	17,493	17,628	5,147	4,763	2,792	2,377
< 250K Pop.	9,067	9,215	2,399	2,526	1,175	1,248
Nonmetro	14,604	14,339	3,531	3,581	1,922	1,808
Urbanized	6,379	6,187	1,563	1,616	854	806
Less Urbanized	6,752	6,793	1,667	1,718	929	861
Completely Rural	1,473	1,359	301	246	139	140

\*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

<sup>a</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 2.66B (REVISED) Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics:**  
**Percentages, 2007 and 2008**

<b>Geographic Characteristic</b>	<b>Lifetime (2007)</b>	<b>Lifetime (2008)</b>	<b>Past Year (2007)</b>	<b>Past Year (2008)</b>	<b>Past Month (2007)</b>	<b>Past Month (2008)</b>
<b>TOTAL</b>	38.3	38.5	10.9	10.7	5.5	5.5
<b>GEOGRAPHIC DIVISION</b>						
Northeast	35.8	36.5	10.0	9.6	4.9	4.6
New England	39.3	38.7	10.9	10.9	5.0	4.5
Middle Atlantic	34.6	35.8	9.7	9.1	4.9	4.7
Midwest	42.1	43.2	11.9	12.7	5.9	6.2
East North Central	41.6	42.1	12.0	13.0	6.1	6.6
West North Central	43.3	45.8	11.6	12.0	5.5	5.5
South	36.7	36.5	10.9	10.1	5.9	5.9
South Atlantic	36.6	36.8	11.0	10.0	6.1	5.9
East South Central	36.8	37.7	10.5	10.7	5.9	6.0
West South Central	36.6	35.5	11.1	10.0	5.4	5.6
West	39.2	38.8	10.6	10.8	5.0	4.7
Mountain	43.1	40.4	12.4	10.6	6.1 <sup>a</sup>	4.6
Pacific	37.5	38.0	9.8	10.9	4.5	4.8
<b>COUNTY TYPE</b>						
Large Metro	37.0	38.0	11.1	11.1	5.3	5.7
Small Metro	39.8	39.4	11.3	10.7	5.9	5.3
250K – 1 Mil. Pop.	39.2	39.1	11.5	10.6	6.3 <sup>a</sup>	5.3
< 250K Pop.	40.9	40.1	10.8	11.0	5.3	5.4
Nonmetro	39.6	38.7	9.6	9.7	5.2	4.9
Urbanized	42.3	40.2	10.4	10.5	5.7	5.2
Less Urbanized	38.3	37.7	9.5	9.5	5.3	4.8
Completely Rural	35.3	37.2	7.2	6.8	3.3	3.8

\*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

<sup>a</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.