

Table 2.67C (REVISED) Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics:
Standard Errors of Numbers in Thousands, 2007 and 2008

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	721	706	957	956	1,019	983
GEOGRAPHIC DIVISION						
Northeast	299	305	375	394	421	410
New England	121	135	175	157	191	187
Middle Atlantic	270	274	330	362	372	364
Midwest	232	230	353	341	372	369
East North Central	186	196	285	286	311	322
West North Central	136	121	197	191	207	197
South	465	461	634	638	631	659
South Atlantic	341	340	481	477	495	505
East South Central	169	154	218	221	213	229
West South Central	250	246	335	321	306	321
West	386	378	454	506	504	531
Mountain	168	138	202	218	205	224
Pacific	350	350	410	457	462	481
COUNTY TYPE						
Large Metro	1,745	1,760	1,514	1,553	1,308	1,345
Small Metro	1,339	1,344	1,114	1,125	931	958
250K – 1 Mil. Pop.	1,119	1,141	926	980	785	831
< 250K Pop.	818	840	672	694	548	570
Nonmetro	882	963	667	756	550	623
Urbanized	617	709	485	572	383	468
Less Urbanized	660	665	504	509	416	416
Completely Rural	410	347	289	248	243	187

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 2.67D (REVISED) Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics:
Standard Errors of Percentages, 2007 and 2008**

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	0.29	0.28	0.39	0.38	0.41	0.39
GEOGRAPHIC DIVISION						
Northeast	0.65	0.66	0.82	0.86	0.92	0.89
New England	1.01	1.12	1.45	1.30	1.59	1.55
Middle Atlantic	0.80	0.80	0.98	1.06	1.10	1.07
Midwest	0.42	0.42	0.64	0.62	0.68	0.67
East North Central	0.48	0.51	0.74	0.74	0.81	0.84
West North Central	0.82	0.73	1.19	1.15	1.26	1.19
South	0.52	0.51	0.71	0.70	0.70	0.72
South Atlantic	0.72	0.71	1.01	0.99	1.04	1.05
East South Central	1.15	1.03	1.48	1.48	1.45	1.54
West South Central	0.90	0.88	1.21	1.14	1.11	1.14
West	0.67	0.65	0.79	0.87	0.88	0.92
Mountain	0.98	0.78	1.17	1.24	1.18	1.27
Pacific	0.88	0.87	1.03	1.14	1.16	1.20
COUNTY TYPE						
Large Metro	0.41	0.41	0.52	0.54	0.59	0.56
Small Metro	0.50	0.46	0.65	0.67	0.66	0.71
250K – 1 Mil. Pop.	0.60	0.61	0.78	0.82	0.80	0.87
< 250K Pop.	0.88	0.68	1.16	1.14	1.15	1.23
Nonmetro	0.63	0.63	0.86	0.94	0.85	0.89
Urbanized	0.92	0.85	1.50	1.27	1.40	1.28
Less Urbanized	0.94	1.02	1.17	1.43	1.24	1.34
Completely Rural	2.31	2.15	2.90	3.57	2.75	3.15

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 2.67P (REVISED) Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics:
***P* Values from Tests of Differences of Percentages, 2008 Versus 2007**

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	0.8816		0.6130		0.4775	
GEOGRAPHIC DIVISION						
Northeast	0.4009		0.2988		0.7488	
New England	0.7952		0.4772		0.6645	
Middle Atlantic	0.4186		0.4192		0.5651	
Midwest	0.0949		0.4536		0.6386	
East North Central	0.0265		0.3484		0.5121	
West North Central	0.8506		0.9474		0.8332	
South	0.5908		1.0000		0.5729	
South Atlantic	0.4545		0.9737		0.8385	
East South Central	0.0631		0.2213		0.3948	
West South Central	0.2723		0.4347		0.7871	
West	0.4246		0.3897		0.4183	
Mountain	0.1671		0.6321		0.9606	
Pacific	0.8032		0.4702		0.3689	
COUNTY TYPE						
Large Metro	0.2414		0.9570		0.8250	
Small Metro	0.6002		0.6407		0.8326	
250K – 1 Mil. Pop.	0.7706		0.9352		0.7615	
< 250K Pop.	0.1438		0.4936		0.4562	
Nonmetro	0.2269		0.5480		0.2214	
Urbanized	0.7242		0.8502		0.1925	
Less Urbanized	0.3476		0.5024		0.4788	
Completely Rural	0.4451		0.8575		0.5381	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 2.68C (REVISED) Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2007 and 2008

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	112	108	107	101	87	79
GEOGRAPHIC DIVISION						
Northeast	45	44	44	42	35	32
New England	19	21	18	21	16	17
Middle Atlantic	40	38	40	36	31	28
Midwest	44	41	40	40	32	31
East North Central	35	34	33	32	26	26
West North Central	27	24	21	23	18	16
South	71	68	66	61	51	42
South Atlantic	54	49	51	44	40	33
East South Central	23	23	22	21	17	14
West South Central	41	40	37	35	29	26
West	60	58	57	56	49	46
Mountain	27	28	27	27	19	19
Pacific	54	51	51	49	44	42
COUNTY TYPE						
Large Metro	135	127	116	109	76	72
Small Metro	92	89	79	76	53	48
250K – 1 Mil. Pop.	73	69	63	58	42	36
< 250K Pop.	57	61	47	50	32	31
Nonmetro	64	65	53	54	39	31
Urbanized	41	46	34	39	25	23
Less Urbanized	45	47	41	36	27	20
Completely Rural	29	21	20	19	15	10

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 2.68D (REVISED) Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Percentages, 2007 and 2008

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	0.44	0.44	0.42	0.41	0.34	0.32
GEOGRAPHIC DIVISION						
Northeast	1.00	1.00	0.98	0.95	0.78	0.74
New England	1.67	1.88	1.54	1.82	1.35	1.49
Middle Atlantic	1.22	1.18	1.21	1.12	0.95	0.85
Midwest	0.79	0.75	0.70	0.72	0.57	0.55
East North Central	0.88	0.88	0.83	0.83	0.66	0.66
West North Central	1.59	1.43	1.28	1.41	1.09	1.00
South	0.78	0.75	0.73	0.67	0.56	0.47
South Atlantic	1.17	1.08	1.10	0.96	0.86	0.73
East South Central	1.56	1.55	1.46	1.43	1.12	0.93
West South Central	1.35	1.33	1.24	1.17	0.95	0.86
West	1.00	0.97	0.95	0.94	0.80	0.77
Mountain	1.50	1.54	1.48	1.47	1.08	1.03
Pacific	1.27	1.22	1.20	1.18	1.04	1.00
COUNTY TYPE						
Large Metro	0.63	0.63	0.59	0.59	0.48	0.46
Small Metro	0.82	0.71	0.75	0.68	0.63	0.54
250K – 1 Mil. Pop.	1.03	0.86	0.93	0.83	0.75	0.60
< 250K Pop.	1.23	1.28	1.15	1.19	1.06	1.09
Nonmetro	1.00	1.03	0.92	0.92	0.77	0.60
Urbanized	1.39	1.70	1.29	1.53	1.13	0.96
Less Urbanized	1.41	1.43	1.41	1.23	1.10	0.85
Completely Rural	3.78	3.07	2.53	3.01	2.56	1.94

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 2.68P (REVISED) Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2008 Versus 2007

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	0.1352		0.1533		0.0061	
GEOGRAPHIC DIVISION						
Northeast	0.5182		0.7875		0.1804	
New England	0.7554		0.4780		0.9329	
Middle Atlantic	0.3727		0.5027		0.1166	
Midwest	0.7916		0.9638		0.0606	
East North Central	0.5253		0.5295		0.1265	
West North Central	0.6790		0.4222		0.2680	
South	0.0365		0.0094		0.0796	
South Atlantic	0.3036		0.2138		0.1304	
East South Central	0.3132		0.0214		0.1032	
West South Central	0.0807		0.1126		0.9128	
West	0.6665		0.5793		0.3726	
Mountain	0.3318		0.3724		0.2288	
Pacific	0.9662		0.8899		0.1290	
COUNTY TYPE						
Large Metro	0.2917		0.5155		0.1624	
Small Metro	0.7169		0.5891		0.1560	
250K – 1 Mil. Pop.	0.6487		0.7706		0.4599	
< 250K Pop.	0.9742		0.5350		0.1638	
Nonmetro	0.2140		0.0986		0.0074	
Urbanized	0.6032		0.3744		0.1393	
Less Urbanized	0.4688		0.1893		0.0159	
Completely Rural	0.2270		0.4956		0.6965	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 2.69C (REVISED) Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2007 and 2008

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	125	126	147	144	170	163
GEOGRAPHIC DIVISION						
Northeast	50	45	58	52	69	64
New England	21	22	24	22	32	29
Middle Atlantic	45	40	53	48	60	57
Midwest	37	39	45	50	63	61
East North Central	30	33	36	40	51	50
West North Central	22	21	27	28	36	34
South	74	75	91	91	104	103
South Atlantic	53	58	66	68	77	80
East South Central	31	24	34	30	37	36
West South Central	43	41	54	51	56	53
West	83	78	91	82	96	92
Mountain	54	41	55	42	51	43
Pacific	62	66	71	70	81	80
COUNTY TYPE						
Large Metro	325	329	316	312	261	263
Small Metro	242	233	233	218	208	187
250K – 1 Mil. Pop.	189	182	184	170	165	143
< 250K Pop.	161	159	152	149	134	131
Nonmetro	167	202	156	195	133	158
Urbanized	119	173	107	168	86	137
Less Urbanized	123	110	118	102	103	83
Completely Rural	46	34	41	31	35	23

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 2.69D (REVISED) Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Percentages, 2007 and 2008

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	0.38	0.38	0.45	0.44	0.52	0.49
GEOGRAPHIC DIVISION						
Northeast	0.84	0.76	0.98	0.87	1.16	1.06
New England	1.40	1.42	1.57	1.44	2.08	1.88
Middle Atlantic	1.02	0.90	1.20	1.07	1.37	1.28
Midwest	0.51	0.53	0.62	0.68	0.86	0.84
East North Central	0.60	0.65	0.72	0.79	1.02	0.99
West North Central	0.96	0.93	1.20	1.24	1.60	1.52
South	0.63	0.64	0.78	0.77	0.89	0.87
South Atlantic	0.89	0.98	1.12	1.14	1.30	1.35
East South Central	1.66	1.30	1.80	1.62	1.98	1.93
West South Central	1.09	1.04	1.37	1.30	1.44	1.36
West	1.06	0.98	1.16	1.03	1.22	1.16
Mountain	2.28	1.74	2.34	1.77	2.18	1.83
Pacific	1.13	1.19	1.29	1.26	1.47	1.45
COUNTY TYPE						
Large Metro	0.49	0.50	0.60	0.59	0.70	0.68
Small Metro	0.81	0.69	0.89	0.75	0.97	0.89
250K – 1 Mil. Pop.	1.07	0.91	1.13	0.95	1.19	1.09
< 250K Pop.	1.10	0.98	1.42	1.24	1.65	1.48
Nonmetro	0.85	0.95	0.96	1.14	1.18	1.19
Urbanized	1.16	1.22	1.31	1.54	1.50	1.64
Less Urbanized	1.40	1.49	1.61	1.68	2.03	1.72
Completely Rural	3.20	2.56	3.30	3.20	4.00	3.34

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 2.69P (REVISED) Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2008 Versus 2007

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	0.4409		0.8694		0.8178	
GEOGRAPHIC DIVISION						
Northeast	0.1601		0.1085		0.6642	
New England	0.7418		0.9073		0.7510	
Middle Atlantic	0.1564		0.0851		0.5119	
Midwest	0.1257		0.0423		0.4118	
East North Central	0.1697		0.0191		0.0949	
West North Central	0.4966		0.8056		0.3753	
South	0.7391		0.9615		0.6743	
South Atlantic	0.2756		0.4925		0.1270	
East South Central	0.0283		0.0471		0.0567	
West South Central	0.4870		0.6825		0.8938	
West	0.1390		0.6894		0.7924	
Mountain	0.2596		0.4717		0.6525	
Pacific	0.3167		0.9935		0.9669	
COUNTY TYPE						
Large Metro	0.2265		0.5506		0.4599	
Small Metro	0.9115		0.8360		0.8630	
250K – 1 Mil. Pop.	0.9117		0.8513		0.9657	
< 250K Pop.	0.9757		0.5003		0.8336	
Nonmetro	0.7045		0.6482		0.5907	
Urbanized	0.7805		0.6718		0.1183	
Less Urbanized	0.7826		0.2255		0.7182	
Completely Rural	0.9322		0.7025		0.3623	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 2.70C (REVISED) Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics:
Standard Errors of Numbers in Thousands, 2007 and 2008

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	679	661	933	931	986	958
GEOGRAPHIC DIVISION						
Northeast	289	283	362	381	407	401
New England	115	124	173	151	191	178
Middle Atlantic	261	254	316	350	357	359
Midwest	212	215	346	329	358	362
East North Central	169	182	282	275	301	312
West North Central	121	113	189	186	196	199
South	442	439	625	621	613	648
South Atlantic	323	326	485	471	484	504
East South Central	156	151	203	211	200	218
West South Central	242	231	323	309	295	313
West	356	353	434	486	487	508
Mountain	149	118	190	208	198	210
Pacific	326	331	392	441	445	461
COUNTY TYPE						
Large Metro	1,621	1,645	1,400	1,443	1,229	1,264
Small Metro	1,241	1,235	1,023	1,021	865	889
250K – 1 Mil. Pop.	1,027	1,046	841	888	721	766
< 250K Pop.	754	755	616	612	510	523
Nonmetro	798	859	593	663	499	557
Urbanized	537	624	416	488	336	411
Less Urbanized	597	585	448	446	374	374
Completely Rural	363	310	250	219	218	169

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 2.70D (REVISED) Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics:
Standard Errors of Percentages, 2007 and 2008**

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	0.36	0.34	0.49	0.49	0.52	0.50
GEOGRAPHIC DIVISION						
Northeast	0.81	0.79	1.02	1.07	1.15	1.12
New England	1.23	1.33	1.85	1.61	2.04	1.90
Middle Atlantic	1.00	0.96	1.21	1.33	1.36	1.36
Midwest	0.51	0.51	0.82	0.78	0.86	0.86
East North Central	0.58	0.62	0.96	0.93	1.03	1.06
West North Central	0.97	0.89	1.51	1.46	1.56	1.57
South	0.64	0.63	0.90	0.89	0.89	0.92
South Atlantic	0.87	0.87	1.31	1.26	1.31	1.34
East South Central	1.37	1.31	1.78	1.83	1.76	1.89
West South Central	1.17	1.09	1.56	1.46	1.42	1.48
West	0.82	0.80	1.00	1.11	1.12	1.16
Mountain	1.13	0.88	1.45	1.55	1.51	1.57
Pacific	1.08	1.08	1.30	1.45	1.47	1.51
COUNTY TYPE						
Large Metro	0.50	0.50	0.66	0.68	0.74	0.71
Small Metro	0.60	0.56	0.82	0.86	0.86	0.90
250K – 1 Mil. Pop.	0.71	0.75	0.99	1.06	1.03	1.11
< 250K Pop.	1.10	0.84	1.49	1.47	1.46	1.56
Nonmetro	0.77	0.78	1.07	1.17	1.05	1.11
Urbanized	1.10	0.94	1.88	1.55	1.76	1.65
Less Urbanized	1.15	1.28	1.45	1.77	1.50	1.66
Completely Rural	2.71	2.57	3.44	4.38	3.25	3.70

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 2.70P (REVISED) Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics:
P Values from Tests of Differences of Percentages, 2008 Versus 2007

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	0.7397		0.6147		0.3972	
GEOGRAPHIC DIVISION						
Northeast	0.5251		0.4477		0.7818	
New England	0.9452		0.5737		0.6438	
Middle Atlantic	0.5046		0.5697		0.5935	
Midwest	0.0848		0.5563		0.7804	
East North Central	0.0258		0.4970		0.7040	
West North Central	0.9071		0.9645		0.9214	
South	0.7504		0.8383		0.4822	
South Atlantic	0.5715		0.9076		0.6350	
East South Central	0.1060		0.2270		0.4908	
West South Central	0.4195		0.5639		0.8140	
West	0.7271		0.5003		0.4249	
Mountain	0.3777		0.8737		0.7531	
Pacific	0.9920		0.5032		0.3146	
COUNTY TYPE						
Large Metro	0.1173		0.8624		0.7912	
Small Metro	0.6395		0.5658		0.7705	
250K – 1 Mil. Pop.	0.6618		0.9627		0.7527	
< 250K Pop.	0.1089		0.3535		0.3703	
Nonmetro	0.1240		0.3625		0.1526	
Urbanized	0.5147		0.7643		0.1988	
Less Urbanized	0.2888		0.3011		0.3147	
Completely Rural	0.3297		0.8571		0.5954	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 2.71C (REVISED) Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics:
Standard Errors of Numbers in Thousands, 2007 and 2008

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	694	682	955	950	1,011	977
GEOGRAPHIC DIVISION						
Northeast	292	292	371	392	415	408
New England	117	132	174	157	192	186
Middle Atlantic	264	261	325	360	365	363
Midwest	217	218	352	337	373	369
East North Central	173	186	285	281	311	320
West North Central	125	114	195	189	207	197
South	456	449	639	639	630	659
South Atlantic	331	328	488	476	498	504
East South Central	165	152	214	219	209	230
West South Central	248	241	335	322	301	321
West	365	364	446	501	498	525
Mountain	161	131	198	216	203	219
Pacific	330	339	403	453	456	477
COUNTY TYPE						
Large Metro	1,698	1,723	1,476	1,520	1,287	1,329
Small Metro	1,301	1,304	1,083	1,090	916	941
250K – 1 Mil. Pop.	1,087	1,109	900	952	774	817
< 250K Pop.	789	809	650	668	536	560
Nonmetro	853	936	645	734	540	610
Urbanized	596	686	470	551	375	457
Less Urbanized	636	642	486	492	407	408
Completely Rural	391	335	274	239	234	182

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 2.71D (REVISED) Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics:
Standard Errors of Percentages, 2007 and 2008**

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	0.31	0.30	0.43	0.42	0.45	0.43
GEOGRAPHIC DIVISION						
Northeast	0.71	0.70	0.89	0.94	1.00	0.98
New England	1.08	1.20	1.60	1.43	1.77	1.70
Middle Atlantic	0.87	0.85	1.06	1.17	1.20	1.18
Midwest	0.44	0.44	0.72	0.68	0.76	0.75
East North Central	0.50	0.54	0.83	0.81	0.91	0.93
West North Central	0.85	0.77	1.32	1.26	1.39	1.32
South	0.56	0.55	0.79	0.78	0.78	0.80
South Atlantic	0.77	0.76	1.14	1.10	1.16	1.16
East South Central	1.24	1.13	1.61	1.64	1.58	1.71
West South Central	1.01	0.96	1.36	1.28	1.22	1.28
West	0.71	0.70	0.87	0.97	0.97	1.01
Mountain	1.04	0.83	1.28	1.37	1.31	1.39
Pacific	0.92	0.94	1.13	1.26	1.28	1.32
COUNTY TYPE						
Large Metro	0.44	0.44	0.58	0.59	0.65	0.62
Small Metro	0.53	0.49	0.71	0.74	0.73	0.78
250K – 1 Mil. Pop.	0.63	0.65	0.86	0.91	0.89	0.96
< 250K Pop.	0.94	0.73	1.27	1.27	1.26	1.35
Nonmetro	0.68	0.70	0.96	1.05	0.95	0.98
Urbanized	0.96	0.82	1.64	1.36	1.55	1.42
Less Urbanized	1.02	1.16	1.31	1.60	1.38	1.49
Completely Rural	2.53	2.36	3.23	3.99	3.03	3.41

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 2.71P (REVISED) Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics:
***P* Values from Tests of Differences of Percentages, 2008 Versus 2007**

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	0.8485		0.6126		0.4321	
GEOGRAPHIC DIVISION						
Northeast	0.4025		0.3234		0.7270	
New England	0.8981		0.5699		0.6251	
Middle Atlantic	0.3871		0.4118		0.5217	
Midwest	0.0545		0.3950		0.6821	
East North Central	0.0167		0.3359		0.5392	
West North Central	0.9922		0.9245		0.8062	
South	0.7185		0.8556		0.5375	
South Atlantic	0.4682		0.9781		0.8022	
East South Central	0.0495		0.1579		0.3544	
West South Central	0.3689		0.5231		0.8087	
West	0.5227		0.4693		0.4098	
Mountain	0.2474		0.7518		0.8351	
Pacific	0.8724		0.5168		0.3242	
COUNTY TYPE						
Large Metro	0.1775		0.9572		0.8724	
Small Metro	0.6177		0.6367		0.7867	
250K – 1 Mil. Pop.	0.7059		0.9778		0.7363	
< 250K Pop.	0.1174		0.4415		0.3768	
Nonmetro	0.1492		0.4367		0.1484	
Urbanized	0.5627		0.7450		0.1429	
Less Urbanized	0.3220		0.4349		0.3853	
Completely Rural	0.3428		0.9082		0.5613	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 2.72C (REVISED) Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2007 and 2008

Geographic Characteristic	Alcohol Use (2007)	Alcohol Use (2008)	Binge Alcohol Use (2007)	Binge Alcohol Use (2008)	Heavy Alcohol Use (2007)	Heavy Alcohol Use (2008)
TOTAL	1,019	983	770	735	429	414
GEOGRAPHIC DIVISION						
Northeast	421	410	321	307	174	183
New England	191	187	155	139	91	79
Middle Atlantic	372	364	279	274	149	165
Midwest	372	369	325	313	174	183
East North Central	311	322	254	270	149	149
West North Central	207	197	184	164	93	111
South	631	659	467	484	264	269
South Atlantic	495	505	365	371	206	199
East South Central	213	229	143	160	83	99
West South Central	306	321	252	251	136	148
West	504	531	387	370	224	209
Mountain	205	224	171	168	121	95
Pacific	462	481	344	331	187	188
COUNTY TYPE						
Large Metro	1,308	1,345	685	733	340	326
Small Metro	931	958	523	470	261	227
250K – 1 Mil. Pop.	785	831	434	395	227	180
< 250K Pop.	548	570	305	290	130	148
Nonmetro	550	623	353	385	165	215
Urbanized	383	468	263	287	130	150
Less Urbanized	416	416	253	258	102	135
Completely Rural	243	187	130	110	35	76

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 2.72D (REVISED) Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2007 and 2008

Geographic Characteristic	Alcohol Use (2007)	Alcohol Use (2008)	Binge Alcohol Use (2007)	Binge Alcohol Use (2008)	Heavy Alcohol Use (2007)	Heavy Alcohol Use (2008)
TOTAL	0.41	0.39	0.31	0.29	0.17	0.17
GEOGRAPHIC DIVISION						
Northeast	0.92	0.89	0.70	0.67	0.38	0.40
New England	1.59	1.55	1.29	1.16	0.75	0.66
Middle Atlantic	1.10	1.07	0.83	0.80	0.44	0.49
Midwest	0.68	0.67	0.59	0.57	0.32	0.33
East North Central	0.81	0.84	0.66	0.70	0.39	0.39
West North Central	1.26	1.19	1.11	0.99	0.56	0.67
South	0.70	0.72	0.52	0.53	0.29	0.30
South Atlantic	1.04	1.05	0.77	0.77	0.43	0.41
East South Central	1.45	1.54	0.97	1.08	0.56	0.67
West South Central	1.11	1.14	0.91	0.90	0.49	0.53
West	0.88	0.92	0.68	0.64	0.39	0.36
Mountain	1.18	1.27	0.99	0.95	0.70	0.54
Pacific	1.16	1.20	0.86	0.82	0.47	0.47
COUNTY TYPE						
Large Metro	0.59	0.56	0.44	0.43	0.25	0.23
Small Metro	0.66	0.71	0.54	0.50	0.31	0.28
250K – 1 Mil. Pop.	0.80	0.87	0.62	0.61	0.39	0.33
< 250K Pop.	1.15	1.23	0.98	0.92	0.50	0.57
Nonmetro	0.85	0.89	0.69	0.72	0.37	0.47
Urbanized	1.40	1.28	1.06	1.16	0.67	0.77
Less Urbanized	1.24	1.34	1.01	1.03	0.48	0.62
Completely Rural	2.75	3.15	2.02	2.33	0.65	1.78

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 2.72P (REVISED) Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2008 Versus 2007

Geographic Characteristic	Alcohol Use (2007)	Alcohol Use (2008)	Binge Alcohol Use (2007)	Binge Alcohol Use (2008)	Heavy Alcohol Use (2007)	Heavy Alcohol Use (2008)
TOTAL	0.4775		0.9626		0.7231	
GEOGRAPHIC DIVISION						
Northeast	0.7488		0.3122		0.2275	
New England	0.6645		0.9892		0.4966	
Middle Atlantic	0.5651		0.2507		0.0830	
Midwest	0.6386		0.5882		0.6718	
East North Central	0.5121		0.8326		0.9840	
West North Central	0.8332		0.5144		0.4865	
South	0.5729		0.6155		0.5847	
South Atlantic	0.8385		0.9745		0.9266	
East South Central	0.3948		0.9146		0.4737	
West South Central	0.7871		0.3944		0.6388	
West	0.4183		0.3628		0.1578	
Mountain	0.9606		0.4162		0.3552	
Pacific	0.3689		0.5228		0.2737	
COUNTY TYPE						
Large Metro	0.8250		0.3116		0.3907	
Small Metro	0.8326		0.2544		0.1962	
250K – 1 Mil. Pop.	0.7615		0.0583		0.0194	
< 250K Pop.	0.4562		0.5784		0.2763	
Nonmetro	0.2214		0.7215		0.3351	
Urbanized	0.1925		0.4158		0.9972	
Less Urbanized	0.4788		0.7180		0.3143	
Completely Rural	0.5381		0.7815		0.3070	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 2.73C (REVISED) Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2007 and 2008

Geographic Characteristic	Alcohol Use (2007)	Alcohol Use (2008)	Binge Alcohol Use (2007)	Binge Alcohol Use (2008)	Heavy Alcohol Use (2007)	Heavy Alcohol Use (2008)
TOTAL	87	79	69	57	37	30
GEOGRAPHIC DIVISION						
Northeast	35	32	28	27	14	13
New England	16	17	12	15	6	7
Middle Atlantic	31	28	25	22	12	11
Midwest	32	31	27	23	18	12
East North Central	26	26	21	20	13	10
West North Central	18	16	16	12	12	6
South	51	42	42	33	22	17
South Atlantic	40	33	32	24	16	13
East South Central	17	14	14	12	6	4
West South Central	29	26	24	21	13	11
West	49	46	38	32	19	18
Mountain	19	19	16	15	7	9
Pacific	44	42	34	28	17	16
COUNTY TYPE						
Large Metro	76	72	55	48	26	23
Small Metro	53	48	39	36	18	18
250K – 1 Mil. Pop.	42	36	31	27	14	15
< 250K Pop.	32	31	23	23	11	9
Nonmetro	39	31	32	25	18	12
Urbanized	25	23	18	18	9	9
Less Urbanized	27	20	23	17	12	7
Completely Rural	15	10	14	7	12	2

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 2.73D (REVISED) Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Percentages, 2007 and 2008

Geographic Characteristic	Alcohol Use (2007)	Alcohol Use (2008)	Binge Alcohol Use (2007)	Binge Alcohol Use (2008)	Heavy Alcohol Use (2007)	Heavy Alcohol Use (2008)
TOTAL	0.34	0.32	0.27	0.23	0.15	0.12
GEOGRAPHIC DIVISION						
Northeast	0.78	0.74	0.63	0.62	0.31	0.30
New England	1.35	1.49	1.07	1.36	0.56	0.66
Middle Atlantic	0.95	0.85	0.76	0.68	0.37	0.33
Midwest	0.57	0.55	0.47	0.42	0.31	0.21
East North Central	0.66	0.66	0.54	0.51	0.33	0.26
West North Central	1.09	1.00	0.95	0.74	0.71	0.36
South	0.56	0.47	0.47	0.37	0.24	0.19
South Atlantic	0.86	0.73	0.70	0.53	0.35	0.29
East South Central	1.12	0.93	0.96	0.80	0.40	0.28
West South Central	0.95	0.86	0.80	0.69	0.42	0.36
West	0.80	0.77	0.63	0.54	0.31	0.31
Mountain	1.08	1.03	0.87	0.85	0.41	0.48
Pacific	1.04	1.00	0.79	0.68	0.41	0.39
COUNTY TYPE						
Large Metro	0.48	0.46	0.36	0.33	0.19	0.17
Small Metro	0.63	0.54	0.49	0.44	0.24	0.24
250K – 1 Mil. Pop.	0.75	0.60	0.58	0.50	0.28	0.31
< 250K Pop.	1.06	1.09	0.84	0.85	0.46	0.36
Nonmetro	0.77	0.60	0.66	0.53	0.43	0.27
Urbanized	1.13	0.96	0.87	0.87	0.51	0.47
Less Urbanized	1.10	0.85	1.00	0.75	0.59	0.35
Completely Rural	2.56	1.94	2.62	1.55	2.63	0.59

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 2.73P (REVISED) Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 17, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2008 Versus 2007

Geographic Characteristic	Alcohol Use (2007)	Alcohol Use (2008)	Binge Alcohol Use (2007)	Binge Alcohol Use (2008)	Heavy Alcohol Use (2007)	Heavy Alcohol Use (2008)
TOTAL	0.0061		0.0132		0.0610	
GEOGRAPHIC DIVISION						
Northeast	0.1804		0.8058		0.5109	
New England	0.9329		0.4585		0.3868	
Middle Atlantic	0.1166		0.4764		0.1909	
Midwest	0.0606		0.0320		0.0137	
East North Central	0.1265		0.1932		0.1602	
West North Central	0.2680		0.0584		0.0251	
South	0.0796		0.0736		0.1895	
South Atlantic	0.1304		0.1339		0.7198	
East South Central	0.1032		0.2571		0.1284	
West South Central	0.9128		0.6343		0.2851	
West	0.3726		0.4124		0.6964	
Mountain	0.2288		0.8684		0.3588	
Pacific	0.1290		0.3048		0.9787	
COUNTY TYPE						
Large Metro	0.1624		0.0741		0.1950	
Small Metro	0.1560		0.5776		0.6759	
250K – 1 Mil. Pop.	0.4599		0.6270		0.2859	
< 250K Pop.	0.1638		0.7748		0.3916	
Nonmetro	0.0074		0.0263		0.0085	
Urbanized	0.1393		0.4386		0.4209	
Less Urbanized	0.0159		0.0639		0.0310	
Completely Rural	0.6965		0.2415		0.1442	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 2.74C (REVISED) Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2007 and 2008

Geographic Characteristic	Alcohol Use (2007)	Alcohol Use (2008)	Binge Alcohol Use (2007)	Binge Alcohol Use (2008)	Heavy Alcohol Use (2007)	Heavy Alcohol Use (2008)
TOTAL	170	163	169	167	114	124
GEOGRAPHIC DIVISION						
Northeast	69	64	73	65	50	53
New England	32	29	35	29	27	29
Middle Atlantic	60	57	64	58	42	44
Midwest	63	61	62	60	46	45
East North Central	51	50	52	46	39	37
West North Central	36	34	34	37	27	25
South	104	103	99	108	66	83
South Atlantic	77	80	74	89	50	71
East South Central	37	36	36	34	22	26
West South Central	56	53	56	49	36	32
West	96	92	92	89	64	59
Mountain	51	43	42	42	27	31
Pacific	81	80	82	78	58	49
COUNTY TYPE						
Large Metro	261	263	205	196	101	100
Small Metro	208	187	159	144	83	73
250K – 1 Mil. Pop.	165	143	124	110	68	57
< 250K Pop.	134	131	102	98	47	48
Nonmetro	133	158	107	133	47	87
Urbanized	86	137	70	116	35	79
Less Urbanized	103	83	83	67	34	38
Completely Rural	35	23	26	17	11	7

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 2.74D (REVISED) Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Percentages, 2007 and 2008

Geographic Characteristic	Alcohol Use (2007)	Alcohol Use (2008)	Binge Alcohol Use (2007)	Binge Alcohol Use (2008)	Heavy Alcohol Use (2007)	Heavy Alcohol Use (2008)
TOTAL	0.52	0.49	0.52	0.51	0.35	0.38
GEOGRAPHIC DIVISION						
Northeast	1.16	1.06	1.24	1.09	0.85	0.88
New England	2.08	1.88	2.28	1.85	1.78	1.84
Middle Atlantic	1.37	1.28	1.47	1.31	0.96	1.00
Midwest	0.86	0.84	0.85	0.82	0.63	0.62
East North Central	1.02	0.99	1.04	0.92	0.77	0.74
West North Central	1.60	1.52	1.52	1.64	1.18	1.10
South	0.89	0.87	0.85	0.92	0.56	0.70
South Atlantic	1.30	1.35	1.25	1.49	0.85	1.20
East South Central	1.98	1.93	1.93	1.83	1.19	1.41
West South Central	1.44	1.36	1.43	1.26	0.91	0.82
West	1.22	1.16	1.17	1.13	0.82	0.74
Mountain	2.18	1.83	1.78	1.78	1.13	1.31
Pacific	1.47	1.45	1.48	1.41	1.06	0.89
COUNTY TYPE						
Large Metro	0.70	0.68	0.72	0.68	0.49	0.47
Small Metro	0.97	0.89	0.94	0.85	0.62	0.60
250K – 1 Mil. Pop.	1.19	1.09	1.15	1.02	0.78	0.72
< 250K Pop.	1.65	1.48	1.53	1.47	1.03	1.08
Nonmetro	1.18	1.19	1.16	1.27	0.72	1.24
Urbanized	1.50	1.64	1.54	2.05	1.09	2.23
Less Urbanized	2.03	1.72	1.93	1.61	1.07	1.24
Completely Rural	4.00	3.34	3.39	2.95	2.70	1.85

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 2.74P (REVISED) Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 to 25, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2008 Versus 2007

Geographic Characteristic	Alcohol Use (2007)	Alcohol Use (2008)	Binge Alcohol Use (2007)	Binge Alcohol Use (2008)	Heavy Alcohol Use (2007)	Heavy Alcohol Use (2008)
TOTAL	0.8178		0.3249		0.7105	
GEOGRAPHIC DIVISION						
Northeast	0.6642		0.8565		0.5093	
New England	0.7510		0.1786		0.2524	
Middle Atlantic	0.5119		0.3635		0.1314	
Midwest	0.4118		0.8950		0.2525	
East North Central	0.0949		0.4654		0.2300	
West North Central	0.3753		0.4656		0.8124	
South	0.6743		0.1149		0.8353	
South Atlantic	0.1270		0.0545		0.5421	
East South Central	0.0567		0.8272		0.2078	
West South Central	0.8938		0.7441		0.8158	
West	0.7924		0.9619		0.0734	
Mountain	0.6525		0.5215		0.9675	
Pacific	0.9669		0.7046		0.0411	
COUNTY TYPE						
Large Metro	0.4599		0.4585		0.6143	
Small Metro	0.8630		0.4764		0.0702	
250K – 1 Mil. Pop.	0.9657		0.5959		0.0116	
< 250K Pop.	0.8336		0.5797		0.7470	
Nonmetro	0.5907		0.9425		0.5341	
Urbanized	0.1183		0.5249		0.5111	
Less Urbanized	0.7182		0.6760		0.7232	
Completely Rural	0.3623		0.2467		0.4694	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 2.75C (REVISED) Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2007 and 2008

Geographic Characteristic	Alcohol Use (2007)	Alcohol Use (2008)	Binge Alcohol Use (2007)	Binge Alcohol Use (2008)	Heavy Alcohol Use (2007)	Heavy Alcohol Use (2008)
TOTAL	986	958	714	703	402	391
GEOGRAPHIC DIVISION						
Northeast	407	401	300	297	160	170
New England	191	178	147	133	84	71
Middle Atlantic	357	359	260	266	137	154
Midwest	358	362	305	300	164	176
East North Central	301	312	244	260	141	143
West North Central	196	199	169	158	86	106
South	613	648	436	469	250	255
South Atlantic	484	504	342	360	196	185
East South Central	200	218	131	154	81	91
West South Central	295	313	233	247	128	142
West	487	508	353	350	208	199
Mountain	198	210	160	156	116	86
Pacific	445	461	313	314	173	181
COUNTY TYPE						
Large Metro	1,229	1,264	616	676	317	309
Small Metro	865	889	469	432	237	215
250K – 1 Mil. Pop.	721	766	394	362	209	171
< 250K Pop.	510	523	261	257	115	134
Nonmetro	499	557	314	335	151	189
Urbanized	336	411	227	243	118	124
Less Urbanized	374	374	223	227	93	124
Completely Rural	218	169	110	101	28	74

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 2.75D (REVISED) Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2007 and 2008

Geographic Characteristic	Alcohol Use (2007)	Alcohol Use (2008)	Binge Alcohol Use (2007)	Binge Alcohol Use (2008)	Heavy Alcohol Use (2007)	Heavy Alcohol Use (2008)
TOTAL	0.52	0.50	0.38	0.37	0.21	0.20
GEOGRAPHIC DIVISION						
Northeast	1.15	1.12	0.84	0.83	0.45	0.47
New England	2.04	1.90	1.58	1.42	0.90	0.76
Middle Atlantic	1.36	1.36	0.99	1.01	0.52	0.59
Midwest	0.86	0.86	0.73	0.71	0.39	0.42
East North Central	1.03	1.06	0.83	0.88	0.48	0.49
West North Central	1.56	1.57	1.34	1.25	0.68	0.84
South	0.89	0.92	0.63	0.67	0.36	0.36
South Atlantic	1.31	1.34	0.92	0.96	0.53	0.49
East South Central	1.76	1.89	1.15	1.34	0.71	0.79
West South Central	1.42	1.48	1.13	1.17	0.62	0.67
West	1.12	1.16	0.81	0.80	0.48	0.45
Mountain	1.51	1.57	1.22	1.17	0.88	0.64
Pacific	1.47	1.51	1.04	1.03	0.57	0.59
COUNTY TYPE						
Large Metro	0.74	0.71	0.53	0.54	0.31	0.29
Small Metro	0.86	0.90	0.67	0.63	0.38	0.35
250K – 1 Mil. Pop.	1.03	1.11	0.79	0.76	0.49	0.42
< 250K Pop.	1.46	1.56	1.17	1.13	0.58	0.68
Nonmetro	1.05	1.11	0.83	0.87	0.45	0.56
Urbanized	1.76	1.65	1.32	1.44	0.83	0.90
Less Urbanized	1.50	1.66	1.19	1.24	0.57	0.74
Completely Rural	3.25	3.70	2.28	2.69	0.73	2.13

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 2.75P (REVISED) Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 26 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2008 Versus 2007

Geographic Characteristic	Alcohol Use (2007)	Alcohol Use (2008)	Binge Alcohol Use (2007)	Binge Alcohol Use (2008)	Heavy Alcohol Use (2007)	Heavy Alcohol Use (2008)
TOTAL	0.3972		0.6697		0.5465	
GEOGRAPHIC DIVISION						
Northeast	0.7818		0.3327		0.2596	
New England	0.6438		0.8542		0.6238	
Middle Atlantic	0.5935		0.3157		0.1260	
Midwest	0.7804		0.6940		0.7272	
East North Central	0.7040		1.0000		0.8359	
West North Central	0.9214		0.4951		0.3873	
South	0.4822		0.3299		0.4728	
South Atlantic	0.6350		0.6083		0.7721	
East South Central	0.4908		0.8711		0.6106	
West South Central	0.8140		0.3093		0.5179	
West	0.4249		0.3732		0.3297	
Mountain	0.7531		0.3069		0.3072	
Pacific	0.3146		0.6076		0.5845	
COUNTY TYPE						
Large Metro	0.7912		0.2373		0.4695	
Small Metro	0.7705		0.4119		0.4374	
250K – 1 Mil. Pop.	0.7527		0.0928		0.0760	
< 250K Pop.	0.3703		0.3952		0.2180	
Nonmetro	0.1526		0.9218		0.2530	
Urbanized	0.1988		0.3737		0.8876	
Less Urbanized	0.3147		0.4458		0.2089	
Completely Rural	0.5954		0.9310		0.1914	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 2.76C (REVISED) Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2007 and 2008

Geographic Characteristic	Alcohol Use (2007)	Alcohol Use (2008)	Binge Alcohol Use (2007)	Binge Alcohol Use (2008)	Heavy Alcohol Use (2007)	Heavy Alcohol Use (2008)
TOTAL	1,011	977	766	737	426	414
GEOGRAPHIC DIVISION						
Northeast	415	408	317	309	173	184
New England	192	186	152	139	90	79
Middle Atlantic	365	363	277	276	148	167
Midwest	373	369	326	311	173	182
East North Central	311	320	254	268	148	147
West North Central	207	197	184	163	92	110
South	630	659	463	482	262	268
South Atlantic	498	504	364	369	206	197
East South Central	209	230	142	160	82	99
West South Central	301	321	247	252	134	148
West	498	525	384	369	222	210
Mountain	203	219	172	166	120	95
Pacific	456	477	342	331	185	189
COUNTY TYPE						
Large Metro	1,287	1,329	674	726	339	324
Small Metro	916	941	514	461	259	225
250K – 1 Mil. Pop.	774	817	427	389	226	179
< 250K Pop.	536	560	298	284	129	146
Nonmetro	540	610	346	377	162	214
Urbanized	375	457	256	281	129	149
Less Urbanized	407	408	248	253	100	135
Completely Rural	234	182	123	108	32	76

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 2.76D (REVISED) Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2007 and 2008

Geographic Characteristic	Alcohol Use (2007)	Alcohol Use (2008)	Binge Alcohol Use (2007)	Binge Alcohol Use (2008)	Heavy Alcohol Use (2007)	Heavy Alcohol Use (2008)
TOTAL	0.45	0.43	0.34	0.33	0.19	0.18
GEOGRAPHIC DIVISION						
Northeast	1.00	0.98	0.77	0.74	0.42	0.44
New England	1.77	1.70	1.40	1.27	0.83	0.72
Middle Atlantic	1.20	1.18	0.91	0.90	0.49	0.54
Midwest	0.76	0.75	0.66	0.63	0.35	0.37
East North Central	0.91	0.93	0.74	0.78	0.43	0.43
West North Central	1.39	1.32	1.24	1.09	0.62	0.74
South	0.78	0.80	0.57	0.59	0.32	0.33
South Atlantic	1.16	1.16	0.85	0.85	0.48	0.45
East South Central	1.58	1.71	1.07	1.19	0.62	0.74
West South Central	1.22	1.28	1.00	1.00	0.55	0.59
West	0.97	1.01	0.75	0.71	0.43	0.41
Mountain	1.31	1.39	1.11	1.06	0.78	0.60
Pacific	1.28	1.32	0.96	0.92	0.52	0.52
COUNTY TYPE						
Large Metro	0.65	0.62	0.48	0.48	0.28	0.25
Small Metro	0.73	0.78	0.59	0.55	0.34	0.31
250K – 1 Mil. Pop.	0.89	0.96	0.69	0.67	0.44	0.37
< 250K Pop.	1.26	1.35	1.08	1.02	0.55	0.63
Nonmetro	0.95	0.98	0.76	0.79	0.40	0.53
Urbanized	1.55	1.42	1.17	1.29	0.73	0.87
Less Urbanized	1.38	1.49	1.11	1.13	0.53	0.68
Completely Rural	3.03	3.41	2.20	2.51	0.69	1.95

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 2.76P (REVISED) Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2008 Versus 2007

Geographic Characteristic	Alcohol Use (2007)	Alcohol Use (2008)	Binge Alcohol Use (2007)	Binge Alcohol Use (2008)	Heavy Alcohol Use (2007)	Heavy Alcohol Use (2008)
TOTAL	0.4321		0.8638		0.6499	
GEOGRAPHIC DIVISION						
Northeast	0.7270		0.3193		0.2189	
New England	0.6251		0.9396		0.4440	
Middle Atlantic	0.5217		0.2392		0.0726	
Midwest	0.6821		0.6711		0.5443	
East North Central	0.5392		0.8824		0.9118	
West North Central	0.8062		0.5927		0.3809	
South	0.5375		0.5482		0.5390	
South Atlantic	0.8022		0.9127		0.9218	
East South Central	0.3544		0.8543		0.4290	
West South Central	0.8087		0.3817		0.5862	
West	0.4098		0.3854		0.1412	
Mountain	0.8351		0.3925		0.3130	
Pacific	0.3242		0.5656		0.2645	
COUNTY TYPE						
Large Metro	0.8724		0.2831		0.3651	
Small Metro	0.7867		0.2578		0.1774	
250K – 1 Mil. Pop.	0.7363		0.0551		0.0136	
< 250K Pop.	0.3768		0.5557		0.2472	
Nonmetro	0.1484		0.8641		0.2370	
Urbanized	0.1429		0.4617		0.9517	
Less Urbanized	0.3853		0.6055		0.2334	
Completely Rural	0.5613		0.8627		0.2319	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.