

**Table 6.10A (REVISED) Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Past Year Substance Dependence or Abuse:
Numbers in Thousands, 2008**

Substance	Total	Dependence	No Dependence	Dependence or Abuse	No Dependence or Abuse
ILLCIT DRUGS¹	8,331	718	7,613	888	7,442
Marijuana	8,331	301	8,030	451	7,880
ILLCIT DRUGS OTHER THAN MARIJUANA¹	8,331	506	7,825	584	7,747
ALCOHOL	8,331	1,069	7,262	1,533	6,798
BOTH ILLCIT DRUGS AND ALCOHOL¹	8,331	268	8,063	469	7,862
ILLCIT DRUGS OR ALCOHOL¹	8,331	1,519	6,812	1,952	6,379

*Low precision; no estimate reported.

NOTE: Some 2008 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Estimates for SMI presented in this table may differ from previously published SMI estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates are based on data from the original questions not including methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2008.

Table 6.10B (REVISED) Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Past Year Substance Dependence or Abuse: Percentages, 2008

Substance	Total	Dependence	No Dependence	Dependence or Abuse	No Dependence or Abuse
ILLCIT DRUGS¹	3.7	17.0	3.5	15.2	3.4
Marijuana	3.7	13.4	3.6	13.3	3.6
ILLCIT DRUGS OTHER THAN MARIJUANA¹	3.7	21.9	3.5	18.4	3.5
ALCOHOL	3.7	12.8	3.4	8.9	3.3
BOTH ILLCIT DRUGS AND ALCOHOL¹	3.7	22.8	3.6	17.8	3.5
ILLCIT DRUGS OR ALCOHOL¹	3.7	13.3	3.2	9.5	3.1

*Low precision; no estimate reported.

NOTE: Some 2008 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Estimates for SMI presented in this table may differ from previously published SMI estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates are based on data from the original questions not including methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2008.

Table 6.11A (REVISED) Substance Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Serious Mental Illness: Numbers in Thousands, 2008

Dependence/Abuse	Total¹	Serious Mental Illness	No Serious Mental Illness
DEPENDENCE OR ABUSE²			
Illicit Drugs ³	5,857	888	4,968
Marijuana	3,382	451	2,930
Illicit Drugs Other Than Marijuana ³	3,170	584	2,585
Alcohol	17,258	1,533	15,725
Both Illicit Drugs and Alcohol ³	2,635	469	2,165
Illicit Drugs or Alcohol ³	20,480	1,952	18,528
DEPENDENCE²			
Illicit Drugs ³	4,235	718	3,516
Marijuana	2,246	301	1,945
Illicit Drugs Other Than Marijuana ³	2,309	506	1,803
Alcohol	8,356	1,069	7,286
Both Illicit Drugs and Alcohol ³	1,176	268	907
Illicit Drugs or Alcohol ³	11,414	1,519	9,895

*Low precision; no estimate reported.

NOTE: Some 2008 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Estimates for SMI presented in this table may differ from previously published SMI estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

¹ Estimates in the Total column represent persons aged 18 or older, including those with unknown past year SMI information.

² Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates are based on data from the original questions not including methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2008.

Table 6.11B (REVISED) Substance Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Serious Mental Illness: Percentages, 2008

Dependence/Abuse	Total¹	Serious Mental Illness	No Serious Mental Illness
DEPENDENCE OR ABUSE²			
Illicit Drugs ³	2.6	10.7	2.3
Marijuana	1.5	5.4	1.4
Illicit Drugs Other Than Marijuana ³	1.4	7.0	1.2
Alcohol	7.7	18.4	7.3
Both Illicit Drugs and Alcohol ³	1.2	5.6	1.0
Illicit Drugs or Alcohol ³	9.1	23.4	8.6
DEPENDENCE²			
Illicit Drugs ³	1.9	8.6	1.6
Marijuana	1.0	3.6	0.9
Illicit Drugs Other Than Marijuana ³	1.0	6.1	0.8
Alcohol	3.7	12.8	3.4
Both Illicit Drugs and Alcohol ³	0.5	3.2	0.4
Illicit Drugs or Alcohol ³	5.1	18.2	4.6

*Low precision; no estimate reported.

NOTE: Some 2008 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Estimates for SMI presented in this table may differ from previously published SMI estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

¹ Estimates in the Total column represent persons aged 18 or older, including those with unknown past year SMI information.

² Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates are based on data from the original questions not including methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2008.

Table 6.12A (REVISED) Illicit Drug Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Serious Mental Illness, Gender, and Age Group: Numbers in Thousands, 2008

Gender/Age	Total ¹	Serious Mental Illness	No Serious Mental Illness
TOTAL	5,857	888	4,968
18-25	2,591	293	2,298
26-49	2,918	553	2,366
50 or Older	348	43	305
MALE	3,561	341	3,220
18-25	1,520	100	1,420
26-49	1,834	229	1,605
50 or Older	207	*	195
FEMALE	2,296	547	1,748
18-25	1,071	193	878
26-49	1,084	323	761
50 or Older	141	31	110

*Low precision; no estimate reported.

NOTE: Some 2008 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, based on data from original questions not including methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Estimates for SMI presented in this table may differ from previously published SMI estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

¹ Estimates in the Total column represent persons aged 18 or older, including those with unknown past year SMI information.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2008.

Table 6.12B (REVISED) Illicit Drug Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Serious Mental Illness, Gender, and Age Group: Percentages, 2008

Gender/Age	Total ¹	Serious Mental Illness	No Serious Mental Illness
TOTAL	2.6	10.7	2.3
18-25	7.9	23.7	7.3
26-49	2.9	11.6	2.5
50 or Older	0.4	1.8	0.3
MALE	3.3	11.9	3.0
18-25	9.2	26.8	8.8
26-49	3.7	13.2	3.4
50 or Older	0.5	*	0.5
FEMALE	2.0	10.0	1.6
18-25	6.5	22.3	5.7
26-49	2.1	10.7	1.6
50 or Older	0.3	2.0	0.2

*Low precision; no estimate reported.

NOTE: Some 2008 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, based on data from original questions not including methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Estimates for SMI presented in this table may differ from previously published SMI estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

¹ Estimates in the Total column represent persons aged 18 or older, including those with unknown past year SMI information.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2008.

Table 6.13A (REVISED) Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Serious Mental Illness, Gender, and Age Group: Numbers in Thousands, 2008

Gender/Age	Total ¹	Serious Mental Illness	No Serious Mental Illness
TOTAL	17,258	1,533	15,725
18-25	5,726	472	5,254
26-49	8,785	928	7,856
50 or Older	2,747	133	2,615
MALE	11,245	686	10,559
18-25	3,463	186	3,277
26-49	5,899	464	5,435
50 or Older	1,884	*	1,848
FEMALE	6,013	846	5,166
18-25	2,263	286	1,977
26-49	2,886	464	2,422
50 or Older	864	97	767

*Low precision; no estimate reported.

NOTE: Some 2008 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Estimates for SMI presented in this table may differ from previously published SMI estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

¹ Estimates in the Total column represent persons aged 18 or older, including those with unknown past year SMI information.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2008.

Table 6.13B (REVISED) Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Serious Mental Illness, Gender, and Age Group: Percentages, 2008

Gender/Age	Total ¹	Serious Mental Illness	No Serious Mental Illness
TOTAL	7.7	18.4	7.3
18-25	17.4	38.1	16.6
26-49	8.8	19.5	8.3
50 or Older	3.0	5.7	2.9
MALE	10.4	23.9	10.0
18-25	20.9	49.7	20.2
26-49	11.9	26.8	11.4
50 or Older	4.4	*	4.4
FEMALE	5.2	15.5	4.7
18-25	13.8	33.1	12.7
26-49	5.7	15.3	5.1
50 or Older	1.7	6.2	1.6

*Low precision; no estimate reported.

NOTE: Some 2008 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Estimates for SMI presented in this table may differ from previously published SMI estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

¹ Estimates in the Total column represent persons aged 18 or older, including those with unknown past year SMI information.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2008.