

Table 6.18C (REVISED) Location of Outpatient Mental Health Treatment/Counseling among Persons Aged 18 or Older with Past Year Serious Mental Illness Who Received Outpatient Mental Health Treatment in the Past Year, by Age Group: Standard Errors of Numbers in Thousands, 2008

Location of Treatment/Counseling ¹	Total	Aged 18-25	Aged 26-49	Aged 50+
Outpatient Mental Health Clinic or Center	105	21	73	*
Office of a Private Therapist, Psychologist, Psychiatrist, Social Worker, or Counselor - Not Part of a Clinic	167	24	112	*
Doctor's Office - Not Part of a Clinic	105	11	79	*
Outpatient Medical Clinic	65	9	49	*
Partial Day Hospital or Day Treatment Program	30	8	25	*
School or University Setting/Clinic/Center ²	10	7	7	*
Some Other Place ³	17	2	4	*

*Low precision; no estimate reported.

NOTE: Some 2008 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Outpatient Mental Health Treatment/Counseling is defined as having received outpatient care for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information were excluded. Estimates were based only on responses to items in the Adult Mental Health Service Utilization module.

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Estimates for SMI presented in this table may differ from previously published SMI estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

¹ Respondents could indicate multiple locations; thus, these response categories are not mutually exclusive.

² Respondents were permitted to specify other locations for receiving outpatient mental health treatment/counseling. This location is one of the most commonly reported other locations for receiving outpatient treatment/counseling.

³ Respondents with unknown or invalid responses to the other-specify question on Some Other Place Received Outpatient Mental Health Treatment/Counseling were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2008.

Table 6.18D (REVISED) Location of Outpatient Mental Health Treatment/Counseling among Persons Aged 18 or Older with Past Year Serious Mental Illness Who Received Outpatient Mental Health Treatment in the Past Year, by Age Group: Standard Errors of Percentages, 2008

Location of Treatment/Counseling ¹	Total	Aged 18-25	Aged 26-49	Aged 50+
Outpatient Mental Health Clinic or Center	2.52	3.74	2.85	*
Office of a Private Therapist, Psychologist, Psychiatrist, Social Worker, or Counselor - Not Part of a Clinic	2.66	3.77	3.23	*
Doctor's Office - Not Part of a Clinic	2.45	2.55	2.95	*
Outpatient Medical Clinic	1.64	2.01	2.06	*
Partial Day Hospital or Day Treatment Program	0.80	1.92	1.11	*
School or University Setting/Clinic/Center ²	0.26	1.69	0.29	*
Some Other Place ³	0.45	0.57	0.16	*

*Low precision; no estimate reported.

NOTE: Some 2008 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Outpatient Mental Health Treatment/Counseling is defined as having received outpatient care for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information were excluded. Estimates were based only on responses to items in the Adult Mental Health Service Utilization module.

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Estimates for SMI presented in this table may differ from previously published SMI estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

¹ Respondents could indicate multiple locations; thus, these response categories are not mutually exclusive.

² Respondents were permitted to specify other locations for receiving outpatient mental health treatment/counseling. This location is one of the most commonly reported other locations for receiving outpatient treatment/counseling.

³ Respondents with unknown or invalid responses to the other-specify question on Some Other Place Received Outpatient Mental Health Treatment/Counseling were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2008.