

Table 6.14C (REVISED) Received Mental Health Treatment/Counseling in the Past Year among Persons Aged 18 or Older, by Past Year Serious Mental Illness and Demographic Characteristics: Standard Errors of Numbers in Thousands, 2008

Demographic Characteristic	Total¹	Serious Mental Illness	No Serious Mental Illness
TOTAL	646	262	669
AGE			
18-25	92	37	98
26-49	340	177	320
50 or Older	523	192	547
GENDER			
Male	374	144	373
Female	503	217	515
HISPANIC ORIGIN AND RACE			
Not Hispanic or Latino	616	251	645
White	658	234	600
Black or African American	204	59	196
American Indian or Alaska Native	32	*	30
Native Hawaiian or Other Pacific Islander	*	*	*
Asian	96	*	71
Two or More Races	96	*	92
Hispanic or Latino	167	*	161
EDUCATION			
< High School	290	75	276
High School Graduate	368	139	340
Some College	330	136	295
College Graduate	399	141	373
CURRENT EMPLOYMENT			
Full-Time	447	156	409
Part-Time	257	98	240
Unemployed	117	*	99
Other ²	488	184	437

*Low precision; no estimate reported.

NOTE: Some 2008 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Mental Health Treatment/Counseling is defined as having received inpatient care or outpatient care or having used prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information were excluded. Estimates were based only on responses to items in the Adult Mental Health Service Utilization module.

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Estimates for SMI presented in this table may differ from previously published SMI estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

¹ Estimates in the Total column represent persons aged 18 or older, including those with unknown past year SMI information.

² The Other Employment category includes retired persons, disabled persons, homemakers, students, or other persons not in the labor force.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2008.

Table 6.14D (REVISED) Received Mental Health Treatment/Counseling in the Past Year among Persons Aged 18 or Older, by Past Year Serious Mental Illness and Demographic Characteristics: Standard Errors of Percentages, 2008

Demographic Characteristic	Total¹	Serious Mental Illness	No Serious Mental Illness
TOTAL	0.29	1.76	0.28
AGE			
18-25	0.28	2.28	0.27
26-49	0.34	2.08	0.31
50 or Older	0.57	4.33	0.56
GENDER			
Male	0.35	3.06	0.33
Female	0.43	2.14	0.43
HISPANIC ORIGIN AND RACE			
Not Hispanic or Latino	0.32	1.84	0.31
White	0.37	1.99	0.36
Black or African American	0.73	5.68	0.73
American Indian or Alaska Native	3.11	*	3.05
Native Hawaiian or Other Pacific Islander	*	*	*
Asian	0.94	*	0.72
Two or More Races	3.42	*	3.50
Hispanic or Latino	0.55	*	0.53
EDUCATION			
< High School	0.77	4.61	0.77
High School Graduate	0.50	3.19	0.48
Some College	0.53	2.99	0.50
College Graduate	0.57	3.06	0.55
CURRENT EMPLOYMENT			
Full-Time	0.33	2.71	0.31
Part-Time	0.77	3.97	0.75
Unemployed	1.23	*	1.13
Other ²	0.67	3.19	0.65

*Low precision; no estimate reported.

NOTE: Some 2008 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Mental Health Treatment/Counseling is defined as having received inpatient care or outpatient care or having used prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information were excluded. Estimates were based only on responses to items in the Adult Mental Health Service Utilization module.

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Estimates for SMI presented in this table may differ from previously published SMI estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

¹ Estimates in the Total column represent persons aged 18 or older, including those with unknown past year SMI information.

² The Other Employment category includes retired persons, disabled persons, homemakers, students, or other persons not in the labor force.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2008.

Table 6.15C (REVISED) Received Mental Health Treatment/Counseling in the Past Year among Persons Aged 18 or Older, by Past Year Serious Mental Illness and Geographic Characteristics: Standard Errors of Numbers in Thousands, 2008

Geographic Characteristic	Total¹	Serious Mental Illness	No Serious Mental Illness
TOTAL	646	262	669
GEOGRAPHIC REGION			
Northeast	293	117	288
New England	122	*	120
Middle Atlantic	266	107	262
Midwest	240	109	246
East North Central	198	86	195
West North Central	142	*	152
South	392	150	408
South Atlantic	311	114	331
East South Central	146	*	151
West South Central	180	82	177
West	354	134	390
Mountain	160	68	175
Pacific	316	115	349
COUNTY TYPE			
Large Metro	571	187	522
Small Metro	404	149	375
250K – 1 Mil. Pop.	349	126	323
< 250K Pop.	219	81	205
Nonmetro	294	104	272
Urbanized	199	*	173
Less Urbanized	212	65	204
Completely Rural	70	*	66

*Low precision; no estimate reported.

NOTE: Some 2008 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Mental Health Treatment/Counseling is defined as having received inpatient care or outpatient care or having used prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information were excluded. Estimates were based only on responses to items in the Adult Mental Health Service Utilization module.

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Estimates for SMI presented in this table may differ from previously published SMI estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

¹ Estimates in the Total column represent persons aged 18 or older, including those with unknown past year SMI information.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2008.

Table 6.15D (REVISED) Received Mental Health Treatment/Counseling in the Past Year among Persons Aged 18 or Older, by Past Year Serious Mental Illness and Geographic Characteristics: Standard Errors of Percentages, 2008

Geographic Characteristic	Total¹	Serious Mental Illness	No Serious Mental Illness
TOTAL	0.29	1.76	0.28
GEOGRAPHIC REGION			
Northeast	0.70	3.91	0.68
New England	1.12	*	1.06
Middle Atlantic	0.87	4.78	0.84
Midwest	0.49	2.96	0.48
East North Central	0.58	3.37	0.54
West North Central	0.95	*	0.97
South	0.48	3.26	0.47
South Atlantic	0.72	4.41	0.71
East South Central	1.09	*	1.11
West South Central	0.72	5.73	0.67
West	0.68	3.84	0.67
Mountain	1.01	5.61	0.99
Pacific	0.88	5.18	0.87
COUNTY TYPE			
Large Metro	0.41	2.55	0.40
Small Metro	0.52	3.09	0.51
250K – 1 Mil. Pop.	0.65	3.44	0.64
< 250K Pop.	0.84	5.76	0.82
Nonmetro	0.70	3.95	0.69
Urbanized	1.09	*	1.02
Less Urbanized	1.06	5.44	1.07
Completely Rural	1.57	*	1.61

*Low precision; no estimate reported.

NOTE: Some 2008 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Mental Health Treatment/Counseling is defined as having received inpatient care or outpatient care or having used prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information were excluded. Estimates were based only on responses to items in the Adult Mental Health Service Utilization module.

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Estimates for SMI presented in this table may differ from previously published SMI estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

¹ Estimates in the Total column represent persons aged 18 or older, including those with unknown past year SMI information.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2008.