

Table 6.8C (REVISED) Types of Illicit Drug Use in the Past Year among Persons Aged 18 or Older, by Past Year Serious Mental Illness: Standard Errors of Numbers in Thousands, 2008

Drug	Total¹	Serious Mental Illness	No Serious Mental Illness
ILLCIT DRUGS²	545	134	569
Marijuana and Hashish	466	116	481
Cocaine	207	61	193
Crack	94	36	88
Heroin	58	21	53
Hallucinogens	124	39	116
LSD	47	18	42
PCP	10	2	10
Ecstasy	92	33	85
Inhalants	108	15	108
Nonmedical Use of Psychotherapeutics ^{3,4}	345	102	335
Pain Relievers	288	87	276
OxyContin [®]	87	24	84
Tranquilizers	199	66	187
Stimulants ⁴	120	46	112
Methamphetamine ⁴	75	29	70
Sedatives	59	30	49
ILLCIT DRUGS OTHER THAN MARIJUANA²	400	114	389

*Low precision; no estimate reported.

NOTE: Some 2008 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Estimates for SMI presented in this table may differ from previously published SMI estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

¹ Estimates in the Total column represent persons aged 18 or older, including those with unknown past year SMI information.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2008.

Table 6.8D (REVISED) Types of Illicit Drug Use in the Past Year among Persons Aged 18 or Older, by Past Year Serious Mental Illness: Standard Errors of Percentages, 2008

Drug	Total ¹	Serious Mental Illness	No Serious Mental Illness
ILLCIT DRUGS²	0.24	1.52	0.24
Marijuana and Hashish	0.21	1.36	0.21
Cocaine	0.09	0.73	0.09
Crack	0.04	0.44	0.04
Heroin	0.03	0.25	0.02
Hallucinogens	0.06	0.46	0.05
LSD	0.02	0.22	0.02
PCP	0.00	0.03	0.00
Ecstasy	0.04	0.39	0.04
Inhalants	0.05	0.18	0.05
Nonmedical Use of Psychotherapeutics ^{3,4}	0.15	1.19	0.15
Pain Relievers	0.13	1.02	0.13
OxyContin [®]	0.04	0.28	0.04
Tranquilizers	0.09	0.78	0.09
Stimulants ⁴	0.05	0.56	0.05
Methamphetamine ⁴	0.03	0.35	0.03
Sedatives	0.03	0.36	0.02
ILLCIT DRUGS OTHER THAN MARIJUANA²	0.18	1.30	0.18

*Low precision; no estimate reported.

NOTE: Some 2008 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Estimates for SMI presented in this table may differ from previously published SMI estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

¹ Estimates in the Total column represent persons aged 18 or older, including those with unknown past year SMI information.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2008.

Table 6.9C (REVISED) Tobacco Product and Alcohol Use in the Past Year and Past Month among Persons Aged 18 or Older, by Past Year Serious Mental Illness: Standard Errors of Numbers in Thousands, 2008

Substance	Total ¹	Serious Mental Illness	No Serious Mental Illness
PAST YEAR			
Tobacco Products ²	923	228	1,044
Cigarettes	855	218	926
Smokeless Tobacco	345	83	343
Cigars	488	94	512
Alcohol	950	256	1,820
PAST MONTH			
Tobacco Products ²	872	220	953
Cigarettes	809	212	856
Smokeless Tobacco	311	60	308
Cigars	363	70	363
Pipe Tobacco	146	33	140
Alcohol	977	231	1,604
Binge Alcohol Use ³	737	144	867
Heavy Alcohol Use ³	414	101	414

*Low precision; no estimate reported.

NOTE: Some 2008 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Estimates for SMI presented in this table may differ from previously published SMI estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

¹ Estimates in the Total column represent persons aged 18 or older, including those with unknown past year SMI information.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2008.

Table 6.9D (REVISED) Tobacco Product and Alcohol Use in the Past Year and Past Month among Persons Aged 18 or Older, by Past Year Serious Mental Illness: Standard Errors of Percentages, 2008

Substance	Total ¹	Serious Mental Illness	No Serious Mental Illness
PAST YEAR			
Tobacco Products ²	0.41	1.96	0.41
Cigarettes	0.38	1.93	0.38
Smokeless Tobacco	0.15	0.97	0.16
Cigars	0.22	1.11	0.22
Alcohol	0.42	1.71	0.43
PAST MONTH			
Tobacco Products ²	0.39	1.91	0.39
Cigarettes	0.36	1.90	0.36
Smokeless Tobacco	0.14	0.71	0.14
Cigars	0.16	0.82	0.16
Pipe Tobacco	0.06	0.39	0.06
Alcohol	0.43	1.97	0.44
Binge Alcohol Use ³	0.33	1.57	0.34
Heavy Alcohol Use ³	0.18	1.17	0.19

*Low precision; no estimate reported.

NOTE: Some 2008 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Estimates for SMI presented in this table may differ from previously published SMI estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

¹ Estimates in the Total column represent persons aged 18 or older, including those with unknown past year SMI information.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2008.