

Table 7.28A Tobacco Product Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2007 and 2008

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2007	2008	2007	2008	2007	2008	2007	2008
TOBACCO PRODUCTS²	11,554	11,899	20,350	19,707	16,102	16,793	22,933	22,469
Cigarettes	9,877	10,032	17,154	16,643	13,135	13,615	19,903	19,490
Smokeless Tobacco	2,099	2,163	2,291	2,401	1,408	1,564	2,253	2,542
Cigars	2,983	3,073	4,617	4,454	2,964	3,235	2,700	2,365
Pipe Tobacco	396	343	520	510	533	435	597	588

*Low precision; no estimate reported.

^a Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 7.28B Tobacco Product Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Percentages, 2007 and 2008

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2007	2008	2007	2008	2007	2008	2007	2008
TOBACCO PRODUCTS²	67.9	68.8	49.9	48.3	23.3	23.7	18.9	18.6
Cigarettes	58.1	58.0	42.1	40.8	19.0	19.2	16.4	16.1
Smokeless Tobacco	12.3	12.5	5.6	5.9	2.0	2.2	1.9	2.1
Cigars	17.5	17.8	11.3	10.9	4.3	4.6	2.2	2.0
Pipe Tobacco	2.3	2.0	1.3	1.2	0.8	0.6	0.5	0.5

*Low precision; no estimate reported.

^a Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 7.29A Tobacco Product Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2007 and 2008

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2007	2008	2007	2008	2007	2008	2007	2008
TOBACCO PRODUCTS²	490 ^b	378	977	890	380	363	1,281	1,213
Cigarettes	422 ^b	319	798	739	291	271	970	931
Smokeless Tobacco	129	93	183	160	56	38	248	256
Cigars	216	177	357	322	105	136	393 ^a	309
Pipe Tobacco	49	38	55	80	22	16	55	43

*Low precision; no estimate reported.

^a Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 7.29B Tobacco Product Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Percentages, 2007 and 2008

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2007	2008	2007	2008	2007	2008	2007	2008
TOBACCO PRODUCTS²	82.9	77.0	52.6	52.4	24.2	25.1	6.0	5.7
Cigarettes	71.5	65.0	43.0	43.5	18.5	18.7	4.6	4.4
Smokeless Tobacco	21.8	19.0	9.9	9.4	3.6	2.6	1.2	1.2
Cigars	36.5	36.1	19.2	18.9	6.7	9.4	1.8 ^a	1.5
Pipe Tobacco	8.4	7.6	3.0 ^a	4.7	1.4	1.1	0.3	0.2

*Low precision; no estimate reported.

^a Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 7.30A Tobacco Product Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2007 and 2008

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2007	2008	2007	2008	2007	2008	2007	2008
TOBACCO PRODUCTS²	3,643	3,544	5,004	5,041	2,205	2,234	2,841	2,810
Cigarettes	3,140	3,080	4,307	4,408	1,915	1,837	2,474	2,433
Smokeless Tobacco	758	694	571	632	116 ^a	176	276	270
Cigars	1,260	1,184	1,477	1,351	528	613	608	573
Pipe Tobacco	119	145	168	151	53	70	54 ^a	91

*Low precision; no estimate reported.

^a Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 7.30B Tobacco Product Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Percentages, 2007 and 2008

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2007	2008	2007	2008	2007	2008	2007	2008
TOBACCO PRODUCTS²	75.6	74.5	56.5	57.6	34.7	33.7	22.4	22.0
Cigarettes	65.1	64.7	48.6	50.4	30.1	27.7	19.5	19.0
Smokeless Tobacco	15.7	14.6	6.4	7.2	1.8 ^a	2.7	2.2	2.1
Cigars	26.1	24.9	16.7	15.4	8.3	9.2	4.8	4.5
Pipe Tobacco	2.5	3.0	1.9	1.7	0.8	1.0	0.4 ^a	0.7

*Low precision; no estimate reported.

^a Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 7.31A Tobacco Product Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2007 and 2008

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2007	2008	2007	2008	2007	2008	2007	2008
TOBACCO PRODUCTS²	7,420	7,976	14,369	13,776	13,517	14,196	18,811	18,446
Cigarettes	6,315	6,633	12,049	11,496	10,929	11,507	16,459	16,126
Smokeless Tobacco	1,212	1,376	1,537	1,609	1,235	1,350	1,729	2,016
Cigars	1,507	1,712	2,783	2,781	2,331	2,486	1,699	1,483
Pipe Tobacco	227	161	297	280	459	349	488	454

*Low precision; no estimate reported.

^a Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 7.31B Tobacco Product Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Percentages, 2007 and 2008

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2007	2008	2007	2008	2007	2008	2007	2008
TOBACCO PRODUCTS²	64.0	66.2	47.8	45.4	22.1	22.6	21.6	21.3
Cigarettes	54.4	55.1	40.1	37.9	17.9	18.3	18.9	18.6
Smokeless Tobacco	10.4	11.4	5.1	5.3	2.0	2.1	2.0	2.3
Cigars	13.0	14.2	9.3	9.2	3.8	4.0	1.9	1.7
Pipe Tobacco	2.0	1.3	1.0	0.9	0.8	0.6	0.6	0.5

*Low precision; no estimate reported.

^a Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.