

Table 7.11C Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2007 and 2008

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2007	2008	2007	2008	2007	2008	2007	2008
ILLCIT DRUGS²	219	195	232	227	203	231	195	200
Marijuana and Hashish	192	180	188	203	166	206	140	155
Cocaine	95	86	75	73	53	40	58	28
Crack	42	35	32	31	37	22	47	19
Heroin	12	22	7	27	28	16	18	10
Hallucinogens	38	44	45	37	32	23	30	27
LSD	10	12	26	11	17	4	7	9
PCP	2	4	8	6	2	2	9	*
Ecstasy	30	36	31	32	30	13	14	12
Inhalants	45	13	28	46	22	46	35	27
Nonmedical Use of Psychotherapeutics ^{3,4}	110	90	123	117	131	93	139	135
Pain Relievers	100	84	87	103	109	78	126	112
OxyContin [®]	17	26	24	32	10	18	43	36
Tranquilizers	55	44	62	66	66	48	70	71
Stimulants ⁴	48	36	49	42	37	27	55	40
Methamphetamine ⁴	42	17	40	32	28	11	48	35
Sedatives	20	18	51	18	23	15	33	23
ILLCIT DRUGS OTHER THAN MARIJUANA²	141	119	148	142	143	111	152	139

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 7.11D Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2007 and 2008

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2007	2008	2007	2008	2007	2008	2007	2008
ILLCIT DRUGS²	1.03	1.03	0.53	0.53	0.29	0.32	0.16	0.16
Marijuana and Hashish	0.97	0.97	0.45	0.48	0.24	0.28	0.11	0.13
Cocaine	0.54	0.49	0.18	0.18	0.08	0.06	0.05	0.02
Crack	0.25	0.20	0.08	0.08	0.05	0.03	0.04	0.02
Heroin	0.07	0.13	0.02	0.07	0.04	0.02	0.01	0.01
Hallucinogens	0.22	0.26	0.11	0.09	0.05	0.03	0.02	0.02
LSD	0.06	0.07	0.06	0.03	0.02	0.01	0.01	0.01
PCP	0.01	0.02	0.02	0.02	0.00	0.00	0.01	*
Ecstasy	0.17	0.21	0.08	0.08	0.04	0.02	0.01	0.01
Inhalants	0.26	0.08	0.07	0.11	0.03	0.06	0.03	0.02
Nonmedical Use of Psychotherapeutics ^{3,4}	0.61	0.52	0.29	0.28	0.19	0.13	0.12	0.11
Pain Relievers	0.56	0.48	0.21	0.25	0.16	0.11	0.10	0.09
OxyContin [®]	0.10	0.15	0.06	0.08	0.01	0.03	0.04	0.03
Tranquilizers	0.32	0.26	0.15	0.16	0.10	0.07	0.06	0.06
Stimulants ⁴	0.28	0.21	0.12	0.10	0.05	0.04	0.05	0.03
Methamphetamine ⁴	0.25	0.10	0.10	0.08	0.04	0.02	0.04	0.03
Sedatives	0.12	0.10	0.13	0.04	0.03	0.02	0.03	0.02
ILLCIT DRUGS OTHER THAN MARIJUANA²	0.75	0.67	0.35	0.34	0.21	0.16	0.13	0.11

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 7.11P Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: *P* Values from Tests of Differences of Percentages, 2008 Versus 2007

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2007	2008	2007	2008	2007	2008	2007	2008
ILLCIT DRUGS²	0.1844		0.6155		0.1613		0.4308	
Marijuana and Hashish	0.5811		0.3480		0.0631		1.0000	
Cocaine	0.6454		0.6340		0.2189		0.0174	
Crack	0.2492		0.8056		0.1650		0.0164	
Heroin	0.0352		0.0361		0.2393		0.4782	
Hallucinogens	0.2079		0.9503		0.8748		0.8485	
LSD	0.3199		0.7816		0.5668		0.3832	
PCP	0.2879		0.5607		0.9651		*	
Ecstasy	0.1490		0.5335		0.2050		0.7663	
Inhalants	0.1124		0.2451		0.2831		0.6178	
Nonmedical Use of Psychotherapeutics ^{3,4}	0.0648		0.8541		0.1091		0.3313	
Pain Relievers	0.0614		0.5392		0.2393		0.3478	
OxyContin [®]	0.8238		0.5937		0.1976		0.8818	
Tranquilizers	0.0956		0.3299		0.7870		0.9856	
Stimulants ⁴	0.7607		0.9470		0.0440		0.5216	
Methamphetamine ⁴	0.0835		0.4474		0.0140		0.6662	
Sedatives	0.4071		0.3733		0.5360		0.5045	
ILLCIT DRUGS OTHER THAN MARIJUANA²	0.1087		0.9400		0.2025		0.1415	

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 7.12C Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2007 and 2008

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2007	2008	2007	2008	2007	2008	2007	2008
ILLICIT DRUGS²	27	26	37	35	24	27	51	41
Marijuana and Hashish	25	24	34	32	21	25	34	31
Cocaine	7	9	7	7	4	4	7	5
Crack	3	1	3	2	1	*	3	*
Heroin	1	3	*	5	1	*	*	3
Hallucinogens	9	11	11	14	5	8	8	10
LSD	3	4	5	7	2	4	4	6
PCP	1	3	3	1	1	2	3	1
Ecstasy	6	8	7	10	3	2	5	9
Inhalants	9	8	10	12	7	7	21	18
Nonmedical Use of Psychotherapeutics ^{3,4}	16	15	19	17	14	9	29	24
Pain Relievers	16	13	16	15	13	7	25	23
OxyContin [®]	5	4	4	5	3	1	4	5
Tranquilizers	10	7	10	9	5	4	11	10
Stimulants ⁴	7	8	7	7	3	4	9	6
Methamphetamine ⁴	5	4	3	4	1	*	5	2
Sedatives	2	5	3	3	1	2	8	6
ILLICIT DRUGS OTHER THAN MARIJUANA²	18	19	22	23	15	14	38	30

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 7.12D Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2007 and 2008

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2007	2008	2007	2008	2007	2008	2007	2008
ILLCIT DRUGS²	2.85	2.76	1.53	1.63	1.40	1.57	0.23	0.18
Marijuana and Hashish	2.87	2.98	1.49	1.57	1.22	1.50	0.16	0.14
Cocaine	1.20	1.68	0.39	0.43	0.23	0.26	0.03	0.02
Crack	0.56	0.28	0.15	0.13	0.09	*	0.01	*
Heroin	0.16	0.51	*	0.31	0.09	*	*	0.01
Hallucinogens	1.50	2.12	0.58	0.78	0.32	0.53	0.04	0.05
LSD	0.53	0.75	0.28	0.38	0.13	0.24	0.02	0.03
PCP	0.12	0.54	0.17	0.09	0.09	0.16	0.01	0.01
Ecstasy	1.01	1.57	0.38	0.56	0.21	0.13	0.02	0.04
Inhalants	1.49	1.48	0.52	0.68	0.42	0.51	0.10	0.08
Nonmedical Use of Psychotherapeutics ^{3,4}	2.41	2.67	0.94	0.96	0.84	0.60	0.14	0.11
Pain Relievers	2.33	2.31	0.82	0.87	0.83	0.51	0.12	0.10
OxyContin [®]	0.88	0.80	0.21	0.30	0.18	0.07	0.02	0.02
Tranquilizers	1.64	1.48	0.53	0.55	0.30	0.28	0.05	0.05
Stimulants ⁴	1.08	1.55	0.40	0.41	0.19	0.27	0.04	0.03
Methamphetamine ⁴	0.79	0.88	0.18	0.21	0.08	*	0.02	0.01
Sedatives	0.29	0.92	0.14	0.18	0.09	0.14	0.04	0.03
ILLCIT DRUGS OTHER THAN MARIJUANA²	2.58	2.96	1.07	1.25	0.92	0.90	0.17	0.13

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 7.12P Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: *P* Values from Tests of Differences of Percentages, 2008 Versus 2007

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2007	2008	2007	2008	2007	2008	2007	2008
ILLCIT DRUGS²	0.0376		0.0448		0.0540		0.0227	
Marijuana and Hashish	0.3194		0.0883		0.0185		0.1763	
Cocaine	0.1342		0.9801		0.9128		0.1875	
Crack	0.3731		0.8157		*		*	
Heroin	0.2308		*		*		*	
Hallucinogens	0.0750		0.0959		0.1695		0.1209	
LSD	0.2519		0.4072		0.4164		0.2851	
PCP	0.0565		0.6993		0.3302		0.6611	
Ecstasy	0.1205		0.0768		0.4691		0.1305	
Inhalants	0.5759		0.7861		0.6988		0.5822	
Nonmedical Use of Psychotherapeutics ^{3,4}	0.3375		0.6818		0.1047		0.0466	
Pain Relievers	0.9041		0.9180		0.0241		0.3467	
OxyContin [®]	0.7882		0.1809		0.0963		0.4265	
Tranquilizers	0.2731		0.9332		0.9171		0.6942	
Stimulants ⁴	0.1105		0.7418		0.7356		0.0222	
Methamphetamine ⁴	0.3696		0.3442		*		0.0618	
Sedatives	0.0994		0.4203		0.6289		0.2927	
ILLCIT DRUGS OTHER THAN MARIJUANA²	0.0320		0.2997		0.7320		0.0558	

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 7.13C Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2007 and 2008

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2007	2008	2007	2008	2007	2008	2007	2008
ILLCIT DRUGS²	90	91	91	85	46	55	45	45
Marijuana and Hashish	84	87	81	77	43	51	39	39
Cocaine	27	29	24	20	9	9	7	8
Crack	6	7	5	6	5	1	5	5
Heroin	3	10	7	6	6	2	4	6
Hallucinogens	25	28	21	23	8	13	9	10
LSD	9	11	6	8	1	3	6	4
PCP	2	3	2	0	*	*	5	*
Ecstasy	15	23	17	17	5	8	6	8
Inhalants	9	9	12	12	6	8	9	3
Nonmedical Use of Psychotherapeutics ^{3,4}	48	46	45	42	24	25	23	26
Pain Relievers	43	39	37	36	19	21	19	24
OxyContin [®]	10	10	12	8	7	6	5	7
Tranquilizers	28	24	20	23	14	16	13	11
Stimulants ⁴	19	21	20	18	13	6	10	9
Methamphetamine ⁴	11	9	10	7	10	2	6	7
Sedatives	9	6	6	7	2	9	6	5
ILLCIT DRUGS OTHER THAN MARIJUANA²	56	57	53	49	26	29	26	28

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 7.13D Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2007 and 2008

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2007	2008	2007	2008	2007	2008	2007	2008
ILLCIT DRUGS²	1.17	1.16	0.78	0.79	0.68	0.74	0.34	0.35
Marijuana and Hashish	1.17	1.21	0.72	0.75	0.64	0.70	0.30	0.31
Cocaine	0.54	0.59	0.26	0.23	0.15	0.14	0.05	0.07
Crack	0.13	0.16	0.06	0.06	0.08	0.01	0.04	0.04
Heroin	0.06	0.21	0.08	0.07	0.09	0.02	0.04	0.04
Hallucinogens	0.51	0.56	0.24	0.25	0.13	0.19	0.07	0.07
LSD	0.18	0.23	0.07	0.10	0.01	0.04	0.05	0.03
PCP	0.04	0.07	0.02	0.00	*	*	0.04	*
Ecstasy	0.31	0.47	0.19	0.20	0.09	0.12	0.05	0.06
Inhalants	0.20	0.20	0.14	0.14	0.09	0.12	0.07	0.03
Nonmedical Use of Psychotherapeutics ^{3,4}	0.87	0.84	0.47	0.45	0.37	0.36	0.18	0.20
Pain Relievers	0.80	0.74	0.40	0.40	0.30	0.30	0.15	0.19
OxyContin [®]	0.21	0.21	0.14	0.10	0.10	0.10	0.04	0.05
Tranquilizers	0.56	0.49	0.22	0.25	0.21	0.25	0.10	0.09
Stimulants ⁴	0.38	0.41	0.23	0.20	0.20	0.10	0.08	0.07
Methamphetamine ⁴	0.22	0.18	0.11	0.08	0.15	0.04	0.05	0.06
Sedatives	0.19	0.12	0.07	0.08	0.03	0.13	0.05	0.04
ILLCIT DRUGS OTHER THAN MARIJUANA²	0.98	0.98	0.54	0.51	0.39	0.42	0.20	0.22

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 7.13P Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: *P* Values from Tests of Differences of Percentages, 2008 Versus 2007

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2007	2008	2007	2008	2007	2008	2007	2008
ILLCIT DRUGS²	0.3254		0.5122		0.0517		0.6617	
Marijuana and Hashish	0.4631		0.6858		0.0328		0.6154	
Cocaine	0.7538		0.1475		0.3631		0.6238	
Crack	0.8346		0.8108		0.1457		0.9954	
Heroin	0.0039		0.7779		0.0786		0.5760	
Hallucinogens	0.2741		0.5367		0.2559		0.8599	
LSD	0.2840		0.2789		0.1477		0.6465	
PCP	0.9846		0.3748		*		*	
Ecstasy	0.0097		0.8085		0.7920		0.9230	
Inhalants	0.8719		0.7774		0.9943		0.0494	
Nonmedical Use of Psychotherapeutics ^{3,4}	0.8529		0.8144		0.9800		0.9431	
Pain Relievers	0.7898		0.7642		0.5746		0.9778	
OxyContin [®]	0.9705		0.1215		0.6955		0.5263	
Tranquilizers	0.5609		0.4980		0.7915		0.7620	
Stimulants ⁴	0.5430		0.5918		0.0478		0.7118	
Methamphetamine ⁴	0.1723		0.4354		0.0816		0.6627	
Sedatives	0.0495		0.8461		0.2548		0.7369	
ILLCIT DRUGS OTHER THAN MARIJUANA²	0.9619		0.4423		0.7781		0.8397	

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 7.14C Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2007 and 2008

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2007	2008	2007	2008	2007	2008	2007	2008
ILLCIT DRUGS²	191	167	206	208	196	220	184	190
Marijuana and Hashish	162	152	164	186	160	196	133	148
Cocaine	91	79	70	68	52	40	57	26
Crack	42	34	32	31	36	22	46	19
Heroin	12	19	*	26	27	16	17	8
Hallucinogens	27	36	38	26	31	18	28	23
LSD	3	2	24	*	16	*	*	5
PCP	*	*	7	6	*	*	6	*
Ecstasy	25	26	24	25	30	10	11	*
Inhalants	42	6	23	42	20	45	26	19
Nonmedical Use of Psychotherapeutics ^{3,4}	98	76	114	104	126	89	136	129
Pain Relievers	89	72	77	94	106	75	122	107
OxyContin [®]	12	24	21	31	7	17	43	36
Tranquilizers	49	36	60	61	65	45	69	70
Stimulants ⁴	43	28	43	36	34	26	53	39
Methamphetamine ⁴	41	14	39	30	26	10	47	34
Sedatives	18	16	51	16	23	12	32	22
ILLCIT DRUGS OTHER THAN MARIJUANA²	130	106	136	130	137	106	145	134

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 7.14D Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2007 and 2008

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2007	2008	2007	2008	2007	2008	2007	2008
ILLCIT DRUGS²	1.38	1.30	0.66	0.65	0.31	0.34	0.21	0.22
Marijuana and Hashish	1.24	1.19	0.54	0.59	0.26	0.30	0.15	0.17
Cocaine	0.76	0.65	0.23	0.22	0.08	0.06	0.07	0.03
Crack	0.36	0.28	0.11	0.10	0.06	0.04	0.05	0.02
Heroin	0.10	0.16	*	0.08	0.04	0.03	0.02	0.01
Hallucinogens	0.23	0.30	0.12	0.09	0.05	0.03	0.03	0.03
LSD	0.03	0.02	0.08	*	0.03	*	*	0.01
PCP	*	*	0.02	0.02	*	*	0.01	*
Ecstasy	0.22	0.22	0.08	0.08	0.05	0.02	0.01	*
Inhalants	0.36	0.05	0.08	0.14	0.03	0.07	0.03	0.02
Nonmedical Use of Psychotherapeutics ^{3,4}	0.81	0.63	0.37	0.34	0.20	0.14	0.16	0.15
Pain Relievers	0.74	0.60	0.25	0.30	0.17	0.12	0.14	0.12
OxyContin [®]	0.11	0.20	0.07	0.10	0.01	0.03	0.05	0.04
Tranquilizers	0.42	0.30	0.20	0.20	0.11	0.07	0.08	0.08
Stimulants ⁴	0.37	0.24	0.14	0.12	0.06	0.04	0.06	0.04
Methamphetamine ⁴	0.35	0.12	0.13	0.10	0.04	0.02	0.05	0.04
Sedatives	0.15	0.13	0.17	0.05	0.04	0.02	0.04	0.03
ILLCIT DRUGS OTHER THAN MARIJUANA²	1.05	0.86	0.44	0.42	0.22	0.17	0.17	0.15

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 7.14P Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: *P* Values from Tests of Differences of Percentages, 2008 Versus 2007

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2007	2008	2007	2008	2007	2008	2007	2008
ILLCIT DRUGS²	0.3292		0.4787		0.3795		0.6800	
Marijuana and Hashish	0.9399		0.2541		0.2306		0.8874	
Cocaine	0.6258		0.3311		0.2905		0.0193	
Crack	0.2358		0.7451		0.2339		0.0191	
Heroin	0.3855		*		0.4169		0.2760	
Hallucinogens	0.5675		0.3502		0.3371		0.3947	
LSD	0.5974		*		*		*	
PCP	*		0.7515		*		*	
Ecstasy	0.8292		0.8999		0.1872		*	
Inhalants	0.1736		0.1548		0.2736		0.7332	
Nonmedical Use of Psychotherapeutics ^{3,4}	0.0407		0.9506		0.1368		0.5364	
Pain Relievers	0.0834		0.3763		0.2841		0.4311	
OxyContin [®]	0.6204		0.3116		0.1299		0.8777	
Tranquilizers	0.2228		0.3960		0.7136		0.8970	
Stimulants ⁴	0.4254		0.7441		0.1335		0.8397	
Methamphetamine ⁴	0.1473		0.4881		0.0525		0.7389	
Sedatives	0.7466		0.3207		0.2817		0.6246	
ILLCIT DRUGS OTHER THAN MARIJUANA²	0.0832		0.7469		0.1898		0.3066	

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 7.15C Types of Illicit Drug Use in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Drinking Age Groups: Standard Errors of Numbers in Thousands, 2007 and 2008

Drug	Total (2007)	Total (2008)	Aged 12-20 (2007)	Aged 12-20 (2008)	Aged 21+ (2007)	Aged 21+ (2008)
ILLICIT DRUGS¹	403	393	114	105	377	371
Marijuana or Hashish	330	353	105	99	304	333
Cocaine (Including Crack)	134	127	26	27	131	124
Heroin	31	38	6	9	31	37
Hallucinogens	68	63	29	36	61	53
Inhalants	57	66	19	22	54	62
Pain Relievers ²	173	160	50	43	165	151
Tranquilizers ²	107	93	27	25	105	90
Stimulants ^{2,3}	77	64	21	21	75	59
Methamphetamine ³	65	41	11	7	64	39
Sedatives ²	60	30	7	6	60	29
ILLICIT DRUGS OTHER THAN MARIJUANA¹	266	224	65	63	255	214

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from the methamphetamine use items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated rows include data from methamphetamine use items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 7.15D Types of Illicit Drug Use in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Drinking Age Groups: Standard Errors of Percentages, 2007 and 2008

Drug	Total (2007)	Total (2008)	Aged 12-20 (2007)	Aged 12-20 (2008)	Aged 21+ (2007)	Aged 21+ (2008)
ILLICIT DRUGS¹	0.30	0.29	0.75	0.78	0.31	0.30
Marijuana or Hashish	0.25	0.26	0.72	0.76	0.26	0.27
Cocaine (Including Crack)	0.10	0.10	0.23	0.26	0.11	0.10
Heroin	0.02	0.03	0.05	0.09	0.03	0.03
Hallucinogens	0.05	0.05	0.26	0.34	0.05	0.04
Inhalants	0.04	0.05	0.18	0.21	0.05	0.05
Pain Relievers ²	0.13	0.12	0.43	0.41	0.14	0.13
Tranquilizers ²	0.08	0.07	0.25	0.25	0.09	0.08
Stimulants ^{2,3}	0.06	0.05	0.19	0.20	0.06	0.05
Methamphetamine ³	0.05	0.03	0.11	0.07	0.05	0.03
Sedatives ²	0.05	0.02	0.07	0.06	0.05	0.02
ILLICIT DRUGS OTHER THAN MARIJUANA¹	0.20	0.17	0.53	0.57	0.21	0.18

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from the methamphetamine use items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated rows include data from methamphetamine use items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 7.15P Types of Illicit Drug Use in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Drinking Age Groups: *P* Values from Tests of Differences of Percentages, 2008 Versus 2007

Drug	Total (2007)	Total (2008)	Aged 12-20 (2007)	Aged 12-20 (2008)	Aged 21+ (2007)	Aged 21+ (2008)
ILLICIT DRUGS¹	0.7437		0.0236		0.7981	
Marijuana or Hashish	0.2314		0.0184		0.2661	
Cocaine (Including Crack)	0.5900		0.9094		0.6490	
Heroin	0.1442		0.0911		0.2511	
Hallucinogens	0.5385		0.0055		0.6394	
Inhalants	0.6462		0.8503		0.5641	
Pain Relievers ²	0.1565		0.6308		0.2617	
Tranquilizers ²	0.6653		0.8725		0.7791	
Stimulants ^{2,3}	0.2479		0.5344		0.2051	
Methamphetamine ³	0.0117		0.2356		0.0190	
Sedatives ²	0.1874		0.9202		0.1941	
ILLICIT DRUGS OTHER THAN MARIJUANA¹	0.1179		0.3983		0.1224	

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from the methamphetamine use items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated rows include data from methamphetamine use items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 7.16C Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Age Drinking Groups: Standard Errors of Numbers in Thousands, 2007 and 2008

Drug¹	Total (2007)	Total (2008)	Aged 12-20 (2007)	Aged 12-20 (2008)	Aged 21+ (2007)	Aged 21+ (2008)
ILLCIT DRUGS²	230	252	68	68	217	240
Marijuana or Hashish	209	236	67	65	194	224
Cocaine (Including Crack)	76	66	12	14	75	64
Heroin	18	22	4	2	17	22
Hallucinogens	24	13	9	9	23	8
Inhalants	6	14	5	8	*	11
Pain Relievers ³	58	62	13	12	57	61
Tranquilizers ³	18	37	5	5	17	37
Stimulants ³	10	21	4	4	9	21
Methamphetamine ⁴	37	15	7	3	36	15
Sedatives ³	18	10	2	2	18	9
ILLCIT DRUGS OTHER THAN MARIJUANA²	103	99	22	22	102	96

*Low precision; no estimate reported.

¹ Estimates are based on nonimputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from the methamphetamine use items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Refers to nonmedical use; does not include over-the-counter drugs.

⁴ Estimates in the designated rows include data from methamphetamine use items added in 2005 and 2006.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 7.16D Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Age Drinking Groups: Standard Errors of Percentages, 2007 and 2008

Drug¹	Total (2007)	Total (2008)	Aged 12-20 (2007)	Aged 12-20 (2008)	Aged 21+ (2007)	Aged 21+ (2008)
ILLCIT DRUGS²	0.18	0.19	0.55	0.60	0.19	0.20
Marijuana or Hashish	0.16	0.18	0.55	0.58	0.17	0.18
Cocaine (Including Crack)	0.06	0.05	0.12	0.14	0.06	0.05
Heroin	0.01	0.02	0.03	0.02	0.01	0.02
Hallucinogens	0.02	0.01	0.08	0.09	0.02	0.01
Inhalants	0.00	0.01	0.05	0.08	*	0.01
Pain Relievers ³	0.05	0.05	0.12	0.12	0.05	0.05
Tranquilizers ³	0.01	0.03	0.05	0.05	0.01	0.03
Stimulants ³	0.01	0.02	0.04	0.04	0.01	0.02
Methamphetamine ⁴	0.03	0.01	0.06	0.03	0.03	0.01
Sedatives ³	0.01	0.01	0.01	0.02	0.02	0.01
ILLCIT DRUGS OTHER THAN MARIJUANA²	0.08	0.08	0.20	0.21	0.09	0.08

*Low precision; no estimate reported.

¹ Estimates are based on nonimputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from the methamphetamine use items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Refers to nonmedical use; does not include over-the-counter drugs.

⁴ Estimates in the designated rows include data from methamphetamine use items added in 2005 and 2006.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 7.16P Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Age Drinking Groups: *P* Values from Tests of Differences of Percentages, 2008 Versus 2007

Drug¹	Total (2007)	Total (2008)	Aged 12-20 (2007)	Aged 12-20 (2008)	Aged 21+ (2007)	Aged 21+ (2008)
ILLCIT DRUGS²	0.6910		0.1531		0.7083	
Marijuana or Hashish	0.3064		0.1321		0.3047	
Cocaine (Including Crack)	0.4243		0.4036		0.3453	
Heroin	0.6131		0.4581		0.5311	
Hallucinogens	0.1772		0.6251		0.0969	
Inhalants	0.3296		0.3584		*	
Pain Relievers ³	0.7556		0.9900		0.7195	
Tranquilizers ³	0.0507		0.4877		0.0343	
Stimulants ³	0.4504		0.7958		0.3846	
Methamphetamine ⁴	0.0500		0.4954		0.0630	
Sedatives ³	0.0932		0.9842		0.0919	
ILLCIT DRUGS OTHER THAN MARIJUANA²	0.4485		0.7892		0.4407	

*Low precision; no estimate reported.

¹ Estimates are based on nonimputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from the methamphetamine use items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Refers to nonmedical use; does not include over-the-counter drugs.

⁴ Estimates in the designated rows include data from methamphetamine use items added in 2005 and 2006.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 7.17C Types of Illicit Drugs Used in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, 2007 and 2008

Drug	Total (2007)	Total (2008)	Aged 12-17 (2007)	Aged 12-17 (2008)	Aged 18-25 (2007)	Aged 18-25 (2008)	Aged 26+ (2007)	Aged 26+ (2008)
ILLCIT DRUGS¹	403	393	52	53	153	149	347	354
Marijuana or Hashish	330	353	47	49	140	138	279	315
Cocaine (Including Crack)	134	127	11	12	38	38	127	119
Heroin	31	38	2	6	10	12	30	36
Hallucinogens	68	63	15	19	34	38	56	48
Inhalants	57	66	15	16	17	18	52	62
Pain Relievers ²	173	160	25	21	62	59	157	142
Tranquilizers ²	107	93	16	13	37	38	101	83
Stimulants ^{2,3}	77	64	10	13	30	28	70	54
Methamphetamine ³	65	41	6	6	18	12	62	37
Sedatives ²	60	30	3	6	12	14	59	26
ILLCIT DRUGS OTHER THAN MARIJUANA¹	266	224	32	33	87	84	243	202

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from the methamphetamine use items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated rows include data from methamphetamine use items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 7.17D Types of Illicit Drugs Used in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, 2007 and 2008

Drug	Total (2007)	Total (2008)	Aged 12-17 (2007)	Aged 12-17 (2008)	Aged 18-25 (2007)	Aged 18-25 (2008)	Aged 26+ (2007)	Aged 26+ (2008)
ILLICIT DRUGS¹	0.30	0.29	1.03	1.11	0.53	0.55	0.32	0.32
Marijuana or Hashish	0.25	0.26	0.98	1.07	0.52	0.53	0.27	0.29
Cocaine (Including Crack)	0.10	0.10	0.26	0.33	0.18	0.19	0.12	0.11
Heroin	0.02	0.03	0.04	0.16	0.05	0.06	0.03	0.03
Hallucinogens	0.05	0.05	0.37	0.50	0.17	0.19	0.05	0.05
Inhalants	0.04	0.05	0.37	0.43	0.08	0.09	0.05	0.06
Pain Relievers ²	0.13	0.12	0.59	0.57	0.28	0.28	0.15	0.13
Tranquilizers ²	0.08	0.07	0.39	0.35	0.18	0.19	0.10	0.08
Stimulants ^{2,3}	0.06	0.05	0.26	0.34	0.15	0.14	0.07	0.05
Methamphetamine ³	0.05	0.03	0.15	0.16	0.09	0.06	0.06	0.04
Sedatives ²	0.05	0.02	0.09	0.16	0.06	0.07	0.06	0.02
ILLICIT DRUGS OTHER THAN MARIJUANA¹	0.20	0.17	0.73	0.83	0.38	0.37	0.23	0.19

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from the methamphetamine use items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated rows include data from methamphetamine use items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 7.17P Types of Illicit Drugs Used in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: *P* Values from Tests of Differences of Percentages, 2008 Versus 2007

Drug	Total (2007)	Total (2008)	Aged 12-17 (2007)	Aged 12-17 (2008)	Aged 18-25 (2007)	Aged 18-25 (2008)	Aged 26+ (2007)	Aged 26+ (2008)
ILLCIT DRUGS¹	0.7437		0.0032		0.6647		0.6277	
Marijuana or Hashish	0.2314		0.0110		0.9383		0.1711	
Cocaine (Including Crack)	0.5900		0.3928		0.1478		0.8644	
Heroin	0.1442		0.1726		0.1776		0.3583	
Hallucinogens	0.5385		0.0066		0.1648		0.4832	
Inhalants	0.6462		0.9815		0.7412		0.4459	
Pain Relievers ²	0.1565		0.2175		0.7447		0.2915	
Tranquilizers ²	0.6653		0.4335		1.0000		0.8077	
Stimulants ^{2,3}	0.2479		0.2547		0.9594		0.1713	
Methamphetamine ³	0.0117		0.3469		0.0252		0.0375	
Sedatives ²	0.1874		0.0714		0.5711		0.1668	
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.1179		0.1571		0.5309		0.1521	

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from the methamphetamine use items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated rows include data from methamphetamine use items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 7.18C Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, 2007 and 2008

Drug¹	Total (2007)	Total (2008)	Aged 12-17 (2007)	Aged 12-17 (2008)	Aged 18-25 (2007)	Aged 18-25 (2008)	Aged 26+ (2007)	Aged 26+ (2008)
ILLCIT DRUGS²	230	252	32	33	94	95	200	229
Marijuana or Hashish	209	236	31	32	91	91	178	212
Cocaine (Including Crack)	76	66	5	6	17	20	74	63
Heroin	18	22	2	2	7	5	16	21
Hallucinogens	24	13	6	5	11	10	21	6
Inhalants	6	14	4	5	4	7	*	11
Pain Relievers ³	58	62	8	7	24	19	52	58
Tranquilizers ³	18	37	4	2	12	11	12	36
Stimulants ³	10	21	2	3	7	7	7	20
Methamphetamine ⁴	37	15	2	2	10	5	35	14
Sedatives ³	18	10	2	2	5	*	17	9
ILLCIT DRUGS OTHER THAN MARIJUANA²	103	99	12	11	33	31	98	93

*Low precision; no estimate reported.

¹ Estimates are based on non-imputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from the methamphetamine use items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Refers to nonmedical use; does not include over-the-counter drugs.

⁴ Estimates in the designated rows include data from methamphetamine use items added in 2005 and 2006.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 7.18D Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, 2007 and 2008

Drug¹	Total (2007)	Total (2008)	Aged 12-17 (2007)	Aged 12-17 (2008)	Aged 18-25 (2007)	Aged 18-25 (2008)	Aged 26+ (2007)	Aged 26+ (2008)
ILLICIT DRUGS²	0.18	0.19	0.75	0.84	0.40	0.41	0.19	0.21
Marijuana or Hashish	0.16	0.18	0.74	0.82	0.39	0.40	0.17	0.20
Cocaine (Including Crack)	0.06	0.05	0.13	0.18	0.09	0.10	0.07	0.06
Heroin	0.01	0.02	0.04	0.05	0.03	0.03	0.02	0.02
Hallucinogens	0.02	0.01	0.16	0.14	0.06	0.05	0.02	0.01
Inhalants	0.00	0.01	0.09	0.14	0.02	0.03	*	0.01
Pain Relievers ³	0.05	0.05	0.20	0.19	0.12	0.10	0.05	0.06
Tranquilizers ³	0.01	0.03	0.11	0.06	0.06	0.06	0.01	0.03
Stimulants ³	0.01	0.02	0.04	0.07	0.03	0.03	0.01	0.02
Methamphetamine ⁴	0.03	0.01	0.05	0.06	0.05	0.02	0.03	0.01
Sedatives ³	0.01	0.01	0.04	0.05	0.03	*	0.02	0.01
ILLICIT DRUGS OTHER THAN MARIJUANA²	0.08	0.08	0.30	0.30	0.16	0.15	0.09	0.09

*Low precision; no estimate reported.

¹ Estimates are based on non-imputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from the methamphetamine use items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Refers to nonmedical use; does not include over-the-counter drugs.

⁴ Estimates in the designated rows include data from methamphetamine use items added in 2005 and 2006.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 7.18P Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: *P* Values from Tests of Differences of Percentages, 2008 Versus 2007

Drug¹	Total (2007)	Total (2008)	Aged 12-17 (2007)	Aged 12-17 (2008)	Aged 18-25 (2007)	Aged 18-25 (2008)	Aged 26+ (2007)	Aged 26+ (2008)
ILLICIT DRUGS²	0.6910		0.4140		0.7841		0.4943	
Marijuana or Hashish	0.3064		0.4890		0.9454		0.1747	
Cocaine (Including Crack)	0.4243		0.3464		0.7857		0.3370	
Heroin	0.6131		0.9274		0.8719		0.6202	
Hallucinogens	0.1772		0.7705		0.8737		0.1710	
Inhalants	0.3296		0.4490		0.6124		*	
Pain Relievers ³	0.7556		0.6187		0.2774		0.4571	
Tranquilizers ³	0.0507		0.1081		0.9557		0.0187	
Stimulants ³	0.4504		0.5438		0.7058		0.5460	
Methamphetamine ⁴	0.0500		0.6447		0.0908		0.1144	
Sedatives ³	0.0932		0.9878		*		0.2886	
ILLICIT DRUGS OTHER THAN MARIJUANA²	0.4485		0.8600		0.5428		0.5784	

*Low precision; no estimate reported.

¹ Estimates are based on non-imputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from the methamphetamine use items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Refers to nonmedical use; does not include over-the-counter drugs.

⁴ Estimates in the designated rows include data from methamphetamine use items added in 2005 and 2006.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.