

Table 1.6C Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2017 and 2018

| Age Category | Lifetime (2017) | Lifetime (2018) | Past Year (2017) | Past Year (2018) | Past Month (2017) | Past Month (2018) |
|---------------------|----------------------------|----------------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| TOTAL | 895 | 930 | 656 | 719 | 515 | 574 |
| 12-20 | 245 | 223 | 206 | 188 | 144 | 141 |
| 21 or Older | 1,357 | 1,378 | 714 | 787 | 530 | 582 |
| 12-17 | 106 | 105 | 93 | 93 | 66 | 67 |
| 12-13 | 45 | 46 | 31 | 36 | 18 | 18 |
| 14-15 | 66 | 72 | 52 | 55 | 36 | 36 |
| 16-17 | 94 | 96 | 83 | 81 | 56 | 57 |
| 16-25 | 396 | 382 | 319 | 307 | 229 | 228 |
| 18 or Older | 881 | 904 | 646 | 714 | 508 | 570 |
| 18-25 | 185 | 188 | 186 | 182 | 160 | 163 |
| 18-20 | 203 | 181 | 171 | 156 | 127 | 121 |
| 21-25 | 281 | 278 | 221 | 219 | 170 | 159 |
| 26 or Older | 865 | 873 | 577 | 648 | 462 | 520 |
| 26-49 | 462 | 494 | 379 | 401 | 310 | 321 |
| 26-29 | 309 | 319 | 211 | 232 | 168 | 171 |
| 30-34 | 309 | 311 | 201 | 214 | 150 | 157 |
| 35-39 | 333 | 315 | 191 | 189 | 145 | 147 |
| 40-44 | 279 | 273 | 167 | 169 | 122 | 135 |
| 45-49 | 288 | 268 | 149 | 141 | 113 | 112 |
| 50 or Older | 704 | 719 | 411 | 470 | 323 | 373 |
| 50-54 | 453 | 446 | 227 | 225 | 168 | 160 |
| 55-59 | 459 | 460 | 227 | 244 | 181 | 177 |
| 60-64 | 407 | 463 | 181 | 219 | 134 | 170 |
| 65 or Older | 448 | 467 | 217 | 243 | 150 | 192 |

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Illicit Drug Use in Lifetime includes the misuse of prescription psychotherapeutics even though those estimates are not reported due to potential underreporting in lifetime prescription psychotherapeutics estimates.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 1.6D Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Detailed Age Category: Standard Errors of Percentages, 2017 and 2018

| Age Category | Lifetime (2017) | Lifetime (2018) | Past Year (2017) | Past Year (2018) | Past Month (2017) | Past Month (2018) |
|--------------------|--------------------|--------------------|---------------------|---------------------|----------------------|----------------------|
| TOTAL | 0.33 | 0.34 | 0.24 | 0.26 | 0.19 | 0.21 |
| 12-20 | 0.46 | 0.43 | 0.43 | 0.40 | 0.33 | 0.33 |
| 21 or Older | 0.37 | 0.38 | 0.27 | 0.30 | 0.21 | 0.23 |
| 12-17 | 0.43 | 0.42 | 0.37 | 0.37 | 0.26 | 0.27 |
| 12-13 | 0.55 | 0.54 | 0.38 | 0.44 | 0.22 | 0.22 |
| 14-15 | 0.65 | 0.75 | 0.56 | 0.61 | 0.41 | 0.42 |
| 16-17 | 0.82 | 0.87 | 0.78 | 0.80 | 0.59 | 0.60 |
| 16-25 | 0.49 | 0.48 | 0.48 | 0.46 | 0.40 | 0.41 |
| 18 or Older | 0.36 | 0.36 | 0.26 | 0.29 | 0.21 | 0.23 |
| 18-25 | 0.54 | 0.55 | 0.54 | 0.54 | 0.47 | 0.48 |
| 18-20 | 0.91 | 0.83 | 0.88 | 0.78 | 0.75 | 0.73 |
| 21-25 | 0.65 | 0.67 | 0.67 | 0.68 | 0.60 | 0.58 |
| 26 or Older | 0.41 | 0.41 | 0.27 | 0.30 | 0.22 | 0.24 |
| 26-49 | 0.46 | 0.49 | 0.38 | 0.40 | 0.31 | 0.32 |
| 26-29 | 1.03 | 0.96 | 0.92 | 0.98 | 0.79 | 0.81 |
| 30-34 | 0.84 | 0.87 | 0.75 | 0.78 | 0.63 | 0.61 |
| 35-39 | 0.95 | 0.94 | 0.75 | 0.76 | 0.63 | 0.64 |
| 40-44 | 0.94 | 0.97 | 0.75 | 0.77 | 0.58 | 0.65 |
| 45-49 | 0.99 | 1.02 | 0.68 | 0.67 | 0.54 | 0.55 |
| 50 or Older | 0.63 | 0.63 | 0.36 | 0.41 | 0.29 | 0.33 |
| 50-54 | 1.41 | 1.41 | 0.99 | 1.00 | 0.78 | 0.75 |
| 55-59 | 1.36 | 1.38 | 0.97 | 1.03 | 0.79 | 0.78 |
| 60-64 | 1.40 | 1.45 | 0.86 | 0.98 | 0.65 | 0.79 |
| 65 or Older | 0.90 | 0.91 | 0.44 | 0.47 | 0.30 | 0.38 |

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Illicit Drug Use in Lifetime includes the misuse of prescription psychotherapeutics even though those estimates are not reported due to potential underreporting in lifetime prescription psychotherapeutics estimates.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 1.6P Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2018 Versus 2017

| Age Category | Lifetime (2017) | Lifetime (2018) | Past Year (2017) | Past Year (2018) | Past Month (2017) | Past Month (2018) |
|---------------------|----------------------------|----------------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| TOTAL | 0.5420 | | 0.2452 | | 0.0839 | |
| 12-20 | 0.9759 | | 0.5979 | | 0.6341 | |
| 21 or Older | 0.5268 | | 0.2898 | | 0.0953 | |
| 12-17 | 1.0000 | | 0.5029 | | 0.6747 | |
| 12-13 | 0.8606 | | 0.1808 | | 0.9015 | |
| 14-15 | 0.1457 | | 0.1667 | | 0.5123 | |
| 16-17 | 0.3286 | | 0.5296 | | 0.9353 | |
| 16-25 | 0.0381 | | 0.2616 | | 0.6650 | |
| 18 or Older | 0.5037 | | 0.2875 | | 0.0970 | |
| 18-25 | 0.0571 | | 0.3217 | | 0.6344 | |
| 18-20 | 0.7849 | | 0.9808 | | 0.8529 | |
| 21-25 | 0.0405 | | 0.2104 | | 0.4655 | |
| 26 or Older | 0.8062 | | 0.1093 | | 0.0374 | |
| 26-49 | 0.4077 | | 0.0133 | | 0.0005 | |
| 26-29 | 0.1165 | | 0.0194 | | 0.1534 | |
| 30-34 | 0.3240 | | 0.4321 | | 0.0968 | |
| 35-39 | 0.6949 | | 0.4697 | | 0.0687 | |
| 40-44 | 0.6174 | | 0.0472 | | 0.0234 | |
| 45-49 | 0.0904 | | 0.7719 | | 0.3259 | |
| 50 or Older | 0.7799 | | 0.9725 | | 0.7984 | |
| 50-54 | 0.9745 | | 0.9486 | | 0.6033 | |
| 55-59 | 0.2649 | | 0.9308 | | 0.2633 | |
| 60-64 | 0.3640 | | 0.7215 | | 0.3629 | |
| 65 or Older | 0.3131 | | 0.9347 | | 0.5803 | |

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Illicit Drug Use in Lifetime includes the misuse of prescription psychotherapeutics even though those estimates are not reported due to potential underreporting in lifetime prescription psychotherapeutics estimates.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 1.7C Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2017 and 2018

| Age Category | Lifetime (2017) | Lifetime (2018) | Past Year (2017) | Past Year (2018) | Past Month (2017) | Past Month (2018) |
|---------------------|----------------------------|----------------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| TOTAL | 904 | 931 | 595 | 669 | 478 | 538 |
| 12-20 | 214 | 204 | 188 | 180 | 137 | 138 |
| 21 or Older | 1,306 | 1,334 | 631 | 712 | 490 | 540 |
| 12-17 | 91 | 92 | 83 | 83 | 61 | 63 |
| 12-13 | 22 | 20 | 18 | 18 | 11 | 13 |
| 14-15 | 50 | 54 | 45 | 48 | 32 | 32 |
| 16-17 | 82 | 85 | 76 | 75 | 53 | 54 |
| 16-25 | 373 | 363 | 300 | 294 | 218 | 220 |
| 18 or Older | 889 | 902 | 586 | 662 | 474 | 533 |
| 18-25 | 189 | 189 | 182 | 181 | 156 | 160 |
| 18-20 | 185 | 173 | 160 | 151 | 120 | 119 |
| 21-25 | 271 | 265 | 207 | 206 | 160 | 149 |
| 26 or Older | 866 | 870 | 518 | 595 | 423 | 483 |
| 26-49 | 476 | 499 | 361 | 375 | 287 | 300 |
| 26-29 | 300 | 309 | 195 | 217 | 156 | 163 |
| 30-34 | 304 | 307 | 180 | 198 | 139 | 151 |
| 35-39 | 323 | 301 | 173 | 171 | 133 | 134 |
| 40-44 | 265 | 260 | 146 | 153 | 115 | 127 |
| 45-49 | 271 | 258 | 126 | 119 | 103 | 98 |
| 50 or Older | 697 | 713 | 350 | 426 | 293 | 351 |
| 50-54 | 429 | 426 | 194 | 183 | 149 | 150 |
| 55-59 | 448 | 451 | 189 | 200 | 153 | 160 |
| 60-64 | 400 | 447 | 154 | 202 | 127 | 160 |
| 65 or Older | 435 | 467 | 168 | 219 | 128 | 177 |

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 1.7D Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Detailed Age Category: Standard Errors of Percentages, 2017 and 2018

| Age Category | Lifetime (2017) | Lifetime (2018) | Past Year (2017) | Past Year (2018) | Past Month (2017) | Past Month (2018) |
|---------------------|----------------------------|----------------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| TOTAL | 0.33 | 0.34 | 0.22 | 0.24 | 0.18 | 0.20 |
| 12-20 | 0.44 | 0.43 | 0.40 | 0.40 | 0.32 | 0.32 |
| 21 or Older | 0.38 | 0.38 | 0.24 | 0.27 | 0.20 | 0.22 |
| 12-17 | 0.37 | 0.37 | 0.33 | 0.33 | 0.24 | 0.25 |
| 12-13 | 0.27 | 0.25 | 0.23 | 0.23 | 0.14 | 0.16 |
| 14-15 | 0.55 | 0.60 | 0.50 | 0.55 | 0.37 | 0.38 |
| 16-17 | 0.78 | 0.81 | 0.73 | 0.75 | 0.56 | 0.57 |
| 16-25 | 0.50 | 0.50 | 0.47 | 0.46 | 0.39 | 0.40 |
| 18 or Older | 0.36 | 0.36 | 0.24 | 0.27 | 0.19 | 0.21 |
| 18-25 | 0.55 | 0.56 | 0.53 | 0.53 | 0.46 | 0.47 |
| 18-20 | 0.87 | 0.82 | 0.83 | 0.79 | 0.73 | 0.73 |
| 21-25 | 0.69 | 0.67 | 0.66 | 0.66 | 0.57 | 0.55 |
| 26 or Older | 0.41 | 0.41 | 0.24 | 0.28 | 0.20 | 0.22 |
| 26-49 | 0.47 | 0.49 | 0.36 | 0.37 | 0.29 | 0.30 |
| 26-29 | 1.02 | 1.00 | 0.87 | 0.97 | 0.75 | 0.79 |
| 30-34 | 0.89 | 0.89 | 0.71 | 0.75 | 0.60 | 0.60 |
| 35-39 | 0.96 | 0.93 | 0.72 | 0.71 | 0.58 | 0.58 |
| 40-44 | 0.93 | 0.98 | 0.68 | 0.71 | 0.55 | 0.61 |
| 45-49 | 0.98 | 1.04 | 0.60 | 0.59 | 0.50 | 0.49 |
| 50 or Older | 0.62 | 0.63 | 0.31 | 0.37 | 0.26 | 0.31 |
| 50-54 | 1.44 | 1.39 | 0.88 | 0.84 | 0.70 | 0.71 |
| 55-59 | 1.37 | 1.37 | 0.83 | 0.86 | 0.68 | 0.71 |
| 60-64 | 1.42 | 1.48 | 0.74 | 0.92 | 0.62 | 0.75 |
| 65 or Older | 0.88 | 0.91 | 0.34 | 0.43 | 0.26 | 0.35 |

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 1.7P Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2018 Versus 2017

| Age Category | Lifetime (2017) | Lifetime (2018) | Past Year (2017) | Past Year (2018) | Past Month (2017) | Past Month (2018) |
|---------------------|----------------------------|----------------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| TOTAL | 0.8341 | | 0.0068 | | 0.0280 | |
| 12-20 | 0.6748 | | 0.4533 | | 0.3583 | |
| 21 or Older | 0.9126 | | 0.0092 | | 0.0394 | |
| 12-17 | 0.8526 | | 0.8555 | | 0.5707 | |
| 12-13 | 0.5841 | | 0.6491 | | 0.3389 | |
| 14-15 | 0.2343 | | 0.2460 | | 0.7822 | |
| 16-17 | 0.7187 | | 0.5393 | | 0.7396 | |
| 16-25 | 0.0931 | | 0.7324 | | 0.9463 | |
| 18 or Older | 0.8921 | | 0.0072 | | 0.0341 | |
| 18-25 | 0.0852 | | 0.8925 | | 0.9866 | |
| 18-20 | 0.8220 | | 0.4512 | | 0.5045 | |
| 21-25 | 0.0291 | | 0.4501 | | 0.5830 | |
| 26 or Older | 0.5886 | | 0.0016 | | 0.0140 | |
| 26-49 | 0.6198 | | 0.0001 | | 0.0002 | |
| 26-29 | 0.2079 | | 0.0103 | | 0.1238 | |
| 30-34 | 0.5893 | | 0.0788 | | 0.0119 | |
| 35-39 | 0.9150 | | 0.0517 | | 0.0613 | |
| 40-44 | 0.7097 | | 0.0116 | | 0.0474 | |
| 45-49 | 0.0771 | | 0.9043 | | 0.5753 | |
| 50 or Older | 0.3173 | | 0.3948 | | 0.9699 | |
| 50-54 | 0.3981 | | 0.8740 | | 0.9216 | |
| 55-59 | 0.4504 | | 0.8573 | | 0.5673 | |
| 60-64 | 0.3434 | | 0.3201 | | 0.5395 | |
| 65 or Older | 0.2013 | | 0.4621 | | 0.7098 | |

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 1.8C Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2017 and 2018

| Age Category | Lifetime (2017) | Lifetime (2018) | Past Year (2017) | Past Year (2018) | Past Month (2017) | Past Month (2018) |
|---------------------|----------------------------|----------------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| TOTAL | 618 | 688 | 217 | 205 | 130 | 123 |
| 12-20 | 57 | 58 | 48 | 48 | 28 | 24 |
| 21 or Older | 698 | 773 | 214 | 200 | 128 | 122 |
| 12-17 | 18 | 21 | 16 | 18 | 7 | 5 |
| 12-13 | 4 | 4 | 4 | 3 | 1 | 0 |
| 14-15 | 6 | 10 | 5 | 9 | 0 | * |
| 16-17 | 16 | 19 | 14 | 15 | 7 | 5 |
| 16-25 | 139 | 143 | 96 | 98 | 49 | 47 |
| 18 or Older | 611 | 682 | 218 | 203 | 130 | 123 |
| 18-25 | 124 | 123 | 89 | 89 | 47 | 46 |
| 18-20 | 54 | 56 | 46 | 44 | 27 | 23 |
| 21-25 | 125 | 125 | 86 | 81 | 44 | 40 |
| 26 or Older | 595 | 666 | 192 | 177 | 123 | 117 |
| 26-49 | 334 | 324 | 140 | 146 | 90 | 100 |
| 26-29 | 154 | 164 | 85 | 89 | 51 | 61 |
| 30-34 | 172 | 173 | 76 | 69 | 47 | 37 |
| 35-39 | 160 | 173 | 60 | 62 | 42 | 41 |
| 40-44 | 155 | 148 | 41 | 42 | 28 | 30 |
| 45-49 | 167 | 141 | 51 | 41 | 32 | 30 |
| 50 or Older | 493 | 580 | 126 | 106 | 89 | 71 |
| 50-54 | 280 | 286 | 73 | 57 | 49 | 28 |
| 55-59 | 303 | 314 | 75 | 44 | 58 | 34 |
| 60-64 | 251 | 301 | 34 | 64 | 14 | 57 |
| 65 or Older | 237 | 288 | 57 | 14 | 45 | 6 |

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 1.8D Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Detailed Age Category: Standard Errors of Percentages, 2017 and 2018

| Age Category | Lifetime (2017) | Lifetime (2018) | Past Year (2017) | Past Year (2018) | Past Month (2017) | Past Month (2018) |
|--------------------|--------------------|--------------------|---------------------|---------------------|----------------------|----------------------|
| TOTAL | 0.23 | 0.25 | 0.08 | 0.07 | 0.05 | 0.05 |
| 12-20 | 0.14 | 0.15 | 0.12 | 0.12 | 0.07 | 0.06 |
| 21 or Older | 0.26 | 0.29 | 0.09 | 0.08 | 0.05 | 0.05 |
| 12-17 | 0.07 | 0.09 | 0.06 | 0.07 | 0.03 | 0.02 |
| 12-13 | 0.06 | 0.05 | 0.06 | 0.04 | 0.01 | 0.01 |
| 14-15 | 0.07 | 0.12 | 0.06 | 0.11 | 0.01 | * |
| 16-17 | 0.19 | 0.22 | 0.16 | 0.17 | 0.08 | 0.06 |
| 16-25 | 0.29 | 0.30 | 0.21 | 0.21 | 0.11 | 0.11 |
| 18 or Older | 0.25 | 0.27 | 0.09 | 0.08 | 0.05 | 0.05 |
| 18-25 | 0.36 | 0.36 | 0.26 | 0.26 | 0.14 | 0.13 |
| 18-20 | 0.39 | 0.40 | 0.34 | 0.33 | 0.21 | 0.18 |
| 21-25 | 0.51 | 0.51 | 0.37 | 0.35 | 0.20 | 0.19 |
| 26 or Older | 0.28 | 0.31 | 0.09 | 0.08 | 0.06 | 0.05 |
| 26-49 | 0.33 | 0.32 | 0.14 | 0.14 | 0.09 | 0.10 |
| 26-29 | 0.75 | 0.80 | 0.44 | 0.47 | 0.28 | 0.33 |
| 30-34 | 0.71 | 0.68 | 0.34 | 0.30 | 0.21 | 0.17 |
| 35-39 | 0.70 | 0.72 | 0.29 | 0.28 | 0.20 | 0.19 |
| 40-44 | 0.70 | 0.70 | 0.20 | 0.22 | 0.14 | 0.15 |
| 45-49 | 0.75 | 0.67 | 0.25 | 0.21 | 0.16 | 0.15 |
| 50 or Older | 0.44 | 0.51 | 0.11 | 0.09 | 0.08 | 0.06 |
| 50-54 | 1.13 | 1.18 | 0.35 | 0.28 | 0.23 | 0.13 |
| 55-59 | 1.17 | 1.22 | 0.34 | 0.20 | 0.26 | 0.16 |
| 60-64 | 1.11 | 1.28 | 0.17 | 0.31 | 0.07 | 0.28 |
| 65 or Older | 0.48 | 0.56 | 0.11 | 0.03 | 0.09 | 0.01 |

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 1.8P Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2018 Versus 2017

| Age Category | Lifetime (2017) | Lifetime (2018) | Past Year (2017) | Past Year (2018) | Past Month (2017) | Past Month (2018) |
|---------------------|----------------------------|----------------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| TOTAL | 0.4885 | | 0.1298 | | 0.1917 | |
| 12-20 | 0.6462 | | 0.7411 | | 0.3834 | |
| 21 or Older | 0.4473 | | 0.1389 | | 0.2574 | |
| 12-17 | 0.8788 | | 0.5124 | | 0.0698 | |
| 12-13 | 0.9012 | | 0.5265 | | 0.8860 | |
| 14-15 | 0.2282 | | 0.3430 | | * | |
| 16-17 | 0.7144 | | 0.2827 | | 0.0839 | |
| 16-25 | 0.1824 | | 0.1819 | | 0.0220 | |
| 18 or Older | 0.4604 | | 0.1400 | | 0.2201 | |
| 18-25 | 0.2023 | | 0.2435 | | 0.0365 | |
| 18-20 | 0.7370 | | 0.8412 | | 0.5943 | |
| 21-25 | 0.1518 | | 0.2463 | | 0.0415 | |
| 26 or Older | 0.6001 | | 0.3099 | | 0.6532 | |
| 26-49 | 0.7645 | | 0.9539 | | 0.9410 | |
| 26-29 | 0.3545 | | 0.4064 | | 0.1581 | |
| 30-34 | 0.8913 | | 0.5868 | | 0.0111 | |
| 35-39 | 0.2157 | | 0.7780 | | 0.9553 | |
| 40-44 | 0.3021 | | 0.4390 | | 0.2239 | |
| 45-49 | 0.0032 | | 0.4176 | | 0.8844 | |
| 50 or Older | 0.4320 | | 0.1018 | | 0.4913 | |
| 50-54 | 0.2637 | | 0.1745 | | 0.1476 | |
| 55-59 | 0.3824 | | 0.0427 | | 0.2935 | |
| 60-64 | 0.5949 | | 0.0354 | | 0.0097 | |
| 65 or Older | 0.1716 | | 0.0395 | | 0.0950 | |

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 1.9C Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2017 and 2018

| Age Category | Lifetime (2017) | Lifetime (2018) | Past Year (2017) | Past Year (2018) | Past Month (2017) | Past Month (2018) |
|--------------------|--------------------|--------------------|---------------------|---------------------|----------------------|----------------------|
| TOTAL | 322 | 330 | 99 | 85 | 74 | 72 |
| 12-20 | 15 | 16 | 9 | 10 | 5 | 6 |
| 21 or Older | 333 | 336 | 99 | 85 | 74 | 72 |
| 12-17 | 8 | 5 | 7 | 3 | * | * |
| 12-13 | 0 | 1 | * | * | * | * |
| 14-15 | 3 | 1 | 3 | * | * | * |
| 16-17 | 8 | 5 | 6 | 3 | 1 | * |
| 16-25 | 39 | 31 | 22 | 15 | 7 | 9 |
| 18 or Older | 322 | 329 | 99 | 85 | 74 | 72 |
| 18-25 | 37 | 32 | 20 | 16 | 7 | 9 |
| 18-20 | 13 | 16 | 7 | 9 | 5 | 6 |
| 21-25 | 36 | 27 | 19 | 12 | 5 | 6 |
| 26 or Older | 316 | 325 | 96 | 83 | 74 | 71 |
| 26-49 | 176 | 170 | 54 | 47 | 40 | 35 |
| 26-29 | 57 | 50 | 19 | 20 | 15 | 14 |
| 30-34 | 78 | 72 | 29 | 16 | 23 | 8 |
| 35-39 | 84 | 84 | 18 | 25 | 17 | 21 |
| 40-44 | 74 | 86 | 16 | 17 | 10 | 12 |
| 45-49 | 89 | 78 | 29 | 21 | 22 | 17 |
| 50 or Older | 254 | 278 | 83 | 68 | 65 | 61 |
| 50-54 | 153 | 159 | 57 | 21 | 41 | 18 |
| 55-59 | 152 | 160 | 49 | 36 | 45 | 27 |
| 60-64 | 121 | 133 | 27 | 53 | 8 | 51 |
| 65 or Older | 103 | 101 | 23 | 6 | 20 | 6 |

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 1.9D Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Detailed Age Category: Standard Errors of Percentages, 2017 and 2018

| Age Category | Lifetime (2017) | Lifetime (2018) | Past Year (2017) | Past Year (2018) | Past Month (2017) | Past Month (2018) |
|---------------------|----------------------------|----------------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| TOTAL | 0.12 | 0.12 | 0.04 | 0.03 | 0.03 | 0.03 |
| 12-20 | 0.04 | 0.04 | 0.02 | 0.03 | 0.01 | 0.02 |
| 21 or Older | 0.14 | 0.14 | 0.04 | 0.04 | 0.03 | 0.03 |
| 12-17 | 0.03 | 0.02 | 0.03 | 0.01 | * | * |
| 12-13 | 0.01 | 0.01 | * | * | * | * |
| 14-15 | 0.03 | 0.02 | 0.03 | * | * | * |
| 16-17 | 0.09 | 0.06 | 0.07 | 0.03 | 0.01 | * |
| 16-25 | 0.09 | 0.07 | 0.05 | 0.04 | 0.02 | 0.02 |
| 18 or Older | 0.13 | 0.13 | 0.04 | 0.03 | 0.03 | 0.03 |
| 18-25 | 0.11 | 0.09 | 0.06 | 0.05 | 0.02 | 0.03 |
| 18-20 | 0.10 | 0.12 | 0.05 | 0.07 | 0.04 | 0.05 |
| 21-25 | 0.16 | 0.13 | 0.09 | 0.06 | 0.02 | 0.03 |
| 26 or Older | 0.15 | 0.15 | 0.05 | 0.04 | 0.03 | 0.03 |
| 26-49 | 0.18 | 0.17 | 0.05 | 0.05 | 0.04 | 0.03 |
| 26-29 | 0.32 | 0.28 | 0.11 | 0.11 | 0.08 | 0.08 |
| 30-34 | 0.35 | 0.32 | 0.13 | 0.07 | 0.11 | 0.04 |
| 35-39 | 0.39 | 0.38 | 0.09 | 0.11 | 0.08 | 0.10 |
| 40-44 | 0.36 | 0.43 | 0.08 | 0.09 | 0.05 | 0.06 |
| 45-49 | 0.43 | 0.40 | 0.14 | 0.11 | 0.11 | 0.09 |
| 50 or Older | 0.23 | 0.24 | 0.07 | 0.06 | 0.06 | 0.05 |
| 50-54 | 0.71 | 0.73 | 0.27 | 0.10 | 0.20 | 0.09 |
| 55-59 | 0.67 | 0.70 | 0.22 | 0.17 | 0.20 | 0.12 |
| 60-64 | 0.60 | 0.63 | 0.13 | 0.26 | 0.04 | 0.25 |
| 65 or Older | 0.21 | 0.20 | 0.05 | 0.01 | 0.04 | 0.01 |

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 1.9P Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2018 Versus 2017

| Age Category | Lifetime (2017) | Lifetime (2018) | Past Year (2017) | Past Year (2018) | Past Month (2017) | Past Month (2018) |
|---------------------|----------------------------|----------------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| TOTAL | 0.3009 | | 0.1858 | | 0.7196 | |
| 12-20 | 0.9968 | | 0.7486 | | 1.0000 | |
| 21 or Older | 0.2944 | | 0.1946 | | 0.7179 | |
| 12-17 | 0.1072 | | 0.1001 | | * | |
| 12-13 | 0.9534 | | * | | * | |
| 14-15 | 0.6042 | | * | | * | |
| 16-17 | 0.1245 | | 0.1828 | | * | |
| 16-25 | 0.0333 | | 0.2602 | | 0.7560 | |
| 18 or Older | 0.3087 | | 0.2168 | | 0.7237 | |
| 18-25 | 0.0643 | | 0.4335 | | 0.6918 | |
| 18-20 | 0.4458 | | 0.4969 | | 0.9136 | |
| 21-25 | 0.0205 | | 0.2256 | | 0.6580 | |
| 26 or Older | 0.3876 | | 0.2626 | | 0.6871 | |
| 26-49 | 0.5867 | | 0.7582 | | 0.4334 | |
| 26-29 | 0.2990 | | 0.8177 | | 0.8394 | |
| 30-34 | 0.4889 | | 0.0158 | | 0.0106 | |
| 35-39 | 0.5013 | | 0.2851 | | 0.8192 | |
| 40-44 | 0.1462 | | 0.4804 | | 0.3962 | |
| 45-49 | 0.5306 | | 0.9515 | | 0.8355 | |
| 50 or Older | 0.4949 | | 0.2725 | | 0.9788 | |
| 50-54 | 0.7177 | | 0.0156 | | 0.1324 | |
| 55-59 | 0.9236 | | 0.3693 | | 0.3499 | |
| 60-64 | 0.9733 | | 0.0714 | | 0.0109 | |
| 65 or Older | 0.2988 | | 0.2859 | | 0.4943 | |

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 1.10C Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2017 and 2018

| Age Category | Lifetime (2017) | Lifetime (2018) | Past Year (2017) | Past Year (2018) | Past Month (2017) | Past Month (2018) |
|---------------------|----------------------------|----------------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| TOTAL | 646 | 726 | 177 | 215 | 91 | 106 |
| 12-20 | 86 | 83 | 67 | 66 | 36 | 32 |
| 21 or Older | 734 | 802 | 163 | 203 | 82 | 102 |
| 12-17 | 39 | 39 | 34 | 31 | 19 | 20 |
| 12-13 | 13 | 11 | 11 | 9 | 6 | 5 |
| 14-15 | 23 | 20 | 19 | 18 | 13 | 10 |
| 16-17 | 30 | 30 | 27 | 23 | 14 | 15 |
| 16-25 | 173 | 181 | 108 | 114 | 52 | 51 |
| 18 or Older | 639 | 714 | 173 | 214 | 89 | 105 |
| 18-25 | 140 | 145 | 95 | 102 | 49 | 48 |
| 18-20 | 75 | 73 | 57 | 57 | 30 | 26 |
| 21-25 | 146 | 146 | 84 | 86 | 37 | 40 |
| 26 or Older | 612 | 674 | 138 | 170 | 74 | 91 |
| 26-49 | 371 | 362 | 114 | 150 | 68 | 79 |
| 26-29 | 178 | 186 | 69 | 94 | 35 | 52 |
| 30-34 | 188 | 185 | 73 | 72 | 48 | 39 |
| 35-39 | 187 | 193 | 52 | 59 | 28 | 37 |
| 40-44 | 174 | 166 | 33 | 42 | 21 | 24 |
| 45-49 | 161 | 150 | 29 | 34 | 10 | 16 |
| 50 or Older | 469 | 553 | 61 | 76 | 30 | 46 |
| 50-54 | 239 | 245 | 32 | 45 | 20 | 33 |
| 55-59 | 263 | 289 | 41 | 21 | 22 | 12 |
| 60-64 | 239 | 279 | 19 | 44 | 6 | 29 |
| 65 or Older | 239 | 280 | 25 | 30 | * | 8 |

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 1.10D Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Detailed Age Category: Standard Errors of Percentages, 2017 and 2018

| Age Category | Lifetime (2017) | Lifetime (2018) | Past Year (2017) | Past Year (2018) | Past Month (2017) | Past Month (2018) |
|---------------------|----------------------------|----------------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| TOTAL | 0.24 | 0.27 | 0.07 | 0.08 | 0.03 | 0.04 |
| 12-20 | 0.21 | 0.21 | 0.17 | 0.17 | 0.09 | 0.08 |
| 21 or Older | 0.27 | 0.30 | 0.07 | 0.08 | 0.03 | 0.04 |
| 12-17 | 0.16 | 0.15 | 0.14 | 0.13 | 0.08 | 0.08 |
| 12-13 | 0.16 | 0.14 | 0.13 | 0.11 | 0.07 | 0.07 |
| 14-15 | 0.27 | 0.24 | 0.22 | 0.21 | 0.15 | 0.13 |
| 16-17 | 0.33 | 0.35 | 0.31 | 0.26 | 0.16 | 0.18 |
| 16-25 | 0.34 | 0.36 | 0.23 | 0.25 | 0.12 | 0.12 |
| 18 or Older | 0.26 | 0.29 | 0.07 | 0.09 | 0.04 | 0.04 |
| 18-25 | 0.41 | 0.43 | 0.28 | 0.30 | 0.14 | 0.14 |
| 18-20 | 0.52 | 0.51 | 0.42 | 0.42 | 0.23 | 0.19 |
| 21-25 | 0.56 | 0.56 | 0.36 | 0.37 | 0.17 | 0.19 |
| 26 or Older | 0.29 | 0.31 | 0.06 | 0.08 | 0.03 | 0.04 |
| 26-49 | 0.37 | 0.36 | 0.11 | 0.15 | 0.07 | 0.08 |
| 26-29 | 0.85 | 0.87 | 0.37 | 0.51 | 0.19 | 0.29 |
| 30-34 | 0.74 | 0.70 | 0.33 | 0.32 | 0.22 | 0.17 |
| 35-39 | 0.76 | 0.77 | 0.25 | 0.27 | 0.13 | 0.17 |
| 40-44 | 0.76 | 0.76 | 0.17 | 0.21 | 0.11 | 0.12 |
| 45-49 | 0.73 | 0.72 | 0.15 | 0.18 | 0.05 | 0.08 |
| 50 or Older | 0.42 | 0.49 | 0.05 | 0.07 | 0.03 | 0.04 |
| 50-54 | 0.98 | 1.06 | 0.15 | 0.22 | 0.09 | 0.16 |
| 55-59 | 1.11 | 1.16 | 0.19 | 0.09 | 0.10 | 0.06 |
| 60-64 | 1.06 | 1.20 | 0.10 | 0.21 | 0.03 | 0.14 |
| 65 or Older | 0.48 | 0.55 | 0.05 | 0.06 | * | 0.02 |

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 1.10P Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2018 Versus 2017

| Age Category | Lifetime (2017) | Lifetime (2018) | Past Year (2017) | Past Year (2018) | Past Month (2017) | Past Month (2018) |
|---------------------|----------------------------|----------------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| TOTAL | 0.2799 | | 0.0923 | | 0.1673 | |
| 12-20 | 0.4534 | | 0.0913 | | 0.4092 | |
| 21 or Older | 0.2354 | | 0.0155 | | 0.0765 | |
| 12-17 | 0.0118 | | 0.0032 | | 0.9118 | |
| 12-13 | 0.3313 | | 0.2826 | | 0.5189 | |
| 14-15 | 0.3145 | | 0.4615 | | 0.6017 | |
| 16-17 | 0.0322 | | 0.0037 | | 0.6120 | |
| 16-25 | 0.1322 | | 0.2534 | | 0.9806 | |
| 18 or Older | 0.2245 | | 0.0264 | | 0.1543 | |
| 18-25 | 0.2717 | | 0.6760 | | 0.8941 | |
| 18-20 | 0.6771 | | 0.7732 | | 0.3308 | |
| 21-25 | 0.1766 | | 0.7593 | | 0.5998 | |
| 26 or Older | 0.1245 | | 0.0019 | | 0.0782 | |
| 26-49 | 0.1539 | | 0.0016 | | 0.1588 | |
| 26-29 | 0.0964 | | 0.0039 | | 0.0507 | |
| 30-34 | 0.4575 | | 0.4590 | | 0.4631 | |
| 35-39 | 0.6828 | | 0.3442 | | 0.4768 | |
| 40-44 | 0.0206 | | 0.0893 | | 0.5658 | |
| 45-49 | 0.4812 | | 0.3469 | | 0.4506 | |
| 50 or Older | 0.3851 | | 0.4282 | | 0.2970 | |
| 50-54 | 0.6395 | | 0.3220 | | 0.3210 | |
| 55-59 | 0.1606 | | 0.0860 | | 0.5750 | |
| 60-64 | 0.5536 | | 0.1692 | | 0.3218 | |
| 65 or Older | 0.1027 | | 0.3853 | | * | |

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 1.11C Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2017 and 2018

| Age Category | Lifetime (2017) | Lifetime (2018) | Past Year (2017) | Past Year (2018) | Past Month (2017) | Past Month (2018) |
|---------------------|----------------------------|----------------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| TOTAL | 489 | 478 | 95 | 112 | 55 | 62 |
| 12-20 | 96 | 97 | 41 | 48 | 22 | 26 |
| 21 or Older | 511 | 508 | 89 | 103 | 49 | 57 |
| 12-17 | 67 | 70 | 34 | 41 | 18 | 23 |
| 12-13 | 38 | 35 | 21 | 23 | 13 | 11 |
| 14-15 | 39 | 46 | 20 | 26 | 11 | 15 |
| 16-17 | 43 | 46 | 19 | 21 | 7 | 12 |
| 16-25 | 120 | 122 | 50 | 48 | 23 | 24 |
| 18 or Older | 482 | 473 | 90 | 105 | 51 | 58 |
| 18-25 | 105 | 100 | 44 | 43 | 22 | 20 |
| 18-20 | 62 | 61 | 24 | 27 | 13 | 12 |
| 21-25 | 90 | 91 | 41 | 34 | 18 | 17 |
| 26 or Older | 458 | 463 | 77 | 97 | 46 | 55 |
| 26-49 | 275 | 288 | 69 | 72 | 42 | 46 |
| 26-29 | 120 | 122 | 40 | 32 | 18 | 21 |
| 30-34 | 132 | 125 | 35 | 37 | 22 | 11 |
| 35-39 | 126 | 150 | 28 | 30 | 24 | 19 |
| 40-44 | 135 | 124 | 20 | 24 | 15 | 18 |
| 45-49 | 128 | 119 | 14 | 26 | 11 | 22 |
| 50 or Older | 341 | 355 | 34 | 63 | 20 | 31 |
| 50-54 | 183 | 200 | 26 | 40 | 20 | 21 |
| 55-59 | 186 | 206 | 5 | 43 | * | 20 |
| 60-64 | 151 | 182 | 14 | 18 | * | 2 |
| 65 or Older | 132 | 153 | 15 | 18 | * | 9 |

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 1.11D Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Detailed Age Category: Standard Errors of Percentages, 2017 and 2018

| Age Category | Lifetime (2017) | Lifetime (2018) | Past Year (2017) | Past Year (2018) | Past Month (2017) | Past Month (2018) |
|---------------------|----------------------------|----------------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| TOTAL | 0.18 | 0.17 | 0.03 | 0.04 | 0.02 | 0.02 |
| 12-20 | 0.24 | 0.25 | 0.11 | 0.13 | 0.06 | 0.07 |
| 21 or Older | 0.20 | 0.20 | 0.04 | 0.04 | 0.02 | 0.02 |
| 12-17 | 0.27 | 0.28 | 0.14 | 0.16 | 0.07 | 0.09 |
| 12-13 | 0.46 | 0.42 | 0.27 | 0.29 | 0.16 | 0.14 |
| 14-15 | 0.42 | 0.51 | 0.23 | 0.31 | 0.13 | 0.18 |
| 16-17 | 0.48 | 0.52 | 0.21 | 0.24 | 0.08 | 0.14 |
| 16-25 | 0.26 | 0.26 | 0.11 | 0.11 | 0.05 | 0.06 |
| 18 or Older | 0.20 | 0.19 | 0.04 | 0.04 | 0.02 | 0.02 |
| 18-25 | 0.31 | 0.29 | 0.13 | 0.13 | 0.06 | 0.06 |
| 18-20 | 0.46 | 0.45 | 0.19 | 0.20 | 0.10 | 0.09 |
| 21-25 | 0.39 | 0.38 | 0.18 | 0.16 | 0.08 | 0.08 |
| 26 or Older | 0.22 | 0.22 | 0.04 | 0.04 | 0.02 | 0.03 |
| 26-49 | 0.27 | 0.29 | 0.07 | 0.07 | 0.04 | 0.05 |
| 26-29 | 0.61 | 0.63 | 0.22 | 0.18 | 0.10 | 0.12 |
| 30-34 | 0.56 | 0.53 | 0.16 | 0.17 | 0.10 | 0.05 |
| 35-39 | 0.56 | 0.62 | 0.13 | 0.13 | 0.12 | 0.09 |
| 40-44 | 0.62 | 0.60 | 0.10 | 0.12 | 0.08 | 0.09 |
| 45-49 | 0.59 | 0.58 | 0.07 | 0.13 | 0.05 | 0.11 |
| 50 or Older | 0.30 | 0.31 | 0.03 | 0.06 | 0.02 | 0.03 |
| 50-54 | 0.82 | 0.90 | 0.12 | 0.19 | 0.09 | 0.10 |
| 55-59 | 0.81 | 0.88 | 0.02 | 0.20 | * | 0.09 |
| 60-64 | 0.73 | 0.83 | 0.07 | 0.09 | * | 0.01 |
| 65 or Older | 0.27 | 0.30 | 0.03 | 0.03 | * | 0.02 |

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 1.11P Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2018 Versus 2017

| Age Category | Lifetime (2017) | Lifetime (2018) | Past Year (2017) | Past Year (2018) | Past Month (2017) | Past Month (2018) |
|---------------------|----------------------------|----------------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| TOTAL | 0.4012 | | 0.0972 | | 0.5133 | |
| 12-20 | 0.8681 | | 0.1433 | | 0.7310 | |
| 21 or Older | 0.3678 | | 0.2662 | | 0.5691 | |
| 12-17 | 0.8779 | | 0.0884 | | 0.2771 | |
| 12-13 | 0.5701 | | 0.1225 | | 0.7087 | |
| 14-15 | 0.0633 | | 0.0732 | | 0.3842 | |
| 16-17 | 0.1542 | | 0.5678 | | 0.1067 | |
| 16-25 | 0.1191 | | 0.2408 | | 0.6087 | |
| 18 or Older | 0.4092 | | 0.2676 | | 0.7647 | |
| 18-25 | 0.2677 | | 0.3176 | | 0.1904 | |
| 18-20 | 0.6526 | | 0.9308 | | 0.2722 | |
| 21-25 | 0.0969 | | 0.1986 | | 0.4290 | |
| 26 or Older | 0.5502 | | 0.0629 | | 0.3646 | |
| 26-49 | 0.8957 | | 0.3984 | | 0.8761 | |
| 26-29 | 0.9622 | | 0.2410 | | 0.7658 | |
| 30-34 | 0.5219 | | 0.5077 | | 0.1506 | |
| 35-39 | 0.1547 | | 0.3817 | | 0.7307 | |
| 40-44 | 0.7935 | | 0.4205 | | 0.5725 | |
| 45-49 | 0.1025 | | 0.1217 | | 0.1867 | |
| 50 or Older | 0.5260 | | 0.0612 | | 0.1402 | |
| 50-54 | 0.9584 | | 0.2467 | | 0.6201 | |
| 55-59 | 0.7197 | | 0.0982 | | * | |
| 60-64 | 0.8025 | | 0.8124 | | * | |
| 65 or Older | 0.3886 | | 0.8390 | | * | |

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 1.12C Methamphetamine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2017 and 2018

| Age Category | Lifetime (2017) | Lifetime (2018) | Past Year (2017) | Past Year (2018) | Past Month (2017) | Past Month (2018) |
|---------------------|----------------------------|----------------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| TOTAL | 402 | 413 | 113 | 123 | 78 | 100 |
| 12-20 | 32 | 29 | 23 | 23 | 17 | 11 |
| 21 or Older | 422 | 425 | 111 | 120 | 75 | 99 |
| 12-17 | 13 | 12 | 9 | 9 | 5 | 6 |
| 12-13 | 2 | 3 | 2 | 2 | 1 | 2 |
| 14-15 | 7 | 5 | 5 | 5 | 3 | 3 |
| 16-17 | 11 | 11 | 8 | 7 | 4 | 4 |
| 16-25 | 60 | 54 | 41 | 31 | 28 | 15 |
| 18 or Older | 401 | 411 | 113 | 122 | 77 | 100 |
| 18-25 | 58 | 52 | 39 | 30 | 27 | 15 |
| 18-20 | 28 | 26 | 20 | 21 | 16 | 9 |
| 21-25 | 52 | 47 | 33 | 21 | 22 | 11 |
| 26 or Older | 395 | 405 | 104 | 118 | 72 | 98 |
| 26-49 | 228 | 230 | 74 | 89 | 47 | 64 |
| 26-29 | 81 | 80 | 31 | 39 | 19 | 20 |
| 30-34 | 108 | 108 | 39 | 51 | 21 | 42 |
| 35-39 | 114 | 111 | 43 | 39 | 30 | 27 |
| 40-44 | 98 | 109 | 24 | 30 | 15 | 18 |
| 45-49 | 107 | 92 | 25 | 32 | 19 | 25 |
| 50 or Older | 308 | 322 | 65 | 83 | 53 | 75 |
| 50-54 | 156 | 174 | 43 | 38 | 29 | 38 |
| 55-59 | 179 | 178 | 36 | 37 | 31 | 28 |
| 60-64 | 161 | 171 | 33 | 51 | 31 | 44 |
| 65 or Older | 118 | 146 | 12 | 38 | 5 | 38 |

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 1.12D Methamphetamine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Detailed Age Category: Standard Errors of Percentages, 2017 and 2018

| Age Category | Lifetime (2017) | Lifetime (2018) | Past Year (2017) | Past Year (2018) | Past Month (2017) | Past Month (2018) |
|---------------------|----------------------------|----------------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| TOTAL | 0.15 | 0.15 | 0.04 | 0.04 | 0.03 | 0.04 |
| 12-20 | 0.08 | 0.08 | 0.06 | 0.06 | 0.05 | 0.03 |
| 21 or Older | 0.17 | 0.17 | 0.05 | 0.05 | 0.03 | 0.04 |
| 12-17 | 0.05 | 0.05 | 0.04 | 0.04 | 0.02 | 0.02 |
| 12-13 | 0.03 | 0.03 | 0.02 | 0.02 | 0.01 | 0.02 |
| 14-15 | 0.09 | 0.06 | 0.06 | 0.06 | 0.03 | 0.04 |
| 16-17 | 0.12 | 0.13 | 0.09 | 0.09 | 0.05 | 0.05 |
| 16-25 | 0.14 | 0.13 | 0.09 | 0.07 | 0.06 | 0.04 |
| 18 or Older | 0.16 | 0.17 | 0.05 | 0.05 | 0.03 | 0.04 |
| 18-25 | 0.17 | 0.15 | 0.11 | 0.09 | 0.08 | 0.04 |
| 18-20 | 0.22 | 0.20 | 0.16 | 0.16 | 0.13 | 0.07 |
| 21-25 | 0.23 | 0.22 | 0.15 | 0.10 | 0.10 | 0.05 |
| 26 or Older | 0.19 | 0.19 | 0.05 | 0.05 | 0.03 | 0.05 |
| 26-49 | 0.23 | 0.23 | 0.07 | 0.09 | 0.05 | 0.06 |
| 26-29 | 0.44 | 0.43 | 0.17 | 0.22 | 0.11 | 0.11 |
| 30-34 | 0.47 | 0.46 | 0.18 | 0.23 | 0.10 | 0.19 |
| 35-39 | 0.53 | 0.49 | 0.20 | 0.18 | 0.14 | 0.12 |
| 40-44 | 0.47 | 0.54 | 0.12 | 0.15 | 0.07 | 0.09 |
| 45-49 | 0.51 | 0.46 | 0.12 | 0.16 | 0.09 | 0.13 |
| 50 or Older | 0.27 | 0.28 | 0.06 | 0.07 | 0.05 | 0.07 |
| 50-54 | 0.70 | 0.79 | 0.20 | 0.18 | 0.14 | 0.18 |
| 55-59 | 0.78 | 0.77 | 0.16 | 0.17 | 0.14 | 0.13 |
| 60-64 | 0.76 | 0.79 | 0.16 | 0.25 | 0.15 | 0.22 |
| 65 or Older | 0.24 | 0.29 | 0.02 | 0.07 | 0.01 | 0.07 |

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 1.12P Methamphetamine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2018 Versus 2017

| Age Category | Lifetime (2017) | Lifetime (2018) | Past Year (2017) | Past Year (2018) | Past Month (2017) | Past Month (2018) |
|---------------------|----------------------------|----------------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| TOTAL | 0.8849 | | 0.1865 | | 0.0858 | |
| 12-20 | 0.3478 | | 0.9321 | | 0.2246 | |
| 21 or Older | 0.8345 | | 0.1693 | | 0.0522 | |
| 12-17 | 0.3731 | | 0.6811 | | 0.9177 | |
| 12-13 | 0.9956 | | 0.8860 | | 0.6742 | |
| 14-15 | 0.2309 | | 0.9036 | | 0.9534 | |
| 16-17 | 0.7412 | | 0.7074 | | 0.9678 | |
| 16-25 | 0.0528 | | 0.0505 | | 0.0529 | |
| 18 or Older | 0.8775 | | 0.1775 | | 0.0862 | |
| 18-25 | 0.0592 | | 0.0551 | | 0.0485 | |
| 18-20 | 0.4968 | | 0.9507 | | 0.1658 | |
| 21-25 | 0.0839 | | 0.0123 | | 0.1548 | |
| 26 or Older | 0.6947 | | 0.0400 | | 0.0243 | |
| 26-49 | 0.8008 | | 0.0164 | | 0.0136 | |
| 26-29 | 0.9911 | | 0.3919 | | 0.5984 | |
| 30-34 | 0.8913 | | 0.0842 | | 0.0140 | |
| 35-39 | 0.8414 | | 0.8548 | | 0.8719 | |
| 40-44 | 0.1511 | | 0.1356 | | 0.2399 | |
| 45-49 | 0.0644 | | 0.1576 | | 0.1680 | |
| 50 or Older | 0.5035 | | 0.5633 | | 0.3697 | |
| 50-54 | 0.7784 | | 0.5334 | | 0.7795 | |
| 55-59 | 0.6809 | | 0.7069 | | 0.9040 | |
| 60-64 | 0.8802 | | 0.3974 | | 0.7156 | |
| 65 or Older | 0.3602 | | 0.3792 | | 0.1860 | |

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 1.13C Any Use of Pain Relievers in Past Year and Misuse of Pain Relievers in Past Year and Past Month among Persons Aged 12 or Older, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2017 and 2018

| Age Category | Any Use in Past Year (2017) | Any Use in Past Year (2018) | Misuse in Past Year (2017) | Misuse in Past Year (2018) | Misuse in Past Month (2017) | Misuse in Past Month (2018) |
|--------------------|--------------------------------|--------------------------------|-------------------------------|-------------------------------|--------------------------------|--------------------------------|
| TOTAL | 781 | 787 | 274 | 286 | 165 | 158 |
| 12-20 | 167 | 156 | 67 | 61 | 33 | 31 |
| 21 or Older | 1,058 | 996 | 278 | 288 | 162 | 157 |
| 12-17 | 94 | 90 | 41 | 42 | 22 | 21 |
| 12-13 | 46 | 46 | 15 | 17 | 8 | 6 |
| 14-15 | 64 | 57 | 24 | 21 | 13 | 10 |
| 16-17 | 70 | 69 | 32 | 32 | 16 | 16 |
| 16-25 | 228 | 215 | 103 | 81 | 50 | 39 |
| 18 or Older | 774 | 776 | 270 | 284 | 163 | 157 |
| 18-25 | 156 | 148 | 88 | 72 | 46 | 37 |
| 18-20 | 122 | 114 | 52 | 45 | 24 | 23 |
| 21-25 | 166 | 155 | 83 | 60 | 42 | 29 |
| 26 or Older | 754 | 765 | 249 | 275 | 154 | 154 |
| 26-49 | 400 | 387 | 172 | 173 | 96 | 98 |
| 26-29 | 188 | 191 | 81 | 76 | 43 | 40 |
| 30-34 | 213 | 209 | 86 | 85 | 56 | 46 |
| 35-39 | 231 | 221 | 80 | 85 | 36 | 45 |
| 40-44 | 229 | 205 | 73 | 73 | 39 | 42 |
| 45-49 | 236 | 203 | 77 | 66 | 41 | 39 |
| 50 or Older | 652 | 636 | 193 | 220 | 123 | 117 |
| 50-54 | 340 | 353 | 100 | 110 | 57 | 54 |
| 55-59 | 338 | 362 | 96 | 115 | 70 | 60 |
| 60-64 | 341 | 361 | 91 | 95 | 48 | 51 |
| 65 or Older | 434 | 438 | 109 | 111 | 68 | 60 |

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 1.13D Any Use of Pain Relievers in Past Year and Misuse of Pain Relievers in Past Year and Past Month among Persons Aged 12 or Older, by Detailed Age Category: Standard Errors of Percentages, 2017 and 2018

| Age Category | Any Use in Past Year (2017) | Any Use in Past Year (2018) | Misuse in Past Year (2017) | Misuse in Past Year (2018) | Misuse in Past Month (2017) | Misuse in Past Month (2018) |
|--------------------|--------------------------------|--------------------------------|-------------------------------|-------------------------------|--------------------------------|--------------------------------|
| TOTAL | 0.29 | 0.29 | 0.10 | 0.10 | 0.06 | 0.06 |
| 12-20 | 0.35 | 0.34 | 0.17 | 0.16 | 0.09 | 0.08 |
| 21 or Older | 0.33 | 0.33 | 0.12 | 0.12 | 0.07 | 0.07 |
| 12-17 | 0.38 | 0.36 | 0.16 | 0.17 | 0.09 | 0.09 |
| 12-13 | 0.55 | 0.55 | 0.20 | 0.22 | 0.11 | 0.08 |
| 14-15 | 0.67 | 0.60 | 0.27 | 0.25 | 0.16 | 0.12 |
| 16-17 | 0.68 | 0.70 | 0.36 | 0.37 | 0.18 | 0.19 |
| 16-25 | 0.39 | 0.37 | 0.22 | 0.18 | 0.11 | 0.09 |
| 18 or Older | 0.31 | 0.31 | 0.11 | 0.11 | 0.07 | 0.06 |
| 18-25 | 0.45 | 0.44 | 0.26 | 0.21 | 0.13 | 0.11 |
| 18-20 | 0.70 | 0.69 | 0.39 | 0.33 | 0.19 | 0.18 |
| 21-25 | 0.58 | 0.56 | 0.35 | 0.27 | 0.19 | 0.14 |
| 26 or Older | 0.35 | 0.36 | 0.12 | 0.13 | 0.07 | 0.07 |
| 26-49 | 0.40 | 0.38 | 0.17 | 0.17 | 0.10 | 0.10 |
| 26-29 | 0.86 | 0.89 | 0.43 | 0.41 | 0.23 | 0.22 |
| 30-34 | 0.81 | 0.78 | 0.39 | 0.36 | 0.26 | 0.20 |
| 35-39 | 0.91 | 0.83 | 0.37 | 0.38 | 0.17 | 0.21 |
| 40-44 | 0.90 | 0.89 | 0.36 | 0.37 | 0.19 | 0.21 |
| 45-49 | 0.92 | 0.85 | 0.37 | 0.34 | 0.20 | 0.20 |
| 50 or Older | 0.58 | 0.56 | 0.17 | 0.19 | 0.11 | 0.10 |
| 50-54 | 1.27 | 1.39 | 0.46 | 0.52 | 0.27 | 0.26 |
| 55-59 | 1.25 | 1.36 | 0.43 | 0.52 | 0.32 | 0.28 |
| 60-64 | 1.36 | 1.39 | 0.46 | 0.45 | 0.24 | 0.25 |
| 65 or Older | 0.88 | 0.86 | 0.22 | 0.22 | 0.14 | 0.12 |

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 1.13P Any Use of Pain Relievers in Past Year and Misuse of Pain Relievers in Past Year and Past Month among Persons Aged 12 or Older, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2018 Versus 2017

| Age Category | Any Use in Past Year (2017) | Any Use in Past Year (2018) | Misuse in Past Year (2017) | Misuse in Past Year (2018) | Misuse in Past Month (2017) | Misuse in Past Month (2018) |
|--------------------|--------------------------------|--------------------------------|-------------------------------|-------------------------------|--------------------------------|--------------------------------|
| TOTAL | 0.0000 | | 0.0040 | | 0.0822 | |
| 12-20 | 0.0114 | | 0.0764 | | 0.1303 | |
| 21 or Older | 0.0001 | | 0.0110 | | 0.1431 | |
| 12-17 | 0.1375 | | 0.2358 | | 0.0899 | |
| 12-13 | 0.9650 | | 0.6733 | | 0.4473 | |
| 14-15 | 0.0187 | | 0.4647 | | 0.1078 | |
| 16-17 | 0.8832 | | 0.1963 | | 0.4407 | |
| 16-25 | 0.0000 | | 0.0000 | | 0.0052 | |
| 18 or Older | 0.0001 | | 0.0061 | | 0.1260 | |
| 18-25 | 0.0000 | | 0.0000 | | 0.0072 | |
| 18-20 | 0.0218 | | 0.1609 | | 0.5844 | |
| 21-25 | 0.0000 | | 0.0000 | | 0.0052 | |
| 26 or Older | 0.0024 | | 0.1707 | | 0.3722 | |
| 26-49 | 0.0000 | | 0.1261 | | 0.7124 | |
| 26-29 | 0.1641 | | 0.4935 | | 0.5246 | |
| 30-34 | 0.0165 | | 0.3217 | | 0.0617 | |
| 35-39 | 0.0116 | | 0.6604 | | 0.3408 | |
| 40-44 | 0.0379 | | 0.8726 | | 0.5255 | |
| 45-49 | 0.3784 | | 0.1471 | | 0.6195 | |
| 50 or Older | 0.2555 | | 0.6653 | | 0.3878 | |
| 50-54 | 0.7649 | | 0.9624 | | 0.8397 | |
| 55-59 | 0.6051 | | 0.4254 | | 0.5863 | |
| 60-64 | 0.1014 | | 0.4038 | | 0.7767 | |
| 65 or Older | 0.6218 | | 0.4426 | | 0.5727 | |

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 1.14C Any Use of Stimulants in Past Year and Misuse of Stimulants in Past Year and Past Month among Persons Aged 12 or Older, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2017 and 2018

| Age Category | Any Use in Past Year (2017) | Any Use in Past Year (2018) | Misuse in Past Year (2017) | Misuse in Past Year (2018) | Misuse in Past Month (2017) | Misuse in Past Month (2018) |
|--------------------|--------------------------------|--------------------------------|-------------------------------|-------------------------------|--------------------------------|--------------------------------|
| TOTAL | 369 | 375 | 191 | 178 | 109 | 103 |
| 12-20 | 114 | 96 | 70 | 56 | 34 | 25 |
| 21 or Older | 372 | 370 | 183 | 165 | 101 | 101 |
| 12-17 | 62 | 62 | 34 | 27 | 17 | 16 |
| 12-13 | 27 | 31 | 8 | 7 | 4 | 2 |
| 14-15 | 35 | 36 | 18 | 16 | 10 | 8 |
| 16-17 | 46 | 45 | 28 | 22 | 13 | 14 |
| 16-25 | 164 | 158 | 113 | 105 | 60 | 53 |
| 18 or Older | 366 | 368 | 190 | 175 | 108 | 102 |
| 18-25 | 129 | 127 | 99 | 93 | 56 | 49 |
| 18-20 | 93 | 74 | 60 | 49 | 29 | 20 |
| 21-25 | 122 | 122 | 88 | 85 | 47 | 46 |
| 26 or Older | 343 | 329 | 153 | 134 | 89 | 85 |
| 26-49 | 234 | 227 | 132 | 124 | 78 | 77 |
| 26-29 | 125 | 117 | 89 | 76 | 46 | 44 |
| 30-34 | 109 | 109 | 65 | 65 | 44 | 42 |
| 35-39 | 93 | 109 | 51 | 58 | 32 | 44 |
| 40-44 | 89 | 91 | 44 | 36 | 27 | 13 |
| 45-49 | 97 | 80 | 36 | 33 | 22 | 19 |
| 50 or Older | 242 | 245 | 66 | 45 | 38 | 37 |
| 50-54 | 134 | 121 | 40 | 30 | 32 | 18 |
| 55-59 | 113 | 121 | 44 | 15 | 16 | 15 |
| 60-64 | 107 | 110 | 22 | 23 | 11 | 23 |
| 65 or Older | 144 | 138 | 9 | 20 | 7 | 16 |

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 1.14D Any Use of Stimulants in Past Year and Misuse of Stimulants in Past Year and Past Month among Persons Aged 12 or Older, by Detailed Age Category: Standard Errors of Percentages, 2017 and 2018

| Age Category | Any Use in Past Year (2017) | Any Use in Past Year (2018) | Misuse in Past Year (2017) | Misuse in Past Year (2018) | Misuse in Past Month (2017) | Misuse in Past Month (2018) |
|--------------------|--------------------------------|--------------------------------|-------------------------------|-------------------------------|--------------------------------|--------------------------------|
| TOTAL | 0.14 | 0.14 | 0.07 | 0.06 | 0.04 | 0.04 |
| 12-20 | 0.27 | 0.24 | 0.18 | 0.14 | 0.09 | 0.07 |
| 21 or Older | 0.15 | 0.15 | 0.08 | 0.07 | 0.04 | 0.04 |
| 12-17 | 0.25 | 0.25 | 0.14 | 0.11 | 0.07 | 0.06 |
| 12-13 | 0.34 | 0.38 | 0.10 | 0.09 | 0.05 | 0.03 |
| 14-15 | 0.39 | 0.41 | 0.21 | 0.19 | 0.11 | 0.09 |
| 16-17 | 0.50 | 0.50 | 0.32 | 0.25 | 0.16 | 0.16 |
| 16-25 | 0.32 | 0.31 | 0.24 | 0.22 | 0.14 | 0.12 |
| 18 or Older | 0.15 | 0.15 | 0.08 | 0.07 | 0.04 | 0.04 |
| 18-25 | 0.38 | 0.37 | 0.29 | 0.27 | 0.16 | 0.14 |
| 18-20 | 0.60 | 0.50 | 0.42 | 0.35 | 0.22 | 0.15 |
| 21-25 | 0.48 | 0.49 | 0.38 | 0.36 | 0.21 | 0.21 |
| 26 or Older | 0.16 | 0.15 | 0.07 | 0.06 | 0.04 | 0.04 |
| 26-49 | 0.23 | 0.22 | 0.13 | 0.12 | 0.08 | 0.08 |
| 26-29 | 0.63 | 0.61 | 0.47 | 0.42 | 0.25 | 0.25 |
| 30-34 | 0.47 | 0.47 | 0.29 | 0.29 | 0.20 | 0.19 |
| 35-39 | 0.42 | 0.48 | 0.24 | 0.26 | 0.15 | 0.20 |
| 40-44 | 0.44 | 0.44 | 0.22 | 0.18 | 0.13 | 0.07 |
| 45-49 | 0.46 | 0.41 | 0.18 | 0.17 | 0.11 | 0.10 |
| 50 or Older | 0.21 | 0.22 | 0.06 | 0.04 | 0.03 | 0.03 |
| 50-54 | 0.62 | 0.57 | 0.19 | 0.14 | 0.15 | 0.09 |
| 55-59 | 0.50 | 0.55 | 0.20 | 0.07 | 0.07 | 0.07 |
| 60-64 | 0.52 | 0.52 | 0.11 | 0.11 | 0.05 | 0.11 |
| 65 or Older | 0.29 | 0.27 | 0.02 | 0.04 | 0.01 | 0.03 |

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 1.14P Any Use of Stimulants in Past Year and Misuse of Stimulants in Past Year and Past Month among Persons Aged 12 or Older, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2018 Versus 2017

| Age Category | Any Use in Past Year (2017) | Any Use in Past Year (2018) | Misuse in Past Year (2017) | Misuse in Past Year (2018) | Misuse in Past Month (2017) | Misuse in Past Month (2018) |
|--------------------|--------------------------------|--------------------------------|-------------------------------|-------------------------------|--------------------------------|--------------------------------|
| TOTAL | 0.1563 | | 0.0023 | | 0.1976 | |
| 12-20 | 0.0541 | | 0.0106 | | 0.0099 | |
| 21 or Older | 0.3509 | | 0.0236 | | 0.6132 | |
| 12-17 | 0.7198 | | 0.0597 | | 0.9043 | |
| 12-13 | 0.2205 | | 0.8666 | | 0.6056 | |
| 14-15 | 0.2385 | | 0.6183 | | 0.4553 | |
| 16-17 | 0.6904 | | 0.0547 | | 0.6435 | |
| 16-25 | 0.0025 | | 0.0044 | | 0.1087 | |
| 18 or Older | 0.1728 | | 0.0062 | | 0.1991 | |
| 18-25 | 0.0026 | | 0.0154 | | 0.0712 | |
| 18-20 | 0.0202 | | 0.0392 | | 0.0018 | |
| 21-25 | 0.0441 | | 0.1378 | | 0.8848 | |
| 26 or Older | 0.7850 | | 0.0911 | | 0.6887 | |
| 26-49 | 0.8825 | | 0.2798 | | 0.7076 | |
| 26-29 | 0.4670 | | 0.1496 | | 0.9631 | |
| 30-34 | 0.5005 | | 0.3149 | | 0.9098 | |
| 35-39 | 0.1972 | | 0.9916 | | 0.6869 | |
| 40-44 | 0.7746 | | 0.1343 | | 0.0515 | |
| 45-49 | 0.1738 | | 0.3209 | | 0.8273 | |
| 50 or Older | 0.6391 | | 0.0969 | | 0.9163 | |
| 50-54 | 0.2468 | | 0.1847 | | 0.2116 | |
| 55-59 | 0.4255 | | 0.1361 | | 0.7475 | |
| 60-64 | 0.5298 | | 0.5371 | | 0.4429 | |
| 65 or Older | 0.9003 | | 0.3335 | | 0.3891 | |

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 1.15C Any Use of Tranquilizers or Sedatives in Past Year and Misuse of Tranquilizers or Sedatives in Past Year and Past Month among Persons Aged 12 or Older, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2017 and 2018

| Age Category | Any Use in Past Year (2017) | Any Use in Past Year (2018) | Misuse in Past Year (2017) | Misuse in Past Year (2018) | Misuse in Past Month (2017) | Misuse in Past Month (2018) |
|--------------------|--------------------------------|--------------------------------|-------------------------------|-------------------------------|--------------------------------|--------------------------------|
| TOTAL | 671 | 672 | 227 | 239 | 112 | 125 |
| 12-20 | 97 | 93 | 58 | 57 | 30 | 26 |
| 21 or Older | 758 | 767 | 221 | 234 | 109 | 125 |
| 12-17 | 56 | 56 | 33 | 34 | 16 | 12 |
| 12-13 | 25 | 26 | 9 | 8 | 4 | 3 |
| 14-15 | 31 | 33 | 18 | 21 | 10 | 6 |
| 16-17 | 42 | 41 | 27 | 25 | 13 | 10 |
| 16-25 | 150 | 131 | 92 | 81 | 47 | 37 |
| 18 or Older | 663 | 663 | 222 | 237 | 111 | 124 |
| 18-25 | 119 | 110 | 81 | 73 | 45 | 35 |
| 18-20 | 76 | 70 | 47 | 45 | 25 | 23 |
| 21-25 | 113 | 103 | 70 | 61 | 39 | 29 |
| 26 or Older | 639 | 652 | 201 | 223 | 100 | 120 |
| 26-49 | 306 | 305 | 135 | 140 | 69 | 76 |
| 26-29 | 122 | 128 | 65 | 69 | 33 | 35 |
| 30-34 | 140 | 150 | 70 | 75 | 32 | 36 |
| 35-39 | 159 | 156 | 58 | 66 | 34 | 36 |
| 40-44 | 157 | 148 | 54 | 46 | 24 | 19 |
| 45-49 | 156 | 155 | 52 | 48 | 31 | 24 |
| 50 or Older | 558 | 569 | 145 | 172 | 75 | 92 |
| 50-54 | 246 | 277 | 72 | 85 | 40 | 40 |
| 55-59 | 257 | 275 | 79 | 106 | 50 | 25 |
| 60-64 | 246 | 285 | 53 | 63 | 25 | 47 |
| 65 or Older | 388 | 378 | 85 | 80 | 31 | 62 |

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 1.15D Any Use of Tranquilizers or Sedatives in Past Year and Misuse of Tranquilizers or Sedatives in Past Year and Past Month among Persons Aged 12 or Older, by Detailed Age Category: Standard Errors of Percentages, 2017 and 2018

| Age Category | Any Use in Past Year (2017) | Any Use in Past Year (2018) | Misuse in Past Year (2017) | Misuse in Past Year (2018) | Misuse in Past Month (2017) | Misuse in Past Month (2018) |
|--------------------|--------------------------------|--------------------------------|-------------------------------|-------------------------------|--------------------------------|--------------------------------|
| TOTAL | 0.25 | 0.25 | 0.08 | 0.09 | 0.04 | 0.05 |
| 12-20 | 0.23 | 0.24 | 0.15 | 0.15 | 0.08 | 0.07 |
| 21 or Older | 0.28 | 0.28 | 0.09 | 0.10 | 0.05 | 0.05 |
| 12-17 | 0.23 | 0.22 | 0.13 | 0.14 | 0.06 | 0.05 |
| 12-13 | 0.32 | 0.32 | 0.11 | 0.11 | 0.05 | 0.03 |
| 14-15 | 0.36 | 0.38 | 0.21 | 0.24 | 0.11 | 0.07 |
| 16-17 | 0.47 | 0.46 | 0.30 | 0.29 | 0.15 | 0.12 |
| 16-25 | 0.29 | 0.27 | 0.20 | 0.18 | 0.11 | 0.09 |
| 18 or Older | 0.27 | 0.27 | 0.09 | 0.10 | 0.04 | 0.05 |
| 18-25 | 0.35 | 0.32 | 0.24 | 0.21 | 0.13 | 0.10 |
| 18-20 | 0.51 | 0.51 | 0.35 | 0.34 | 0.19 | 0.17 |
| 21-25 | 0.45 | 0.42 | 0.30 | 0.27 | 0.18 | 0.13 |
| 26 or Older | 0.30 | 0.30 | 0.09 | 0.10 | 0.05 | 0.06 |
| 26-49 | 0.31 | 0.30 | 0.13 | 0.14 | 0.07 | 0.08 |
| 26-29 | 0.62 | 0.65 | 0.35 | 0.38 | 0.18 | 0.19 |
| 30-34 | 0.60 | 0.61 | 0.32 | 0.33 | 0.15 | 0.16 |
| 35-39 | 0.69 | 0.65 | 0.28 | 0.30 | 0.16 | 0.17 |
| 40-44 | 0.71 | 0.71 | 0.27 | 0.24 | 0.12 | 0.10 |
| 45-49 | 0.70 | 0.71 | 0.26 | 0.24 | 0.15 | 0.12 |
| 50 or Older | 0.50 | 0.50 | 0.13 | 0.15 | 0.07 | 0.08 |
| 50-54 | 1.03 | 1.21 | 0.34 | 0.41 | 0.19 | 0.19 |
| 55-59 | 1.07 | 1.14 | 0.36 | 0.48 | 0.23 | 0.12 |
| 60-64 | 1.12 | 1.23 | 0.26 | 0.30 | 0.13 | 0.23 |
| 65 or Older | 0.78 | 0.74 | 0.17 | 0.16 | 0.06 | 0.12 |

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 1.15P Any Use of Tranquilizers or Sedatives in Past Year and Misuse of Tranquilizers or Sedatives in Past Year and Past Month among Persons Aged 12 or Older, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2018 Versus 2017

| Age Category | Any Use in Past Year (2017) | Any Use in Past Year (2018) | Misuse in Past Year (2017) | Misuse in Past Year (2018) | Misuse in Past Month (2017) | Misuse in Past Month (2018) |
|--------------------|--------------------------------|--------------------------------|-------------------------------|-------------------------------|--------------------------------|--------------------------------|
| TOTAL | 0.5396 | | 0.2833 | | 0.3214 | |
| 12-20 | 0.0182 | | 0.3769 | | 0.0934 | |
| 21 or Older | 0.7606 | | 0.3708 | | 0.5364 | |
| 12-17 | 0.0834 | | 0.5158 | | 0.0117 | |
| 12-13 | 0.3247 | | 0.6977 | | 0.6686 | |
| 14-15 | 0.7015 | | 0.4557 | | 0.3210 | |
| 16-17 | 0.0435 | | 0.1962 | | 0.0196 | |
| 16-25 | 0.0044 | | 0.0050 | | 0.0013 | |
| 18 or Older | 0.6150 | | 0.3154 | | 0.4875 | |
| 18-25 | 0.0231 | | 0.0110 | | 0.0075 | |
| 18-20 | 0.0604 | | 0.4586 | | 0.6475 | |
| 21-25 | 0.1386 | | 0.0068 | | 0.0038 | |
| 26 or Older | 0.9006 | | 0.9332 | | 0.8114 | |
| 26-49 | 0.3520 | | 0.6031 | | 0.8467 | |
| 26-29 | 0.5984 | | 0.8702 | | 0.8767 | |
| 30-34 | 0.4330 | | 0.6034 | | 0.9476 | |
| 35-39 | 0.0540 | | 0.3087 | | 0.3597 | |
| 40-44 | 0.3281 | | 0.6772 | | 0.3412 | |
| 45-49 | 0.9366 | | 0.5603 | | 0.3324 | |
| 50 or Older | 0.7450 | | 0.6067 | | 0.6313 | |
| 50-54 | 0.3615 | | 0.8720 | | 0.4957 | |
| 55-59 | 0.9558 | | 0.9825 | | 0.0808 | |
| 60-64 | 0.7016 | | 0.8874 | | 0.0562 | |
| 65 or Older | 0.8488 | | 0.4282 | | 0.1621 | |

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 1.16C Any Use of Tranquilizers in Past Year and Misuse of Tranquilizers in Past Year and Past Month among Persons Aged 12 or Older, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2017 and 2018

| Age Category | Any Use in Past Year (2017) | Any Use in Past Year (2018) | Misuse in Past Year (2017) | Misuse in Past Year (2018) | Misuse in Past Month (2017) | Misuse in Past Month (2018) |
|--------------------|--------------------------------|--------------------------------|-------------------------------|-------------------------------|--------------------------------|--------------------------------|
| TOTAL | 615 | 604 | 209 | 226 | 102 | 119 |
| 12-20 | 90 | 83 | 57 | 56 | 28 | 25 |
| 21 or Older | 671 | 666 | 201 | 220 | 99 | 118 |
| 12-17 | 50 | 48 | 32 | 33 | 15 | 12 |
| 12-13 | 18 | 19 | 8 | 6 | 4 | 3 |
| 14-15 | 29 | 28 | 18 | 20 | 9 | 6 |
| 16-17 | 39 | 36 | 26 | 24 | 13 | 10 |
| 16-25 | 140 | 122 | 91 | 80 | 46 | 37 |
| 18 or Older | 607 | 599 | 205 | 223 | 101 | 118 |
| 18-25 | 112 | 102 | 80 | 72 | 44 | 35 |
| 18-20 | 73 | 64 | 46 | 45 | 24 | 22 |
| 21-25 | 102 | 95 | 69 | 61 | 38 | 28 |
| 26 or Older | 585 | 588 | 182 | 211 | 89 | 114 |
| 26-49 | 286 | 287 | 127 | 133 | 64 | 74 |
| 26-29 | 112 | 117 | 63 | 67 | 32 | 34 |
| 30-34 | 128 | 139 | 65 | 72 | 31 | 36 |
| 35-39 | 146 | 144 | 52 | 60 | 28 | 35 |
| 40-44 | 139 | 131 | 50 | 43 | 23 | 17 |
| 45-49 | 141 | 139 | 49 | 46 | 30 | 24 |
| 50 or Older | 510 | 510 | 125 | 161 | 65 | 86 |
| 50-54 | 221 | 247 | 58 | 81 | 36 | 38 |
| 55-59 | 234 | 251 | 71 | 101 | 43 | 25 |
| 60-64 | 208 | 237 | 40 | 52 | 15 | 46 |
| 65 or Older | 357 | 342 | 76 | 74 | 31 | 56 |

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 1.16D Any Use of Tranquilizers in Past Year and Misuse of Tranquilizers in Past Year and Past Month among Persons Aged 12 or Older, by Detailed Age Category: Standard Errors of Percentages, 2017 and 2018

| Age Category | Any Use in Past Year (2017) | Any Use in Past Year (2018) | Misuse in Past Year (2017) | Misuse in Past Year (2018) | Misuse in Past Month (2017) | Misuse in Past Month (2018) |
|--------------------|--------------------------------|--------------------------------|-------------------------------|-------------------------------|--------------------------------|--------------------------------|
| TOTAL | 0.23 | 0.22 | 0.08 | 0.08 | 0.04 | 0.04 |
| 12-20 | 0.22 | 0.21 | 0.15 | 0.15 | 0.07 | 0.07 |
| 21 or Older | 0.26 | 0.25 | 0.08 | 0.09 | 0.04 | 0.05 |
| 12-17 | 0.20 | 0.19 | 0.13 | 0.13 | 0.06 | 0.05 |
| 12-13 | 0.23 | 0.23 | 0.10 | 0.07 | 0.05 | 0.03 |
| 14-15 | 0.34 | 0.32 | 0.21 | 0.23 | 0.11 | 0.07 |
| 16-17 | 0.43 | 0.41 | 0.30 | 0.28 | 0.15 | 0.11 |
| 16-25 | 0.27 | 0.25 | 0.20 | 0.18 | 0.11 | 0.09 |
| 18 or Older | 0.25 | 0.24 | 0.08 | 0.09 | 0.04 | 0.05 |
| 18-25 | 0.33 | 0.30 | 0.23 | 0.21 | 0.13 | 0.10 |
| 18-20 | 0.50 | 0.47 | 0.34 | 0.33 | 0.18 | 0.17 |
| 21-25 | 0.41 | 0.39 | 0.30 | 0.27 | 0.18 | 0.13 |
| 26 or Older | 0.27 | 0.27 | 0.09 | 0.10 | 0.04 | 0.05 |
| 26-49 | 0.28 | 0.28 | 0.13 | 0.13 | 0.06 | 0.07 |
| 26-29 | 0.57 | 0.61 | 0.34 | 0.37 | 0.18 | 0.19 |
| 30-34 | 0.55 | 0.58 | 0.30 | 0.31 | 0.14 | 0.16 |
| 35-39 | 0.66 | 0.62 | 0.25 | 0.27 | 0.13 | 0.16 |
| 40-44 | 0.64 | 0.64 | 0.25 | 0.22 | 0.12 | 0.09 |
| 45-49 | 0.65 | 0.65 | 0.24 | 0.23 | 0.15 | 0.12 |
| 50 or Older | 0.45 | 0.45 | 0.11 | 0.14 | 0.06 | 0.08 |
| 50-54 | 0.96 | 1.10 | 0.27 | 0.39 | 0.17 | 0.18 |
| 55-59 | 0.99 | 1.06 | 0.32 | 0.46 | 0.20 | 0.12 |
| 60-64 | 0.97 | 1.04 | 0.20 | 0.25 | 0.07 | 0.22 |
| 65 or Older | 0.72 | 0.67 | 0.15 | 0.14 | 0.06 | 0.11 |

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 1.16P Any Use of Tranquilizers in Past Year and Misuse of Tranquilizers in Past Year and Past Month among Persons Aged 12 or Older, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2018 Versus 2017

| Age Category | Any Use in Past Year (2017) | Any Use in Past Year (2018) | Misuse in Past Year (2017) | Misuse in Past Year (2018) | Misuse in Past Month (2017) | Misuse in Past Month (2018) |
|--------------------|-----------------------------------|-----------------------------------|----------------------------------|----------------------------------|-----------------------------------|-----------------------------------|
| TOTAL | 0.3529 | | 0.4415 | | 0.5128 | |
| 12-20 | 0.0072 | | 0.3073 | | 0.0780 | |
| 21 or Older | 0.5634 | | 0.5990 | | 0.8054 | |
| 12-17 | 0.0540 | | 0.4647 | | 0.0153 | |
| 12-13 | 0.6377 | | 0.3544 | | 0.9187 | |
| 14-15 | 0.9884 | | 0.3026 | | 0.3616 | |
| 16-17 | 0.0208 | | 0.1341 | | 0.0187 | |
| 16-25 | 0.0017 | | 0.0019 | | 0.0008 | |
| 18 or Older | 0.4201 | | 0.4995 | | 0.7178 | |
| 18-25 | 0.0116 | | 0.0054 | | 0.0055 | |
| 18-20 | 0.0321 | | 0.3911 | | 0.5454 | |
| 21-25 | 0.1152 | | 0.0034 | | 0.0036 | |
| 26 or Older | 0.7065 | | 0.6785 | | 0.4776 | |
| 26-49 | 0.2227 | | 0.5071 | | 0.8174 | |
| 26-29 | 0.6035 | | 0.8966 | | 0.8613 | |
| 30-34 | 0.5075 | | 0.4666 | | 0.8293 | |
| 35-39 | 0.0779 | | 0.2391 | | 0.0878 | |
| 40-44 | 0.1452 | | 0.5195 | | 0.3391 | |
| 45-49 | 0.7471 | | 0.7549 | | 0.4103 | |
| 50 or Older | 0.8264 | | 1.0000 | | 0.4687 | |
| 50-54 | 0.1992 | | 0.7217 | | 0.5165 | |
| 55-59 | 0.7539 | | 0.8416 | | 0.3332 | |
| 60-64 | 0.8059 | | 0.4676 | | 0.0227 | |
| 65 or Older | 0.6130 | | 0.3638 | | 0.3661 | |

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 1.17C Any Use of Sedatives in Past Year and Misuse of Sedatives in Past Year and Past Month among Persons Aged 12 or Older, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2017 and 2018

| Age Category | Any Use in Past Year (2017) | Any Use in Past Year (2018) | Misuse in Past Year (2017) | Misuse in Past Year (2018) | Misuse in Past Month (2017) | Misuse in Past Month (2018) |
|--------------------|--------------------------------|--------------------------------|-------------------------------|-------------------------------|--------------------------------|--------------------------------|
| TOTAL | 421 | 442 | 117 | 96 | 55 | 43 |
| 12-20 | 51 | 48 | 18 | 18 | 10 | 8 |
| 21 or Older | 447 | 465 | 116 | 92 | 54 | 43 |
| 12-17 | 36 | 32 | 13 | 13 | 6 | 4 |
| 12-13 | 18 | 18 | 5 | 7 | 2 | 1 |
| 14-15 | 19 | 21 | 8 | 7 | 4 | 3 |
| 16-17 | 23 | 21 | 9 | 9 | 4 | 3 |
| 16-25 | 73 | 63 | 26 | 27 | 13 | 11 |
| 18 or Older | 420 | 437 | 116 | 94 | 55 | 43 |
| 18-25 | 64 | 58 | 24 | 24 | 13 | 11 |
| 18-20 | 35 | 34 | 12 | 13 | 8 | 7 |
| 21-25 | 56 | 48 | 20 | 20 | 10 | 9 |
| 26 or Older | 413 | 430 | 112 | 89 | 52 | 41 |
| 26-49 | 180 | 188 | 58 | 60 | 30 | 22 |
| 26-29 | 70 | 69 | 24 | 26 | 10 | 9 |
| 30-34 | 77 | 84 | 29 | 30 | 11 | 11 |
| 35-39 | 82 | 92 | 29 | 32 | 24 | 15 |
| 40-44 | 84 | 95 | 26 | 23 | 10 | 8 |
| 45-49 | 97 | 96 | 22 | 17 | 8 | 4 |
| 50 or Older | 370 | 386 | 94 | 66 | 43 | 35 |
| 50-54 | 157 | 172 | 55 | 27 | 28 | 13 |
| 55-59 | 167 | 165 | 43 | 36 | 25 | 14 |
| 60-64 | 171 | 161 | 44 | 36 | 20 | 11 |
| 65 or Older | 262 | 271 | 40 | 32 | * | 27 |

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 1.17D Any Use of Sedatives in Past Year and Misuse of Sedatives in Past Year and Past Month among Persons Aged 12 or Older, by Detailed Age Category: Standard Errors of Percentages, 2017 and 2018

| Age Category | Any Use in Past Year (2017) | Any Use in Past Year (2018) | Misuse in Past Year (2017) | Misuse in Past Year (2018) | Misuse in Past Month (2017) | Misuse in Past Month (2018) |
|--------------------|--------------------------------|--------------------------------|-------------------------------|-------------------------------|--------------------------------|--------------------------------|
| TOTAL | 0.15 | 0.16 | 0.04 | 0.04 | 0.02 | 0.02 |
| 12-20 | 0.13 | 0.13 | 0.05 | 0.05 | 0.03 | 0.02 |
| 21 or Older | 0.18 | 0.18 | 0.05 | 0.04 | 0.02 | 0.02 |
| 12-17 | 0.14 | 0.13 | 0.05 | 0.05 | 0.02 | 0.02 |
| 12-13 | 0.24 | 0.23 | 0.06 | 0.09 | 0.02 | 0.01 |
| 14-15 | 0.22 | 0.25 | 0.10 | 0.08 | 0.04 | 0.03 |
| 16-17 | 0.26 | 0.24 | 0.10 | 0.10 | 0.05 | 0.04 |
| 16-25 | 0.16 | 0.15 | 0.06 | 0.06 | 0.03 | 0.03 |
| 18 or Older | 0.17 | 0.18 | 0.05 | 0.04 | 0.02 | 0.02 |
| 18-25 | 0.19 | 0.17 | 0.07 | 0.07 | 0.04 | 0.03 |
| 18-20 | 0.26 | 0.26 | 0.09 | 0.10 | 0.06 | 0.05 |
| 21-25 | 0.24 | 0.22 | 0.09 | 0.10 | 0.05 | 0.04 |
| 26 or Older | 0.19 | 0.20 | 0.05 | 0.04 | 0.02 | 0.02 |
| 26-49 | 0.18 | 0.19 | 0.06 | 0.06 | 0.03 | 0.02 |
| 26-29 | 0.38 | 0.37 | 0.13 | 0.15 | 0.06 | 0.05 |
| 30-34 | 0.35 | 0.36 | 0.13 | 0.13 | 0.05 | 0.05 |
| 35-39 | 0.38 | 0.41 | 0.14 | 0.15 | 0.11 | 0.07 |
| 40-44 | 0.42 | 0.48 | 0.13 | 0.12 | 0.05 | 0.04 |
| 45-49 | 0.46 | 0.47 | 0.11 | 0.09 | 0.04 | 0.02 |
| 50 or Older | 0.33 | 0.34 | 0.08 | 0.06 | 0.04 | 0.03 |
| 50-54 | 0.71 | 0.81 | 0.26 | 0.13 | 0.13 | 0.06 |
| 55-59 | 0.73 | 0.73 | 0.20 | 0.17 | 0.11 | 0.07 |
| 60-64 | 0.82 | 0.77 | 0.22 | 0.17 | 0.10 | 0.06 |
| 65 or Older | 0.53 | 0.53 | 0.08 | 0.06 | * | 0.05 |

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 1.17P Any Use of Sedatives in Past Year and Misuse of Sedatives in Past Year and Past Month among Persons Aged 12 or Older, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2018 Versus 2017

| Age Category | Any Use in Past Year (2017) | Any Use in Past Year (2018) | Misuse in Past Year (2017) | Misuse in Past Year (2018) | Misuse in Past Month (2017) | Misuse in Past Month (2018) |
|--------------------|--------------------------------|--------------------------------|-------------------------------|-------------------------------|--------------------------------|--------------------------------|
| TOTAL | 0.9097 | | 0.0626 | | 0.1101 | |
| 12-20 | 0.2681 | | 0.9403 | | 0.5466 | |
| 21 or Older | 0.9915 | | 0.0548 | | 0.1326 | |
| 12-17 | 0.2043 | | 0.5886 | | 0.1299 | |
| 12-13 | 0.4996 | | 0.5694 | | 0.3765 | |
| 14-15 | 0.3933 | | 0.3318 | | 0.5045 | |
| 16-17 | 0.0293 | | 0.7289 | | 0.2417 | |
| 16-25 | 0.0802 | | 0.8262 | | 0.4044 | |
| 18 or Older | 0.9798 | | 0.0691 | | 0.1456 | |
| 18-25 | 0.2776 | | 0.9125 | | 0.6117 | |
| 18-20 | 0.7185 | | 0.4978 | | 0.7991 | |
| 21-25 | 0.2861 | | 0.5889 | | 0.3909 | |
| 26 or Older | 0.9184 | | 0.0629 | | 0.1683 | |
| 26-49 | 0.8673 | | 0.7394 | | 0.2106 | |
| 26-29 | 0.8143 | | 0.7534 | | 0.5567 | |
| 30-34 | 0.7818 | | 0.5627 | | 0.9448 | |
| 35-39 | 0.6256 | | 0.7285 | | 0.3188 | |
| 40-44 | 0.2460 | | 0.9494 | | 0.4925 | |
| 45-49 | 0.8170 | | 0.3745 | | 0.4839 | |
| 50 or Older | 0.9886 | | 0.0113 | | 0.4215 | |
| 50-54 | 0.8110 | | 0.0074 | | 0.1304 | |
| 55-59 | 0.8755 | | 0.4319 | | 0.3198 | |
| 60-64 | 0.1195 | | 0.1910 | | 0.7646 | |
| 65 or Older | 0.3543 | | 0.8661 | | * | |

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 1.18C Any Use and Misuse of Benzodiazepines in Past Year among Persons Aged 12 or Older, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2017 and 2018

| Age Category | Any Use in Past Year (2017) | Any Use in Past Year (2018) | Misuse in Past Year (2017) | Misuse in Past Year (2018) |
|---------------------|--|--|---------------------------------------|---------------------------------------|
| TOTAL | 553 | 587 | 203 | 218 |
| 12-20 | 83 | 76 | 56 | 55 |
| 21 or Older | 588 | 633 | 195 | 213 |
| 12-17 | 45 | 41 | 32 | 32 |
| 12-13 | 16 | 12 | 8 | 5 |
| 14-15 | 26 | 25 | 17 | 19 |
| 16-17 | 35 | 32 | 26 | 24 |
| 16-25 | 132 | 116 | 88 | 79 |
| 18 or Older | 545 | 582 | 199 | 217 |
| 18-25 | 108 | 98 | 79 | 71 |
| 18-20 | 69 | 61 | 45 | 44 |
| 21-25 | 98 | 91 | 67 | 60 |
| 26 or Older | 524 | 569 | 177 | 203 |
| 26-49 | 266 | 265 | 123 | 129 |
| 26-29 | 106 | 109 | 61 | 65 |
| 30-34 | 120 | 126 | 62 | 71 |
| 35-39 | 130 | 134 | 51 | 59 |
| 40-44 | 129 | 120 | 47 | 43 |
| 45-49 | 129 | 126 | 50 | 44 |
| 50 or Older | 450 | 506 | 129 | 155 |
| 50-54 | 195 | 219 | 53 | 80 |
| 55-59 | 216 | 224 | 68 | 94 |
| 60-64 | 184 | 232 | 41 | 52 |
| 65 or Older | 307 | 329 | 84 | 71 |

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 1.18D Any Use and Misuse of Benzodiazepines in Past Year among Persons Aged 12 or Older, by Detailed Age Category: Standard Errors of Percentages, 2017 and 2018

| Age Category | Any Use in Past Year (2017) | Any Use in Past Year (2018) | Misuse in Past Year (2017) | Misuse in Past Year (2018) |
|---------------------|--|--|---------------------------------------|---------------------------------------|
| TOTAL | 0.20 | 0.21 | 0.07 | 0.08 |
| 12-20 | 0.21 | 0.20 | 0.14 | 0.15 |
| 21 or Older | 0.23 | 0.24 | 0.08 | 0.09 |
| 12-17 | 0.18 | 0.16 | 0.13 | 0.13 |
| 12-13 | 0.20 | 0.16 | 0.10 | 0.06 |
| 14-15 | 0.30 | 0.29 | 0.20 | 0.22 |
| 16-17 | 0.39 | 0.36 | 0.29 | 0.28 |
| 16-25 | 0.26 | 0.24 | 0.19 | 0.18 |
| 18 or Older | 0.22 | 0.23 | 0.08 | 0.09 |
| 18-25 | 0.32 | 0.29 | 0.23 | 0.21 |
| 18-20 | 0.48 | 0.44 | 0.34 | 0.33 |
| 21-25 | 0.40 | 0.38 | 0.29 | 0.27 |
| 26 or Older | 0.25 | 0.26 | 0.08 | 0.09 |
| 26-49 | 0.26 | 0.26 | 0.12 | 0.13 |
| 26-29 | 0.55 | 0.57 | 0.33 | 0.36 |
| 30-34 | 0.52 | 0.52 | 0.28 | 0.31 |
| 35-39 | 0.59 | 0.58 | 0.24 | 0.27 |
| 40-44 | 0.61 | 0.60 | 0.24 | 0.22 |
| 45-49 | 0.59 | 0.60 | 0.25 | 0.22 |
| 50 or Older | 0.40 | 0.44 | 0.11 | 0.14 |
| 50-54 | 0.86 | 0.99 | 0.25 | 0.39 |
| 55-59 | 0.93 | 0.96 | 0.31 | 0.43 |
| 60-64 | 0.87 | 1.03 | 0.21 | 0.25 |
| 65 or Older | 0.62 | 0.64 | 0.17 | 0.14 |

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 1.18P Any Use of and Misuse of Benzodiazepines in Past Year among Persons Aged 12 or Older, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2018 Versus 2017

| Age Category | Any Use in Past Year (2017) | Any Use in Past Year (2018) | Misuse in Past Year (2017) | Misuse in Past Year (2018) |
|---------------------|--|--|---------------------------------------|---------------------------------------|
| TOTAL | 0.2040 | | 0.3660 | |
| 12-20 | 0.0591 | | 0.2198 | |
| 21 or Older | 0.2986 | | 0.5442 | |
| 12-17 | 0.1550 | | 0.3697 | |
| 12-13 | 0.1055 | | 0.1657 | |
| 14-15 | 0.5070 | | 0.3960 | |
| 16-17 | 0.1031 | | 0.1642 | |
| 16-25 | 0.0007 | | 0.0026 | |
| 18 or Older | 0.2292 | | 0.4330 | |
| 18-25 | 0.0025 | | 0.0064 | |
| 18-20 | 0.1191 | | 0.3139 | |
| 21-25 | 0.0067 | | 0.0061 | |
| 26 or Older | 0.5161 | | 0.7705 | |
| 26-49 | 0.2169 | | 0.4914 | |
| 26-29 | 0.6224 | | 0.9264 | |
| 30-34 | 0.7936 | | 0.4750 | |
| 35-39 | 0.0848 | | 0.2191 | |
| 40-44 | 0.2822 | | 0.8195 | |
| 45-49 | 0.9227 | | 0.5122 | |
| 50 or Older | 0.9535 | | 0.8518 | |
| 50-54 | 0.2608 | | 0.4600 | |
| 55-59 | 0.8168 | | 1.0000 | |
| 60-64 | 0.9022 | | 0.5442 | |
| 65 or Older | 0.4822 | | 0.1963 | |

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 1.19C Misuse of Opioids in Past Year and Past Month among Persons Aged 12 or Older, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2017 and 2018

| Age Category | Misuse in Past Year (2017) | Misuse in Past Year (2018) | Misuse in Past Month (2017) | Misuse in Past Month (2018) |
|---------------------|---------------------------------------|---------------------------------------|--|--|
| TOTAL | 279 | 294 | 176 | 162 |
| 12-20 | 68 | 62 | 34 | 32 |
| 21 or Older | 284 | 297 | 174 | 161 |
| 12-17 | 41 | 42 | 22 | 22 |
| 12-13 | 15 | 17 | 8 | 6 |
| 14-15 | 24 | 22 | 13 | 11 |
| 16-17 | 32 | 32 | 16 | 17 |
| 16-25 | 103 | 82 | 58 | 41 |
| 18 or Older | 276 | 293 | 175 | 161 |
| 18-25 | 88 | 73 | 53 | 38 |
| 18-20 | 52 | 46 | 25 | 24 |
| 21-25 | 84 | 60 | 51 | 30 |
| 26 or Older | 255 | 283 | 161 | 157 |
| 26-49 | 174 | 177 | 101 | 103 |
| 26-29 | 84 | 79 | 47 | 45 |
| 30-34 | 87 | 87 | 57 | 48 |
| 35-39 | 82 | 87 | 39 | 48 |
| 40-44 | 74 | 74 | 39 | 42 |
| 45-49 | 77 | 66 | 41 | 40 |
| 50 or Older | 197 | 229 | 128 | 118 |
| 50-54 | 104 | 117 | 60 | 54 |
| 55-59 | 100 | 119 | 75 | 61 |
| 60-64 | 91 | 95 | 48 | 51 |
| 65 or Older | 109 | 112 | 68 | 61 |

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 1.19D Misuse of Opioids in Past Year and Past Month among Persons Aged 12 or Older, by Detailed Age Category: Standard Errors of Percentages, 2017 and 2018

| Age Category | Misuse in Past Year (2017) | Misuse in Past Year (2018) | Misuse in Past Month (2017) | Misuse in Past Month (2018) |
|---------------------|---------------------------------------|---------------------------------------|--|--|
| TOTAL | 0.10 | 0.11 | 0.06 | 0.06 |
| 12-20 | 0.17 | 0.16 | 0.09 | 0.08 |
| 21 or Older | 0.12 | 0.12 | 0.07 | 0.07 |
| 12-17 | 0.16 | 0.17 | 0.09 | 0.09 |
| 12-13 | 0.20 | 0.22 | 0.11 | 0.08 |
| 14-15 | 0.27 | 0.25 | 0.16 | 0.13 |
| 16-17 | 0.36 | 0.37 | 0.18 | 0.19 |
| 16-25 | 0.22 | 0.18 | 0.13 | 0.10 |
| 18 or Older | 0.11 | 0.12 | 0.07 | 0.06 |
| 18-25 | 0.26 | 0.21 | 0.16 | 0.11 |
| 18-20 | 0.39 | 0.34 | 0.19 | 0.18 |
| 21-25 | 0.35 | 0.27 | 0.22 | 0.14 |
| 26 or Older | 0.12 | 0.13 | 0.08 | 0.07 |
| 26-49 | 0.17 | 0.18 | 0.10 | 0.10 |
| 26-29 | 0.45 | 0.42 | 0.26 | 0.25 |
| 30-34 | 0.39 | 0.37 | 0.26 | 0.21 |
| 35-39 | 0.38 | 0.38 | 0.19 | 0.22 |
| 40-44 | 0.36 | 0.37 | 0.20 | 0.21 |
| 45-49 | 0.37 | 0.34 | 0.20 | 0.21 |
| 50 or Older | 0.18 | 0.20 | 0.11 | 0.10 |
| 50-54 | 0.48 | 0.55 | 0.28 | 0.26 |
| 55-59 | 0.44 | 0.54 | 0.34 | 0.28 |
| 60-64 | 0.46 | 0.46 | 0.24 | 0.25 |
| 65 or Older | 0.22 | 0.22 | 0.14 | 0.12 |

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 1.19P Misuse of Opioids in Past Year and Past Month among Persons Aged 12 or Older, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2018 Versus 2017

| Age Category | Misuse in Past Year (2017) | Misuse in Past Year (2018) | Misuse in Past Month (2017) | Misuse in Past Month (2018) |
|---------------------|---------------------------------------|---------------------------------------|--|--|
| TOTAL | 0.0040 | | 0.0302 | |
| 12-20 | 0.1063 | | 0.2249 | |
| 21 or Older | 0.0100 | | 0.0494 | |
| 12-17 | 0.2491 | | 0.1421 | |
| 12-13 | 0.6527 | | 0.5424 | |
| 14-15 | 0.4945 | | 0.1729 | |
| 16-17 | 0.1948 | | 0.4794 | |
| 16-25 | 0.0000 | | 0.0037 | |
| 18 or Older | 0.0061 | | 0.0456 | |
| 18-25 | 0.0000 | | 0.0047 | |
| 18-20 | 0.2212 | | 0.7240 | |
| 21-25 | 0.0000 | | 0.0029 | |
| 26 or Older | 0.1581 | | 0.1909 | |
| 26-49 | 0.1131 | | 0.5919 | |
| 26-29 | 0.3192 | | 0.4714 | |
| 30-34 | 0.4837 | | 0.0501 | |
| 35-39 | 0.5597 | | 0.4419 | |
| 40-44 | 0.8144 | | 0.5252 | |
| 45-49 | 0.1732 | | 0.6179 | |
| 50 or Older | 0.6522 | | 0.2015 | |
| 50-54 | 0.9699 | | 0.5396 | |
| 55-59 | 0.5326 | | 0.3191 | |
| 60-64 | 0.4453 | | 0.7767 | |
| 65 or Older | 0.4700 | | 0.6267 | |

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

**Table 1.20C Use of Illicit Drugs Other Than Marijuana in Past Year and Past Month among Persons Aged 12 or Older, by Detailed Age Category:
Standard Errors of Numbers in Thousands, 2017 and 2018**

| Age Category | Past Year (2017) | Past Year (2018) | Past Month (2017) | Past Month (2018) |
|---------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| TOTAL | 416 | 441 | 271 | 269 |
| 12-20 | 122 | 111 | 66 | 60 |
| 21 or Older | 426 | 451 | 263 | 270 |
| 12-17 | 65 | 69 | 37 | 38 |
| 12-13 | 28 | 31 | 15 | 13 |
| 14-15 | 37 | 39 | 22 | 22 |
| 16-17 | 50 | 48 | 25 | 28 |
| 16-25 | 185 | 182 | 111 | 95 |
| 18 or Older | 410 | 438 | 269 | 269 |
| 18-25 | 137 | 144 | 96 | 85 |
| 18-20 | 96 | 88 | 55 | 47 |
| 21-25 | 145 | 140 | 94 | 77 |
| 26 or Older | 365 | 398 | 244 | 252 |
| 26-49 | 261 | 274 | 174 | 182 |
| 26-29 | 146 | 150 | 87 | 97 |
| 30-34 | 140 | 134 | 91 | 83 |
| 35-39 | 122 | 128 | 81 | 83 |
| 40-44 | 103 | 97 | 60 | 58 |
| 45-49 | 106 | 96 | 60 | 65 |
| 50 or Older | 264 | 298 | 170 | 176 |
| 50-54 | 140 | 145 | 90 | 87 |
| 55-59 | 147 | 163 | 106 | 81 |
| 60-64 | 114 | 130 | 61 | 90 |
| 65 or Older | 151 | 138 | 87 | 94 |

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Illicit Drugs Other Than Marijuana excludes respondents who used only marijuana but includes those who used marijuana in addition to other illicit drugs.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

**Table 1.20D Use of Illicit Drugs Other Than Marijuana in Past Year and Past Month among Persons Aged 12 or Older, by Detailed Age Category:
Standard Errors of Percentages, 2017 and 2018**

| Age Category | Past Year (2017) | Past Year (2018) | Past Month (2017) | Past Month (2018) |
|---------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| TOTAL | 0.15 | 0.16 | 0.10 | 0.10 |
| 12-20 | 0.29 | 0.28 | 0.17 | 0.16 |
| 21 or Older | 0.17 | 0.18 | 0.11 | 0.11 |
| 12-17 | 0.26 | 0.28 | 0.15 | 0.15 |
| 12-13 | 0.35 | 0.38 | 0.20 | 0.17 |
| 14-15 | 0.41 | 0.46 | 0.26 | 0.26 |
| 16-17 | 0.54 | 0.53 | 0.28 | 0.32 |
| 16-25 | 0.34 | 0.35 | 0.23 | 0.21 |
| 18 or Older | 0.17 | 0.18 | 0.11 | 0.11 |
| 18-25 | 0.40 | 0.42 | 0.28 | 0.25 |
| 18-20 | 0.62 | 0.60 | 0.39 | 0.35 |
| 21-25 | 0.53 | 0.54 | 0.40 | 0.34 |
| 26 or Older | 0.17 | 0.19 | 0.11 | 0.12 |
| 26-49 | 0.26 | 0.27 | 0.17 | 0.18 |
| 26-29 | 0.71 | 0.75 | 0.45 | 0.52 |
| 30-34 | 0.60 | 0.55 | 0.41 | 0.35 |
| 35-39 | 0.56 | 0.55 | 0.38 | 0.37 |
| 40-44 | 0.49 | 0.48 | 0.30 | 0.29 |
| 45-49 | 0.50 | 0.48 | 0.29 | 0.33 |
| 50 or Older | 0.23 | 0.26 | 0.15 | 0.15 |
| 50-54 | 0.64 | 0.68 | 0.42 | 0.42 |
| 55-59 | 0.65 | 0.72 | 0.48 | 0.37 |
| 60-64 | 0.56 | 0.62 | 0.31 | 0.44 |
| 65 or Older | 0.30 | 0.27 | 0.18 | 0.18 |

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Illicit Drugs Other Than Marijuana excludes respondents who used only marijuana but includes those who used marijuana in addition to other illicit drugs.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 1.20P Use of Illicit Drugs Other Than Marijuana in Past Year and Past Month among Persons Aged 12 or Older, by Detailed Age Category:
P Values from Tests of Differences of Percentages, 2018 Versus 2017

| Age Category | Past Year (2017) | Past Year (2018) | Past Month (2017) | Past Month (2018) |
|---------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| TOTAL | 0.0400 | | 0.3365 | |
| 12-20 | 0.4268 | | 0.0693 | |
| 21 or Older | 0.0597 | | 0.5930 | |
| 12-17 | 0.6782 | | 0.6455 | |
| 12-13 | 0.5509 | | 0.3210 | |
| 14-15 | 0.7252 | | 0.6772 | |
| 16-17 | 0.2197 | | 0.8223 | |
| 16-25 | 0.0011 | | 0.0081 | |
| 18 or Older | 0.0452 | | 0.3689 | |
| 18-25 | 0.0023 | | 0.0047 | |
| 18-20 | 0.3926 | | 0.0418 | |
| 21-25 | 0.0020 | | 0.0540 | |
| 26 or Older | 0.3237 | | 0.9003 | |
| 26-49 | 0.7571 | | 0.6366 | |
| 26-29 | 0.6585 | | 0.3401 | |
| 30-34 | 0.8928 | | 0.1242 | |
| 35-39 | 0.7534 | | 0.9950 | |
| 40-44 | 0.7821 | | 0.3922 | |
| 45-49 | 0.3905 | | 0.2479 | |
| 50 or Older | 0.3217 | | 0.7947 | |
| 50-54 | 0.6573 | | 0.5565 | |
| 55-59 | 0.9699 | | 0.1286 | |
| 60-64 | 0.9910 | | 0.1197 | |
| 65 or Older | 0.2154 | | 0.6486 | |

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Illicit Drugs Other Than Marijuana excludes respondents who used only marijuana but includes those who used marijuana in addition to other illicit drugs.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.