

Table 1.97A Specific Hallucinogen, Inhalant, Needle, Heroin, and Other Drug Use in Lifetime among Persons Aged 12 or Older, by Age Group: Numbers in Thousands, 2017 and 2018

Drug or Method of Administration	Aged 12+ (2017)	Aged 12+ (2018)	Aged 12-17 (2017)	Aged 12-17 (2018)	Aged 18+ (2017)	Aged 18+ (2018)	Aged 18-25 (2017)	Aged 18-25 (2018)	Aged 26+ (2017)	Aged 26+ (2018)
HALLUCINOGENS	42,072	43,255	700 ^a	565	41,372	42,690	5,859	5,597	35,513	37,093
PCP	6,098	6,085	46	30	6,053	6,055	261	215	5,792	5,839
LSD	26,069	27,339	364	326	25,705	27,013	3,125	3,338	22,581	23,675
Peyote	5,586	5,884	20	26	5,566	5,859	183	167	5,382	5,692
Mescaline	7,565	7,425	8	17	7,557	7,408	217	202	7,341	7,207
Psilocybin/Mushrooms	24,002	25,138	244	207	23,758	24,930	2,987	3,013	20,771	21,917
Ecstasy	19,173	19,949	257	209	18,915	19,740	4,105 ^a	3,586	14,811 ^a	16,155
Ketamine	3,463	3,622	46 ^a	19	3,417	3,602	632 ^a	505	2,785	3,098
DMT, AMT, or 5-MeO-DIPT/Foxy	2,649	2,817	70	55	2,579	2,762	727	700	1,852	2,062
Salvia divinorum	4,845	5,043	43	28	4,802	5,014	1,192 ^a	916	3,610 ^a	4,098
INHALANTS	25,187	24,783	2,144	2,125	23,043	22,658	3,242	3,053	19,801	19,605
Amyl Nitrite, Poppers, Locker Room Odorizers, or Rush	6,868	6,810	115	98	6,752	6,712	536	504	6,216	6,208
Correction Fluid, Degreaser, or Cleaning Fluid	1,438	1,391	178	207	1,260	1,185	185 ^a	121	1,075	1,064
Gasoline or Lighter Fluid	3,080	2,828	397	418	2,683	2,409	327	335	2,356	2,074
Glue, Shoe Polish, or Toluene	3,120	2,984	434	485	2,686	2,499	343	263	2,343	2,236
Halothane, Ether, or Other Anesthetics	869	798	21	29	848	769	86	52	761	717
Lacquer Thinner or Other Paint Solvents	1,532	1,325	122	130	1,409	1,195	166	127	1,243	1,068
Lighter Gases	774	782	113	130	660	652	114	76	546	576
Nitrous Oxide or Whippits	12,813	12,496	90	122	12,723	12,374	1,283	1,127	11,439	11,247
Felt-Tip Pens or Felt-Tip/Magic Markers	6,390	6,155	1,607	1,577	4,782	4,578	1,365	1,305	3,417	3,273
Spray Paints	1,914	1,785	369	369	1,545	1,417	311	236	1,235	1,181
Computer Cleaner/Air Duster	2,623	2,931	199	190	2,424	2,741	744 ^a	532	1,681 ^a	2,209
Other Aerosol Sprays	1,326	1,281	179	146	1,147	1,134	190	141	958	993
NEEDLE USE¹	4,442	4,447	38	18	4,404	4,430	437 ^a	307	3,967	4,123
Heroin Needle Use ¹	2,508	2,360	17	7	2,491	2,353	335 ^a	195	2,155	2,158
Cocaine Needle Use ¹	2,250	2,240	*	4	2,248	2,236	153	100	2,095	2,136
Methamphetamine Needle Use ¹	1,949	2,178	9	7	1,940	2,171	180	143	1,761	2,028
HEROIN	5,295	5,108	32	16	5,262	5,093	602 ^a	459	4,661	4,633
Smoke Heroin ¹	2,000	2,122	10	9	1,989	2,113	294	235	1,695	1,878
Sniff or Snort Heroin ¹	3,869	3,598	5	7	3,863	3,590	421 ^a	310	3,442	3,281
OTHER DRUGS										
GHB	1,512	1,439	7	17	1,506	1,422	116	78	1,390	1,344
Nonprescription Cough or Cold Medicine	10,867	10,809	602	638	10,265	10,171	1,827	1,655	8,438	8,516

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

^a The difference between this estimate and the 2018 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

¹ These estimates may be inconsistent with estimates presented in other detailed tables because NSDUH data are not edited for consistency across discrete sections of the interview, except with regard to skip logic.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 1.97B Specific Hallucinogen, Inhalant, Needle, Heroin, and Other Drug Use in Lifetime among Persons Aged 12 or Older, by Age Group: Percentages, 2017 and 2018

Drug or Method of Administration	Aged 12+ (2017)	Aged 12+ (2018)	Aged 12-17 (2017)	Aged 12-17 (2018)	Aged 18+ (2017)	Aged 18+ (2018)	Aged 18-25 (2017)	Aged 18-25 (2018)	Aged 26+ (2017)	Aged 26+ (2018)
HALLUCINOGENS	15.5	15.8	2.8 ^a	2.3	16.7	17.2	17.1	16.4	16.7	17.3
PCP	2.2	2.2	0.2	0.1	2.4	2.4	0.8	0.6	2.7	2.7
LSD	9.6	10.0	1.5	1.3	10.4	10.9	9.1	9.8	10.6	11.0
Peyote	2.1	2.1	0.1	0.1	2.3	2.4	0.5	0.5	2.5	2.6
Mescaline	2.8	2.7	0.0	0.1	3.1	3.0	0.6	0.6	3.4	3.4
Psilocybin/Mushrooms	8.8	9.2	1.0	0.8	9.6	10.0	8.7	8.9	9.8	10.2
Ecstasy	7.0	7.3	1.0	0.8	7.7	7.9	12.0 ^a	10.5	7.0 ^a	7.5
Ketamine	1.3	1.3	0.2 ^a	0.1	1.4	1.4	1.8	1.5	1.3	1.4
DMT, AMT, or 5-MeO-DIPT/Foxy	1.0	1.0	0.3	0.2	1.0	1.1	2.1	2.1	0.9	1.0
Salvia divinorum	1.8	1.8	0.2	0.1	1.9	2.0	3.5 ^a	2.7	1.7	1.9
INHALANTS	9.3	9.1	8.6	8.5	9.3	9.1	9.5	9.0	9.3	9.1
Amyl Nitrite, Poppers, Locker Room Odorizers, or Rush	2.5	2.5	0.5	0.4	2.7	2.7	1.6	1.5	2.9	2.9
Correction Fluid, Degreaser, or Cleaning Fluid	0.5	0.5	0.7	0.8	0.5	0.5	0.5 ^a	0.4	0.5	0.5
Gasoline or Lighter Fluid	1.1	1.0	1.6	1.7	1.1	1.0	1.0	1.0	1.1	1.0
Glue, Shoe Polish, or Toluene	1.1	1.1	1.7	1.9	1.1	1.0	1.0	0.8	1.1	1.0
Halothane, Ether, or Other Anesthetics	0.3	0.3	0.1	0.1	0.3	0.3	0.3	0.2	0.4	0.3
Lacquer Thinner or Other Paint Solvents	0.6	0.5	0.5	0.5	0.6	0.5	0.5	0.4	0.6	0.5
Lighter Gases	0.3	0.3	0.5	0.5	0.3	0.3	0.3	0.2	0.3	0.3
Nitrous Oxide or Whippits	4.7	4.6	0.4	0.5	5.1	5.0	3.7	3.3	5.4	5.2
Felt-Tip Pens or Felt-Tip/Magic Markers	2.3	2.2	6.4	6.3	1.9	1.8	4.0	3.8	1.6	1.5
Spray Paints	0.7	0.7	1.5	1.5	0.6	0.6	0.9	0.7	0.6	0.5
Computer Cleaner/Air Duster	1.0	1.1	0.8	0.8	1.0	1.1	2.2 ^a	1.6	0.8 ^a	1.0
Other Aerosol Sprays	0.5	0.5	0.7	0.6	0.5	0.5	0.6	0.4	0.4	0.5
NEEDLE USE¹	1.6	1.6	0.2	0.1	1.8	1.8	1.3 ^a	0.9	1.9	1.9
Heroin Needle Use ¹	0.9	0.9	0.1	0.0	1.0	0.9	1.0 ^a	0.6	1.0	1.0
Cocaine Needle Use ¹	0.8	0.8	*	0.0	0.9	0.9	0.4	0.3	1.0	1.0
Methamphetamine Needle Use ¹	0.7	0.8	0.0	0.0	0.8	0.9	0.5	0.4	0.8	0.9
HEROIN	1.9	1.9	0.1	0.1	2.1	2.0	1.8 ^a	1.3	2.2	2.2
Smoke Heroin ¹	0.7	0.8	0.0	0.0	0.8	0.8	0.9	0.7	0.8	0.9
Sniff or Snort Heroin ¹	1.4	1.3	0.0	0.0	1.6	1.4	1.2 ^a	0.9	1.6	1.5
OTHER DRUGS										
GHB	0.6	0.5	0.0	0.1	0.6	0.6	0.3	0.2	0.7	0.6
Nonprescription Cough or Cold Medicine	4.0	3.9	2.4	2.6	4.2	4.1	5.3	4.9	4.0	4.0

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

^a The difference between this estimate and the 2018 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

¹ These estimates may be inconsistent with estimates presented in other detailed tables because NSDUH data are not edited for consistency across discrete sections of the interview, except with regard to skip logic.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.