

**Table 1.97C Specific Hallucinogen, Inhalant, Needle, Heroin, and Other Drug Use in Lifetime among Persons Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, 2017 and 2018**

Drug or Method of Administration	Aged 12+ (2017)	Aged 12+ (2018)	Aged 12-17 (2017)	Aged 12-17 (2018)	Aged 18+ (2017)	Aged 18+ (2018)	Aged 18-25 (2017)	Aged 18-25 (2018)	Aged 26+ (2017)	Aged 26+ (2018)
<b>HALLUCINOGENS</b>	646	726	39	39	639	714	140	145	612	674
PCP	263	313	10	7	262	312	30	25	258	309
LSD	519	591	27	30	516	584	106	107	496	567
Peyote	247	280	6	7	246	279	25	23	244	276
Mescaline	318	332	4	7	317	330	29	27	315	326
Psilocybin/Mushrooms	511	542	21	23	510	539	104	109	486	513
Ecstasy	412	444	25	24	409	442	120	118	385	415
Ketamine	168	179	10	6	168	178	45	42	160	171
DMT, AMT, or 5-MeO-DIPT/Foxy	140	147	13	11	139	147	54	50	127	138
Salvia divinorum	184	188	10	6	184	187	65	61	171	173
<b>INHALANTS</b>	489	478	67	70	482	473	105	100	458	463
Amyl Nitrite, Poppers, Locker Room Odorizers, or Rush	287	285	16	14	287	284	44	44	277	282
Correction Fluid, Degreaser, or Cleaning Fluid	101	104	20	22	99	102	26	17	95	100
Gasoline or Lighter Fluid	166	143	30	30	165	139	33	33	161	134
Glue, Shoe Polish, or Toluene	173	184	30	35	170	180	36	30	166	178
Halothane, Ether, or Other Anesthetics	94	90	6	8	94	90	16	12	92	89
Lacquer Thinner or Other Paint Solvents	124	117	17	17	123	117	23	17	121	115
Lighter Gases	75	83	16	19	75	82	19	14	71	80
Nitrous Oxide or Whippits	364	346	13	19	364	345	65	69	356	339
Felt-Tip Pens or Felt-Tip/Magic Markers	193	189	60	61	182	176	68	62	164	167
Spray Paints	125	142	30	29	122	139	34	28	116	135
Computer Cleaner/Air Duster	121	143	22	23	118	142	47	42	103	134
Other Aerosol Sprays	107	102	21	17	105	101	24	21	100	99
<b>NEEDLE USE<sup>1</sup></b>	214	232	9	6	214	232	36	29	208	228
Heroin Needle Use <sup>1</sup>	167	165	6	3	166	165	34	23	160	162
Cocaine Needle Use <sup>1</sup>	150	179	*	2	150	179	24	17	146	177
Methamphetamine Needle Use <sup>1</sup>	142	167	5	4	141	167	25	20	138	165
<b>HEROIN</b>	239	264	8	5	238	264	46	38	230	261
Smoke Heroin <sup>1</sup>	144	145	4	4	143	145	33	27	138	144
Sniff or Snort Heroin <sup>1</sup>	201	217	2	3	201	217	38	31	193	213
<b>OTHER DRUGS</b>										
GHB	113	114	3	8	113	114	22	15	110	114
Nonprescription Cough or Cold Medicine	322	324	35	41	320	322	81	71	309	311

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

<sup>1</sup> These estimates may be inconsistent with estimates presented in other detailed tables because NSDUH data are not edited for consistency across discrete sections of the interview, except with regard to skip logic.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

**Table 1.97D Specific Hallucinogen, Inhalant, Needle, Heroin, and Other Drug Use in Lifetime among Persons Aged 12 or Older, by Age Group: Standard Errors of Percentages, 2017 and 2018**

Drug or Method of Administration	Aged 12+ (2017)	Aged 12+ (2018)	Aged 12-17 (2017)	Aged 12-17 (2018)	Aged 18+ (2017)	Aged 18+ (2018)	Aged 18-25 (2017)	Aged 18-25 (2018)	Aged 26+ (2017)	Aged 26+ (2018)
<b>HALLUCINOGENS</b>	0.24	0.27	0.16	0.15	0.26	0.29	0.41	0.43	0.29	0.31
PCP	0.10	0.11	0.04	0.03	0.11	0.13	0.09	0.07	0.12	0.14
LSD	0.19	0.22	0.11	0.12	0.21	0.23	0.31	0.31	0.23	0.26
Peyote	0.09	0.10	0.02	0.03	0.10	0.11	0.07	0.07	0.11	0.13
Mescaline	0.12	0.12	0.02	0.03	0.13	0.13	0.08	0.08	0.15	0.15
Psilocybin/Mushrooms	0.19	0.20	0.08	0.09	0.21	0.22	0.30	0.32	0.23	0.24
Ecstasy	0.15	0.16	0.10	0.10	0.17	0.18	0.35	0.35	0.18	0.19
Ketamine	0.06	0.07	0.04	0.03	0.07	0.07	0.13	0.12	0.08	0.08
DMT, AMT, or 5-MeO-DIPT/Foxy	0.05	0.05	0.05	0.04	0.06	0.06	0.16	0.15	0.06	0.06
Salvia divinorum	0.07	0.07	0.04	0.03	0.07	0.08	0.19	0.18	0.08	0.08
<b>INHALANTS</b>	0.18	0.17	0.27	0.28	0.20	0.19	0.31	0.29	0.22	0.22
Amyl Nitrite, Poppers, Locker Room Odorizers, or Rush	0.11	0.10	0.06	0.06	0.12	0.11	0.13	0.13	0.13	0.13
Correction Fluid, Degreaser, or Cleaning Fluid	0.04	0.04	0.08	0.09	0.04	0.04	0.07	0.05	0.04	0.05
Gasoline or Lighter Fluid	0.06	0.05	0.12	0.12	0.07	0.06	0.10	0.10	0.08	0.06
Glue, Shoe Polish, or Toluene	0.06	0.07	0.12	0.14	0.07	0.07	0.10	0.09	0.08	0.08
Halothane, Ether, or Other Anesthetics	0.03	0.03	0.02	0.03	0.04	0.04	0.05	0.03	0.04	0.04
Lacquer Thinner or Other Paint Solvents	0.05	0.04	0.07	0.07	0.05	0.05	0.07	0.05	0.06	0.05
Lighter Gases	0.03	0.03	0.07	0.08	0.03	0.03	0.06	0.04	0.03	0.04
Nitrous Oxide or Whippits	0.13	0.13	0.05	0.07	0.15	0.14	0.19	0.20	0.17	0.16
Felt-Tip Pens or Felt-Tip/Magic Markers	0.07	0.07	0.24	0.24	0.07	0.07	0.20	0.18	0.08	0.08
Spray Paints	0.05	0.05	0.12	0.12	0.05	0.06	0.10	0.08	0.05	0.06
Computer Cleaner/Air Duster	0.04	0.05	0.09	0.09	0.05	0.06	0.14	0.12	0.05	0.06
Other Aerosol Sprays	0.04	0.04	0.08	0.07	0.04	0.04	0.07	0.06	0.05	0.05
<b>NEEDLE USE<sup>1</sup></b>	0.08	0.08	0.04	0.02	0.09	0.09	0.11	0.08	0.10	0.11
Heroin Needle Use <sup>1</sup>	0.06	0.06	0.02	0.01	0.07	0.07	0.10	0.07	0.08	0.08
Cocaine Needle Use <sup>1</sup>	0.06	0.07	*	0.01	0.06	0.07	0.07	0.05	0.07	0.08
Methamphetamine Needle Use <sup>1</sup>	0.05	0.06	0.02	0.01	0.06	0.07	0.07	0.06	0.06	0.08
<b>HEROIN</b>	0.09	0.10	0.03	0.02	0.10	0.11	0.13	0.11	0.11	0.12
Smoke Heroin <sup>1</sup>	0.05	0.05	0.02	0.02	0.06	0.06	0.10	0.08	0.06	0.07
Sniff or Snort Heroin <sup>1</sup>	0.07	0.08	0.01	0.01	0.08	0.09	0.11	0.09	0.09	0.10
<b>OTHER DRUGS</b>										
GHB	0.04	0.04	0.01	0.03	0.05	0.05	0.06	0.04	0.05	0.05
Nonprescription Cough or Cold Medicine	0.12	0.12	0.14	0.16	0.13	0.13	0.24	0.21	0.15	0.14

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

<sup>1</sup> These estimates may be inconsistent with estimates presented in other detailed tables because NSDUH data are not edited for consistency across discrete sections of the interview, except with regard to skip logic.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

**Table 1.97P Specific Hallucinogen, Inhalant, Needle, Heroin, and Other Drug Use in Lifetime among Persons Aged 12 or Older, by Age Group: *P* Values from Tests of Differences of Percentages, 2018 Versus 2017**

Drug or Method of Administration	Aged 12+ (2017)	Aged 12+ (2018)	Aged 12-17 (2017)	Aged 12-17 (2018)	Aged 18+ (2017)	Aged 18+ (2018)	Aged 18-25 (2017)	Aged 18-25 (2018)	Aged 26+ (2017)	Aged 26+ (2018)
<b>HALLUCINOGENS</b>	0.2799		0.0118		0.2245		0.2717		0.1245	
PCP	0.9002		0.1909		0.9220		0.2821		0.9882	
LSD	0.1231		0.3397		0.1145		0.1080		0.2032	
Peyote	0.4547		0.5589		0.4704		0.6629		0.4600	
Mescaline	0.6660		0.2498		0.6397		0.7314		0.6364	
Psilocybin/Mushrooms	0.1557		0.2261		0.1468		0.7431		0.1495	
Ecstasy	0.2692		0.1600		0.2420		0.0035		0.0335	
Ketamine	0.5670		0.0234		0.4997		0.0512		0.2118	
DMT, AMT, or 5-MeO-DIPT/Foxy	0.4525		0.4114		0.4123		0.7716		0.3095	
Salvia divinorum	0.5196		0.1875		0.4912		0.0025		0.0655	
<b>INHALANTS</b>	0.4012		0.8779		0.4092		0.2677		0.5502	
Amyl Nitrite, Poppers, Locker Room Odorizers, or Rush	0.8076		0.4193		0.8313		0.6561		0.8698	
Correction Fluid, Degreaser, or Cleaning Fluid	0.6974		0.3300		0.5493		0.0393		0.8745	
Gasoline or Lighter Fluid	0.2048		0.6098		0.1627		0.8135		0.1405	
Glue, Shoe Polish, or Toluene	0.5288		0.2583		0.3951		0.0890		0.5909	
Halothane, Ether, or Other Anesthetics	0.5691		0.3984		0.5252		0.0839		0.6984	
Lacquer Thinner or Other Paint Solvents	0.1965		0.7378		0.1839		0.1727		0.2624	
Lighter Gases	0.9748		0.4905		0.9058		0.1092		0.8181	
Nitrous Oxide or Whippits	0.4220		0.1459		0.3762		0.1195		0.5408	
Felt-Tip Pens or Felt-Tip/Magic Markers	0.2954		0.7427		0.3319		0.5933		0.4422	
Spray Paints	0.4627		0.9928		0.4574		0.0938		0.7193	
Computer Cleaner/Air Duster	0.1104		0.7752		0.0979		0.0016		0.0022	
Other Aerosol Sprays	0.7133		0.2239		0.8839		0.1373		0.8475	
<b>NEEDLE USE<sup>1</sup></b>	0.9436		0.0586		0.9878		0.0056		0.6948	
Heroin Needle Use <sup>1</sup>	0.4909		0.1562		0.5119		0.0008		0.9399	
Cocaine Needle Use <sup>1</sup>	0.9201		*		0.9035		0.0670		0.9234	
Methamphetamine Needle Use <sup>1</sup>	0.3287		0.7725		0.3269		0.2676		0.2489	
<b>HEROIN</b>	0.5306		0.0775		0.5539		0.0199		0.8362	
Smoke Heroin <sup>1</sup>	0.5941		0.8483		0.5959		0.1909		0.4065	
Sniff or Snort Heroin <sup>1</sup>	0.3112		0.6241		0.3026		0.0226		0.4956	
<b>OTHER DRUGS</b>										
GHB	0.5804		0.1936		0.5275		0.1628		0.6879	
Nonprescription Cough or Cold Medicine	0.7829		0.4948		0.7127		0.1433		1.0000	

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

<sup>1</sup> The estimates that are used in the comparisons may be inconsistent with estimates presented in other detailed tables because NSDUH data are not edited for consistency across discrete sections of the interview, except with regard to skip logic.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.