

**Table 10.36C Had Serious Thoughts of Suicide in Past Year among Persons Aged 18 or Older, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2008-2018**

Age Category	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018
<b>TOTAL</b>	299	299	311	299	305	340	281	280	267	289	282
<b>18-25</b>	74	66	75	85	79	85	87	91	96	108	100
18-20	52	47	56	61	62	62	61	62	78	75	70
21-25	60	51	59	61	60	62	69	74	72	87	85
<b>26 or Older</b>	292	292	296	284	294	327	263	267	247	262	257
26-49	193	200	201	169	208	207	167	165	154	163	170
26-29	82	80	94	78	90	88	71	81	74	88	90
30-34	74	92	86	63	84	106	78	73	78	74	80
35-39	93	103	72	86	94	89	68	71	72	74	75
40-44	88	81	88	83	97	88	73	68	60	73	69
45-49	94	98	111	93	100	85	69	70	70	63	63
50 or Older	210	215	214	229	214	271	194	209	193	210	189
50-54	121	118	111	131	103	209	106	102	108	77	96
55-59	118	118	114	124	95	117	109	98	89	107	86
60-64	77	67	93	84	109	83	80	79	74	84	82
65 or Older	104	107	124	106	123	109	102	121	109	136	102

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Some 2008 to 2010 estimates may differ from previously published estimates due to updates (see Section 3.3.5 in Chapter 3 of the *2018 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

NOTE: Respondents with unknown suicide information were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2018.

**Table 10.36D Had Serious Thoughts of Suicide in Past Year among Persons Aged 18 or Older, by Detailed Age Category: Standard Errors of Percentages, 2008-2018**

Age Category	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018
<b>TOTAL</b>	0.13	0.13	0.14	0.13	0.13	0.14	0.12	0.12	0.11	0.12	0.11
<b>18-25</b>	0.23	0.20	0.22	0.25	0.23	0.24	0.25	0.26	0.28	0.32	0.30
18-20	0.37	0.32	0.39	0.41	0.41	0.44	0.44	0.45	0.56	0.53	0.49
21-25	0.29	0.25	0.27	0.28	0.27	0.28	0.31	0.32	0.31	0.38	0.37
<b>26 or Older</b>	0.15	0.15	0.15	0.14	0.15	0.16	0.13	0.13	0.12	0.12	0.12
26-49	0.19	0.20	0.20	0.17	0.21	0.21	0.17	0.17	0.16	0.16	0.17
26-29	0.47	0.47	0.54	0.44	0.51	0.51	0.41	0.46	0.42	0.47	0.49
30-34	0.39	0.46	0.43	0.32	0.42	0.51	0.36	0.34	0.36	0.34	0.35
35-39	0.45	0.50	0.37	0.44	0.47	0.46	0.34	0.36	0.34	0.35	0.34
40-44	0.41	0.39	0.42	0.40	0.47	0.41	0.36	0.33	0.31	0.36	0.35
45-49	0.41	0.43	0.48	0.42	0.46	0.42	0.33	0.33	0.34	0.31	0.32
50 or Older	0.23	0.23	0.22	0.23	0.21	0.26	0.18	0.19	0.17	0.19	0.17
50-54	0.55	0.52	0.50	0.57	0.43	0.88	0.44	0.43	0.49	0.37	0.46
55-59	0.65	0.62	0.58	0.65	0.46	0.56	0.52	0.46	0.41	0.49	0.40
60-64	0.51	0.43	0.56	0.46	0.63	0.47	0.46	0.41	0.39	0.42	0.40
65 or Older	0.28	0.28	0.32	0.27	0.29	0.25	0.23	0.26	0.23	0.28	0.20

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Some 2008 to 2010 estimates may differ from previously published estimates due to updates (see Section 3.3.5 in Chapter 3 of the *2018 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

NOTE: Respondents with unknown suicide information were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2018.

**Table 10.36P Had Serious Thoughts of Suicide in Past Year among Persons Aged 18 or Older, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2018 Versus 2008-2017**

Age Category	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018
<b>TOTAL</b>	0.0004	0.0005	0.0030	0.0001	0.0045	0.0228	0.0128	0.0652	0.0618	0.9377	
<b>18-25</b>	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.2198	
18-20	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0000	0.0011	0.2055	0.6304	
21-25	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0492	
<b>26 or Older</b>	0.6004	0.8339	0.8828	0.4211	0.9047	0.8682	0.7430	0.8435	0.8485	0.7454	
26-49	0.0083	0.2207	0.0305	0.0001	0.0715	0.0129	0.0115	0.0381	0.0727	0.1372	
26-29	0.0001	0.0000	0.0001	0.0000	0.0001	0.0003	0.0000	0.0298	0.0036	0.3001	
30-34	0.0046	0.0594	0.0078	0.0000	0.0140	0.1643	0.0532	0.1007	0.6041	0.0455	
35-39	0.6382	0.2210	0.0519	0.8317	0.5809	0.4623	0.4971	0.4756	0.7883	0.3800	
40-44	0.7057	0.4324	0.6163	0.6546	0.3242	0.4476	0.2087	0.9157	0.1958	0.3517	
45-49	0.1549	0.0766	0.0016	0.4415	0.0234	0.1288	0.8232	0.6534	0.4718	0.9246	
50 or Older	0.3713	0.5002	0.0997	0.0619	0.2228	0.0417	0.0117	0.0583	0.2354	0.1108	
50-54	0.7304	0.6634	0.8807	0.1331	0.5616	0.0546	0.2482	0.3705	0.2973	0.4645	
55-59	0.3768	0.1575	0.1834	0.3029	0.8437	0.2335	0.0476	0.5425	0.6683	0.2411	
60-64	0.8192	0.6233	0.4249	0.8456	0.1113	0.9688	0.3763	0.2165	0.9027	0.5582	
65 or Older	0.9365	0.3948	0.5928	0.7829	0.2418	0.8998	0.6630	0.3229	0.6392	0.0745	

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Some 2008 to 2010 estimates that are used in the comparisons may differ from previously published estimates due to updates (see Section 3.3.5 in Chapter 3 of the *2018 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

NOTE: Respondents with unknown suicide information were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2018.

**Table 10.39C Made Any Suicide Plans in Past Year among Persons Aged 18 or Older, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2008-2018**

Age Category	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018
<b>TOTAL</b>	155	157	168	157	156	166	138	138	127	148	155
<b>18-25</b>	39	41	41	44	47	50	49	54	56	67	58
18-20	29	29	28	28	37	35	36	41	40	48	39
21-25	27	29	31	35	31	35	38	39	42	49	43
<b>26 or Older</b>	148	150	164	148	152	156	130	127	114	131	147
26-49	99	94	91	102	120	120	85	78	85	90	100
26-29	40	34	32	54	50	55	35	40	35	43	58
30-34	44	34	42	26	32	62	35	38	47	38	42
35-39	51	49	29	47	62	51	42	32	42	39	49
40-44	43	46	49	50	58	45	36	31	37	40	33
45-49	47	46	44	45	59	46	36	34	35	41	38
50 or Older	108	117	134	111	101	98	98	101	78	94	105
50-54	82	69	74	72	47	47	41	65	45	41	65
55-59	47	56	61	68	43	58	54	40	34	48	53
60-64	29	37	50	32	56	37	54	34	43	42	41
65 or Older	49	37	81	42	56	54	54	56	36	64	41

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Some 2008 to 2010 estimates may differ from previously published estimates due to updates (see Section 3.3.5 in Chapter 3 of the *2018 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

NOTE: Respondents were asked whether they made any suicide plans in the past 12 months only if they had serious thoughts of suicide in the past 12 months. Respondents who had no suicide thoughts were categorized as not making any suicide plans in the past 12 months. Respondents with unknown suicide information were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2018.

**Table 10.39D Made Any Suicide Plans in Past Year among Persons Aged 18 or Older, by Detailed Age Category: Standard Errors of Percentages, 2008-2018**

Age Category	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018
<b>TOTAL</b>	0.07	0.07	0.07	0.07	0.07	0.07	0.06	0.06	0.05	0.06	0.06
<b>18-25</b>	0.12	0.12	0.12	0.13	0.14	0.14	0.14	0.16	0.16	0.20	0.17
18-20	0.22	0.21	0.20	0.20	0.28	0.26	0.27	0.31	0.31	0.36	0.30
21-25	0.14	0.14	0.15	0.16	0.15	0.16	0.17	0.18	0.19	0.23	0.20
<b>26 or Older</b>	0.08	0.08	0.08	0.07	0.08	0.08	0.06	0.06	0.05	0.06	0.07
26-49	0.10	0.10	0.09	0.10	0.12	0.12	0.09	0.08	0.09	0.09	0.10
26-29	0.23	0.20	0.19	0.32	0.29	0.32	0.21	0.23	0.20	0.24	0.32
30-34	0.24	0.17	0.21	0.13	0.16	0.30	0.17	0.18	0.22	0.17	0.19
35-39	0.25	0.24	0.15	0.24	0.32	0.26	0.21	0.17	0.21	0.19	0.22
40-44	0.20	0.22	0.24	0.25	0.29	0.21	0.18	0.15	0.19	0.20	0.17
45-49	0.21	0.20	0.20	0.21	0.28	0.23	0.18	0.16	0.17	0.20	0.20
50 or Older	0.12	0.12	0.14	0.11	0.10	0.09	0.09	0.09	0.07	0.08	0.09
50-54	0.37	0.31	0.34	0.31	0.20	0.20	0.17	0.28	0.21	0.20	0.31
55-59	0.26	0.30	0.32	0.36	0.21	0.28	0.26	0.19	0.15	0.22	0.25
60-64	0.19	0.24	0.30	0.17	0.33	0.21	0.31	0.18	0.23	0.21	0.20
65 or Older	0.13	0.10	0.21	0.11	0.13	0.13	0.12	0.12	0.08	0.13	0.08

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Some 2008 to 2010 estimates may differ from previously published estimates due to updates (see Section 3.3.5 in Chapter 3 of the *2018 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

NOTE: Respondents were asked whether they made any suicide plans in the past 12 months only if they had serious thoughts of suicide in the past 12 months. Respondents who had no suicide thoughts were categorized as not making any suicide plans in the past 12 months. Respondents with unknown suicide information were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2018.

**Table 10.39P Made Any Suicide Plans in Past Year among Persons Aged 18 or Older, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2018 Versus 2008-2017**

Age Category	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018
<b>TOTAL</b>	0.0008	0.0002	0.0186	0.0011	0.0328	0.0409	0.0078	0.0117	0.0120	0.6434	
<b>18-25</b>	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0015	0.0173	0.2928	
18-20	0.0002	0.0000	0.0000	0.0000	0.0638	0.0625	0.0099	0.3431	0.4703	0.0294	
21-25	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0006	0.0132	0.6159	
<b>26 or Older</b>	0.1694	0.0826	0.7003	0.2158	0.4719	0.4036	0.2857	0.1030	0.0711	0.2824	
26-49	0.0061	0.0022	0.0022	0.0162	0.3517	0.2894	0.0139	0.0027	0.2106	0.1088	
26-29	0.0008	0.0000	0.0000	0.0525	0.0080	0.0243	0.0008	0.0261	0.0058	0.0253	
30-34	0.3279	0.0110	0.2909	0.0006	0.0105	0.8479	0.1032	0.2808	0.5066	0.2612	
35-39	0.2979	0.2650	0.0097	0.7641	0.8595	0.7159	0.6196	0.0510	0.8279	0.0783	
40-44	0.7517	0.3714	0.3494	0.9150	0.2282	0.6643	0.7875	0.4044	0.7541	0.4261	
45-49	0.8292	0.7673	0.6649	0.8866	0.0732	0.4981	0.7153	0.8004	0.7280	0.2098	
50 or Older	0.8156	0.9959	0.0923	0.7350	0.7923	0.8069	0.4993	0.6705	0.1708	0.9598	
50-54	0.7365	0.8662	0.4727	0.5670	0.3410	0.3940	0.3204	0.5400	0.6340	0.5576	
55-59	0.5249	0.7673	0.7485	0.9876	0.1395	0.9244	0.7624	0.3204	0.0582	0.4197	
60-64	0.9807	0.8021	0.8130	0.4931	0.4571	0.8259	0.3278	0.8189	0.9050	0.7327	
65 or Older	0.9956	0.7972	0.0473	0.8027	0.3724	0.6812	0.0967	0.4126	0.4025	0.2617	

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Some 2008 to 2010 estimates that are used in the comparisons may differ from previously published estimates due to updates (see Section 3.3.5 in Chapter 3 of the *2018 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

NOTE: Respondents were asked whether they made any suicide plans in the past 12 months only if they had serious thoughts of suicide in the past 12 months. Respondents who had no suicide thoughts were categorized as not making any suicide plans in the past 12 months. Respondents with unknown suicide information were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2018.

**Table 10.41C Attempted Suicide in Past Year among Persons Aged 18 or Older, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2008-2018**

Age Category	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018
<b>TOTAL</b>	105	91	108	110	103	118	76	100	96	97	101
<b>18-25</b>	32	31	30	35	41	35	36	44	45	48	44
18-20	25	20	21	22	33	26	22	32	35	36	31
21-25	18	24	22	26	24	22	30	32	30	33	32
<b>26 or Older</b>	100	85	103	103	95	112	67	90	84	87	91
26-49	67	63	59	64	75	83	54	57	53	53	59
26-29	30	23	14	32	35	32	21	24	25	23	24
30-34	36	25	28	20	25	52	22	28	29	20	25
35-39	22	36	16	30	33	36	25	21	27	32	36
40-44	29	23	32	31	33	34	23	28	18	17	22
45-49	29	31	33	29	38	26	30	25	20	23	23
50 or Older	76	57	81	81	64	72	40	73	65	68	66
50-54	67	24	31	54	26	25	25	45	27	36	44
55-59	28	31	30	57	29	46	31	36	41	26	41
60-64	16	28	38	2	23	32	14	27	25	25	10
65 or Older	18	31	58	18	45	38	24	37	34	40	27

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Some 2008 to 2010 estimates may differ from previously published estimates due to updates (see Section 3.3.5 in Chapter 3 of the *2018 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

NOTE: Respondents were asked whether they attempted suicide in the past 12 months only if they had serious thoughts of suicide in the past 12 months. Respondents who had no suicide thoughts were categorized as not attempting suicide in the past 12 months. Respondents with unknown suicide information were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2018.

**Table 10.41D Attempted Suicide in Past Year among Persons Aged 18 or Older, by Detailed Age Category: Standard Errors of Percentages, 2008-2018**

Age Category	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018
<b>TOTAL</b>	0.05	0.04	0.05	0.05	0.04	0.05	0.03	0.04	0.04	0.04	0.04
<b>18-25</b>	0.10	0.09	0.09	0.10	0.12	0.10	0.10	0.13	0.13	0.14	0.13
18-20	0.19	0.14	0.15	0.16	0.25	0.19	0.17	0.24	0.27	0.28	0.24
21-25	0.09	0.12	0.11	0.13	0.12	0.10	0.14	0.15	0.14	0.15	0.15
<b>26 or Older</b>	0.05	0.04	0.05	0.05	0.05	0.06	0.03	0.04	0.04	0.04	0.04
26-49	0.07	0.06	0.06	0.07	0.08	0.09	0.06	0.06	0.05	0.05	0.06
26-29	0.18	0.14	0.09	0.19	0.21	0.19	0.13	0.14	0.14	0.13	0.14
30-34	0.20	0.13	0.14	0.11	0.13	0.26	0.10	0.13	0.14	0.09	0.11
35-39	0.11	0.18	0.08	0.15	0.17	0.19	0.13	0.11	0.13	0.16	0.17
40-44	0.13	0.11	0.15	0.15	0.16	0.16	0.12	0.14	0.09	0.09	0.11
45-49	0.13	0.14	0.15	0.13	0.18	0.13	0.15	0.12	0.10	0.11	0.12
50 or Older	0.08	0.06	0.08	0.08	0.06	0.07	0.04	0.07	0.06	0.06	0.06
50-54	0.31	0.11	0.14	0.24	0.11	0.11	0.10	0.20	0.13	0.17	0.21
55-59	0.16	0.17	0.16	0.30	0.14	0.22	0.15	0.17	0.19	0.12	0.19
60-64	0.11	0.18	0.23	0.01	0.14	0.18	0.08	0.14	0.13	0.13	0.05
65 or Older	0.05	0.08	0.15	0.05	0.11	0.09	0.05	0.08	0.07	0.08	0.05

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Some 2008 to 2010 estimates may differ from previously published estimates due to updates (see Section 3.3.5 in Chapter 3 of the *2018 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

NOTE: Respondents were asked whether they attempted suicide in the past 12 months only if they had serious thoughts of suicide in the past 12 months. Respondents who had no suicide thoughts were categorized as not attempting suicide in the past 12 months. Respondents with unknown suicide information were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2018.



**Table 10.41P Attempted Suicide in Past Year among Persons Aged 18 or Older, by Detailed Age Category: P Values from Tests of Differences of Percentages, 2018 Versus 2008-2017**

Age Category	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018
<b>TOTAL</b>	0.1167	0.0307	0.1402	0.1361	0.5853	0.6561	0.0233	0.9398	0.4802	0.7475	
<b>18-25</b>	0.0000	0.0000	0.0000	0.0000	0.0119	0.0001	0.0001	0.1348	0.5082	0.9403	
18-20	0.0496	0.0002	0.0010	0.0056	0.3880	0.1021	0.0009	0.3498	0.5528	0.2631	
21-25	0.0000	0.0004	0.0003	0.0007	0.0045	0.0000	0.0204	0.2800	0.1383	0.2180	
<b>26 or Older</b>	0.8867	0.6673	0.9734	0.9117	0.7680	0.3705	0.4992	0.6491	0.5482	0.7033	
26-49	0.4201	0.5554	0.1736	0.4449	0.9470	0.3854	0.8244	0.9501	0.8282	0.2982	
26-29	0.9088	0.5984	0.0368	0.6619	0.9465	0.8949	0.9416	0.7076	0.4332	0.8109	
30-34	0.7011	0.4388	0.9128	0.2655	0.4892	0.4078	0.4625	0.6796	0.9315	0.2671	
35-39	0.0295	0.6516	0.0117	0.3787	0.2547	0.8151	0.1176	0.1034	0.4429	0.1880	
40-44	0.4912	0.4878	0.9275	0.7709	0.2663	0.2421	0.5864	0.4611	0.4931	0.3138	
45-49	0.3912	0.2309	0.2530	0.7721	0.3904	0.5864	0.1195	0.7233	0.6336	0.2703	
50 or Older	0.7794	0.7618	0.3243	0.7093	0.7080	0.7458	0.4073	0.5417	0.5271	0.7130	
50-54	0.4366	0.4650	0.6336	0.5611	0.4953	0.3662	0.5540	0.5179	0.4454	0.7135	
55-59	0.2063	0.2701	0.3407	0.9024	0.2192	0.8931	0.3017	0.3669	0.3512	0.2037	
60-64	0.3221	0.4032	0.3578	0.2511	0.0713	0.1670	0.7611	0.2398	0.6703	0.5886	
65 or Older	0.5213	0.5253	0.0590	0.5545	0.1878	0.8055	0.9223	0.6622	0.6858	0.2385	

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Some 2008 to 2010 estimates that are used in the comparisons may differ from previously published estimates due to updates (see Section 3.3.5 in Chapter 3 of the *2018 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

NOTE: Respondents were asked whether they attempted suicide in the past 12 months only if they had serious thoughts of suicide in the past 12 months. Respondents who had no suicide thoughts were categorized as not attempting suicide in the past 12 months. Respondents with unknown suicide information were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2018.