

Table 11.12A Co-Occurring Substance Use Disorder, Illicit Drug Use Disorder, or Alcohol Use Disorder and Major Depressive Episode (MDE) or MDE with Severe Impairment in Past Year among Persons Aged 12 to 17, by Gender: Numbers in Thousands, 2015-2018

Substance Use Disorder (SUD), Illicit Drug Use Disorder (IDUD), or Alcohol Use Disorder (AUD) and MDE/Gender	2015	2016	2017	2018
SUD and MDE	350	333	345	358
Male	87	72 ^a	101	114
Female	263	261	244	244
SUD and MDE with Severe Impairment	303	278	276	288
Male	70	56	81	82
Female	232	222	195	207
IDUD and MDE	241	236	260	279
Male	57	59	80	87
Female	184	177	180	193
IDUD and MDE with Severe Impairment	219	202	215	227
Male	48	44	66	60
Female	172	158	149	167
AUD and MDE	202	182	164	159
Male	51	27	47	50
Female	151	156 ^a	118	108
AUD and MDE with Severe Impairment	172	152	124	132
Male	43	22	34	39
Female	129	130	90	93

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

NOTE: Respondents with unknown impairment data were excluded.

^a The difference between this estimate and the 2018 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2015-2018.

Table 11.12B Co-Occurring Substance Use Disorder, Illicit Drug Use Disorder, or Alcohol Use Disorder and Major Depressive Episode (MDE) or MDE with Severe Impairment in Past Year among Persons Aged 12 to 17, by Gender: Percentages, 2015-2018

Substance Use Disorder (SUD), Illicit Drug Use Disorder (IDUD), or Alcohol Use Disorder (AUD) and MDE/Gender	2015	2016	2017	2018
SUD and MDE	1.4	1.4	1.4	1.5
Male	0.7	0.6 ^a	0.8	0.9
Female	2.2	2.2	2.1	2.1
SUD and MDE with Severe Impairment	1.2	1.1	1.1	1.2
Male	0.6	0.5	0.7	0.7
Female	2.0	1.9	1.6	1.8
IDUD and MDE	1.0	1.0	1.1	1.2
Male	0.5	0.5	0.6	0.7
Female	1.6	1.5	1.5	1.6
IDUD and MDE with Severe Impairment	0.9	0.8	0.9	0.9
Male	0.4	0.4	0.5	0.5
Female	1.4	1.3	1.3	1.4
AUD and MDE	0.8	0.8	0.7	0.7
Male	0.4	0.2	0.4	0.4
Female	1.3	1.3	1.0	0.9
AUD and MDE with Severe Impairment	0.7	0.6	0.5	0.5
Male	0.3	0.2	0.3	0.3
Female	1.1	1.1	0.8	0.8

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

NOTE: Respondents with unknown impairment data were excluded.

^a The difference between this estimate and the 2018 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2015-2018.