

Table 11.6C Substance Use in Past Year and Past Month among Persons Aged 12 to 17 with Major Depressive Episode in Past Year: Standard Errors of Numbers in Thousands, 2004-2018

Substance	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018
PAST YEAR USE															
Illicit Drugs	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	46	47	46	54
Marijuana	35	32	32	31	32	30	29	32	31	36	39	39	40	39	46
Cocaine	11	12	11	11	12	8	9	9	10	7	11	9	8	8	10
Crack	6	5	6	7	5	4	4	5	3	2	5	*	3	2	0
Heroin	4	4	4	4	5	4	4	4	4	4	6	3	2	3	3
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	18	16	18	16
LSD	9	9	5	5	8	8	6	7	8	7	12	13	9	13	11
PCP	8	5	4	3	5	4	3	5	3	3	5	2	3	5	4
Ecstasy	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	10	12	10	10
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	22	19	17	20
Methamphetamine	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	7	3	5	4
Misuse of Psychotherapeutics	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	29	30	30	31
Pain Relievers	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	23	26	21	24
Stimulants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	21	19	21	17
Tranquilizers or Sedatives	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	15	18	20	20
Tranquilizers	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	14	19	20	19
Sedatives	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	7	8	6	10
Benzodiazepines	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	13	18	20	19
Opioids	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	23	26	21	24
Illicit Drugs Other Than Marijuana ¹	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	37	35	36	38
PAST MONTH USE															
Alcohol	37	30	33	35	33	30	29	32	30	30	36	36	33	35	39
Binge Alcohol Use ²	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	27	25	26	29
Heavy Alcohol Use	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	12	11	9	6
Cigarette Use	30	28	26	27	26	24	22	22	21	22	24	24	23	23	22
Daily Cigarette Use	17	14	14	14	11	11	9	11	9	7	11	12	6	7	7

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section 3.3.5 in Chapter 3 of the *2018 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

¹ Illicit Drugs Other Than Marijuana excludes respondents who used only marijuana but includes those who used marijuana in addition to other illicit drugs.

² Estimates of binge alcohol use include use by those who were heavy alcohol users.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2004-2018.

Table 11.6D Substance Use in Past Year and Past Month among Persons Aged 12 to 17 with Major Depressive Episode in Past Year: Standard Errors of Percentages, 2004-2018

Substance	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018
PAST YEAR USE															
Illicit Drugs	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	1.28	1.25	1.19	1.21
Marijuana	1.29	1.25	1.38	1.33	1.37	1.34	1.35	1.37	1.18	1.22	1.24	1.16	1.14	1.09	1.09
Cocaine	0.47	0.53	0.53	0.54	0.57	0.39	0.45	0.45	0.44	0.26	0.40	0.29	0.27	0.24	0.30
Crack	0.26	0.22	0.30	0.32	0.26	0.19	0.19	0.23	0.15	0.08	0.18	*	0.09	0.05	0.01
Heroin	0.17	0.17	0.22	0.19	0.26	0.19	0.20	0.19	0.19	0.14	0.21	0.09	0.05	0.09	0.07
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.58	0.52	0.55	0.47
LSD	0.41	0.39	0.28	0.22	0.38	0.41	0.29	0.33	0.35	0.26	0.43	0.42	0.30	0.41	0.32
PCP	0.34	0.21	0.20	0.16	0.27	0.20	0.18	0.24	0.12	0.13	0.19	0.06	0.08	0.17	0.11
Ecstasy	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.32	0.37	0.31	0.29
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.70	0.59	0.53	0.56
Methamphetamine	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.22	0.08	0.15	0.11
Misuse of Psychotherapeutics	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.91	0.91	0.88	0.85
Pain Relievers	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.75	0.80	0.64	0.67
Stimulants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.67	0.59	0.64	0.49
Tranquilizers or Sedatives	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.48	0.59	0.61	0.57
Tranquilizers	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.45	0.59	0.60	0.54
Sedatives	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.24	0.27	0.17	0.29
Benzodiazepines	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.44	0.57	0.60	0.53
Opioids	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.75	0.80	0.64	0.67
Illicit Drugs Other Than Marijuana ¹	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	1.12	1.01	1.01	1.00
PAST MONTH USE															
Alcohol	1.33	1.21	1.36	1.41	1.43	1.32	1.38	1.34	1.22	1.04	1.17	1.09	0.98	0.99	1.00
Binge Alcohol Use ²	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.84	0.78	0.78	0.77
Heavy Alcohol Use	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.40	0.35	0.27	0.17
Cigarette Use	1.16	1.15	1.21	1.18	1.15	1.15	1.08	1.01	0.89	0.84	0.82	0.73	0.70	0.67	0.61
Daily Cigarette Use	0.75	0.61	0.68	0.66	0.53	0.54	0.45	0.53	0.41	0.27	0.40	0.40	0.20	0.23	0.21

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section 3.3.5 in Chapter 3 of the *2018 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

¹ Illicit Drugs Other Than Marijuana excludes respondents who used only marijuana but includes those who used marijuana in addition to other illicit drugs.

² Estimates of binge alcohol use include use by those who were heavy alcohol users.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2004-2018.

Table 11.6P Substance Use in Past Year and Past Month among Persons Aged 12 to 17 with Major Depressive Episode in Past Year: P Values from Tests of Differences of Percentages, 2018 Versus 2004-2017

Substance	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018
PAST YEAR USE															
Illicit Drugs	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.5072	0.5638	0.0568	
Marijuana	0.2910	0.9841	0.6907	0.3067	0.4669	0.5420	0.9801	0.6541	0.7620	0.7747	0.5169	0.0668	0.5727	0.0860	
Cocaine	0.0005	0.0001	0.0002	0.0001	0.0023	0.0167	0.0105	0.0017	0.0089	0.8552	0.3421	0.8166	0.7803	0.7250	
Crack	0.0109	0.0112	0.0079	0.0089	0.0241	0.0115	0.0206	0.0249	0.0506	0.2967	0.0293	*	0.1233	0.2700	
Heroin	0.0671	0.0593	0.0428	0.2582	0.1835	0.0382	0.0462	0.0273	0.0499	0.2104	0.0982	0.5147	0.9832	0.3531	
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.1631	0.1896	0.0660	
LSD	0.7638	0.8960	0.0418	0.0102	0.4256	0.4421	0.2045	0.5204	0.9651	0.1073	0.9359	0.5702	0.2604	0.3824	
PCP	0.0310	0.0181	0.0753	0.2661	0.0065	0.0505	0.2226	0.1292	0.1490	0.7154	0.2964	0.5122	0.8312	0.3407	
Ecstasy	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.6809	0.1605	0.7146	
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.1723	0.9709	0.4479	
Methamphetamine	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.3504	0.8624	0.2705	
Misuse of Psychotherapeutics	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.2356	0.0420	0.5632	
Pain Relievers	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.1043	0.0040	0.6752	
Stimulants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.0231	0.2017	0.1072	
Tranquilizers or Sedatives	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.2003	0.4484	0.6644	
Tranquilizers	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.1603	0.5827	0.3869	
Sedatives	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.7337	0.2740	0.3895	
Benzodiazepines	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.1394	0.5979	0.4580	
Opioids	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.1013	0.0034	0.6300	
Illicit Drugs Other Than Marijuana ¹	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.0517	0.1191	0.8325	
PAST MONTH USE															
Alcohol	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0441	0.0062	0.3948	0.7524	0.8500	
Binge Alcohol Use ²	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.3798	0.8630	0.5689	
Heavy Alcohol Use	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.0157	0.0064	0.0672	
Cigarette Use	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0023	0.0640	0.0732	0.5470	
Daily Cigarette Use	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0013	0.0318	0.0110	0.4572	0.7131	

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Some 2006 to 2010 estimates that are used in the comparisons may differ from previously published estimates due to updates (see Section 3.3.5 in Chapter 3 of the *2018 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

¹ Illicit Drugs Other Than Marijuana excludes respondents who used only marijuana but includes those who used marijuana in addition to other illicit drugs.

² Estimates of binge alcohol use that are used in the comparisons include use by those who were heavy alcohol users.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2004-2018.

Table 11.8C Substance Use in Past Year and Past Month among Persons Aged 12 to 17 with No Major Depressive Episode in Past Year: Standard Errors of Numbers in Thousands, 2004-2018

Substance	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018
PAST YEAR USE															
Illicit Drugs	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	90	84	88	87
Marijuana	76	76	77	74	73	78	87	81	79	72	76	75	71	79	76
Cocaine	25	25	25	23	20	21	20	16	17	12	18	18	15	14	13
Crack	8	8	10	9	6	5	3	5	7	3	2	3	3	6	1
Heroin	9	6	7	5	9	6	7	9	8	6	5	6	4	4	3
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	33	27	30	25
LSD	15	15	13	14	16	11	16	14	13	13	21	23	18	19	18
PCP	9	13	10	9	8	9	7	6	10	5	7	10	5	5	3
Ecstasy	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	21	17	18	15
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	33	28	29	33
Methamphetamine	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	7	7	8	8
Misuse of Psychotherapeutics	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	52	46	44	42
Pain Relievers	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	41	36	35	33
Stimulants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	28	27	26	22
Tranquilizers or Sedatives	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	32	26	28	25
Tranquilizers	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	29	25	26	25
Sedatives	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	13	12	12	9
Benzodiazepines	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	29	24	26	25
Opioids	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	42	36	35	33
Illicit Drugs Other Than Marijuana ¹	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	67	55	56	57
PAST MONTH USE															
Alcohol	87	85	84	88	80	84	81	77	83	73	74	66	69	68	65
Binge Alcohol Use ²	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	52	48	50	48
Heavy Alcohol Use	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	20	17	17	16
Cigarette Use	74	68	64	64	55	62	62	56	51	45	45	41	36	35	31
Daily Cigarette Use	35	31	31	31	27	28	27	25	23	19	24	18	14	11	13

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section 3.3.5 in Chapter 3 of the *2018 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

¹ Illicit Drugs Other Than Marijuana excludes respondents who used only marijuana but includes those who used marijuana in addition to other illicit drugs.

² Estimates of binge alcohol use include use by those who were heavy alcohol users.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2004-2018.

Table 11.8D Substance Use in Past Year and Past Month among Persons Aged 12 to 17 with No Major Depressive Episode in Past Year: Standard Errors of Percentages, 2004-2018

Substance	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018
PAST YEAR USE															
Illicit Drugs	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.38	0.36	0.37	0.40
Marijuana	0.31	0.30	0.31	0.30	0.29	0.32	0.36	0.34	0.31	0.31	0.33	0.33	0.31	0.34	0.35
Cocaine	0.11	0.11	0.11	0.10	0.09	0.09	0.09	0.07	0.07	0.05	0.09	0.08	0.07	0.06	0.06
Crack	0.04	0.03	0.04	0.04	0.02	0.02	0.01	0.02	0.03	0.01	0.01	0.01	0.01	0.03	0.01
Heroin	0.04	0.03	0.03	0.02	0.04	0.03	0.03	0.04	0.04	0.03	0.03	0.03	0.02	0.02	0.01
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.15	0.13	0.14	0.12
LSD	0.07	0.06	0.05	0.06	0.07	0.05	0.07	0.06	0.06	0.06	0.10	0.11	0.08	0.09	0.09
PCP	0.04	0.06	0.05	0.04	0.03	0.04	0.03	0.02	0.05	0.02	0.03	0.05	0.02	0.02	0.02
Ecstasy	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.10	0.08	0.08	0.07
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.15	0.13	0.14	0.16
Methamphetamine	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.03	0.03	0.04	0.04
Misuse of Psychotherapeutics	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.23	0.21	0.21	0.20
Pain Relievers	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.19	0.17	0.16	0.16
Stimulants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.13	0.12	0.12	0.11
Tranquilizers or Sedatives	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.15	0.12	0.13	0.12
Tranquilizers	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.14	0.11	0.12	0.12
Sedatives	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.06	0.05	0.06	0.04
Benzodiazepines	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.13	0.11	0.12	0.12
Opioids	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.19	0.17	0.16	0.16
Illicit Drugs Other Than Marijuana ¹	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.30	0.24	0.26	0.27
PAST MONTH USE															
Alcohol	0.33	0.33	0.34	0.35	0.32	0.33	0.33	0.31	0.32	0.31	0.32	0.30	0.31	0.31	0.29
Binge Alcohol Use ²	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.24	0.22	0.23	0.22
Heavy Alcohol Use	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.10	0.08	0.08	0.08
Cigarette Use	0.30	0.29	0.27	0.27	0.24	0.26	0.27	0.24	0.22	0.20	0.20	0.19	0.17	0.16	0.15
Daily Cigarette Use	0.15	0.13	0.14	0.14	0.12	0.12	0.12	0.11	0.10	0.09	0.11	0.08	0.07	0.05	0.06

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section 3.3.5 in Chapter 3 of the *2018 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

¹ Illicit Drugs Other Than Marijuana excludes respondents who used only marijuana but includes those who used marijuana in addition to other illicit drugs.

² Estimates of binge alcohol use include use by those who were heavy alcohol users.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2004-2018.

Table 11.8P Substance Use in Past Year and Past Month among Persons Aged 12 to 17 with No Major Depressive Episode in Past Year: P Values from Tests of Differences of Percentages, 2018 Versus 2004-2017

Substance	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018
PAST YEAR USE															
Illicit Drugs	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.0131	0.2374	0.5299	
Marijuana	0.0000	0.0003	0.0002	0.0179	0.0007	0.0000	0.0000	0.0000	0.0001	0.0047	0.0245	0.1529	0.4375	0.4248	
Cocaine	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0162	0.1846	0.0335	0.0314	0.2330	0.2748	
Crack	0.0000	0.0000	0.0000	0.0000	0.0002	0.0018	0.0108	0.0012	0.1300	0.0632	0.1075	0.1991	0.2916	0.0704	
Heroin	0.0009	0.0090	0.0063	0.0656	0.0244	0.0051	0.0637	0.0011	0.0815	0.0320	0.1644	0.1307	0.3873	0.5904	
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.0074	0.1999	0.0248	
LSD	0.2011	0.0975	0.0072	0.0937	0.9161	0.0323	0.1043	0.1652	0.0976	0.1365	0.4243	0.1843	0.4735	0.1909	
PCP	0.0001	0.0000	0.0004	0.0000	0.0001	0.0061	0.0153	0.0049	0.0073	0.0562	0.0492	0.0396	0.4267	0.6106	
Ecstasy	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.0104	0.1848	0.0960	
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.7369	0.0298	0.2558	
Methamphetamine	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.2916	0.3526	0.8694	
Misuse of Psychotherapeutics	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.0001	0.2673	0.4926	
Pain Relievers	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.0000	0.0328	0.1224	
Stimulants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.0584	0.2554	0.1867	
Tranquilizers or Sedatives	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.4110	0.9194	0.4045	
Tranquilizers	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.6505	0.9755	0.5148	
Sedatives	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.0086	0.1584	0.1791	
Benzodiazepines	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.7723	0.9515	0.3811	
Opioids	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.0000	0.0275	0.1440	
Illicit Drugs Other Than Marijuana ¹	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.0006	0.7616	0.4095	
PAST MONTH USE															
Alcohol	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0740	0.4139	0.0161	
Binge Alcohol Use ²	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.0002	0.5412	0.0526	
Heavy Alcohol Use	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.0287	0.2712	0.3346	
Cigarette Use	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0041	0.0207	
Daily Cigarette Use	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0033	0.2939	0.7311	

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Some 2006 to 2010 estimates that are used in the comparisons may differ from previously published estimates due to updates (see Section 3.3.5 in Chapter 3 of the 2018 National Survey on Drug Use and Health: Methodological Summary and Definitions).

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

¹ Illicit Drugs Other Than Marijuana excludes respondents who used only marijuana but includes those who used marijuana in addition to other illicit drugs.

² Estimates of binge alcohol use that are used in the comparisons include use by those who were heavy alcohol users.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2004-2018.