

Table 6.21A Illicit Drug, Tobacco Product, and Alcohol Use in Past Month among Females Aged 15 to 44, by Pregnancy Status: Numbers in Thousands, 2017 and 2018

Substance	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2017	2018	2017	2018	2017	2018
ILLCIT DRUGS	8,855 ^a	9,686	194	128	8,604 ^a	9,509
TOBACCO PRODUCTS	13,222	13,034	334	271	12,780	12,692
Cigarettes	11,909	11,601	313	258	11,499	11,273
Smokeless Tobacco	383	456	6	8	377	447
Cigars	2,115	2,251	51	39	2,041	2,196
Pipe Tobacco	395	446	*	11	389	434
ALCOHOL	33,967	34,133	261	233	33,528	33,722
Binge Alcohol Use ²	18,073	17,963	118	111	17,879	17,786
Heavy Alcohol Use	3,576	3,587	11	34	3,553	3,522
ILLCIT DRUGS, TOBACCO PRODUCTS, OR ALCOHOL	39,290	39,556	563	487	38,508	38,863
ILLCIT DRUGS OR ALCOHOL	35,839	36,193	379	309	35,275	35,699
TOBACCO PRODUCTS OR ALCOHOL	38,382	38,549	515	446	37,653	37,898

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

^a The difference between this estimate and the 2018 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

¹ The Total column includes respondents with unknown pregnancy status.

² Estimates of binge alcohol use include use by those who were heavy alcohol users.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 6.21B Illicit Drug, Tobacco Product, and Alcohol Use in Past Month among Females Aged 15 to 44, by Pregnancy Status: Percentages, 2017 and 2018

Substance	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2017	2018	2017	2018	2017	2018
ILLCIT DRUGS	13.8 ^a	15.0	8.5 ^a	5.4	14.0 ^a	15.4
TOBACCO PRODUCTS	20.6	20.2	14.7	11.6	20.8	20.6
Cigarettes	18.5	18.0	13.8	11.0	18.7	18.3
Smokeless Tobacco	0.6	0.7	0.3	0.3	0.6	0.7
Cigars	3.3	3.5	2.2	1.7	3.3	3.6
Pipe Tobacco	0.6	0.7	*	0.5	0.6	0.7
ALCOHOL	52.9	53.0	11.5	9.9	54.5	54.7
Binge Alcohol Use ²	28.1	27.9	5.2	4.7	29.0	28.8
Heavy Alcohol Use	5.6	5.6	0.5	1.5	5.8	5.7
ILLCIT DRUGS, TOBACCO PRODUCTS, OR ALCOHOL	61.2	61.5	24.7	20.8	62.5	63.0
ILLCIT DRUGS OR ALCOHOL	55.8	56.2	16.6	13.2	57.3	57.9
TOBACCO PRODUCTS OR ALCOHOL	59.8	59.9	22.6	19.0	61.2	61.5

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

^a The difference between this estimate and the 2018 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

¹ The Total column includes respondents with unknown pregnancy status.

² Estimates of binge alcohol use include use by those who were heavy alcohol users.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.