

**Table 8.1C Any Mental Illness in Past Year among Persons Aged 18 or Older, by Gender and Detailed Age Category: Standard Errors of Numbers in Thousands, 2017 and 2018**

<b>Age Category</b>	<b>Total (2017)</b>	<b>Total (2018)</b>	<b>Male (2017)</b>	<b>Male (2018)</b>	<b>Female (2017)</b>	<b>Female (2018)</b>
<b>TOTAL</b>	673	644	409	395	500	500
<b>18-25</b>	169	151	109	99	115	115
18-20	123	106	72	63	87	84
21-25	180	155	105	94	124	116
<b>26 or Older</b>	625	615	393	380	469	477
26-49	363	351	231	228	251	253
26-29	188	187	119	122	135	132
30-34	182	198	127	125	126	139
35-39	173	183	105	122	131	137
40-44	166	154	112	100	124	116
45-49	168	152	106	89	121	123
50 or Older	473	480	296	302	375	371
50-54	244	244	154	169	183	171
55-59	226	244	142	157	170	192
60-64	186	222	124	127	152	176
65 or Older	299	310	171	174	255	254

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Any Mental Illness (AMI) aligns with DSM-IV criteria and is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder. These mental illness estimates are based on a predictive model and are not direct measures of diagnostic status.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

**Table 8.1D Any Mental Illness in Past Year among Persons Aged 18 or Older, by Gender and Detailed Age Category: Standard Errors of Percentages, 2017 and 2018**

<b>Age Category</b>	<b>Total (2017)</b>	<b>Total (2018)</b>	<b>Male (2017)</b>	<b>Male (2018)</b>	<b>Female (2017)</b>	<b>Female (2018)</b>
<b>TOTAL</b>	0.27	0.26	0.34	0.33	0.39	0.39
<b>18-25</b>	0.49	0.44	0.63	0.58	0.67	0.68
18-20	0.72	0.66	0.94	0.81	1.06	1.06
21-25	0.63	0.57	0.83	0.78	0.84	0.84
<b>26 or Older</b>	0.29	0.29	0.38	0.37	0.42	0.43
26-49	0.36	0.35	0.47	0.46	0.49	0.50
26-29	0.84	0.83	1.18	1.15	1.18	1.25
30-34	0.71	0.74	0.99	0.99	1.02	1.03
35-39	0.74	0.72	0.96	1.02	1.05	1.06
40-44	0.75	0.70	1.04	0.95	1.07	1.04
45-49	0.75	0.69	0.99	0.85	1.05	1.08
50 or Older	0.42	0.42	0.56	0.57	0.63	0.61
50-54	1.05	1.07	1.44	1.58	1.51	1.45
55-59	0.97	1.03	1.29	1.44	1.38	1.50
60-64	0.90	0.98	1.21	1.18	1.40	1.54
65 or Older	0.60	0.60	0.77	0.76	0.93	0.90

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Any Mental Illness (AMI) aligns with DSM-IV criteria and is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder. These mental illness estimates are based on a predictive model and are not direct measures of diagnostic status.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

**Table 8.1P Any Mental Illness in Past Year among Persons Aged 18 or Older, by Gender and Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2018 Versus 2017**

Age Category	Total (2017)	Total (2018)	Male (2017)	Male (2018)	Female (2017)	Female (2018)
<b>TOTAL</b>	0.4580		0.8401		0.4160	
<b>18-25</b>	0.4775		0.3138		0.9253	
18-20	0.8001		0.5882		0.3729	
21-25	0.4588		0.0911		0.6304	
<b>26 or Older</b>	0.5170		0.9747		0.3781	
26-49	0.5182		0.8328		0.4567	
26-29	0.2433		0.2611		0.4994	
30-34	0.0733		0.9219		0.0112	
35-39	0.4919		0.0409		0.3715	
40-44	0.2361		0.3963		0.4874	
45-49	0.1880		0.0264		0.9750	
50 or Older	0.7031		0.8589		0.5248	
50-54	0.5485		0.9185		0.4188	
55-59	0.8933		0.9208		0.8557	
60-64	0.6521		0.5191		0.2830	
65 or Older	0.3306		0.8329		0.2961	

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Any Mental Illness (AMI) aligns with DSM-IV criteria and is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder. The mental illness estimates that are used in the comparisons are based on a predictive model and are not direct measures of diagnostic status.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

**Table 8.4C Serious Mental Illness in Past Year among Persons Aged 18 or Older, by Gender and Detailed Age Category: Standard Errors of Numbers in Thousands, 2017 and 2018**

Age Category	Total (2017)	Total (2018)	Male (2017)	Male (2018)	Female (2017)	Female (2018)
<b>TOTAL</b>	307	300	188	187	230	233
<b>18-25</b>	90	86	56	51	70	70
18-20	58	52	37	29	45	45
21-25	77	74	46	45	60	58
<b>26 or Older</b>	290	282	180	179	215	215
26-49	187	186	113	124	139	141
26-29	89	100	47	72	72	72
30-34	82	95	55	59	62	68
35-39	76	83	43	54	63	65
40-44	86	71	52	45	68	57
45-49	83	78	49	46	66	62
50 or Older	209	211	135	125	152	164
50-54	106	124	68	71	75	93
55-59	120	112	82	75	79	86
60-64	79	97	46	53	64	79
65 or Older	113	108	53	53	94	91

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Serious Mental Illness (SMI) aligns with DSM-IV criteria and is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder. Estimates of SMI are a subset of estimates of any mental illness (AMI) because SMI is limited to persons with AMI that resulted in serious functional impairment. These mental illness estimates are based on a predictive model and are not direct measures of diagnostic status.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

**Table 8.4D Serious Mental Illness in Past Year among Persons Aged 18 or Older, by Gender and Detailed Age Category: Standard Errors of Percentages, 2017 and 2018**

Age Category	Total (2017)	Total (2018)	Male (2017)	Male (2018)	Female (2017)	Female (2018)
<b>TOTAL</b>	0.12	0.12	0.16	0.16	0.18	0.18
<b>18-25</b>	0.26	0.25	0.32	0.30	0.41	0.41
18-20	0.43	0.39	0.55	0.41	0.69	0.70
21-25	0.33	0.34	0.42	0.42	0.51	0.51
<b>26 or Older</b>	0.14	0.13	0.18	0.17	0.19	0.19
26-49	0.19	0.18	0.23	0.25	0.27	0.28
26-29	0.47	0.53	0.53	0.76	0.72	0.77
30-34	0.37	0.41	0.49	0.51	0.55	0.59
35-39	0.36	0.37	0.41	0.49	0.58	0.56
40-44	0.43	0.36	0.53	0.46	0.66	0.56
45-49	0.40	0.38	0.48	0.47	0.62	0.60
50 or Older	0.19	0.19	0.26	0.24	0.25	0.27
50-54	0.49	0.59	0.68	0.70	0.65	0.86
55-59	0.54	0.51	0.76	0.73	0.68	0.72
60-64	0.39	0.47	0.45	0.51	0.64	0.76
65 or Older	0.23	0.21	0.24	0.23	0.34	0.32

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Serious Mental Illness (SMI) aligns with DSM-IV criteria and is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder. Estimates of SMI are a subset of estimates of any mental illness (AMI) because SMI is limited to persons with AMI that resulted in serious functional impairment. These mental illness estimates are based on a predictive model and are not direct measures of diagnostic status.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

**Table 8.4P Serious Mental Illness in Past Year among Persons Aged 18 or Older, by Gender and Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2018 Versus 2017**

<b>Age Category</b>	<b>Total (2017)</b>	<b>Total (2018)</b>	<b>Male (2017)</b>	<b>Male (2018)</b>	<b>Female (2017)</b>	<b>Female (2018)</b>
<b>TOTAL</b>	0.8010		0.6579		0.9820	
<b>18-25</b>	0.5568		0.7968		0.5888	
18-20	0.2815		0.0996		0.9261	
21-25	0.1113		0.0973		0.4579	
<b>26 or Older</b>	0.8991		0.6802		0.8673	
26-49	0.2500		0.0477		0.8423	
26-29	0.0445		0.0109		0.5915	
30-34	0.1075		0.2273		0.2848	
35-39	0.7651		0.1170		0.3945	
40-44	0.0701		0.3899		0.1268	
45-49	0.8869		0.7626		0.7341	
50 or Older	0.4348		0.2116		0.9830	
50-54	0.8319		0.3979		0.2991	
55-59	0.3956		0.4341		0.6267	
60-64	0.9699		0.3614		0.5719	
65 or Older	0.5337		0.8435		0.3905	

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Serious Mental Illness (SMI) aligns with DSM-IV criteria and is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder. Estimates of SMI that are used in the comparisons are a subset of estimates of any mental illness (AMI) because SMI is limited to persons with AMI that resulted in serious functional impairment. The mental illness estimates that are used in the comparisons are based on a predictive model and are not direct measures of diagnostic status.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.