

Table 8.36C Major Depressive Episode in Past Year among Persons Aged 18 or Older, by Gender and Detailed Age Category: Standard Errors of Numbers in Thousands, 2017 and 2018

Age Category	Total (2017)	Total (2018)	Male (2017)	Male (2018)	Female (2017)	Female (2018)
TOTAL	383	379	227	227	296	303
18-25	117	115	72	70	91	94
18-20	83	76	50	43	66	64
21-25	101	102	59	61	78	81
26 or Older	362	356	214	213	275	282
26-49	224	225	132	146	166	166
26-29	104	120	60	81	85	85
30-34	98	114	72	74	70	80
35-39	93	94	48	63	78	72
40-44	100	96	59	59	81	75
45-49	94	87	59	50	72	74
50 or Older	267	274	159	164	211	223
50-54	135	156	83	99	103	111
55-59	143	140	86	89	107	109
60-64	129	122	73	72	106	99
65 or Older	145	153	87	80	117	132

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 8.36D Major Depressive Episode in Past Year among Persons Aged 18 or Older, by Gender and Detailed Age Category: Standard Errors of Percentages, 2017 and 2018

Age Category	Total (2017)	Total (2018)	Male (2017)	Male (2018)	Female (2017)	Female (2018)
TOTAL	0.16	0.15	0.19	0.19	0.23	0.24
18-25	0.34	0.34	0.42	0.42	0.54	0.56
18-20	0.58	0.54	0.72	0.60	0.95	0.93
21-25	0.43	0.44	0.55	0.57	0.64	0.68
26 or Older	0.17	0.17	0.21	0.21	0.25	0.25
26-49	0.23	0.23	0.27	0.30	0.33	0.33
26-29	0.54	0.61	0.67	0.86	0.85	0.90
30-34	0.44	0.49	0.63	0.65	0.62	0.69
35-39	0.43	0.42	0.46	0.58	0.70	0.63
40-44	0.49	0.48	0.60	0.59	0.77	0.73
45-49	0.45	0.43	0.58	0.52	0.68	0.71
50 or Older	0.24	0.24	0.31	0.31	0.36	0.37
50-54	0.63	0.74	0.83	0.98	0.91	1.01
55-59	0.65	0.64	0.81	0.88	0.93	0.91
60-64	0.63	0.58	0.71	0.69	1.04	0.96
65 or Older	0.30	0.30	0.40	0.35	0.43	0.47

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 8.36P Major Depressive Episode in Past Year among Persons Aged 18 or Older, by Gender and Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2018 Versus 2017

Age Category	Total (2017)	Total (2018)	Male (2017)	Male (2018)	Female (2017)	Female (2018)
TOTAL	0.5561		0.9272		0.4104	
18-25	0.1538		0.5068		0.1980	
18-20	0.5097		0.5708		0.1772	
21-25	0.1994		0.2004		0.5514	
26 or Older	0.8162		0.7891		0.6044	
26-49	0.2796		0.2991		0.5535	
26-29	0.0150		0.0230		0.1805	
30-34	0.1653		0.8365		0.0375	
35-39	0.8270		0.2646		0.2349	
40-44	0.3300		0.9016		0.2909	
45-49	0.7048		0.2944		0.8168	
50 or Older	0.6131		0.2335		0.7881	
50-54	0.6040		0.9614		0.4159	
55-59	0.4905		0.6822		0.5227	
60-64	0.3540		0.1595		0.8762	
65 or Older	0.9196		0.5504		0.5966	

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

**Table 8.37C Major Depressive Episode with Severe Impairment in Past Year among Persons Aged 18 or Older, by Gender and Detailed Age Category:
Standard Errors of Numbers in Thousands, 2017 and 2018**

Age Category	Total (2017)	Total (2018)	Male (2017)	Male (2018)	Female (2017)	Female (2018)
TOTAL	297	311	180	184	228	244
18-25	98	93	62	57	76	74
18-20	68	62	44	34	51	51
21-25	79	77	47	48	63	60
26 or Older	280	293	166	175	216	226
26-49	179	189	107	122	135	138
26-29	90	100	49	70	73	69
30-34	77	96	52	64	56	64
35-39	78	76	42	47	65	59
40-44	80	81	45	50	66	62
45-49	72	67	47	41	53	54
50 or Older	206	222	122	125	158	180
50-54	98	133	62	72	69	103
55-59	121	110	71	73	92	84
60-64	87	97	54	50	68	81
65 or Older	106	113	55	56	90	99

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

NOTE: Respondents with unknown impairment data were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

**Table 8.37D Major Depressive Episode with Severe Impairment in Past Year among Persons Aged 18 or Older, by Gender and Detailed Age Category:
Standard Errors of Percentages, 2017 and 2018**

Age Category	Total (2017)	Total (2018)	Male (2017)	Male (2018)	Female (2017)	Female (2018)
TOTAL	0.12	0.13	0.15	0.16	0.18	0.19
18-25	0.29	0.28	0.37	0.34	0.45	0.44
18-20	0.50	0.46	0.65	0.49	0.79	0.78
21-25	0.35	0.34	0.44	0.46	0.55	0.52
26 or Older	0.13	0.14	0.16	0.17	0.20	0.20
26-49	0.18	0.19	0.22	0.25	0.27	0.27
26-29	0.48	0.53	0.55	0.76	0.74	0.73
30-34	0.35	0.41	0.47	0.56	0.51	0.55
35-39	0.37	0.34	0.41	0.44	0.59	0.52
40-44	0.40	0.41	0.47	0.51	0.64	0.61
45-49	0.35	0.34	0.47	0.42	0.52	0.54
50 or Older	0.18	0.20	0.23	0.24	0.27	0.30
50-54	0.46	0.63	0.63	0.71	0.61	0.95
55-59	0.55	0.51	0.67	0.72	0.79	0.71
60-64	0.43	0.47	0.53	0.49	0.67	0.79
65 or Older	0.22	0.22	0.25	0.25	0.33	0.36

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

NOTE: Respondents with unknown impairment data were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 8.37P Major Depressive Episode with Severe Impairment in Past Year among Persons Aged 18 or Older, by Gender and Detailed Age Category:
***P* Values from Tests of Differences of Percentages, 2018 Versus 2017**

Age Category	Total (2017)	Total (2018)	Male (2017)	Male (2018)	Female (2017)	Female (2018)
TOTAL	0.3331		0.4036		0.5435	
18-25	0.4220		0.3434		0.7901	
18-20	0.7758		0.6868		0.4760	
21-25	0.4236		0.1162		0.7928	
26 or Older	0.4136		0.5465		0.5441	
26-49	0.3361		0.1118		0.9875	
26-29	0.0804		0.0537		0.4701	
30-34	0.0939		0.2025		0.2681	
35-39	0.4485		0.6931		0.1938	
40-44	0.4015		0.8092		0.2317	
45-49	0.8027		0.4648		0.3933	
50 or Older	0.7593		0.5381		0.4074	
50-54	0.1087		0.8397		0.0201	
55-59	0.4007		0.8636		0.1745	
60-64	0.7078		0.1559		0.5901	
65 or Older	0.8179		0.9173		0.8313	

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

NOTE: Respondents with unknown impairment data were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.