

**Table 9.5A Major Depressive Episode (MDE) or MDE with Severe Impairment in Past Year among Persons Aged 12 to 17, and Receipt of Treatment for Depression in Past Year among Persons Aged 12 to 17 with MDE or MDE with Severe Impairment in Past Year, by Demographic Characteristics: Numbers in Thousands, 2017 and 2018**

<b>Demographic Characteristic</b>	<b>Had MDE (2017)</b>	<b>Had MDE (2018)</b>	<b>Had MDE with Severe Impairment<sup>1</sup> (2017)</b>	<b>Had MDE with Severe Impairment<sup>1</sup> (2018)</b>	<b>Received Treatment for Depression in Past Year among Persons with MDE<sup>2</sup> (2017)</b>	<b>Received Treatment for Depression in Past Year among Persons with MDE<sup>2</sup> (2018)</b>	<b>Received Treatment for Depression in Past Year among Persons with MDE with Severe Impairment<sup>1,2</sup> (2017)</b>	<b>Received Treatment for Depression in Past Year among Persons with MDE with Severe Impairment<sup>1,2</sup> (2018)</b>
<b>TOTAL</b>	3,214 <sup>a</sup>	3,482	2,265	2,423	1,330	1,432	1,074	1,133
<b>AGE GROUP</b>								
12-13	516 <sup>a</sup>	632	332	382	193	251	153	177
14-15	1,200	1,250	861	891	453	514	364	406
16-17	1,498	1,600	1,072	1,150	684	667	558	550
<b>GENDER</b>								
Male	845	946	581	628	274 <sup>a</sup>	351	208 <sup>a</sup>	274
Female	2,369	2,537	1,684	1,795	1,056	1,081	866	859
<b>HISPANIC ORIGIN AND RACE</b>								
Not Hispanic or Latino	2,419	2,601	1,694	1,841	1,072	1,100	868	882
White	1,785	1,905	1,256	1,354	847	870	679	695
Black or African American	313	334	233	225	110	115	97	90
AIAN	24	23	6	*	*	*	*	*
NHOPI	*	*	*	*	*	*	*	*
Asian	145	189	101	127	*	*	*	*
Two or More Races	138	137	89	109	62	48	*	*
Hispanic or Latino	795	882	571	582	258	332	206	251

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

<sup>a</sup> The difference between this estimate and the 2018 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

<sup>1</sup> Respondents with unknown impairment data were excluded.

<sup>2</sup> Respondents with unknown past year treatment for depression data were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

**Table 9.5B Major Depressive Episode (MDE) or MDE with Severe Impairment in Past Year among Persons Aged 12 to 17, and Receipt of Treatment for Depression in Past Year among Persons Aged 12 to 17 with MDE or MDE with Severe Impairment in Past Year, by Demographic Characteristics: Percentages, 2017 and 2018**

Demographic Characteristic	Had MDE (2017)	Had MDE (2018)	Had MDE with Severe Impairment <sup>1</sup> (2017)	Had MDE with Severe Impairment <sup>1</sup> (2018)	Received Treatment for Depression in Past Year among Persons with MDE <sup>2</sup> (2017)	Received Treatment for Depression in Past Year among Persons with MDE <sup>2</sup> (2018)	Received Treatment for Depression in Past Year among Persons with MDE with Severe Impairment <sup>1,2</sup> (2017)	Received Treatment for Depression in Past Year among Persons with MDE with Severe Impairment <sup>1,2</sup> (2018)
<b>TOTAL</b>	13.3 <sup>a</sup>	14.4	9.4	10.0	41.5	41.4	47.5	46.9
<b>AGE GROUP</b>								
12-13	6.9 <sup>a</sup>	8.4	4.4	5.1	37.6	40.7	46.2	47.2
14-15	14.5	15.3	10.4	10.9	37.9	41.2	42.3	45.6
16-17	17.7	19.0	12.7	13.7	45.8	41.8	52.0	47.8
<b>GENDER</b>								
Male	6.8	7.7	4.7	5.1	32.5	37.5	35.8	43.7
Female	20.0	21.5	14.2	15.2	44.8	42.9	51.5	48.1
<b>HISPANIC ORIGIN AND RACE</b>								
Not Hispanic or Latino	13.1 <sup>a</sup>	14.2	9.2	10.1	44.4	42.6	51.3	48.1
White	14.0	15.1	9.8	10.7	47.5	46.1	54.1	51.6
Black or African American	9.5	10.3	7.1	6.9	35.1	34.6	41.6	40.1
AIAN	16.3	15.2	3.9	*	*	*	*	*
NHOPI	*	*	*	*	*	*	*	*
Asian	11.3	13.6	7.9	9.2	*	*	*	*
Two or More Races	16.9	17.7	10.9	14.2	44.8	34.8	*	*
Hispanic or Latino	13.8	15.1	9.9	10.0	32.7	37.9	36.1	43.2

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

<sup>a</sup> The difference between this estimate and the 2018 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

<sup>1</sup> Respondents with unknown impairment data were excluded.

<sup>2</sup> Respondents with unknown past year treatment for depression data were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.