

LOUISIANA

Table 47 Selected Drug Use, Perceptions of Great Risk, Past Year Substance Use Disorder and Treatment, and Past Year Mental Health Measures in Louisiana, by Age Group: Estimated Numbers (in Thousands), Annual Averages Based on 2016-2017 NSDUHs

Measure	12+	12-17	18-25	26+	18+
ILLICIT DRUGS					
Past Month Illicit Drug Use ^{1,2}	356	25	103	228	331
Past Year Marijuana Use	469	36	142	290	433
Past Month Marijuana Use	291	21	92	178	271
Perceptions of Great Risk from Smoking Marijuana Once a Month	1,115	101	72	942	1,014
First Use of Marijuana ^{3,4}	35	15	15	5	20
Past Month Illicit Drug Use Other Than Marijuana ^{1,2}	133	10	34	90	124
Past Year Cocaine Use	55	1	18	36	54
Perceptions of Great Risk from Using Cocaine Once a Month	2,833	201	334	2,298	2,632
Past Year Heroin Use	8	0	2	6	8
Perceptions of Great Risk from Trying Heroin Once or Twice	3,355	242	412	2,701	3,113
Past Year Methamphetamine Use	17	1	3	13	16
Past Year Misuse of Pain Relievers ²	158	11	34	113	147
ALCOHOL					
Past Month Alcohol Use	1,922	37	267	1,618	1,884
Past Month Binge Alcohol Use ⁵	1,024	19	175	830	1,005
Perceptions of Great Risk from Having Five or More Drinks of an Alcoholic Beverage Once or Twice a Week	1,687	164	192	1,331	1,523
Past Month Alcohol Use ⁶ (Individuals Aged 12 to 20)	105	--	--	--	--
Past Month Binge Alcohol Use ^{5,6} (Individuals Aged 12 to 20)	61	--	--	--	--
TOBACCO PRODUCTS					
Past Month Tobacco Product Use ⁷	1,158	25	170	963	1,132
Past Month Cigarette Use	917	16	133	768	901
Perceptions of Great Risk from Smoking One or More Packs of Cigarettes per Day	2,596	228	311	2,056	2,368
PAST YEAR SUBSTANCE USE DISORDER AND TREATMENT					
Illicit Drug Use Disorder ^{1,2,8}	102	9	33	61	94
Pain Reliever Use Disorder ^{2,8}	27	2	6	19	25
Alcohol Use Disorder ⁸	236	8	49	179	228
Substance Use Disorder ^{1,2,8}	307	14	70	223	293
Needing But Not Receiving Treatment for Illicit Drug Use ^{1,2,9}	91	8	31	52	83
Needing But Not Receiving Treatment for Alcohol Use ⁹	213	7	45	160	205
Needing But Not Receiving Treatment for Substance Use ^{1,2,9}	277	13	67	197	264
PAST YEAR MENTAL HEALTH ISSUES					
Serious Mental Illness ^{4,10}	--	--	27	137	164
Any Mental Illness ^{4,10}	--	--	97	566	663
Received Mental Health Services ¹¹	--	--	57	411	468
Had Serious Thoughts of Suicide ¹²	--	--	40	108	147
Major Depressive Episode ^{4,13}	--	39	43	205	249

-- Not available.

NOTE: Estimates are based on a survey-weighted hierarchical Bayes estimation approach.

NOTE: Estimated numbers appearing as 0 in this table mean that the estimate is greater than 0 but less than 500 because estimated numbers are shown in thousands.

¹ Illicit Drug Use includes the misuse of prescription psychotherapeutics or the use of marijuana, cocaine (including crack), heroin, hallucinogens, inhalants, or methamphetamine. Illicit Drug Use Other Than Marijuana includes the misuse of prescription psychotherapeutics or the use of cocaine (including crack), heroin, hallucinogens, inhalants, or methamphetamine.² Misuse of prescription psychotherapeutics is defined as use in any way not directed by a doctor, including use without a prescription of one's own; use in greater amounts, more often, or longer than told; or use in any other way not directed by a doctor. Prescription psychotherapeutics do not include over-the-counter drugs.³ First Use of Marijuana is defined as follows: *Average annual number of marijuana initiates* = $X_1 \div 2$, where X_1 is the number of marijuana initiates in the past 24 months.⁴ For details, see Section B of the "2016-2017 NSDUH: Guide to State Tables and Summary of Small Area Estimation Methodology" at <https://www.samhsa.gov/data/>.⁵ Binge Alcohol Use is defined as drinking five or more drinks (for males) or four or more drinks (for females) on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.⁶ Underage drinking is defined for individuals aged 12 to 20; therefore, the "12+" estimate reflects that age group and not individuals aged 12 or older.⁷ Tobacco Products include cigarettes, smokeless tobacco (i.e., snuff, dip, chewing tobacco, or "snus"), cigars, or pipe tobacco.⁸ Substance Use Disorder is defined as meeting criteria for illicit drug or alcohol dependence or abuse. Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).⁹ Respondents were classified as needing substance use treatment if they met the criteria for illicit drug or alcohol use disorder as defined in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV) or received treatment for illicit drug or alcohol use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center). Needing But Not Receiving Substance Use Treatment refers to respondents who are classified as needing illicit drug or alcohol treatment, but who did not receive illicit drug or alcohol treatment at a specialty facility.¹⁰ Mental illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder, assessed by the Mental Health Surveillance Study (MHSS) *Structured Clinical Interview for the Diagnostic and Statistical Manual of Mental Disorders—Fourth Edition—Research Version—Axis I Disorders* (MHSS-SCID), which is based on the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Any mental illness (AMI) includes adults with any mental disorder regardless of whether their disorder resulted in functional impairment. Serious mental illness (SMI) includes adults with any mental disorder that resulted in serious functional impairment.¹¹ Mental health services are defined as having received inpatient treatment/counseling or outpatient treatment/counseling or having used prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use.¹² Respondents were asked, "At any time in the past 12 months, did you seriously think about trying to kill yourself?" If they answered "Yes," they were categorized as having serious thoughts of suicide in the past year.¹³ Major depressive episode (MDE) is defined as in the 5th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-5), which specifies a period of at least 2 weeks when an individual experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of specified depression symptoms. There are minor wording differences in the questions in the adult and adolescent MDE modules. Therefore, data from youths aged 12 to 17 were not combined with data from adults aged 18 or older to produce an estimate for those aged 12 or older.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2016 and 2017.

LOUISIANA

Table 48 Selected Drug Use, Perceptions of Great Risk, Past Year Substance Use Disorder and Treatment, and Past Year Mental Health Measures in Louisiana, by Age Group: Percentages, Annual Averages Based on 2016-2017 NSDUHs

Measure	12+	12-17	18-25	26+	18+
ILLCIT DRUGS					
Past Month Illicit Drug Use ^{1,2}	9.29	6.95	21.05	7.64	9.54
Past Year Marijuana Use	12.23	9.92	28.98	9.76	12.48
Past Month Marijuana Use	7.60	5.67	18.80	5.99	7.80
Perceptions of Great Risk from Smoking Marijuana Once a Month	29.08	27.75	14.73	31.63	29.22
First Use of Marijuana ^{3,4}	1.58	4.53	5.81	0.33	1.08
Past Month Illicit Drug Use Other Than Marijuana ^{1,2}	3.48	2.61	6.87	3.03	3.57
Past Year Cocaine Use	1.43	0.28	3.69	1.19	1.55
Perceptions of Great Risk from Using Cocaine Once a Month	73.91	55.03	68.05	77.18	75.88
Past Year Heroin Use	0.22	0.04	0.40	0.21	0.24
Perceptions of Great Risk from Trying Heroin Once or Twice	87.54	66.18	83.82	90.74	89.75
Past Year Methamphetamine Use	0.44	0.14	0.67	0.44	0.47
Past Year Misuse of Pain Relievers ²	4.12	3.00	6.83	3.81	4.24
ALCOHOL					
Past Month Alcohol Use	50.12	10.19	54.27	54.34	54.33
Past Month Binge Alcohol Use ⁵	26.72	5.21	35.73	27.87	28.98
Perceptions of Great Risk from Having Five or More Drinks of an Alcoholic Beverage Once or Twice a Week	43.99	44.76	39.19	44.70	43.91
Past Month Alcohol Use ⁶ (Individuals Aged 12 to 20)	19.51	--	--	--	--
Past Month Binge Alcohol Use ^{5,6} (Individuals Aged 12 to 20)	11.35	--	--	--	--
TOBACCO PRODUCTS					
Past Month Tobacco Product Use ⁷	30.20	6.90	34.52	32.34	32.65
Past Month Cigarette Use	23.92	4.36	27.11	25.79	25.98
Perceptions of Great Risk from Smoking One or More Packs of Cigarettes per Day	67.72	62.51	63.36	69.08	68.26
PAST YEAR SUBSTANCE USE DISORDER AND TREATMENT					
Illicit Drug Use Disorder ^{1,2,8}	2.67	2.42	6.63	2.05	2.70
Pain Reliever Use Disorder ^{2,8}	0.70	0.44	1.15	0.65	0.72
Alcohol Use Disorder ⁸	6.16	2.26	10.03	6.00	6.57
Substance Use Disorder ^{1,2,8}	8.02	3.89	14.25	7.50	8.46
Needing But Not Receiving Treatment for Illicit Drug Use ^{1,2,9}	2.37	2.15	6.39	1.74	2.40
Needing But Not Receiving Treatment for Alcohol Use ⁹	5.55	2.05	9.13	5.39	5.92
Needing But Not Receiving Treatment for Substance Use ^{1,2,9}	7.22	3.49	13.59	6.62	7.61
PAST YEAR MENTAL HEALTH ISSUES					
Serious Mental Illness ^{4,10}	--	--	5.47	4.60	4.72
Any Mental Illness ^{4,10}	--	--	19.67	19.02	19.11
Received Mental Health Services ¹¹	--	--	11.52	13.81	13.49
Had Serious Thoughts of Suicide ¹²	--	--	8.07	3.62	4.25
Major Depressive Episode ^{4,13}	--	10.76	8.80	6.90	7.17

-- Not available.

NOTE: Estimates are based on a survey-weighted hierarchical Bayes estimation approach.

¹ Illicit Drug Use includes the misuse of prescription psychotherapeutics or the use of marijuana, cocaine (including crack), heroin, hallucinogens, inhalants, or methamphetamine. Illicit Drug Use Other Than Marijuana includes the misuse of prescription psychotherapeutics or the use of cocaine (including crack), heroin, hallucinogens, inhalants, or methamphetamine.² Misuse of prescription psychotherapeutics is defined as use in any way not directed by a doctor, including use without a prescription of one's own; use in greater amounts, more often, or longer than told; or use in any other way not directed by a doctor. Prescription psychotherapeutics do not include over-the-counter drugs.³ First Use of Marijuana is defined as follows: $Average\ annual\ initiation\ of\ marijuana\ (\%) = 100 * \{ [X_1 \div (0.5 * X_1 + X_2)] \div 2 \}$, where X_1 is the number of marijuana initiates in the past 24 months and X_2 is the number of individuals who never used marijuana (with the at-risk population defined as $0.5 * X_1 + X_2$). Both of the computation components, X_1 and X_2 , are based on a survey-weighted hierarchical Bayes estimation approach. The age group shown is based on a respondent's age at the time of the interview, not his or her age at first use.⁴ For details, see Section B of the "2016-2017 NSDUH: Guide to State Tables and Summary of Small Area Estimation Methodology" at <https://www.samhsa.gov/data/>.⁵ Binge Alcohol Use is defined as drinking five or more drinks (for males) or four or more drinks (for females) on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.⁶ Underage drinking is defined for individuals aged 12 to 20; therefore, the "12+" estimate reflects that age group and not individuals aged 12 or older.⁷ Tobacco Products include cigarettes, smokeless tobacco (i.e., snuff, dip, chewing tobacco, or "snus"), cigars, or pipe tobacco.⁸ Substance Use Disorder is defined as meeting criteria for illicit drug or alcohol dependence or abuse. Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).⁹ Respondents were classified as needing substance use treatment if they met the criteria for illicit drug or alcohol use disorder as defined in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV) or received treatment for illicit drug or alcohol use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center). Needing But Not Receiving Substance Use Treatment refers to respondents who are classified as needing illicit drug or alcohol treatment, but who did not receive illicit drug or alcohol treatment at a specialty facility.¹⁰ Mental illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder, assessed by the Mental Health Surveillance Study (MHSS) *Structured Clinical Interview for the Diagnostic and Statistical Manual of Mental Disorders—Fourth Edition—Research Version—Axis I Disorders* (MHSS-SCID), which is based on the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Any mental illness (AMI) includes adults with any mental disorder regardless of whether their disorder resulted in functional impairment. Serious mental illness (SMI) includes adults with any mental disorder that resulted in serious functional impairment.¹¹ Mental health services are defined as having received inpatient treatment/counseling or outpatient treatment/counseling or having used prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use.¹² Respondents were asked, "At any time in the past 12 months, did you seriously think about trying to kill yourself?" If they answered "Yes," they were categorized as having serious thoughts of suicide in the past year.¹³ Major depressive episode (MDE) is defined as in the 5th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-5), which specifies a period of at least 2 weeks when an individual experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of specified depression symptoms. There are minor wording differences in the questions in the adult and adolescent MDE modules. Therefore, data from youths aged 12 to 17 were not combined with data from adults aged 18 or older to produce an estimate for those aged 12 or older.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2016 and 2017.