

Table 36.1 Marijuana Use in the Past Year, Marijuana Use in the Past Month, and Average Annual Incidence Estimates (Expressed as Percentages of the At-Risk Population) of First Use of Marijuana in Ohio among Individuals Aged 12 or Older, by Substate Region: Percentages, Annual Averages Based on 2014, 2015, and 2016 NSDUHs

State/Substate Region	Marijuana Use in the Past Year		Marijuana Use in the Past Month		First Use of Marijuana ¹	
	Estimate	95% Confidence Interval	Estimate	95% Confidence Interval	Estimate	95% Confidence Interval
Ohio	12.75	(11.87 - 13.70)	8.14	(7.42 - 8.92)	1.89	(1.72 - 2.06)
Boards 2, 46, 55, and 68	10.31	(8.03 - 13.14)	6.59	(4.94 - 8.73)	1.48	(1.13 - 1.93)
Boards 3, 52, and 85	9.82	(7.48 - 12.78)	6.95	(5.19 - 9.27)	1.53	(1.16 - 2.02)
Boards 4 and 78	10.44	(8.15 - 13.28)	6.56	(4.87 - 8.78)	1.90	(1.44 - 2.49)
Boards 5 and 60	13.74	(10.98 - 17.07)	8.66	(6.62 - 11.24)	1.86	(1.42 - 2.43)
Boards 7, 15, 41, 79, and 84	10.39	(8.02 - 13.35)	6.30	(4.60 - 8.57)	1.26	(0.97 - 1.65)
Boards 8, 13, and 83	9.19	(7.06 - 11.88)	6.36	(4.74 - 8.49)	1.70	(1.30 - 2.21)
Board 9 (Butler)	13.55	(10.82 - 16.85)	8.74	(6.79 - 11.18)	2.00	(1.54 - 2.60)
Board 12	11.69	(9.18 - 14.78)	8.06	(6.10 - 10.59)	1.72	(1.33 - 2.22)
Boards 18 and 47	14.33	(12.34 - 16.57)	8.65	(7.15 - 10.43)	1.92	(1.58 - 2.34)
Boards 20, 32, 54, and 69	9.18	(7.19 - 11.65)	6.36	(4.81 - 8.37)	1.41	(1.10 - 1.82)
Boards 21, 39, 51, 70, and 80	10.02	(7.91 - 12.62)	7.05	(5.34 - 9.24)	1.56	(1.20 - 2.01)
Boards 22, 74, and 87	11.82	(9.49 - 14.63)	7.60	(5.81 - 9.88)	1.91	(1.50 - 2.44)
Boards 23 and 45	10.75	(8.52 - 13.47)	7.10	(5.38 - 9.33)	1.93	(1.51 - 2.46)
Board 25 (Franklin)	16.33	(14.04 - 18.91)	9.95	(8.22 - 12.01)	2.22	(1.83 - 2.68)
Boards 27, 71, and 73	12.27	(9.71 - 15.40)	8.51	(6.45 - 11.13)	1.69	(1.31 - 2.19)
Boards 28, 43, and 67	10.51	(8.34 - 13.17)	6.70	(5.07 - 8.81)	2.14	(1.68 - 2.72)
Board 31 (Hamilton)	17.02	(14.43 - 19.98)	9.81	(7.89 - 12.14)	2.47	(1.98 - 3.09)
Board 48 (Lucas)	14.90	(11.92 - 18.47)	10.05	(7.76 - 12.92)	2.43	(1.87 - 3.15)
Boards 50 and 76	12.69	(10.39 - 15.41)	8.24	(6.47 - 10.43)	1.75	(1.40 - 2.20)
Board 57 (Montgomery)	13.62	(10.93 - 16.86)	8.20	(6.37 - 10.50)	2.02	(1.56 - 2.60)
Board 77 (Summit)	11.75	(9.35 - 14.67)	8.61	(6.67 - 11.06)	2.06	(1.59 - 2.68)

NOTE: For substate region definitions, see the "2014-2016 National Survey on Drug Use and Health Substate Region Definitions" at <https://www.samhsa.gov/data/>.

NOTE: Estimates along with the 95 percent Bayesian confidence (credible) intervals are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques.

¹ Average annual initiation of marijuana (%) = $100 * \{[X_1 \div (0.5 * X_1 + X_2)] \div 2\}$, where X_1 is the number of marijuana initiates in the past 24 months and X_2 is the number of individuals who never used marijuana (with the at-risk population defined as $0.5 * X_1 + X_2$). Both of the computation components, X_1 and X_2 , are based on a survey-weighted hierarchical Bayes estimation approach. The age group is based on a respondent's age at the time of the interview, not his or her age at first use. The Bayesian confidence (credible) intervals presented for Average Annual Estimates of First Use of Marijuana use a simultaneous solution for First Use of Marijuana and Never Used Marijuana outcomes, which had age group-specific random effects that were allowed to be correlated across the two outcomes. The associated Markov Chain Monte Carlo chains were used to calculate the posterior variance. For details, see Section B of the "2015-2016 NSDUH: Guide to State Tables and Summary of Small Area Estimation Methodology" at <https://www.samhsa.gov/data/>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2014, 2015, and 2016.

Table 36.2 Cocaine Use in the Past Year and Heroin Use in the Past Year in Ohio among Individuals Aged 12 or Older, by Substate Region: Percentages, Annual Averages Based on 2014, 2015, and 2016 NSDUHs

State/Substate Region	Cocaine Use in the Past Year		Heroin Use in the Past Year	
	Estimate	95% Confidence Interval	Estimate	95% Confidence Interval
Ohio	1.55	(1.28 - 1.87)	0.37	(0.25 - 0.54)
Boards 2, 46, 55, and 68	1.37	(0.82 - 2.29)	0.37	(0.21 - 0.67)
Boards 3, 52, and 85	1.37	(0.81 - 2.31)	0.36	(0.20 - 0.64)
Boards 4 and 78	1.49	(0.90 - 2.45)	0.37	(0.21 - 0.65)
Boards 5 and 60	1.75	(1.06 - 2.87)	0.40	(0.22 - 0.72)
Boards 7, 15, 41, 79, and 84	1.47	(0.87 - 2.48)	0.38	(0.22 - 0.67)
Boards 8, 13, and 83	1.40	(0.83 - 2.35)	0.31	(0.17 - 0.56)
Board 9 (Butler)	1.76	(1.07 - 2.88)	0.34	(0.18 - 0.62)
Board 12	1.78	(1.08 - 2.93)	0.35	(0.20 - 0.61)
Boards 18 and 47	1.21	(0.77 - 1.89)	0.36	(0.20 - 0.65)
Boards 20, 32, 54, and 69	1.23	(0.75 - 2.01)	0.41	(0.23 - 0.71)
Boards 21, 39, 51, 70, and 80	1.40	(0.88 - 2.24)	0.31	(0.18 - 0.55)
Boards 22, 74, and 87	1.41	(0.87 - 2.28)	0.35	(0.20 - 0.62)
Boards 23 and 45	1.16	(0.70 - 1.92)	0.40	(0.23 - 0.70)
Board 25 (Franklin)	2.15	(1.50 - 3.07)	0.30	(0.17 - 0.53)
Boards 27, 71, and 73	2.03	(1.18 - 3.44)	0.42	(0.24 - 0.73)
Boards 28, 43, and 67	1.27	(0.79 - 2.03)	0.34	(0.19 - 0.60)
Board 31 (Hamilton)	1.96	(1.25 - 3.05)	0.44	(0.25 - 0.78)
Board 48 (Lucas)	1.74	(1.05 - 2.88)	0.40	(0.23 - 0.70)
Boards 50 and 76	1.48	(0.91 - 2.39)	0.42	(0.24 - 0.72)
Board 57 (Montgomery)	1.42	(0.85 - 2.39)	0.40	(0.22 - 0.72)
Board 77 (Summit)	1.33	(0.82 - 2.16)	0.40	(0.22 - 0.72)

NOTE: For substate region definitions, see the "2014-2016 National Survey on Drug Use and Health Substate Region Definitions" at <https://www.samhsa.gov/data/>.

NOTE: Estimates along with the 95 percent Bayesian confidence (credible) intervals are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2014, 2015, and 2016.

Table 36.3 Alcohol Use in the Past Month, Underage Alcohol Use in the Past Month, and Alcohol Use Disorder in the Past Year in Ohio among Individuals Aged 12 or Older or Aged 12 to 20, by Substate Region: Percentages, Annual Averages Based on 2014, 2015, and 2016 NSDUHs

State/Substate Region	Alcohol Use in the Past Month (12 or Older)		Underage Alcohol Use in the Past Month (12 to 20)		Alcohol Use Disorder in the Past Year ¹ (12 or Older)	
	95% Confidence		95% Confidence		95% Confidence	
	Estimate	Interval	Estimate	Interval	Estimate	Interval
Ohio	52.28	(50.78 - 53.77)	21.44	(19.83 - 23.13)	6.30	(5.70 - 6.96)
Boards 2, 46, 55, and 68	52.67	(47.56 - 57.73)	23.30	(19.61 - 27.46)	6.24	(4.70 - 8.22)
Boards 3, 52, and 85	52.06	(46.99 - 57.10)	20.60	(17.11 - 24.58)	5.27	(3.99 - 6.93)
Boards 4 and 78	44.59	(39.30 - 50.02)	19.60	(16.22 - 23.48)	5.19	(3.93 - 6.84)
Boards 5 and 60	46.59	(41.23 - 52.03)	25.65	(21.36 - 30.45)	6.69	(5.16 - 8.64)
Boards 7, 15, 41, 79, and 84	45.03	(39.84 - 50.34)	19.77	(16.32 - 23.74)	5.54	(4.18 - 7.32)
Boards 8, 13, and 83	53.44	(48.38 - 58.43)	19.60	(16.33 - 23.34)	5.35	(4.00 - 7.13)
Board 9 (Butler)	51.61	(46.52 - 56.67)	22.52	(18.83 - 26.70)	6.53	(4.98 - 8.53)
Board 12	52.16	(46.80 - 57.46)	22.71	(18.98 - 26.94)	6.30	(4.78 - 8.26)
Boards 18 and 47	55.76	(52.32 - 59.14)	20.17	(17.49 - 23.14)	7.37	(5.95 - 9.09)
Boards 20, 32, 54, and 69	48.86	(44.02 - 53.72)	19.86	(16.60 - 23.59)	5.70	(4.36 - 7.43)
Boards 21, 39, 51, 70, and 80	46.51	(41.64 - 51.46)	18.27	(15.21 - 21.79)	5.40	(4.13 - 7.05)
Boards 22, 74, and 87	53.32	(48.27 - 58.31)	25.16	(21.50 - 29.21)	5.93	(4.55 - 7.68)
Boards 23 and 45	51.19	(46.45 - 55.91)	19.55	(16.43 - 23.11)	6.11	(4.69 - 7.93)
Board 25 (Franklin)	54.82	(51.23 - 58.36)	21.70	(18.87 - 24.83)	6.19	(4.98 - 7.66)
Boards 27, 71, and 73	42.67	(37.76 - 47.72)	18.06	(15.10 - 21.47)	5.91	(4.51 - 7.73)
Boards 28, 43, and 67	58.49	(54.09 - 62.76)	25.20	(21.70 - 29.06)	6.25	(4.79 - 8.12)
Board 31 (Hamilton)	58.02	(53.97 - 61.97)	23.08	(19.64 - 26.91)	7.47	(5.88 - 9.45)
Board 48 (Lucas)	49.44	(44.32 - 54.57)	20.02	(16.51 - 24.06)	6.51	(4.91 - 8.58)
Boards 50 and 76	50.53	(46.19 - 54.86)	21.69	(18.57 - 25.16)	5.95	(4.60 - 7.66)
Board 57 (Montgomery)	53.59	(48.67 - 58.44)	22.23	(18.69 - 26.23)	6.52	(4.97 - 8.51)
Board 77 (Summit)	55.78	(51.10 - 60.35)	22.93	(19.54 - 26.71)	6.40	(4.97 - 8.20)

NOTE: For substate region definitions, see the "2014-2016 National Survey on Drug Use and Health Substate Region Definitions" at <https://www.samhsa.gov/data/>.

NOTE: Estimates along with the 95 percent Bayesian confidence (credible) intervals are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques.

¹ Alcohol Use Disorder is defined as meeting criteria for alcohol dependence or abuse. Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2014, 2015, and 2016.

Table 36.4 Cigarette Use in the Past Month and Tobacco Product Use in the Past Month in Ohio among Individuals Aged 12 or Older, by Substate Region: Percentages, Annual Averages Based on 2014, 2015, and 2016 NSDUHs

State/Substate Region	Cigarette Use in the Past Month		Tobacco Product Use in the Past Month ¹	
	Estimate	95% Confidence Interval	Estimate	95% Confidence Interval
Ohio	24.27	(23.00 - 25.59)	29.86	(28.54 - 31.22)
Boards 2, 46, 55, and 68	27.59	(23.55 - 32.04)	33.26	(29.14 - 37.64)
Boards 3, 52, and 85	19.36	(16.06 - 23.15)	25.36	(21.77 - 29.31)
Boards 4 and 78	29.12	(24.92 - 33.71)	34.26	(30.05 - 38.73)
Boards 5 and 60	28.94	(24.60 - 33.70)	35.01	(30.86 - 39.39)
Boards 7, 15, 41, 79, and 84	26.45	(22.51 - 30.81)	34.35	(30.43 - 38.50)
Boards 8, 13, and 83	20.82	(17.61 - 24.45)	26.79	(23.30 - 30.58)
Board 9 (Butler)	22.42	(19.04 - 26.20)	28.89	(25.20 - 32.88)
Board 12	22.64	(19.02 - 26.72)	28.37	(24.51 - 32.58)
Boards 18 and 47	24.20	(21.56 - 27.06)	28.78	(25.97 - 31.76)
Boards 20, 32, 54, and 69	28.91	(25.05 - 33.09)	33.51	(29.68 - 37.57)
Boards 21, 39, 51, 70, and 80	23.81	(20.28 - 27.74)	29.45	(25.85 - 33.33)
Boards 22, 74, and 87	28.07	(24.08 - 32.45)	34.22	(30.30 - 38.37)
Boards 23 and 45	24.24	(20.83 - 28.01)	30.30	(26.59 - 34.28)
Board 25 (Franklin)	22.02	(19.22 - 25.10)	26.81	(24.01 - 29.81)
Boards 27, 71, and 73	29.54	(25.28 - 34.18)	35.93	(31.71 - 40.38)
Boards 28, 43, and 67	21.49	(18.31 - 25.06)	27.67	(24.27 - 31.34)
Board 31 (Hamilton)	23.42	(20.29 - 26.87)	28.24	(25.18 - 31.53)
Board 48 (Lucas)	26.55	(22.71 - 30.78)	32.22	(28.38 - 36.31)
Boards 50 and 76	24.67	(21.36 - 28.32)	31.40	(27.91 - 35.12)
Board 57 (Montgomery)	23.63	(20.04 - 27.65)	28.97	(25.34 - 32.89)
Board 77 (Summit)	19.95	(16.84 - 23.47)	25.39	(22.07 - 29.03)

NOTE: For substate region definitions, see the "2014-2016 National Survey on Drug Use and Health Substate Region Definitions" at <https://www.samhsa.gov/data/>.

NOTE: Estimates along with the 95 percent Bayesian confidence (credible) intervals are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., snuff, dip, chewing tobacco, or "snus"), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2014, 2015, and 2016.

Table 36.5 Serious Mental Illness in the Past Year, Any Mental Illness in the Past Year, and Received Mental Health Services in the Past Year in Ohio among Adults Aged 18 or Older, by Substate Region: Percentages, Annual Averages Based on 2014, 2015, and 2016 NSDUHs

State/Substate Region	Serious Mental Illness in the Past Year ¹		Any Mental Illness in the Past Year ¹		Received Mental Health Services in the Past Year ²	
	Estimate	95% Confidence Interval	Estimate	95% Confidence Interval	Estimate	95% Confidence Interval
Ohio	5.11	(4.52 - 5.77)	20.16	(18.96 - 21.43)	16.91	(15.77 - 18.12)
Boards 2, 46, 55, and 68	5.20	(3.93 - 6.85)	20.37	(17.19 - 23.98)	16.90	(14.25 - 19.93)
Boards 3, 52, and 85	4.84	(3.61 - 6.46)	18.49	(15.54 - 21.85)	17.04	(14.30 - 20.19)
Boards 4 and 78	5.57	(4.16 - 7.42)	20.67	(17.43 - 24.33)	16.72	(13.98 - 19.87)
Boards 5 and 60	6.16	(4.76 - 7.94)	21.32	(17.96 - 25.11)	17.83	(14.96 - 21.10)
Boards 7, 15, 41, 79, and 84	5.91	(4.47 - 7.79)	19.86	(16.83 - 23.29)	17.45	(14.66 - 20.64)
Boards 8, 13, and 83	4.80	(3.66 - 6.27)	18.98	(16.00 - 22.37)	16.29	(13.69 - 19.28)
Board 9 (Butler)	5.09	(3.86 - 6.67)	20.29	(17.13 - 23.87)	17.10	(14.45 - 20.13)
Board 12	5.03	(3.83 - 6.57)	20.34	(17.14 - 23.97)	16.34	(13.77 - 19.29)
Boards 18 and 47	4.77	(3.74 - 6.06)	19.89	(17.37 - 22.67)	16.31	(14.10 - 18.78)
Boards 20, 32, 54, and 69	5.38	(4.10 - 7.02)	19.11	(16.12 - 22.50)	16.13	(13.55 - 19.10)
Boards 21, 39, 51, 70, and 80	5.15	(3.94 - 6.70)	20.49	(17.21 - 24.22)	17.37	(14.58 - 20.58)
Boards 22, 74, and 87	5.17	(3.93 - 6.76)	19.98	(16.92 - 23.44)	16.35	(13.67 - 19.44)
Boards 23 and 45	5.39	(4.18 - 6.93)	20.98	(17.89 - 24.44)	18.46	(15.57 - 21.76)
Board 25 (Franklin)	4.64	(3.65 - 5.89)	21.58	(18.73 - 24.74)	16.89	(14.50 - 19.58)
Boards 27, 71, and 73	6.44	(4.87 - 8.46)	21.83	(18.40 - 25.70)	18.04	(15.14 - 21.36)
Boards 28, 43, and 67	4.76	(3.62 - 6.24)	18.56	(15.72 - 21.78)	17.02	(14.38 - 20.03)
Board 31 (Hamilton)	4.83	(3.72 - 6.23)	19.64	(16.92 - 22.67)	16.42	(13.94 - 19.23)
Board 48 (Lucas)	5.29	(4.00 - 6.98)	20.80	(17.63 - 24.39)	17.21	(14.48 - 20.34)
Boards 50 and 76	5.05	(3.90 - 6.51)	19.19	(16.37 - 22.36)	17.56	(14.95 - 20.51)
Board 57 (Montgomery)	5.21	(3.95 - 6.84)	20.98	(17.79 - 24.56)	17.23	(14.57 - 20.25)
Board 77 (Summit)	4.83	(3.68 - 6.32)	19.27	(16.36 - 22.56)	16.07	(13.56 - 18.95)

NOTE: For substate region definitions, see the "2014-2016 National Survey on Drug Use and Health Substate Region Definitions" at <https://www.samhsa.gov/data/>.

NOTE: Estimates along with the 95 percent Bayesian confidence (credible) intervals are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques.

¹ Mental illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder, assessed by the Mental Health Surveillance Study (MHSS) *Structured Clinical Interview for the Diagnostic and Statistical Manual of Mental Disorders—Fourth Edition—Research Version—Axis I Disorders* (MHSS-SCID), which is based on the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Any mental illness (AMI) includes adults with any mental disorder regardless of whether their disorder resulted in functional impairment. Serious mental illness (SMI) includes adults with any mental disorder that resulted in serious functional impairment. For details, see Section B of the "2015-2016 NSDUH: Guide to State Tables and Summary of Small Area Estimation Methodology" at <https://www.samhsa.gov/data/>.

² Mental health services are defined as having received inpatient treatment/counseling or outpatient treatment/counseling or having used prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2014, 2015, and 2016.

Table 36.6 Had Serious Thoughts of Suicide in the Past Year and Major Depressive Episode in the Past Year in Ohio among Adults Aged 18 or Older, by Substate Region: Percentages, Annual Averages Based on 2014, 2015, and 2016 NSDUHs

State/Substate Region	Had Serious Thoughts of Suicide in the Past Year ¹		Major Depressive Episode in the Past Year ²	
	Estimate	95% Confidence Interval	Estimate	95% Confidence Interval
Ohio	4.53	(3.99 - 5.12)	7.54	(6.81 - 8.33)
Boards 2, 46, 55, and 68	4.41	(3.42 - 5.67)	7.13	(5.56 - 9.11)
Boards 3, 52, and 85	4.59	(3.54 - 5.95)	7.04	(5.50 - 8.96)
Boards 4 and 78	4.85	(3.64 - 6.42)	7.29	(5.71 - 9.27)
Boards 5 and 60	5.10	(3.96 - 6.55)	8.27	(6.57 - 10.35)
Boards 7, 15, 41, 79, and 84	4.95	(3.76 - 6.50)	7.75	(6.06 - 9.87)
Boards 8, 13, and 83	4.02	(3.09 - 5.22)	6.91	(5.41 - 8.80)
Board 9 (Butler)	4.71	(3.67 - 6.03)	7.54	(5.92 - 9.56)
Board 12	4.77	(3.67 - 6.18)	7.54	(5.91 - 9.57)
Boards 18 and 47	4.16	(3.32 - 5.21)	7.28	(5.91 - 8.94)
Boards 20, 32, 54, and 69	4.28	(3.32 - 5.50)	7.38	(5.74 - 9.46)
Boards 21, 39, 51, 70, and 80	4.41	(3.40 - 5.70)	7.79	(6.13 - 9.86)
Boards 22, 74, and 87	4.71	(3.64 - 6.07)	7.76	(6.16 - 9.73)
Boards 23 and 45	4.38	(3.40 - 5.62)	8.32	(6.60 - 10.44)
Board 25 (Franklin)	4.40	(3.49 - 5.52)	7.86	(6.46 - 9.53)
Boards 27, 71, and 73	5.02	(3.84 - 6.54)	7.86	(6.18 - 9.94)
Boards 28, 43, and 67	4.62	(3.56 - 5.97)	7.30	(5.78 - 9.19)
Board 31 (Hamilton)	4.12	(3.22 - 5.27)	7.64	(6.09 - 9.54)
Board 48 (Lucas)	4.96	(3.82 - 6.43)	7.46	(5.84 - 9.48)
Boards 50 and 76	4.63	(3.60 - 5.94)	6.82	(5.41 - 8.55)
Board 57 (Montgomery)	4.94	(3.84 - 6.33)	8.01	(6.30 - 10.14)
Board 77 (Summit)	4.60	(3.56 - 5.93)	7.67	(6.01 - 9.75)

NOTE: For substate region definitions, see the "2014-2016 National Survey on Drug Use and Health Substate Region Definitions" at <https://www.samhsa.gov/data/>.


NOTE: Estimates along with the 95 percent Bayesian confidence (credible) intervals are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques.

¹ Respondents were asked, "At any time in the past 12 months, did you seriously think about trying to kill yourself?" If they answered "Yes," they were categorized as having serious thoughts of suicide in the past year.

² Major depressive episode (MDE) is defined as in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV), which specifies a period of at least 2 weeks when an individual experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of specified depression symptoms. For details, see Section B in the "2015-2016 National Survey on Drug Use and Health: Guide to State Tables and Summary of Small Area Estimation Methodology" at <https://www.samhsa.gov/data/>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2014, 2015, and 2016.


Figure 36.1 Marijuana Use in the Past Year among Individuals Aged 12 or Older in Ohio, by Substate Region: Percentages, Annual Averages Based on 2014, 2015, and 2016 NSDUHs



NOTE: The legend's ranges were created by dividing 395 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2014-2016 National Survey on Drug Use and Health Substate Region Definitions" at <https://www.samhsa.gov/data/>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2014, 2015, and 2016.


Figure 36.2 Marijuana Use in the Past Month among Individuals Aged 12 or Older in Ohio, by Substate Region: Percentages, Annual Averages Based on 2014, 2015, and 2016 NSDUHs



NOTE: The legend's ranges were created by dividing 395 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2014-2016 National Survey on Drug Use and Health Substate Region Definitions" at <https://www.samhsa.gov/data/>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2014, 2015, and 2016.


Figure 36.3 First Use of Marijuana among Individuals Aged 12 or Older in Ohio, by Substate Region: Average Annual Incidence Estimates (Expressed as Percentages of the At-Risk Population) Based on 2014, 2015, and 2016 NSDUHs



NOTE: The legend's ranges were created by dividing 395 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2014-2016 National Survey on Drug Use and Health Substate Region Definitions" at <https://www.samhsa.gov/data/>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2014, 2015, and 2016.


Figure 36.4 Cocaine Use in the Past Year among Individuals Aged 12 or Older in Ohio, by Substate Region: Percentages, Annual Averages Based on 2014, 2015, and 2016 NSDUHs



NOTE: The legend's ranges were created by dividing 395 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2014-2016 National Survey on Drug Use and Health Substate Region Definitions" at <https://www.samhsa.gov/data/>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2014, 2015, and 2016.


Figure 36.5 Heroin Use in the Past Year among Individuals Aged 12 or Older in Ohio, by Substate Region: Percentages, Annual Averages Based on 2014, 2015, and 2016 NSDUHs



NOTE: The legend's ranges were created by dividing 395 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2014-2016 National Survey on Drug Use and Health Substate Region Definitions" at <https://www.samhsa.gov/data/>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2014, 2015, and 2016.


Figure 36.6 Alcohol Use in the Past Month among Individuals Aged 12 or Older in Ohio, by Substate Region: Percentages, Annual Averages Based on 2014, 2015, and 2016 NSDUHs



NOTE: The legend's ranges were created by dividing 395 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2014-2016 National Survey on Drug Use and Health Substate Region Definitions" at <https://www.samhsa.gov/data/>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2014, 2015, and 2016.


Figure 36.7 Alcohol Use in the Past Month among Individuals Aged 12 to 20 in Ohio, by Substate Region: Percentages, Annual Averages Based on 2014, 2015, and 2016 NSDUHs



NOTE: The legend's ranges were created by dividing 395 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2014-2016 National Survey on Drug Use and Health Substate Region Definitions" at <https://www.samhsa.gov/data/>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2014, 2015, and 2016.


Figure 36.8 Tobacco Product Use in the Past Month among Individuals Aged 12 or Older in Ohio, by Substate Region: Percentages, Annual Averages Based on 2014, 2015, and 2016 NSDUHs



NOTE: The legend's ranges were created by dividing 395 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2014-2016 National Survey on Drug Use and Health Substate Region Definitions" at <https://www.samhsa.gov/data/>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2014, 2015, and 2016.


Figure 36.9 Cigarette Use in the Past Month among Individuals Aged 12 or Older in Ohio, by Substate Region: Percentages, Annual Averages Based on 2014, 2015, and 2016 NSDUHs



NOTE: The legend's ranges were created by dividing 395 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2014-2016 National Survey on Drug Use and Health Substate Region Definitions" at <https://www.samhsa.gov/data/>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2014, 2015, and 2016.


Figure 36.10 Alcohol Use Disorder in the Past Year among Individuals Aged 12 or Older in Ohio, by Substate Region: Percentages, Annual Averages Based on 2014, 2015, and 2016 NSDUHs



NOTE: The legend's ranges were created by dividing 395 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2014-2016 National Survey on Drug Use and Health Substate Region Definitions" at <https://www.samhsa.gov/data/>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2014, 2015, and 2016.


Figure 36.11 Serious Mental Illness in the Past Year among Adults Aged 18 or Older in Ohio, by Substate Region: Percentages, Annual Averages Based on 2014, 2015, and 2016 NSDUHs



NOTE: The legend's ranges were created by dividing 395 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2014-2016 National Survey on Drug Use and Health Substate Region Definitions" at <https://www.samhsa.gov/data/>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2014, 2015, and 2016.


Figure 36.12 Any Mental Illness in the Past Year among Adults Aged 18 or Older in Ohio, by Substate Region: Percentages, Annual Averages Based on 2014, 2015, and 2016 NSDUHs



NOTE: The legend's ranges were created by dividing 395 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2014-2016 National Survey on Drug Use and Health Substate Region Definitions" at <https://www.samhsa.gov/data/>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2014, 2015, and 2016.


Figure 36.13 Received Mental Health Services in the Past Year among Adults Aged 18 or Older in Ohio, by Substate Region: Percentages, Annual Averages Based on 2014, 2015, and 2016 NSDUHs



NOTE: The legend's ranges were created by dividing 395 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2014-2016 National Survey on Drug Use and Health Substate Region Definitions" at <https://www.samhsa.gov/data/>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2014, 2015, and 2016.


Figure 36.14 Had Serious Thoughts of Suicide in the Past Year among Adults Aged 18 or Older in Ohio, by Substate Region: Percentages, Annual Averages Based on 2014, 2015, and 2016 NSDUHs



NOTE: The legend's ranges were created by dividing 395 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2014-2016 National Survey on Drug Use and Health Substate Region Definitions" at <https://www.samhsa.gov/data/>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2014, 2015, and 2016.

Figure 36.15 Major Depressive Episode in the Past Year among Adults Aged 18 or Older in Ohio, by Substate Region: Percentages, Annual Averages Based on 2014, 2015, and 2016 NSDUHs



NOTE: The legend's ranges were created by dividing 395 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2014-2016 National Survey on Drug Use and Health Substate Region Definitions" at <https://www.samhsa.gov/data/>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2014, 2015, and 2016.