

**Table 1.108A Any Use and Misuse of Tranquilizer Subtypes in Past Year among Persons Aged 12 or Older and Misuse of Tranquilizer Subtypes in Past Year among Past Year Any Users of Tranquilizer Subtypes Aged 12 or Older: Numbers in Thousands, 2018 and 2019**

<b>Tranquilizer Subtype</b>	<b>Any Use in Past Year (2018)</b>	<b>Any Use in Past Year (2019)</b>	<b>Misuse in Past Year among Total Population (2018)</b>	<b>Misuse in Past Year among Total Population (2019)</b>	<b>Misuse in Past Year among Past Year Any Users of Tranquilizer Subtypes (2018)</b>	<b>Misuse in Past Year among Past Year Any Users of Tranquilizer Subtypes (2019)</b>
<b>ANY PRESCRIPTION TRANQUILIZER</b>	37,610 <sup>a</sup>	35,601	5,742	5,220	5,742	5,220
Benzodiazepine Tranquilizers	29,505	28,316	5,379	4,819	5,379	4,819
Alprazolam Products	17,690	16,748	4,171 <sup>a</sup>	3,668	4,171 <sup>a</sup>	3,668
Lorazepam Products	7,630	7,787	772	795	772	795
Clonazepam Products	6,041	5,998	1,029	1,029	1,029	1,029
Diazepam Products	6,361	5,672	879	758	879	758
Muscle Relaxants	6,418	6,049	543	552	543	552
Cyclobenzaprine	5,082	4,955	332	361	332	361
Soma <sup>®</sup>	1,486	1,182	211	223	211	223
Any Other Prescription Tranquilizer <sup>1</sup>	6,054	5,738	94	97	94	97

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

<sup>a</sup> The difference between this estimate and the 2019 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

<sup>1</sup> Includes use or misuse of tranquilizers containing other active ingredients. Reports of misuse of "any other prescription tranquilizer" that correspond only to the specific tranquilizer categories shown in the table are excluded from estimates for Any Other Prescription Tranquilizer and are included instead in the relevant tranquilizer category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

**Table 1.108B Any Use and Misuse of Tranquilizer Subtypes in Past Year among Persons Aged 12 or Older and Misuse of Tranquilizer Subtypes in Past Year among Past Year Any Users of Tranquilizer Subtypes Aged 12 or Older: Percentages, 2018 and 2019**

<b>Tranquilizer Subtype</b>	<b>Any Use in Past Year (2018)</b>	<b>Any Use in Past Year (2019)</b>	<b>Misuse in Past Year among Total Population (2018)</b>	<b>Misuse in Past Year among Total Population (2019)</b>	<b>Misuse in Past Year among Past Year Any Users of Tranquilizer Subtypes (2018)</b>	<b>Misuse in Past Year among Past Year Any Users of Tranquilizer Subtypes (2019)</b>
<b>ANY PRESCRIPTION TRANQUILIZER</b>	13.7 <sup>a</sup>	12.9	2.1	1.9	15.3	14.7
Benzodiazepine Tranquilizers	10.8	10.3	2.0	1.8	18.2	17.0
Alprazolam Products	6.5	6.1	1.5 <sup>a</sup>	1.3	23.6	21.9
Lorazepam Products	2.8	2.8	0.3	0.3	10.1	10.2
Clonazepam Products	2.2	2.2	0.4	0.4	17.0	17.2
Diazepam Products	2.3	2.1	0.3	0.3	13.8	13.4
Muscle Relaxants	2.3	2.2	0.2	0.2	8.5	9.1
Cyclobenzaprine	1.9	1.8	0.1	0.1	6.5	7.3
Soma <sup>®</sup>	0.5	0.4	0.1	0.1	14.2	18.8
Any Other Prescription Tranquilizer <sup>1</sup>	2.2	2.1	0.0	0.0	1.6	1.7

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

<sup>a</sup> The difference between this estimate and the 2019 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

<sup>1</sup> Includes use or misuse of tranquilizers containing other active ingredients. Reports of misuse of "any other prescription tranquilizer" that correspond only to the specific tranquilizer categories shown in the table are excluded from estimates for Any Other Prescription Tranquilizer and are included instead in the relevant tranquilizer category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

**Table 1.109A Any Use and Misuse of Tranquilizer Subtypes in Past Year among Persons Aged 12 to 17 and Misuse of Tranquilizer Subtypes in Past Year among Past Year Any Users of Tranquilizer Subtypes Aged 12 to 17: Numbers in Thousands, 2018 and 2019**

<b>Tranquilizer Subtype</b>	<b>Any Use in Past Year (2018)</b>	<b>Any Use in Past Year (2019)</b>	<b>Misuse in Past Year among Total Population (2018)</b>	<b>Misuse in Past Year among Total Population (2019)</b>	<b>Misuse in Past Year among Past Year Any Users of Tranquilizer Subtypes (2018)</b>	<b>Misuse in Past Year among Past Year Any Users of Tranquilizer Subtypes (2019)</b>
<b>ANY PRESCRIPTION TRANQUILIZER</b>	1,001	1,010	423	392	423	392
Benzodiazepine Tranquilizers	732	700	399	371	399	371
Alprazolam Products	620	613	372	359	372	359
Lorazepam Products	97	84	24	9	*	*
Clonazepam Products	70	83	25	33	*	*
Diazepam Products	84	80	28	16	*	*
Muscle Relaxants	31	45	11	15	*	*
Cyclobenzaprine	18	32	2	4	*	*
Soma®	13	11	10	10	*	*
Any Other Prescription Tranquilizer <sup>1</sup>	299	367	13	16	13	16

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

<sup>a</sup> The difference between this estimate and the 2019 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

<sup>1</sup> Includes use or misuse of tranquilizers containing other active ingredients. Reports of misuse of "any other prescription tranquilizer" that correspond only to the specific tranquilizer categories shown in the table are excluded from estimates for Any Other Prescription Tranquilizer and are included instead in the relevant tranquilizer category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

**Table 1.109B Any Use and Misuse of Tranquilizer Subtypes in Past Year among Persons Aged 12 to 17 and Misuse of Tranquilizer Subtypes in Past Year among Past Year Any Users of Tranquilizer Subtypes Aged 12 to 17: Percentages, 2018 and 2019**

<b>Tranquilizer Subtype</b>	<b>Any Use in Past Year (2018)</b>	<b>Any Use in Past Year (2019)</b>	<b>Misuse in Past Year among Total Population (2018)</b>	<b>Misuse in Past Year among Total Population (2019)</b>	<b>Misuse in Past Year among Past Year Any Users of Tranquilizer Subtypes (2018)</b>	<b>Misuse in Past Year among Past Year Any Users of Tranquilizer Subtypes (2019)</b>
<b>ANY PRESCRIPTION TRANQUILIZER</b>	4.0	4.1	1.7	1.6	42.2	38.8
Benzodiazepine Tranquilizers	2.9	2.8	1.6	1.5	54.6	52.9
Alprazolam Products	2.5	2.5	1.5	1.4	59.9	58.6
Lorazepam Products	0.4	0.3	0.1	0.0	*	*
Clonazepam Products	0.3	0.3	0.1	0.1	*	*
Diazepam Products	0.3	0.3	0.1	0.1	*	*
Muscle Relaxants	0.1	0.2	0.0	0.1	*	*
Cyclobenzaprine	0.1	0.1	0.0	0.0	*	*
Soma®	0.1	0.0	0.0	0.0	*	*
Any Other Prescription Tranquilizer <sup>1</sup>	1.2	1.5	0.1	0.1	4.4	4.3

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

<sup>a</sup> The difference between this estimate and the 2019 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

<sup>1</sup> Includes use or misuse of tranquilizers containing other active ingredients. Reports of misuse of "any other prescription tranquilizer" that correspond only to the specific tranquilizer categories shown in the table are excluded from estimates for Any Other Prescription Tranquilizer and are included instead in the relevant tranquilizer category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

**Table 1.110A Any Use and Misuse of Tranquilizer Subtypes in Past Year among Persons Aged 18 or Older and Misuse of Tranquilizer Subtypes in Past Year among Past Year Any Users of Tranquilizer Subtypes Aged 18 or Older: Numbers in Thousands, 2018 and 2019**

<b>Tranquilizer Subtype</b>	<b>Any Use in Past Year (2018)</b>	<b>Any Use in Past Year (2019)</b>	<b>Misuse in Past Year among Total Population (2018)</b>	<b>Misuse in Past Year among Total Population (2019)</b>	<b>Misuse in Past Year among Past Year Any Users of Tranquilizer Subtypes (2018)</b>	<b>Misuse in Past Year among Past Year Any Users of Tranquilizer Subtypes (2019)</b>
<b>ANY PRESCRIPTION TRANQUILIZER</b>	36,609 <sup>a</sup>	34,591	5,319	4,829	5,319	4,829
Benzodiazepine Tranquilizers	28,773	27,615	4,980	4,448	4,980	4,448
Alprazolam Products	17,070	16,134	3,800 <sup>a</sup>	3,308	3,800 <sup>a</sup>	3,308
Lorazepam Products	7,534	7,703	748	786	748	786
Clonazepam Products	5,971	5,916	1,004	996	1,004	996
Diazepam Products	6,277	5,592	851	742	851	742
Muscle Relaxants	6,387	6,005	531	537	531	537
Cyclobenzaprine	5,064	4,923	330	358	330	358
Soma <sup>®</sup>	1,473	1,170	201	213	201	213
Any Other Prescription Tranquilizer <sup>1</sup>	5,755	5,371	81	81	81	81

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

<sup>a</sup> The difference between this estimate and the 2019 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

<sup>1</sup> Includes use or misuse of tranquilizers containing other active ingredients. Reports of misuse of "any other prescription tranquilizer" that correspond only to the specific tranquilizer categories shown in the table are excluded from estimates for Any Other Prescription Tranquilizer and are included instead in the relevant tranquilizer category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

**Table 1.110B Any Use and Misuse of Tranquilizer Subtypes in Past Year among Persons Aged 18 or Older and Misuse of Tranquilizer Subtypes in Past Year among Past Year Any Users of Tranquilizer Subtypes Aged 18 or Older: Percentages, 2018 and 2019**

<b>Tranquilizer Subtype</b>	<b>Any Use in Past Year (2018)</b>	<b>Any Use in Past Year (2019)</b>	<b>Misuse in Past Year among Total Population (2018)</b>	<b>Misuse in Past Year among Total Population (2019)</b>	<b>Misuse in Past Year among Past Year Any Users of Tranquilizer Subtypes (2018)</b>	<b>Misuse in Past Year among Past Year Any Users of Tranquilizer Subtypes (2019)</b>
<b>ANY PRESCRIPTION TRANQUILIZER</b>	14.7 <sup>a</sup>	13.8	2.1	1.9	14.5	14.0
Benzodiazepine Tranquilizers	11.6	11.0	2.0	1.8	17.3	16.1
Alprazolam Products	6.9	6.4	1.5 <sup>a</sup>	1.3	22.3	20.5
Lorazepam Products	3.0	3.1	0.3	0.3	9.9	10.2
Clonazepam Products	2.4	2.4	0.4	0.4	16.8	16.8
Diazepam Products	2.5	2.2	0.3	0.3	13.5	13.3
Muscle Relaxants	2.6	2.4	0.2	0.2	8.3	8.9
Cyclobenzaprine	2.0	2.0	0.1	0.1	6.5	7.3
Soma <sup>®</sup>	0.6	0.5	0.1	0.1	13.7	18.2
Any Other Prescription Tranquilizer <sup>1</sup>	2.3	2.1	0.0	0.0	1.4	1.5

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

<sup>a</sup> The difference between this estimate and the 2019 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

<sup>1</sup> Includes use or misuse of tranquilizers containing other active ingredients. Reports of misuse of "any other prescription tranquilizer" that correspond only to the specific tranquilizer categories shown in the table are excluded from estimates for Any Other Prescription Tranquilizer and are included instead in the relevant tranquilizer category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

**Table 1.111A Any Use and Misuse of Tranquilizer Subtypes in Past Year among Persons Aged 18 to 25 and Misuse of Tranquilizer Subtypes in Past Year among Past Year Any Users of Tranquilizer Subtypes Aged 18 to 25: Numbers in Thousands, 2018 and 2019**

<b>Tranquilizer Subtype</b>	<b>Any Use in Past Year (2018)</b>	<b>Any Use in Past Year (2019)</b>	<b>Misuse in Past Year among Total Population (2018)</b>	<b>Misuse in Past Year among Total Population (2019)</b>	<b>Misuse in Past Year among Past Year Any Users of Tranquilizer Subtypes (2018)</b>	<b>Misuse in Past Year among Past Year Any Users of Tranquilizer Subtypes (2019)</b>
<b>ANY PRESCRIPTION TRANQUILIZER</b>	3,681	3,419	1,566 <sup>a</sup>	1,313	1,566 <sup>a</sup>	1,313
Benzodiazepine Tranquilizers	3,180 <sup>a</sup>	2,922	1,525 <sup>a</sup>	1,260	1,525 <sup>a</sup>	1,260
Alprazolam Products	2,531 <sup>a</sup>	2,229	1,385 <sup>a</sup>	1,091	1,385 <sup>a</sup>	1,091
Lorazepam Products	593	630	137	132	137	132
Clonazepam Products	639	599	250	214	250	214
Diazepam Products	452	451	149	138	149	138
Muscle Relaxants	364	314	77	59	77	59
Cyclobenzaprine	253	258	35	35	35	35
Soma <sup>®</sup>	117	70	43	33	*	*
Any Other Prescription Tranquilizer <sup>1</sup>	534	515	34	33	34	33

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

<sup>a</sup> The difference between this estimate and the 2019 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

<sup>1</sup> Includes use or misuse of tranquilizers containing other active ingredients. Reports of misuse of "any other prescription tranquilizer" that correspond only to the specific tranquilizer categories shown in the table are excluded from estimates for Any Other Prescription Tranquilizer and are included instead in the relevant tranquilizer category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

**Table 1.111B Any Use and Misuse of Tranquilizer Subtypes in Past Year among Persons Aged 18 to 25 and Misuse of Tranquilizer Subtypes in Past Year among Past Year Any Users of Tranquilizer Subtypes Aged 18 to 25: Percentages, 2018 and 2019**

<b>Tranquilizer Subtype</b>	<b>Any Use in Past Year (2018)</b>	<b>Any Use in Past Year (2019)</b>	<b>Misuse in Past Year among Total Population (2018)</b>	<b>Misuse in Past Year among Total Population (2019)</b>	<b>Misuse in Past Year among Past Year Any Users of Tranquilizer Subtypes (2018)</b>	<b>Misuse in Past Year among Past Year Any Users of Tranquilizer Subtypes (2019)</b>
<b>ANY PRESCRIPTION TRANQUILIZER</b>	10.8	10.1	4.6 <sup>a</sup>	3.9	42.5	38.4
Benzodiazepine Tranquilizers	9.3	8.7	4.5 <sup>a</sup>	3.7	47.9 <sup>a</sup>	43.1
Alprazolam Products	7.4 <sup>a</sup>	6.6	4.1 <sup>a</sup>	3.2	54.7 <sup>a</sup>	49.0
Lorazepam Products	1.7	1.9	0.4	0.4	23.1	21.0
Clonazepam Products	1.9	1.8	0.7	0.6	39.1	35.7
Diazepam Products	1.3	1.3	0.4	0.4	33.0	30.6
Muscle Relaxants	1.1	0.9	0.2	0.2	21.2	18.9
Cyclobenzaprine	0.7	0.8	0.1	0.1	14.0	13.5
Soma <sup>®</sup>	0.3	0.2	0.1	0.1	*	*
Any Other Prescription Tranquilizer <sup>1</sup>	1.6	1.5	0.1	0.1	6.3	6.4

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

<sup>a</sup> The difference between this estimate and the 2019 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

<sup>1</sup> Includes use or misuse of tranquilizers containing other active ingredients. Reports of misuse of "any other prescription tranquilizer" that correspond only to the specific tranquilizer categories shown in the table are excluded from estimates for Any Other Prescription Tranquilizer and are included instead in the relevant tranquilizer category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

**Table 1.112A Any Use and Misuse of Tranquilizer Subtypes in Past Year among Persons Aged 26 or Older and Misuse of Tranquilizer Subtypes in Past Year among Past Year Any Users of Tranquilizer Subtypes Aged 26 or Older: Numbers in Thousands, 2018 and 2019**

<b>Tranquilizer Subtype</b>	<b>Any Use in Past Year (2018)</b>	<b>Any Use in Past Year (2019)</b>	<b>Misuse in Past Year among Total Population (2018)</b>	<b>Misuse in Past Year among Total Population (2019)</b>	<b>Misuse in Past Year among Past Year Any Users of Tranquilizer Subtypes (2018)</b>	<b>Misuse in Past Year among Past Year Any Users of Tranquilizer Subtypes (2019)</b>
<b>ANY PRESCRIPTION TRANQUILIZER</b>	32,928 <sup>a</sup>	31,172	3,753	3,516	3,753	3,516
Benzodiazepine Tranquilizers	25,593	24,693	3,455	3,188	3,455	3,188
Alprazolam Products	14,539	13,906	2,415	2,217	2,415	2,217
Lorazepam Products	6,940	7,074	611	654	611	654
Clonazepam Products	5,332	5,317	754	782	754	782
Diazepam Products	5,825	5,141	701	604	701	604
Muscle Relaxants	6,022	5,690	454	477	454	477
Cyclobenzaprine	4,811	4,665	295	323	295	323
Soma <sup>®</sup>	1,356	1,100	158	179	158	179
Any Other Prescription Tranquilizer <sup>1</sup>	5,221	4,856	47	48	47	48

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

<sup>a</sup> The difference between this estimate and the 2019 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

<sup>1</sup> Includes use or misuse of tranquilizers containing other active ingredients. Reports of misuse of "any other prescription tranquilizer" that correspond only to the specific tranquilizer categories shown in the table are excluded from estimates for Any Other Prescription Tranquilizer and are included instead in the relevant tranquilizer category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

**Table 1.112B Any Use and Misuse of Tranquilizer Subtypes in Past Year among Persons Aged 26 or Older and Misuse of Tranquilizer Subtypes in Past Year among Past Year Any Users of Tranquilizer Subtypes Aged 26 or Older: Percentages, 2018 and 2019**

<b>Tranquilizer Subtype</b>	<b>Any Use in Past Year (2018)</b>	<b>Any Use in Past Year (2019)</b>	<b>Misuse in Past Year among Total Population (2018)</b>	<b>Misuse in Past Year among Total Population (2019)</b>	<b>Misuse in Past Year among Past Year Any Users of Tranquilizer Subtypes (2018)</b>	<b>Misuse in Past Year among Past Year Any Users of Tranquilizer Subtypes (2019)</b>
<b>ANY PRESCRIPTION TRANQUILIZER</b>	15.3 <sup>a</sup>	14.4	1.7	1.6	11.4	11.3
Benzodiazepine Tranquilizers	11.9	11.4	1.6	1.5	13.5	12.9
Alprazolam Products	6.8	6.4	1.1	1.0	16.6	15.9
Lorazepam Products	3.2	3.3	0.3	0.3	8.8	9.2
Clonazepam Products	2.5	2.5	0.4	0.4	14.1	14.7
Diazepam Products	2.7	2.4	0.3	0.3	12.0	11.7
Muscle Relaxants	2.8	2.6	0.2	0.2	7.5	8.4
Cyclobenzaprine	2.2	2.2	0.1	0.1	6.1	6.9
Soma <sup>®</sup>	0.6	0.5	0.1	0.1	11.7	16.3
Any Other Prescription Tranquilizer <sup>1</sup>	2.4	2.2	0.0	0.0	0.9	1.0

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

<sup>a</sup> The difference between this estimate and the 2019 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

<sup>1</sup> Includes use or misuse of tranquilizers containing other active ingredients. Reports of misuse of "any other prescription tranquilizer" that correspond only to the specific tranquilizer categories shown in the table are excluded from estimates for Any Other Prescription Tranquilizer and are included instead in the relevant tranquilizer category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.