

Table 1.98C Any Use and Misuse of Pain Reliever Subtypes in Past Year among Persons Aged 12 or Older and Misuse of Pain Reliever Subtypes in Past Year among Past Year Any Users of Pain Reliever Subtypes Aged 12 or Older: Standard Errors of Numbers in Thousands, 2018 and 2019

Pain Reliever Subtype	Any Use in Past Year (2018)	Any Use in Past Year (2019)	Misuse in Past Year among Total Population (2018)	Misuse in Past Year among Total Population (2019)	Misuse in Past Year among Past Year Any Users of Pain Reliever Subtypes (2018)	Misuse in Past Year among Past Year Any Users of Pain Reliever Subtypes (2019)
ANY PRESCRIPTION PAIN RELIEVER	787	790	286	289	290	298
Hydrocodone Products	670	651	236	209	238	212
Zohydro [®] ER	57	76	8	13	*	*
Oxycodone Products	490	495	158	151	158	153
OxyContin [®]	310	300	88	89	89	89
Tramadol Products	423	434	113	110	113	110
Codeine Products	511	500	145	156	144	156
Morphine Products	250	246	57	59	57	60
Fentanyl Products	140	166	36	44	36	44
Buprenorphine Products	156	135	71	67	71	67
Oxymorphone Products	82	93	37	41	37	41
Demerol [®]	135	95	19	16	19	16
Hydromorphone Products	134	134	38	34	38	34
Methadone	101	101	52	50	51	50
Any Other Prescription Pain Reliever ¹	458	443	73	81	73	81

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

¹ Includes use or misuse of pain relievers containing other active ingredients. Reports of misuse of "any other prescription pain reliever" that correspond only to the specific pain reliever categories shown in the table are excluded from estimates for Any Other Prescription Pain Reliever and are included instead in the relevant pain reliever category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

Table 1.98D Any Use and Misuse of Pain Reliever Subtypes in Past Year among Persons Aged 12 or Older and Misuse of Pain Reliever Subtypes in Past Year among Past Year Any Users of Pain Reliever Subtypes Aged 12 or Older: Standard Errors of Percentages, 2018 and 2019

Pain Reliever Subtype	Any Use in Past Year (2018)	Any Use in Past Year (2019)	Misuse in Past Year among Total Population (2018)	Misuse in Past Year among Total Population (2019)	Misuse in Past Year among Past Year Any Users of Pain Reliever Subtypes (2018)	Misuse in Past Year among Past Year Any Users of Pain Reliever Subtypes (2019)
ANY PRESCRIPTION PAIN RELIEVER	0.29	0.29	0.10	0.11	0.32	0.34
Hydrocodone Products	0.24	0.24	0.09	0.08	0.47	0.47
Zohydro [®] ER	0.02	0.03	0.00	0.00	*	*
Oxycodone Products	0.18	0.18	0.06	0.05	0.56	0.61
OxyContin [®]	0.11	0.11	0.03	0.03	0.95	1.00
Tramadol Products	0.15	0.16	0.04	0.04	0.61	0.63
Codeine Products	0.19	0.18	0.05	0.06	0.55	0.62
Morphine Products	0.09	0.09	0.02	0.02	0.91	0.99
Fentanyl Products	0.05	0.06	0.01	0.02	1.66	1.73
Buprenorphine Products	0.06	0.05	0.03	0.02	2.40	2.47
Oxymorphone Products	0.03	0.03	0.01	0.01	4.01	4.52
Demerol [®]	0.05	0.03	0.01	0.01	1.63	1.77
Hydromorphone Products	0.05	0.05	0.01	0.01	2.10	2.06
Methadone	0.04	0.04	0.02	0.02	4.04	4.10
Any Other Prescription Pain Reliever ¹	0.17	0.16	0.03	0.03	0.30	0.35

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

¹ Includes use or misuse of pain relievers containing other active ingredients. Reports of misuse of "any other prescription pain reliever" that correspond only to the specific pain reliever categories shown in the table are excluded from estimates for Any Other Prescription Pain Reliever and are included instead in the relevant pain reliever category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

Table 1.98P Any Use and Misuse of Pain Reliever Subtypes in Past Year among Persons Aged 12 or Older and Misuse of Pain Reliever Subtypes in Past Year among Past Year Any Users of Pain Reliever Subtypes Aged 12 or Older: *P* Values from Tests of Differences of Percentages, 2019 Versus 2018

Pain Reliever Subtype	Any Use in Past Year (2018)	Any Use in Past Year (2019)	Misuse in Past Year among Total Population (2018)	Misuse in Past Year among Total Population (2019)	Misuse in Past Year among Past Year Any Users of Pain Reliever Subtypes (2018)	Misuse in Past Year among Past Year Any Users of Pain Reliever Subtypes (2019)
ANY PRESCRIPTION PAIN RELIEVER	0.0001		0.4890		0.5310	
Hydrocodone Products	0.0000		0.1514		0.8748	
Zohydro [®] ER	0.7165		0.2490		*	
Oxycodone Products	0.0023		0.3445		0.7350	
OxyContin [®]	0.4364		0.6302		0.9242	
Tramadol Products	0.1270		0.4319		0.7298	
Codeine Products	0.0426		0.9023		0.5707	
Morphine Products	0.2878		0.6860		0.9439	
Fentanyl Products	0.0639		0.9813		0.3902	
Buprenorphine Products	0.6882		0.7250		0.8921	
Oxymorphone Products	0.8922		0.9387		0.8585	
Demerol [®]	0.0833		0.9062		0.6540	
Hydromorphone Products	0.3913		0.3151		0.5215	
Methadone	0.8248		0.8119		0.8742	
Any Other Prescription Pain Reliever ¹	0.0318		0.2054		0.0777	

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

¹ Includes use or misuse of pain relievers containing other active ingredients. Reports of misuse of "any other prescription pain reliever" that correspond only to the specific pain reliever categories shown in the table are excluded from estimates for Any Other Prescription Pain Reliever and are included instead in the relevant pain reliever category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

Table 1.99C Any Use and Misuse of Pain Reliever Subtypes in Past Year among Persons Aged 12 to 17 and Misuse of Pain Reliever Subtypes in Past Year among Past Year Any Users of Pain Reliever Subtypes Aged 12 to 17: Standard Errors of Numbers in Thousands, 2018 and 2019

Pain Reliever Subtype	Any Use in Past Year (2018)	Any Use in Past Year (2019)	Misuse in Past Year among Total Population (2018)	Misuse in Past Year among Total Population (2019)	Misuse in Past Year among Past Year Any Users of Pain Reliever Subtypes (2018)	Misuse in Past Year among Past Year Any Users of Pain Reliever Subtypes (2019)
ANY PRESCRIPTION PAIN RELIEVER	90	94	42	39	43	40
Hydrocodone Products	44	43	21	22	21	22
Zohydro [®] ER	8	9	4	6	*	*
Oxycodone Products	42	41	18	19	18	19
OxyContin [®]	31	27	14	14	14	14
Tramadol Products	30	29	13	13	13	13
Codeine Products	55	60	29	26	29	26
Morphine Products	27	29	9	11	9	11
Fentanyl Products	10	12	8	3	*	*
Buprenorphine Products	12	14	6	5	*	*
Oxymorphone Products	10	10	6	4	*	*
Demerol [®]	6	9	*	3	*	*
Hydromorphone Products	5	5	2	4	*	*
Methadone	8	11	6	7	*	*
Any Other Prescription Pain Reliever ¹	65	69	19	16	19	16

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

¹ Includes use or misuse of pain relievers containing other active ingredients. Reports of misuse of "any other prescription pain reliever" that correspond only to the specific pain reliever categories shown in the table are excluded from estimates for Any Other Prescription Pain Reliever and are included instead in the relevant pain reliever category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

Table 1.99D Any Use and Misuse of Pain Reliever Subtypes in Past Year among Persons Aged 12 to 17 and Misuse of Pain Reliever Subtypes in Past Year among Past Year Any Users of Pain Reliever Subtypes Aged 12 to 17: Standard Errors of Percentages, 2018 and 2019

Pain Reliever Subtype	Any Use in Past Year (2018)	Any Use in Past Year (2019)	Misuse in Past Year among Total Population (2018)	Misuse in Past Year among Total Population (2019)	Misuse in Past Year among Past Year Any Users of Pain Reliever Subtypes (2018)	Misuse in Past Year among Past Year Any Users of Pain Reliever Subtypes (2019)
ANY PRESCRIPTION PAIN RELIEVER	0.36	0.38	0.17	0.16	0.91	0.91
Hydrocodone Products	0.18	0.17	0.09	0.09	2.18	2.50
Zohydro [®] ER	0.03	0.04	0.02	0.03	*	*
Oxycodone Products	0.17	0.16	0.07	0.08	1.98	2.43
OxyContin [®]	0.12	0.11	0.06	0.05	3.42	4.08
Tramadol Products	0.12	0.12	0.05	0.05	3.53	3.40
Codeine Products	0.22	0.24	0.12	0.10	1.75	1.75
Morphine Products	0.11	0.12	0.04	0.05	2.92	3.26
Fentanyl Products	0.04	0.05	0.03	0.01	*	*
Buprenorphine Products	0.05	0.05	0.02	0.02	*	*
Oxymorphone Products	0.04	0.04	0.03	0.02	*	*
Demerol [®]	0.02	0.03	*	0.01	*	*
Hydromorphone Products	0.02	0.02	0.01	0.02	*	*
Methadone	0.03	0.04	0.02	0.03	*	*
Any Other Prescription Pain Reliever ¹	0.26	0.28	0.08	0.06	0.92	0.82

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

¹ Includes use or misuse of pain relievers containing other active ingredients. Reports of misuse of "any other prescription pain reliever" that correspond only to the specific pain reliever categories shown in the table are excluded from estimates for Any Other Prescription Pain Reliever and are included instead in the relevant pain reliever category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

Table 1.99P Any Use and Misuse of Pain Reliever Subtypes in Past Year among Persons Aged 12 to 17 and Misuse of Pain Reliever Subtypes in Past Year among Past Year Any Users of Pain Reliever Subtypes Aged 12 to 17: *P* Values from Tests of Differences of Percentages, 2019 Versus 2018

Pain Reliever Subtype	Any Use in Past Year (2018)	Any Use in Past Year (2019)	Misuse in Past Year among Total Population (2018)	Misuse in Past Year among Total Population (2019)	Misuse in Past Year among Past Year Any Users of Pain Reliever Subtypes (2018)	Misuse in Past Year among Past Year Any Users of Pain Reliever Subtypes (2019)
ANY PRESCRIPTION PAIN RELIEVER	0.2142		0.0214		0.0432	
Hydrocodone Products	0.0900		0.2890		0.7006	
Zohydro [®] ER	0.9334		0.8290		*	
Oxycodone Products	0.0521		0.6036		0.1235	
OxyContin [®]	0.0403		0.4018		0.8564	
Tramadol Products	0.7655		0.3159		0.2006	
Codeine Products	0.3773		0.0150		0.0214	
Morphine Products	0.6899		0.4637		0.5094	
Fentanyl Products	0.2117		0.1241		*	
Buprenorphine Products	0.4962		0.6319		*	
Oxymorphone Products	0.8476		0.1544		*	
Demerol [®]	0.1419		*		*	
Hydromorphone Products	0.5283		0.3570		*	
Methadone	0.1679		0.5087		*	
Any Other Prescription Pain Reliever ¹	0.4286		0.8036		0.9603	

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

¹ Includes use or misuse of pain relievers containing other active ingredients. Reports of misuse of "any other prescription pain reliever" that correspond only to the specific pain reliever categories shown in the table are excluded from estimates for Any Other Prescription Pain Reliever and are included instead in the relevant pain reliever category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

Table 1.100C Any Use and Misuse of Pain Reliever Subtypes in Past Year among Persons Aged 18 or Older and Misuse of Pain Reliever Subtypes in Past Year among Past Year Any Users of Pain Reliever Subtypes Aged 18 or Older: Standard Errors of Numbers in Thousands, 2018 and 2019

Pain Reliever Subtype	Any Use in Past Year (2018)	Any Use in Past Year (2019)	Misuse in Past Year among Total Population (2018)	Misuse in Past Year among Total Population (2019)	Misuse in Past Year among Past Year Any Users of Pain Reliever Subtypes (2018)	Misuse in Past Year among Past Year Any Users of Pain Reliever Subtypes (2019)
ANY PRESCRIPTION PAIN RELIEVER	776	784	284	285	289	294
Hydrocodone Products	666	648	235	208	237	211
Zohydro [®] ER	56	75	*	12	*	*
Oxycodone Products	487	493	157	150	158	152
OxyContin [®]	309	296	87	88	88	88
Tramadol Products	419	430	113	109	113	110
Codeine Products	508	498	142	152	141	153
Morphine Products	249	245	57	58	57	58
Fentanyl Products	140	166	35	44	35	44
Buprenorphine Products	155	134	71	67	71	67
Oxymorphone Products	82	93	36	41	36	41
Demerol [®]	135	95	19	15	19	15
Hydromorphone Products	134	134	38	34	38	34
Methadone	100	100	51	50	51	50
Any Other Prescription Pain Reliever ¹	450	439	70	79	71	79

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

¹ Includes use or misuse of pain relievers containing other active ingredients. Reports of misuse of "any other prescription pain reliever" that correspond only to the specific pain reliever categories shown in the table are excluded from estimates for Any Other Prescription Pain Reliever and are included instead in the relevant pain reliever category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

Table 1.100D Any Use and Misuse of Pain Reliever Subtypes in Past Year among Persons Aged 18 or Older and Misuse of Pain Reliever Subtypes in Past Year among Past Year Any Users of Pain Reliever Subtypes Aged 18 or Older: Standard Errors of Percentages, 2018 and 2019

Pain Reliever Subtype	Any Use in Past Year (2018)	Any Use in Past Year (2019)	Misuse in Past Year among Total Population (2018)	Misuse in Past Year among Total Population (2019)	Misuse in Past Year among Past Year Any Users of Pain Reliever Subtypes (2018)	Misuse in Past Year among Past Year Any Users of Pain Reliever Subtypes (2019)
ANY PRESCRIPTION PAIN RELIEVER	0.31	0.31	0.11	0.11	0.33	0.35
Hydrocodone Products	0.27	0.26	0.09	0.08	0.48	0.47
Zohydro [®] ER	0.02	0.03	*	0.00	*	*
Oxycodone Products	0.20	0.20	0.06	0.06	0.58	0.63
OxyContin [®]	0.12	0.12	0.04	0.04	0.98	1.03
Tramadol Products	0.17	0.17	0.05	0.04	0.62	0.63
Codeine Products	0.20	0.20	0.06	0.06	0.58	0.64
Morphine Products	0.10	0.10	0.02	0.02	0.95	1.02
Fentanyl Products	0.06	0.07	0.01	0.02	1.68	1.78
Buprenorphine Products	0.06	0.05	0.03	0.03	2.46	2.54
Oxymorphone Products	0.03	0.04	0.01	0.02	4.21	4.79
Demerol [®]	0.05	0.04	0.01	0.01	1.66	1.79
Hydromorphone Products	0.05	0.05	0.02	0.01	2.12	2.06
Methadone	0.04	0.04	0.02	0.02	4.13	4.25
Any Other Prescription Pain Reliever ¹	0.18	0.18	0.03	0.03	0.31	0.37

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

¹ Includes use or misuse of pain relievers containing other active ingredients. Reports of misuse of "any other prescription pain reliever" that correspond only to the specific pain reliever categories shown in the table are excluded from estimates for Any Other Prescription Pain Reliever and are included instead in the relevant pain reliever category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

Table 1.100P Any Use and Misuse of Pain Reliever Subtypes in Past Year among Persons Aged 18 or Older and Misuse of Pain Reliever Subtypes in Past Year among Past Year Any Users of Pain Reliever Subtypes Aged 18 or Older: *P* Values from Tests of Differences of Percentages, 2019 Versus 2018

Pain Reliever Subtype	Any Use in Past Year (2018)	Any Use in Past Year (2019)	Misuse in Past Year among Total Population (2018)	Misuse in Past Year among Total Population (2019)	Misuse in Past Year among Past Year Any Users of Pain Reliever Subtypes (2018)	Misuse in Past Year among Past Year Any Users of Pain Reliever Subtypes (2019)
ANY PRESCRIPTION PAIN RELIEVER	0.0001		0.7056		0.3658	
Hydrocodone Products	0.0000		0.1784		0.8397	
Zohydro [®] ER	0.7216		*		*	
Oxycodone Products	0.0036		0.3105		0.8537	
OxyContin [®]	0.5505		0.7243		0.9503	
Tramadol Products	0.1159		0.5030		0.8217	
Codeine Products	0.0521		0.7412		0.3270	
Morphine Products	0.2635		0.5846		0.8416	
Fentanyl Products	0.0785		0.8508		0.5560	
Buprenorphine Products	0.6414		0.7496		0.9575	
Oxymorphone Products	0.8763		0.9103		0.9905	
Demerol [®]	0.0685		0.8141		0.7227	
Hydromorphone Products	0.4023		0.2755		0.4578	
Methadone	0.7240		0.7471		0.8478	
Any Other Prescription Pain Reliever ¹	0.0392		0.1703		0.0678	

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

¹ Includes use or misuse of pain relievers containing other active ingredients. Reports of misuse of "any other prescription pain reliever" that correspond only to the specific pain reliever categories shown in the table are excluded from estimates for Any Other Prescription Pain Reliever and are included instead in the relevant pain reliever category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

Table 1.101C Any Use and Misuse of Pain Reliever Subtypes in Past Year among Persons Aged 18 to 25 and Misuse of Pain Reliever Subtypes in Past Year among Past Year Any Users of Pain Reliever Subtypes Aged 18 to 25: Standard Errors of Numbers in Thousands, 2018 and 2019

Pain Reliever Subtype	Any Use in Past Year (2018)	Any Use in Past Year (2019)	Misuse in Past Year among Total Population (2018)	Misuse in Past Year among Total Population (2019)	Misuse in Past Year among Past Year Any Users of Pain Reliever Subtypes (2018)	Misuse in Past Year among Past Year Any Users of Pain Reliever Subtypes (2019)
ANY PRESCRIPTION PAIN RELIEVER	148	142	72	75	77	80
Hydrocodone Products	108	101	51	52	52	54
Zohydro [®] ER	8	12	2	2	*	*
Oxycodone Products	86	87	47	42	48	42
OxyContin [®]	56	53	30	29	31	29
Tramadol Products	64	61	26	27	26	27
Codeine Products	96	86	39	41	39	42
Morphine Products	48	50	17	15	17	15
Fentanyl Products	23	28	12	11	12	11
Buprenorphine Products	30	29	18	17	18	17
Oxymorphone Products	17	23	14	15	*	*
Demerol [®]	13	13	4	7	*	*
Hydromorphone Products	18	18	11	10	*	*
Methadone	16	15	9	9	*	*
Any Other Prescription Pain Reliever ¹	100	95	23	30	23	30

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

¹ Includes use or misuse of pain relievers containing other active ingredients. Reports of misuse of "any other prescription pain reliever" that correspond only to the specific pain reliever categories shown in the table are excluded from estimates for Any Other Prescription Pain Reliever and are included instead in the relevant pain reliever category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

Table 1.101D Any Use and Misuse of Pain Reliever Subtypes in Past Year among Persons Aged 18 to 25 and Misuse of Pain Reliever Subtypes in Past Year among Past Year Any Users of Pain Reliever Subtypes Aged 18 to 25: Standard Errors of Percentages, 2018 and 2019

Pain Reliever Subtype	Any Use in Past Year (2018)	Any Use in Past Year (2019)	Misuse in Past Year among Total Population (2018)	Misuse in Past Year among Total Population (2019)	Misuse in Past Year among Past Year Any Users of Pain Reliever Subtypes (2018)	Misuse in Past Year among Past Year Any Users of Pain Reliever Subtypes (2019)
ANY PRESCRIPTION PAIN RELIEVER	0.44	0.42	0.21	0.22	0.76	0.83
Hydrocodone Products	0.32	0.30	0.15	0.15	1.10	1.31
Zohydro [®] ER	0.02	0.04	0.01	0.01	*	*
Oxycodone Products	0.25	0.26	0.14	0.12	1.47	1.48
OxyContin [®]	0.16	0.16	0.09	0.09	2.36	2.69
Tramadol Products	0.19	0.18	0.08	0.08	1.85	1.99
Codeine Products	0.28	0.25	0.11	0.12	1.33	1.43
Morphine Products	0.14	0.15	0.05	0.05	2.11	1.98
Fentanyl Products	0.07	0.08	0.03	0.03	5.03	4.10
Buprenorphine Products	0.09	0.09	0.05	0.05	4.79	4.70
Oxymorphone Products	0.05	0.07	0.04	0.04	*	*
Demerol [®]	0.04	0.04	0.01	0.02	*	*
Hydromorphone Products	0.05	0.05	0.03	0.03	*	*
Methadone	0.05	0.04	0.03	0.03	*	*
Any Other Prescription Pain Reliever ¹	0.29	0.28	0.07	0.09	0.76	1.04

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

¹ Includes use or misuse of pain relievers containing other active ingredients. Reports of misuse of "any other prescription pain reliever" that correspond only to the specific pain reliever categories shown in the table are excluded from estimates for Any Other Prescription Pain Reliever and are included instead in the relevant pain reliever category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

Table 1.101P Any Use and Misuse of Pain Reliever Subtypes in Past Year among Persons Aged 18 to 25 and Misuse of Pain Reliever Subtypes in Past Year among Past Year Any Users of Pain Reliever Subtypes Aged 18 to 25: *P* Values from Tests of Differences of Percentages, 2019 Versus 2018

Pain Reliever Subtype	Any Use in Past Year (2018)	Any Use in Past Year (2019)	Misuse in Past Year among Total Population (2018)	Misuse in Past Year among Total Population (2019)	Misuse in Past Year among Past Year Any Users of Pain Reliever Subtypes (2018)	Misuse in Past Year among Past Year Any Users of Pain Reliever Subtypes (2019)
ANY PRESCRIPTION PAIN RELIEVER	0.0166		0.3871		0.8701	
Hydrocodone Products	0.0000		0.4901		0.1498	
Zohydro [®] ER	0.3851		0.6359		*	
Oxycodone Products	0.0207		0.0021		0.0329	
OxyContin [®]	0.0504		0.0484		0.3032	
Tramadol Products	0.4260		0.8436		0.8649	
Codeine Products	0.1890		0.6637		0.9306	
Morphine Products	0.9443		0.9372		0.9087	
Fentanyl Products	0.1483		0.8737		0.5484	
Buprenorphine Products	0.9402		0.9403		0.9799	
Oxymorphone Products	0.2412		0.5664		*	
Demerol [®]	0.8250		0.4783		*	
Hydromorphone Products	0.1228		0.4048		*	
Methadone	0.5998		0.8481		*	
Any Other Prescription Pain Reliever ¹	0.6489		0.2884		0.2269	

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

¹ Includes use or misuse of pain relievers containing other active ingredients. Reports of misuse of "any other prescription pain reliever" that correspond only to the specific pain reliever categories shown in the table are excluded from estimates for Any Other Prescription Pain Reliever and are included instead in the relevant pain reliever category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

Table 1.102C Any Use and Misuse of Pain Reliever Subtypes in Past Year among Persons Aged 26 or Older and Misuse of Pain Reliever Subtypes in Past Year among Past Year Any Users of Pain Reliever Subtypes Aged 26 or Older: Standard Errors of Numbers in Thousands, 2018 and 2019

Pain Reliever Subtype	Any Use in Past Year (2018)	Any Use in Past Year (2019)	Misuse in Past Year among Total Population (2018)	Misuse in Past Year among Total Population (2019)	Misuse in Past Year among Past Year Any Users of Pain Reliever Subtypes (2018)	Misuse in Past Year among Past Year Any Users of Pain Reliever Subtypes (2019)
ANY PRESCRIPTION PAIN RELIEVER	765	770	275	276	279	285
Hydrocodone Products	649	636	225	203	228	207
Zohydro [®] ER	55	74	*	12	*	*
Oxycodone Products	482	481	150	144	150	146
OxyContin [®]	301	292	82	83	82	83
Tramadol Products	413	423	107	106	108	106
Codeine Products	499	491	139	147	138	147
Morphine Products	245	238	53	57	53	57
Fentanyl Products	139	160	34	42	34	42
Buprenorphine Products	152	131	67	63	67	63
Oxymorphone Products	79	91	33	39	33	*
Demerol [®]	134	95	19	14	19	14
Hydromorphone Products	132	131	36	33	36	33
Methadone	99	99	50	49	50	49
Any Other Prescription Pain Reliever ¹	442	430	65	71	65	71

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

¹ Includes use or misuse of pain relievers containing other active ingredients. Reports of misuse of "any other prescription pain reliever" that correspond only to the specific pain reliever categories shown in the table are excluded from estimates for Any Other Prescription Pain Reliever and are included instead in the relevant pain reliever category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

Table 1.102D Any Use and Misuse of Pain Reliever Subtypes in Past Year among Persons Aged 26 or Older and Misuse of Pain Reliever Subtypes in Past Year among Past Year Any Users of Pain Reliever Subtypes Aged 26 or Older: Standard Errors of Percentages, 2018 and 2019

Pain Reliever Subtype	Any Use in Past Year (2018)	Any Use in Past Year (2019)	Misuse in Past Year among Total Population (2018)	Misuse in Past Year among Total Population (2019)	Misuse in Past Year among Past Year Any Users of Pain Reliever Subtypes (2018)	Misuse in Past Year among Past Year Any Users of Pain Reliever Subtypes (2019)
ANY PRESCRIPTION PAIN RELIEVER	0.36	0.36	0.13	0.13	0.36	0.39
Hydrocodone Products	0.30	0.29	0.10	0.09	0.50	0.51
Zohydro [®] ER	0.03	0.03	*	0.01	*	*
Oxycodone Products	0.22	0.22	0.07	0.07	0.62	0.67
OxyContin [®]	0.14	0.13	0.04	0.04	1.04	1.09
Tramadol Products	0.19	0.20	0.05	0.05	0.64	0.67
Codeine Products	0.23	0.23	0.06	0.07	0.63	0.70
Morphine Products	0.11	0.11	0.02	0.03	1.02	1.16
Fentanyl Products	0.06	0.07	0.02	0.02	1.76	1.88
Buprenorphine Products	0.07	0.06	0.03	0.03	2.66	2.74
Oxymorphone Products	0.04	0.04	0.02	0.02	4.46	*
Demerol [®]	0.06	0.04	0.01	0.01	1.70	1.71
Hydromorphone Products	0.06	0.06	0.02	0.02	2.15	2.11
Methadone	0.05	0.05	0.02	0.02	4.41	4.55
Any Other Prescription Pain Reliever ¹	0.21	0.20	0.03	0.03	0.34	0.38

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

¹ Includes use or misuse of pain relievers containing other active ingredients. Reports of misuse of "any other prescription pain reliever" that correspond only to the specific pain reliever categories shown in the table are excluded from estimates for Any Other Prescription Pain Reliever and are included instead in the relevant pain reliever category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

Table 1.102P Any Use and Misuse of Pain Reliever Subtypes in Past Year among Persons Aged 26 or Older and Misuse of Pain Reliever Subtypes in Past Year among Past Year Any Users of Pain Reliever Subtypes Aged 26 or Older: *P* Values from Tests of Differences of Percentages, 2019 Versus 2018

Pain Reliever Subtype	Any Use in Past Year (2018)	Any Use in Past Year (2019)	Misuse in Past Year among Total Population (2018)	Misuse in Past Year among Total Population (2019)	Misuse in Past Year among Past Year Any Users of Pain Reliever Subtypes (2018)	Misuse in Past Year among Past Year Any Users of Pain Reliever Subtypes (2019)
ANY PRESCRIPTION PAIN RELIEVER	0.0004		0.8935		0.3523	
Hydrocodone Products	0.0000		0.2213		0.9879	
Zohydro [®] ER	0.6169		*		*	
Oxycodone Products	0.0096		0.9057		0.3938	
OxyContin [®]	0.7801		0.7241		0.6352	
Tramadol Products	0.1326		0.5122		0.7943	
Codeine Products	0.0844		0.6396		0.3191	
Morphine Products	0.2456		0.5894		0.8327	
Fentanyl Products	0.1208		0.8751		0.6289	
Buprenorphine Products	0.6446		0.7554		0.9534	
Oxymorphone Products	0.9125		0.7191		*	
Demerol [®]	0.0694		0.6329		0.9484	
Hydromorphone Products	0.5158		0.3740		0.5190	
Methadone	0.7788		0.7687		0.8403	
Any Other Prescription Pain Reliever ¹	0.0452		0.2718		0.1335	

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

¹ Includes use or misuse of pain relievers containing other active ingredients. Reports of misuse of "any other prescription pain reliever" that correspond only to the specific pain reliever categories shown in the table are excluded from estimates for Any Other Prescription Pain Reliever and are included instead in the relevant pain reliever category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.