

**Table 10.36C Had Serious Thoughts of Suicide in Past Year among Persons Aged 18 or Older, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2008-2019**

| Age Category       | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | 2016 | 2017 | 2018 | 2019 |
|--------------------|------|------|------|------|------|------|------|------|------|------|------|------|
| <b>TOTAL</b>       | 299  | 299  | 311  | 299  | 305  | 340  | 281  | 280  | 267  | 289  | 282  | 300  |
| <b>18-25</b>       | 74   | 66   | 75   | 85   | 79   | 85   | 87   | 91   | 96   | 108  | 100  | 102  |
| 18-20              | 52   | 47   | 56   | 61   | 62   | 62   | 61   | 62   | 78   | 75   | 70   | 81   |
| 21-25              | 60   | 51   | 59   | 61   | 60   | 62   | 69   | 74   | 72   | 87   | 85   | 86   |
| <b>26 or Older</b> | 292  | 292  | 296  | 284  | 294  | 327  | 263  | 267  | 247  | 262  | 257  | 278  |
| 26-49              | 193  | 200  | 201  | 169  | 208  | 207  | 167  | 165  | 154  | 163  | 170  | 182  |
| 26-29              | 82   | 80   | 94   | 78   | 90   | 88   | 71   | 81   | 74   | 88   | 90   | 98   |
| 30-34              | 74   | 92   | 86   | 63   | 84   | 106  | 78   | 73   | 78   | 74   | 80   | 92   |
| 35-39              | 93   | 103  | 72   | 86   | 94   | 89   | 68   | 71   | 72   | 74   | 75   | 82   |
| 40-44              | 88   | 81   | 88   | 83   | 97   | 88   | 73   | 68   | 60   | 73   | 69   | 69   |
| 45-49              | 94   | 98   | 111  | 93   | 100  | 85   | 69   | 70   | 70   | 63   | 63   | 71   |
| 50 or Older        | 210  | 215  | 214  | 229  | 214  | 271  | 194  | 209  | 193  | 210  | 189  | 210  |
| 50-54              | 121  | 118  | 111  | 131  | 103  | 209  | 106  | 102  | 108  | 77   | 96   | 91   |
| 55-59              | 118  | 118  | 114  | 124  | 95   | 117  | 109  | 98   | 89   | 107  | 86   | 104  |
| 60-64              | 77   | 67   | 93   | 84   | 109  | 83   | 80   | 79   | 74   | 84   | 82   | 91   |
| 65 or Older        | 104  | 107  | 124  | 106  | 123  | 109  | 102  | 121  | 109  | 136  | 102  | 128  |

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Some 2008 to 2010 estimates may differ from previously published estimates due to updates (see Section 3.3.5 in Chapter 3 of the *2019 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

NOTE: Respondents with unknown suicide information were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2019.

**Table 10.36D Had Serious Thoughts of Suicide in Past Year among Persons Aged 18 or Older, by Detailed Age Category: Standard Errors of Percentages, 2008-2019**

| Age Category       | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | 2016 | 2017 | 2018 | 2019 |
|--------------------|------|------|------|------|------|------|------|------|------|------|------|------|
| <b>TOTAL</b>       | 0.13 | 0.13 | 0.14 | 0.13 | 0.13 | 0.14 | 0.12 | 0.12 | 0.11 | 0.12 | 0.11 | 0.12 |
| <b>18-25</b>       | 0.23 | 0.20 | 0.22 | 0.25 | 0.23 | 0.24 | 0.25 | 0.26 | 0.28 | 0.32 | 0.30 | 0.31 |
| 18-20              | 0.37 | 0.32 | 0.39 | 0.41 | 0.41 | 0.44 | 0.44 | 0.45 | 0.56 | 0.53 | 0.49 | 0.57 |
| 21-25              | 0.29 | 0.25 | 0.27 | 0.28 | 0.27 | 0.28 | 0.31 | 0.32 | 0.31 | 0.38 | 0.37 | 0.38 |
| <b>26 or Older</b> | 0.15 | 0.15 | 0.15 | 0.14 | 0.15 | 0.16 | 0.13 | 0.13 | 0.12 | 0.12 | 0.12 | 0.13 |
| 26-49              | 0.19 | 0.20 | 0.20 | 0.17 | 0.21 | 0.21 | 0.17 | 0.17 | 0.16 | 0.16 | 0.17 | 0.18 |
| 26-29              | 0.47 | 0.47 | 0.54 | 0.44 | 0.51 | 0.51 | 0.41 | 0.46 | 0.42 | 0.47 | 0.49 | 0.52 |
| 30-34              | 0.39 | 0.46 | 0.43 | 0.32 | 0.42 | 0.51 | 0.36 | 0.34 | 0.36 | 0.34 | 0.35 | 0.40 |
| 35-39              | 0.45 | 0.50 | 0.37 | 0.44 | 0.47 | 0.46 | 0.34 | 0.36 | 0.34 | 0.35 | 0.34 | 0.37 |
| 40-44              | 0.41 | 0.39 | 0.42 | 0.40 | 0.47 | 0.41 | 0.36 | 0.33 | 0.31 | 0.36 | 0.35 | 0.35 |
| 45-49              | 0.41 | 0.43 | 0.48 | 0.42 | 0.46 | 0.42 | 0.33 | 0.33 | 0.34 | 0.31 | 0.32 | 0.35 |
| 50 or Older        | 0.23 | 0.23 | 0.22 | 0.23 | 0.21 | 0.26 | 0.18 | 0.19 | 0.17 | 0.19 | 0.17 | 0.18 |
| 50-54              | 0.55 | 0.52 | 0.50 | 0.57 | 0.43 | 0.88 | 0.44 | 0.43 | 0.49 | 0.37 | 0.46 | 0.44 |
| 55-59              | 0.65 | 0.62 | 0.58 | 0.65 | 0.46 | 0.56 | 0.52 | 0.46 | 0.41 | 0.49 | 0.40 | 0.49 |
| 60-64              | 0.51 | 0.43 | 0.56 | 0.46 | 0.63 | 0.47 | 0.46 | 0.41 | 0.39 | 0.42 | 0.40 | 0.44 |
| 65 or Older        | 0.28 | 0.28 | 0.32 | 0.27 | 0.29 | 0.25 | 0.23 | 0.26 | 0.23 | 0.28 | 0.20 | 0.24 |

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Some 2008 to 2010 estimates may differ from previously published estimates due to updates (see Section 3.3.5 in Chapter 3 of the *2019 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

NOTE: Respondents with unknown suicide information were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2019.

**Table 10.36P Had Serious Thoughts of Suicide in Past Year among Persons Aged 18 or Older, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2019 Versus 2008-2018**

| Age Category       | 2008   | 2009   | 2010   | 2011   | 2012   | 2013   | 2014   | 2015   | 2016   | 2017   | 2018   | 2019 |
|--------------------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|------|
| <b>TOTAL</b>       | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0049 | 0.0044 |      |
| <b>18-25</b>       | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0029 | 0.0780 |      |
| 18-20              | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0058 | 0.2298 | 0.0977 |      |
| 21-25              | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0053 | 0.4452 |      |
| <b>26 or Older</b> | 0.0075 | 0.0481 | 0.0424 | 0.0026 | 0.0198 | 0.0532 | 0.0360 | 0.0320 | 0.0065 | 0.0336 | 0.0123 |      |
| 26-49              | 0.0000 | 0.0004 | 0.0000 | 0.0000 | 0.0001 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0101 |      |
| 26-29              | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0001 | 0.0000 | 0.0044 | 0.0877 |      |
| 30-34              | 0.0005 | 0.0106 | 0.0009 | 0.0000 | 0.0018 | 0.0399 | 0.0061 | 0.0136 | 0.1638 | 0.0055 | 0.3745 |      |
| 35-39              | 0.1485 | 0.7987 | 0.0032 | 0.4238 | 0.1357 | 0.0916 | 0.0659 | 0.0760 | 0.3729 | 0.0463 | 0.2466 |      |
| 40-44              | 0.7856 | 0.9070 | 0.8951 | 0.2726 | 0.6978 | 0.1612 | 0.5821 | 0.4091 | 0.0393 | 0.8081 | 0.4926 |      |
| 45-49              | 0.8841 | 0.6100 | 0.0499 | 0.6302 | 0.2922 | 0.7975 | 0.2407 | 0.3458 | 0.4686 | 0.1320 | 0.1564 |      |
| 50 or Older        | 0.9948 | 0.8282 | 0.4800 | 0.3488 | 0.7966 | 0.2395 | 0.1521 | 0.3625 | 0.8796 | 0.5764 | 0.2933 |      |
| 50-54              | 0.6300 | 0.5632 | 0.7710 | 0.0977 | 0.6561 | 0.0421 | 0.1760 | 0.2945 | 0.2095 | 0.5612 | 0.8836 |      |
| 55-59              | 0.9638 | 0.6087 | 0.6924 | 0.8607 | 0.2583 | 0.8094 | 0.3524 | 0.7032 | 0.5376 | 0.8833 | 0.2986 |      |
| 60-64              | 0.6479 | 0.4679 | 0.5846 | 0.9495 | 0.1811 | 0.8339 | 0.5572 | 0.3482 | 0.8731 | 0.7727 | 0.7882 |      |
| 65 or Older        | 0.4675 | 0.1136 | 0.8135 | 0.5698 | 0.7537 | 0.3369 | 0.6250 | 0.9282 | 0.6490 | 0.3999 | 0.3412 |      |

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Some 2008 to 2010 estimates that are used in the comparisons may differ from previously published estimates due to updates (see Section 3.3.5 in Chapter 3 of the *2019 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

NOTE: Respondents with unknown suicide information were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2019.

**Table 10.39C Made Any Suicide Plans in Past Year among Persons Aged 18 or Older, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2008-2019**

| Age Category       | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | 2016 | 2017 | 2018 | 2019 |
|--------------------|------|------|------|------|------|------|------|------|------|------|------|------|
| <b>TOTAL</b>       | 155  | 157  | 168  | 157  | 156  | 166  | 138  | 138  | 127  | 148  | 155  | 152  |
| <b>18-25</b>       | 39   | 41   | 41   | 44   | 47   | 50   | 49   | 54   | 56   | 67   | 58   | 62   |
| 18-20              | 29   | 29   | 28   | 28   | 37   | 35   | 36   | 41   | 40   | 48   | 39   | 44   |
| 21-25              | 27   | 29   | 31   | 35   | 31   | 35   | 38   | 39   | 42   | 49   | 43   | 49   |
| <b>26 or Older</b> | 148  | 150  | 164  | 148  | 152  | 156  | 130  | 127  | 114  | 131  | 147  | 142  |
| 26-49              | 99   | 94   | 91   | 102  | 120  | 120  | 85   | 78   | 85   | 90   | 100  | 91   |
| 26-29              | 40   | 34   | 32   | 54   | 50   | 55   | 35   | 40   | 35   | 43   | 58   | 46   |
| 30-34              | 44   | 34   | 42   | 26   | 32   | 62   | 35   | 38   | 47   | 38   | 42   | 46   |
| 35-39              | 51   | 49   | 29   | 47   | 62   | 51   | 42   | 32   | 42   | 39   | 49   | 38   |
| 40-44              | 43   | 46   | 49   | 50   | 58   | 45   | 36   | 31   | 37   | 40   | 33   | 39   |
| 45-49              | 47   | 46   | 44   | 45   | 59   | 46   | 36   | 34   | 35   | 41   | 38   | 37   |
| 50 or Older        | 108  | 117  | 134  | 111  | 101  | 98   | 98   | 101  | 78   | 94   | 105  | 107  |
| 50-54              | 82   | 69   | 74   | 72   | 47   | 47   | 41   | 65   | 45   | 41   | 65   | 40   |
| 55-59              | 47   | 56   | 61   | 68   | 43   | 58   | 54   | 40   | 34   | 48   | 53   | 55   |
| 60-64              | 29   | 37   | 50   | 32   | 56   | 37   | 54   | 34   | 43   | 42   | 41   | 55   |
| 65 or Older        | 49   | 37   | 81   | 42   | 56   | 54   | 54   | 56   | 36   | 64   | 41   | 60   |

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Some 2008 to 2010 estimates may differ from previously published estimates due to updates (see Section 3.3.5 in Chapter 3 of the *2019 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

NOTE: Respondents were asked whether they made any suicide plans in the past 12 months only if they had serious thoughts of suicide in the past 12 months. Respondents who had no suicide thoughts were categorized as not making any suicide plans in the past 12 months. Respondents with unknown suicide information were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2019.

**Table 10.39D Made Any Suicide Plans in Past Year among Persons Aged 18 or Older, by Detailed Age Category: Standard Errors of Percentages, 2008-2019**

| Age Category       | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | 2016 | 2017 | 2018 | 2019 |
|--------------------|------|------|------|------|------|------|------|------|------|------|------|------|
| <b>TOTAL</b>       | 0.07 | 0.07 | 0.07 | 0.07 | 0.07 | 0.07 | 0.06 | 0.06 | 0.05 | 0.06 | 0.06 | 0.06 |
| <b>18-25</b>       | 0.12 | 0.12 | 0.12 | 0.13 | 0.14 | 0.14 | 0.14 | 0.16 | 0.16 | 0.20 | 0.17 | 0.19 |
| 18-20              | 0.22 | 0.21 | 0.20 | 0.20 | 0.28 | 0.26 | 0.27 | 0.31 | 0.31 | 0.36 | 0.30 | 0.33 |
| 21-25              | 0.14 | 0.14 | 0.15 | 0.16 | 0.15 | 0.16 | 0.17 | 0.18 | 0.19 | 0.23 | 0.20 | 0.23 |
| <b>26 or Older</b> | 0.08 | 0.08 | 0.08 | 0.07 | 0.08 | 0.08 | 0.06 | 0.06 | 0.05 | 0.06 | 0.07 | 0.07 |
| 26-49              | 0.10 | 0.10 | 0.09 | 0.10 | 0.12 | 0.12 | 0.09 | 0.08 | 0.09 | 0.09 | 0.10 | 0.09 |
| 26-29              | 0.23 | 0.20 | 0.19 | 0.32 | 0.29 | 0.32 | 0.21 | 0.23 | 0.20 | 0.24 | 0.32 | 0.25 |
| 30-34              | 0.24 | 0.17 | 0.21 | 0.13 | 0.16 | 0.30 | 0.17 | 0.18 | 0.22 | 0.17 | 0.19 | 0.21 |
| 35-39              | 0.25 | 0.24 | 0.15 | 0.24 | 0.32 | 0.26 | 0.21 | 0.17 | 0.21 | 0.19 | 0.22 | 0.18 |
| 40-44              | 0.20 | 0.22 | 0.24 | 0.25 | 0.29 | 0.21 | 0.18 | 0.15 | 0.19 | 0.20 | 0.17 | 0.20 |
| 45-49              | 0.21 | 0.20 | 0.20 | 0.21 | 0.28 | 0.23 | 0.18 | 0.16 | 0.17 | 0.20 | 0.20 | 0.18 |
| 50 or Older        | 0.12 | 0.12 | 0.14 | 0.11 | 0.10 | 0.09 | 0.09 | 0.09 | 0.07 | 0.08 | 0.09 | 0.09 |
| 50-54              | 0.37 | 0.31 | 0.34 | 0.31 | 0.20 | 0.20 | 0.17 | 0.28 | 0.21 | 0.20 | 0.31 | 0.19 |
| 55-59              | 0.26 | 0.30 | 0.32 | 0.36 | 0.21 | 0.28 | 0.26 | 0.19 | 0.15 | 0.22 | 0.25 | 0.26 |
| 60-64              | 0.19 | 0.24 | 0.30 | 0.17 | 0.33 | 0.21 | 0.31 | 0.18 | 0.23 | 0.21 | 0.20 | 0.27 |
| 65 or Older        | 0.13 | 0.10 | 0.21 | 0.11 | 0.13 | 0.13 | 0.12 | 0.12 | 0.08 | 0.13 | 0.08 | 0.11 |

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Some 2008 to 2010 estimates may differ from previously published estimates due to updates (see Section 3.3.5 in Chapter 3 of the *2019 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

NOTE: Respondents were asked whether they made any suicide plans in the past 12 months only if they had serious thoughts of suicide in the past 12 months. Respondents who had no suicide thoughts were categorized as not making any suicide plans in the past 12 months. Respondents with unknown suicide information were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2019.

**Table 10.39P Made Any Suicide Plans in Past Year among Persons Aged 18 or Older, by Detailed Age Category: P Values from Tests of Differences of Percentages, 2019 Versus 2008-2018**

| Age Category       | 2008   | 2009   | 2010   | 2011   | 2012   | 2013   | 2014   | 2015   | 2016   | 2017   | 2018   | 2019 |
|--------------------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|------|
| <b>TOTAL</b>       | 0.0001 | 0.0000 | 0.0038 | 0.0001 | 0.0068 | 0.0093 | 0.0012 | 0.0013 | 0.0018 | 0.3157 | 0.5683 |      |
| <b>18-25</b>       | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.4450 | 0.0536 |      |
| 18-20              | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0015 | 0.0013 | 0.0001 | 0.0251 | 0.0410 | 0.4207 | 0.1598 |      |
| 21-25              | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0003 | 0.0842 | 0.2047 |      |
| <b>26 or Older</b> | 0.2028 | 0.1007 | 0.7857 | 0.2571 | 0.5447 | 0.4686 | 0.3605 | 0.1220 | 0.0962 | 0.3649 | 0.8945 |      |
| 26-49              | 0.0034 | 0.0011 | 0.0011 | 0.0102 | 0.3100 | 0.2510 | 0.0101 | 0.0009 | 0.1824 | 0.0877 | 0.9556 |      |
| 26-29              | 0.0012 | 0.0000 | 0.0000 | 0.0998 | 0.0146 | 0.0461 | 0.0014 | 0.0468 | 0.0103 | 0.0612 | 0.6117 |      |
| 30-34              | 0.1205 | 0.0019 | 0.0976 | 0.0001 | 0.0018 | 0.4747 | 0.0191 | 0.0791 | 0.9790 | 0.0850 | 0.4875 |      |
| 35-39              | 0.9442 | 0.9945 | 0.1574 | 0.3678 | 0.4069 | 0.4457 | 0.4212 | 0.4360 | 0.2509 | 0.5770 | 0.1924 |      |
| 40-44              | 0.3338 | 0.8524 | 0.7917 | 0.4810 | 0.5489 | 0.7912 | 0.6408 | 0.1314 | 0.3311 | 0.9499 | 0.4676 |      |
| 45-49              | 0.5777 | 0.6297 | 0.7229 | 0.5242 | 0.2361 | 0.9683 | 0.6537 | 0.2498 | 0.6189 | 0.6276 | 0.4245 |      |
| 50 or Older        | 0.6714 | 0.8488 | 0.0645 | 0.5925 | 0.9603 | 0.9798 | 0.3950 | 0.5154 | 0.2667 | 0.7820 | 0.8297 |      |
| 50-54              | 0.1592 | 0.3337 | 0.0509 | 0.0636 | 0.7941 | 0.6957 | 0.7770 | 0.0447 | 0.3759 | 0.4589 | 0.2478 |      |
| 55-59              | 0.6498 | 0.8914 | 0.8681 | 0.8795 | 0.2134 | 0.9487 | 0.8976 | 0.4384 | 0.1124 | 0.5463 | 0.8669 |      |
| 60-64              | 0.8254 | 0.9952 | 0.9879 | 0.4127 | 0.6256 | 0.6727 | 0.5213 | 0.6610 | 0.9020 | 0.9556 | 0.8116 |      |
| 65 or Older        | 0.5250 | 0.3417 | 0.1602 | 0.3543 | 0.8669 | 0.7713 | 0.4461 | 0.9559 | 0.1426 | 0.7272 | 0.4335 |      |

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Some 2008 to 2010 estimates that are used in the comparisons may differ from previously published estimates due to updates (see Section 3.3.5 in Chapter 3 of the *2019 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

NOTE: Respondents were asked whether they made any suicide plans in the past 12 months only if they had serious thoughts of suicide in the past 12 months. Respondents who had no suicide thoughts were categorized as not making any suicide plans in the past 12 months. Respondents with unknown suicide information were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2019.

**Table 10.41C Attempted Suicide in Past Year among Persons Aged 18 or Older, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2008-2019**

| Age Category       | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | 2016 | 2017 | 2018 | 2019 |
|--------------------|------|------|------|------|------|------|------|------|------|------|------|------|
| <b>TOTAL</b>       | 105  | 91   | 108  | 110  | 103  | 118  | 76   | 100  | 96   | 97   | 101  | 93   |
| <b>18-25</b>       | 32   | 31   | 30   | 35   | 41   | 35   | 36   | 44   | 45   | 48   | 44   | 45   |
| 18-20              | 25   | 20   | 21   | 22   | 33   | 26   | 22   | 32   | 35   | 36   | 31   | 33   |
| 21-25              | 18   | 24   | 22   | 26   | 24   | 22   | 30   | 32   | 30   | 33   | 32   | 32   |
| <b>26 or Older</b> | 100  | 85   | 103  | 103  | 95   | 112  | 67   | 90   | 84   | 87   | 91   | 83   |
| 26-49              | 67   | 63   | 59   | 64   | 75   | 83   | 54   | 57   | 53   | 53   | 59   | 58   |
| 26-29              | 30   | 23   | 14   | 32   | 35   | 32   | 21   | 24   | 25   | 23   | 24   | 27   |
| 30-34              | 36   | 25   | 28   | 20   | 25   | 52   | 22   | 28   | 29   | 20   | 25   | 30   |
| 35-39              | 22   | 36   | 16   | 30   | 33   | 36   | 25   | 21   | 27   | 32   | 36   | 21   |
| 40-44              | 29   | 23   | 32   | 31   | 33   | 34   | 23   | 28   | 18   | 17   | 22   | 24   |
| 45-49              | 29   | 31   | 33   | 29   | 38   | 26   | 30   | 25   | 20   | 23   | 23   | 22   |
| 50 or Older        | 76   | 57   | 81   | 81   | 64   | 72   | 40   | 73   | 65   | 68   | 66   | 56   |
| 50-54              | 67   | 24   | 31   | 54   | 26   | 25   | 25   | 45   | 27   | 36   | 44   | 19   |
| 55-59              | 28   | 31   | 30   | 57   | 29   | 46   | 31   | 36   | 41   | 26   | 41   | 29   |
| 60-64              | 16   | 28   | 38   | 2    | 23   | 32   | 14   | 27   | 25   | 25   | 10   | 11   |
| 65 or Older        | 18   | 31   | 58   | 18   | 45   | 38   | 24   | 37   | 34   | 40   | 27   | 43   |

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Some 2008 to 2010 estimates may differ from previously published estimates due to updates (see Section 3.3.5 in Chapter 3 of the *2019 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

NOTE: Respondents were asked whether they attempted suicide in the past 12 months only if they had serious thoughts of suicide in the past 12 months. Respondents who had no suicide thoughts were categorized as not attempting suicide in the past 12 months. Respondents with unknown suicide information were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2019.

**Table 10.41D Attempted Suicide in Past Year among Persons Aged 18 or Older, by Detailed Age Category: Standard Errors of Percentages, 2008-2019**

| Age Category       | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | 2016 | 2017 | 2018 | 2019 |
|--------------------|------|------|------|------|------|------|------|------|------|------|------|------|
| <b>TOTAL</b>       | 0.05 | 0.04 | 0.05 | 0.05 | 0.04 | 0.05 | 0.03 | 0.04 | 0.04 | 0.04 | 0.04 | 0.04 |
| <b>18-25</b>       | 0.10 | 0.09 | 0.09 | 0.10 | 0.12 | 0.10 | 0.10 | 0.13 | 0.13 | 0.14 | 0.13 | 0.14 |
| 18-20              | 0.19 | 0.14 | 0.15 | 0.16 | 0.25 | 0.19 | 0.17 | 0.24 | 0.27 | 0.28 | 0.24 | 0.25 |
| 21-25              | 0.09 | 0.12 | 0.11 | 0.13 | 0.12 | 0.10 | 0.14 | 0.15 | 0.14 | 0.15 | 0.15 | 0.16 |
| <b>26 or Older</b> | 0.05 | 0.04 | 0.05 | 0.05 | 0.05 | 0.06 | 0.03 | 0.04 | 0.04 | 0.04 | 0.04 | 0.04 |
| 26-49              | 0.07 | 0.06 | 0.06 | 0.07 | 0.08 | 0.09 | 0.06 | 0.06 | 0.05 | 0.05 | 0.06 | 0.06 |
| 26-29              | 0.18 | 0.14 | 0.09 | 0.19 | 0.21 | 0.19 | 0.13 | 0.14 | 0.14 | 0.13 | 0.14 | 0.15 |
| 30-34              | 0.20 | 0.13 | 0.14 | 0.11 | 0.13 | 0.26 | 0.10 | 0.13 | 0.14 | 0.09 | 0.11 | 0.14 |
| 35-39              | 0.11 | 0.18 | 0.08 | 0.15 | 0.17 | 0.19 | 0.13 | 0.11 | 0.13 | 0.16 | 0.17 | 0.10 |
| 40-44              | 0.13 | 0.11 | 0.15 | 0.15 | 0.16 | 0.16 | 0.12 | 0.14 | 0.09 | 0.09 | 0.11 | 0.12 |
| 45-49              | 0.13 | 0.14 | 0.15 | 0.13 | 0.18 | 0.13 | 0.15 | 0.12 | 0.10 | 0.11 | 0.12 | 0.11 |
| 50 or Older        | 0.08 | 0.06 | 0.08 | 0.08 | 0.06 | 0.07 | 0.04 | 0.07 | 0.06 | 0.06 | 0.06 | 0.05 |
| 50-54              | 0.31 | 0.11 | 0.14 | 0.24 | 0.11 | 0.11 | 0.10 | 0.20 | 0.13 | 0.17 | 0.21 | 0.09 |
| 55-59              | 0.16 | 0.17 | 0.16 | 0.30 | 0.14 | 0.22 | 0.15 | 0.17 | 0.19 | 0.12 | 0.19 | 0.14 |
| 60-64              | 0.11 | 0.18 | 0.23 | 0.01 | 0.14 | 0.18 | 0.08 | 0.14 | 0.13 | 0.13 | 0.05 | 0.05 |
| 65 or Older        | 0.05 | 0.08 | 0.15 | 0.05 | 0.11 | 0.09 | 0.05 | 0.08 | 0.07 | 0.08 | 0.05 | 0.08 |

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Some 2008 to 2010 estimates may differ from previously published estimates due to updates (see Section 3.3.5 in Chapter 3 of the *2019 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

NOTE: Respondents were asked whether they attempted suicide in the past 12 months only if they had serious thoughts of suicide in the past 12 months. Respondents who had no suicide thoughts were categorized as not attempting suicide in the past 12 months. Respondents with unknown suicide information were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2019.

**Table 10.41P Attempted Suicide in Past Year among Persons Aged 18 or Older, by Detailed Age Category: P Values from Tests of Differences of Percentages, 2019 Versus 2008-2018**

| Age Category       | 2008   | 2009   | 2010   | 2011   | 2012   | 2013   | 2014   | 2015   | 2016   | 2017   | 2018   | 2019 |
|--------------------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|------|
| <b>TOTAL</b>       | 0.2488 | 0.0820 | 0.2899 | 0.2825 | 0.9403 | 0.9966 | 0.0826 | 0.6687 | 0.8295 | 0.8565 | 0.6047 |      |
| <b>18-25</b>       | 0.0003 | 0.0000 | 0.0001 | 0.0004 | 0.0638 | 0.0010 | 0.0013 | 0.3624 | 0.9452 | 0.6277 | 0.5636 |      |
| 18-20              | 0.1477 | 0.0021 | 0.0075 | 0.0286 | 0.6556 | 0.2558 | 0.0071 | 0.6143 | 0.3509 | 0.1491 | 0.6883 |      |
| 21-25              | 0.0001 | 0.0032 | 0.0026 | 0.0051 | 0.0252 | 0.0004 | 0.0753 | 0.5402 | 0.3373 | 0.4650 | 0.6450 |      |
| <b>26 or Older</b> | 0.9701 | 0.8067 | 0.8260 | 0.9435 | 0.6132 | 0.2676 | 0.6454 | 0.4989 | 0.6812 | 0.8508 | 0.8308 |      |
| 26-49              | 0.1510 | 0.2164 | 0.0411 | 0.1608 | 0.5221 | 0.7314 | 0.3744 | 0.4621 | 0.3612 | 0.0880 | 0.5049 |      |
| 26-29              | 0.2871 | 0.0667 | 0.0005 | 0.4791 | 0.2584 | 0.2071 | 0.1453 | 0.3436 | 0.5954 | 0.2550 | 0.1886 |      |
| 30-34              | 0.8114 | 0.1404 | 0.4018 | 0.0680 | 0.1632 | 0.7586 | 0.1206 | 0.7038 | 0.4135 | 0.0655 | 0.4120 |      |
| 35-39              | 0.3507 | 0.3748 | 0.1795 | 0.6093 | 0.9088 | 0.2731 | 0.8858 | 0.8464 | 0.4198 | 0.9814 | 0.1303 |      |
| 40-44              | 0.1568 | 0.1359 | 0.5369 | 0.6668 | 0.6987 | 0.6710 | 0.7876 | 0.9547 | 0.1234 | 0.0666 | 0.4007 |      |
| 45-49              | 0.5989 | 0.3730 | 0.3999 | 0.5140 | 0.5585 | 0.8365 | 0.2043 | 0.9955 | 0.9194 | 0.4488 | 0.7035 |      |
| 50 or Older        | 0.3065 | 0.5675 | 0.0792 | 0.2564 | 0.1997 | 0.2400 | 0.8314 | 0.1350 | 0.8141 | 0.1970 | 0.3567 |      |
| 50-54              | 0.0895 | 0.5870 | 0.4338 | 0.0842 | 0.5377 | 0.7938 | 0.4254 | 0.0425 | 0.6846 | 0.0685 | 0.2782 |      |
| 55-59              | 0.8595 | 0.9795 | 0.8486 | 0.3401 | 0.9286 | 0.2339 | 0.9035 | 0.8396 | 0.9166 | 0.9502 | 0.2434 |      |
| 60-64              | 0.2863 | 0.3767 | 0.3382 | 0.3300 | 0.0623 | 0.1528 | 0.6886 | 0.2166 | 0.6238 | 0.5443 | 0.9008 |      |
| 65 or Older        | 0.2474 | 0.9774 | 0.1714 | 0.2610 | 0.4919 | 0.7395 | 0.5559 | 0.8393 | 0.7881 | 0.6733 | 0.5041 |      |

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Some 2008 to 2010 estimates that are used in the comparisons may differ from previously published estimates due to updates (see Section 3.3.5 in Chapter 3 of the *2019 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

NOTE: Respondents were asked whether they attempted suicide in the past 12 months only if they had serious thoughts of suicide in the past 12 months. Respondents who had no suicide thoughts were categorized as not attempting suicide in the past 12 months. Respondents with unknown suicide information were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2019.