

Table 11.11C Substance Use Disorder, Illicit Drug Use Disorder, and Alcohol Use Disorder in Past Year among Persons Aged 12 to 17, by Past Year Major Depressive Episode (MDE) and MDE with Severe Impairment Status and Gender: Standard Errors of Numbers in Thousands, 2015-2019

Substance Use Disorder/ MDE Status/Gender	2015	2016	2017	2018	2019
SUBSTANCE USE DISORDER					
MDE	28	27	27	29	32
Male	14	12	15	16	17
Female	24	24	23	24	27
MDE WITH SEVERE IMPAIRMENT	26	24	25	26	30
Male	12	10	14	14	15
Female	23	22	20	22	25
NO MDE	45	40	37	36	44
Male	34	32	28	28	32
Female	30	24	25	22	31
ILLICIT DRUG USE DISORDER					
MDE	23	23	24	25	28
Male	11	11	14	13	15
Female	20	20	20	21	23
MDE WITH SEVERE IMPAIRMENT	22	21	21	23	25
Male	10	10	12	11	13
Female	20	19	18	20	21
NO MDE	37	35	32	28	42
Male	28	29	24	24	30
Female	23	20	21	16	28
ALCOHOL USE DISORDER					
MDE	22	20	19	20	22
Male	11	8	11	12	12
Female	19	18	16	16	18
MDE WITH SEVERE IMPAIRMENT	20	18	16	18	20
Male	10	7	10	11	11
Female	18	16	13	15	17
NO MDE	33	24	24	25	22
Male	26	18	17	17	15
Female	21	16	17	18	16

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

NOTE: Respondents with unknown impairment data were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2015-2019.

Table 11.11D Substance Use Disorder, Illicit Drug Use Disorder, and Alcohol Use Disorder in Past Year among Persons Aged 12 to 17, by Past Year Major Depressive Episode (MDE) and MDE with Severe Impairment Status and Gender: Standard Errors of Percentages, 2015-2019

Substance Use Disorder/ MDE Status/Gender	2015	2016	2017	2018	2019
SUBSTANCE USE DISORDER					
MDE	0.86	0.83	0.80	0.77	0.79
Male	1.82	1.46	1.67	1.69	1.47
Female	0.99	0.99	0.94	0.90	0.93
MDE WITH SEVERE IMPAIRMENT	1.11	1.06	1.02	0.97	1.02
Male	2.32	1.85	2.19	2.14	1.88
Female	1.30	1.23	1.15	1.15	1.20
NO MDE	0.21	0.19	0.17	0.17	0.21
Male	0.29	0.27	0.23	0.24	0.28
Female	0.31	0.24	0.26	0.24	0.33
ILLICIT DRUG USE DISORDER					
MDE	0.73	0.73	0.70	0.68	0.71
Male	1.46	1.42	1.50	1.33	1.33
Female	0.84	0.85	0.81	0.81	0.82
MDE WITH SEVERE IMPAIRMENT	0.99	0.94	0.88	0.86	0.89
Male	1.93	1.83	1.98	1.69	1.67
Female	1.14	1.08	0.99	1.03	1.04
NO MDE	0.17	0.16	0.15	0.14	0.20
Male	0.24	0.24	0.20	0.21	0.27
Female	0.24	0.20	0.21	0.17	0.30
ALCOHOL USE DISORDER					
MDE	0.70	0.61	0.57	0.55	0.57
Male	1.45	0.98	1.22	1.24	1.11
Female	0.80	0.75	0.67	0.61	0.63
MDE WITH SEVERE IMPAIRMENT	0.90	0.80	0.70	0.73	0.73
Male	1.94	1.29	1.58	1.65	1.39
Female	1.04	0.96	0.79	0.82	0.84
NO MDE	0.15	0.12	0.11	0.12	0.11
Male	0.22	0.15	0.14	0.15	0.14
Female	0.21	0.17	0.18	0.19	0.18

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

NOTE: Respondents with unknown impairment data were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2015-2019.

Table 11.11P Substance Use Disorder, Illicit Drug Use Disorder, and Alcohol Use Disorder in Past Year among Persons Aged 12 to 17, by Past Year Major Depressive Episode (MDE) and MDE with Severe Impairment Status and Gender: *P* Values from Tests of Differences of Percentages, 2019 Versus 2015-2018

Substance Use Disorder/ MDE Status/Gender	2015	2016	2017	2018	2019
SUBSTANCE USE DISORDER					
MDE	0.3724	0.8146	0.8398	0.8406	
Male	0.4612	0.5892	0.4547	0.4523	
Female	0.5461	0.5979	0.8313	0.4645	
MDE WITH SEVERE IMPAIRMENT	0.2353	0.7862	0.8813	0.7128	
Male	0.4953	0.3760	0.6713	0.9250	
Female	0.3203	0.4459	0.6760	0.6360	
NO MDE	0.0278	0.5574	0.1026	0.0021	
Male	0.0337	0.7464	0.2384	0.0700	
Female	0.3722	0.2227	0.2615	0.0087	
ILLICIT DRUG USE DISORDER					
MDE	0.7586	0.5457	0.8601	0.8111	
Male	0.7132	0.5767	0.6312	0.7411	
Female	0.8938	0.7131	0.6263	0.6331	
MDE WITH SEVERE IMPAIRMENT	0.6369	0.7915	0.8747	0.8085	
Male	0.8840	0.3916	0.6974	0.7440	
Female	0.5277	0.8510	0.6864	0.9378	
NO MDE	0.7564	0.2241	0.0361	0.0002	
Male	0.6623	0.7362	0.1054	0.0295	
Female	0.9841	0.1377	0.1594	0.0011	
ALCOHOL USE DISORDER					
MDE	0.0273	0.1386	0.5583	0.9106	
Male	0.2606	0.2921	0.7147	0.8142	
Female	0.0518	0.0249	0.6379	0.7649	
MDE WITH SEVERE IMPAIRMENT	0.0388	0.2057	0.8731	0.8550	
Male	0.1988	0.3187	0.9926	0.8832	
Female	0.0938	0.0541	0.8539	0.7584	
NO MDE	0.0000	0.1677	0.3414	0.9425	
Male	0.0000	0.0568	0.4062	0.8362	
Female	0.0659	1.0000	0.6306	0.9214	

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

NOTE: Respondents with unknown impairment data were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2015-2019.