

Table 11.12A Co-Occurring Substance Use Disorder, Illicit Drug Use Disorder, or Alcohol Use Disorder and Major Depressive Episode (MDE) or MDE with Severe Impairment in Past Year among Persons Aged 12 to 17, by Gender: Numbers in Thousands, 2015-2019

Substance Use Disorder (SUD), Illicit Drug Use Disorder (IDUD), or Alcohol Use Disorder (AUD) and MDE/Gender	2015	2016	2017	2018	2019
SUD and MDE	350	333	345	358	397
Male	87	72	101	114	111
Female	263	261	244	244	286
SUD and MDE with Severe Impairment	303	278	276	288	332
Male	70	56 ^a	81	82	94
Female	232	222	195	207	238
IDUD and MDE	241 ^a	236 ^a	260	279	313
Male	57	59	80	87	92
Female	184	177	180	193	221
IDUD and MDE with Severe Impairment	219	202	215	227	258
Male	48	44	66	60	76
Female	172	158	149	167	182
AUD and MDE	202	182	164	159	176
Male	51	27	47	50	53
Female	151	156	118	108	123
AUD and MDE with Severe Impairment	172	152	124	132	151
Male	43	22	34	39	43
Female	129	130	90	93	107

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

NOTE: Respondents with unknown impairment data were excluded.

^a The difference between this estimate and the 2019 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2015-2019.

Table 11.12B Co-Occurring Substance Use Disorder, Illicit Drug Use Disorder, or Alcohol Use Disorder and Major Depressive Episode (MDE) or MDE with Severe Impairment in Past Year among Persons Aged 12 to 17, by Gender: Percentages, 2015-2019

Substance Use Disorder (SUD), Illicit Drug Use Disorder (IDUD), or Alcohol Use Disorder (AUD) and MDE/Gender	2015	2016	2017	2018	2019
SUD and MDE	1.4	1.4	1.4	1.5	1.7
Male	0.7	0.6	0.8	0.9	0.9
Female	2.2	2.2	2.1	2.1	2.4
SUD and MDE with Severe Impairment	1.2	1.1	1.1	1.2	1.4
Male	0.6	0.5 ^a	0.7	0.7	0.8
Female	2.0	1.9	1.6	1.8	2.0
IDUD and MDE	1.0 ^a	1.0 ^a	1.1	1.2	1.3
Male	0.5	0.5	0.6	0.7	0.7
Female	1.6	1.5	1.5	1.6	1.9
IDUD and MDE with Severe Impairment	0.9	0.8	0.9	0.9	1.1
Male	0.4	0.4 ^a	0.5	0.5	0.6
Female	1.4	1.3	1.3	1.4	1.6
AUD and MDE	0.8	0.8	0.7	0.7	0.7
Male	0.4	0.2	0.4	0.4	0.4
Female	1.3	1.3	1.0	0.9	1.0
AUD and MDE with Severe Impairment	0.7	0.6	0.5	0.5	0.6
Male	0.3	0.2	0.3	0.3	0.4
Female	1.1	1.1	0.8	0.8	0.9

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

NOTE: Respondents with unknown impairment data were excluded.

^a The difference between this estimate and the 2019 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2015-2019.