

Table 11.3A Major Depressive Episode with Severe Impairment in Past Year among Persons Aged 12 to 17, by Demographic Characteristics: Numbers in Thousands, 2006-2019

Demographic Characteristic	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
TOTAL	1,358 ^a	1,371 ^a	1,460 ^a	1,404 ^a	1,350 ^a	1,388 ^a	1,544 ^a	1,868 ^a	1,990 ^a	2,129 ^a	2,168 ^a	2,265 ^a	2,423 ^a	2,671
AGE GROUP														
12-13	211 ^a	200 ^a	239 ^a	235 ^a	232 ^a	218 ^a	285 ^a	314 ^a	375 ^a	388 ^a	354 ^a	332 ^a	382 ^a	530
14-15	518 ^a	500 ^a	505 ^a	521 ^a	479 ^a	487 ^a	590 ^a	752 ^a	707 ^a	826 ^a	789 ^a	861	891	970
16-17	629 ^a	671 ^a	716 ^a	648 ^a	639 ^a	683 ^a	669 ^a	801 ^a	909 ^a	915 ^a	1,025 ^a	1,072	1,150	1,171
GENDER														
Male	335 ^a	386 ^a	359 ^a	391 ^a	395 ^a	397 ^a	373 ^a	435 ^a	461 ^a	477 ^a	539 ^a	581 ^a	628 ^a	736
Female	1,023 ^a	986 ^a	1,101 ^a	1,013 ^a	954 ^a	991 ^a	1,172 ^a	1,432 ^a	1,529 ^a	1,652 ^a	1,629 ^a	1,684 ^a	1,795	1,935
HISPANIC ORIGIN AND RACE														
Not Hispanic or Latino	1,118 ^a	1,141 ^a	1,226 ^a	1,150 ^a	1,093 ^a	1,113 ^a	1,152 ^a	1,425 ^a	1,540 ^a	1,651 ^a	1,703 ^a	1,694 ^a	1,841	1,981
White	871 ^a	873 ^a	944 ^a	858 ^a	853 ^a	799 ^a	883 ^a	1,046 ^a	1,167 ^a	1,258 ^a	1,290	1,256	1,354	1,415
Black or African American	150 ^a	193 ^a	171 ^a	204	157 ^a	183 ^a	164 ^a	207	214	198	196	233	225	256
AIAN	9	4 ^a	8	5 ^a	7 ^a	13	4 ^a	6 ^a	8 ^a	*	8	6 ^a	*	19
NHOPI	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Asian	54 ^a	39 ^a	44 ^a	48 ^a	44 ^a	60 ^a	30 ^a	98	80 ^a	67 ^a	118	101	127	142
Two or More Races	32 ^a	32 ^a	50 ^a	31 ^a	30 ^a	57 ^a	64 ^a	61 ^a	66 ^a	98	86 ^a	89 ^a	109	134
Hispanic or Latino	240 ^a	230 ^a	233 ^a	254 ^a	257 ^a	275 ^a	392 ^a	442 ^a	450 ^a	477 ^a	465 ^a	571 ^a	582	690

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.
NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section 3.3.5 in Chapter 3 of the 2019 National Survey on Drug Use and Health: Methodological Summary and Definitions).
NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.
NOTE: Respondents with unknown impairment data were excluded.

^a The difference between this estimate and the 2019 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2019.

Table 11.3B Major Depressive Episode with Severe Impairment in Past Year among Persons Aged 12 to 17, by Demographic Characteristics: Percentages, 2006-2019

Demographic Characteristic	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
TOTAL	5.5 ^a	5.5 ^a	6.0 ^a	5.8 ^a	5.7 ^a	5.7 ^a	6.3 ^a	7.7 ^a	8.2 ^a	8.8 ^a	9.0 ^a	9.4 ^a	10.0 ^a	11.1
AGE GROUP														
12-13	2.7 ^a	2.5 ^a	3.2 ^a	3.2 ^a	3.0 ^a	2.8 ^a	3.7 ^a	4.1 ^a	4.9 ^a	5.1 ^a	4.7 ^a	4.4 ^a	5.1 ^a	6.8
14-15	6.0 ^a	6.0 ^a	6.1 ^a	6.2 ^a	6.1 ^a	5.9 ^a	7.1 ^a	9.1 ^a	8.5 ^a	9.8 ^a	9.4 ^a	10.4	10.9	11.9
16-17	7.5 ^a	7.9 ^a	8.4 ^a	7.7 ^a	7.7 ^a	8.1 ^a	8.0 ^a	9.7 ^a	10.9 ^a	11.1 ^a	12.4 ^a	12.7 ^a	13.7	14.5
GENDER														
Male	2.6 ^a	3.0 ^a	2.9 ^a	3.2 ^a	3.2 ^a	3.2 ^a	3.0 ^a	3.5 ^a	3.7 ^a	3.8 ^a	4.4 ^a	4.7 ^a	5.1 ^a	6.0
Female	8.4 ^a	8.2 ^a	9.3 ^a	8.6 ^a	8.2 ^a	8.3 ^a	9.8 ^a	12.0 ^a	13.0 ^a	14.0 ^a	13.7 ^a	14.2 ^a	15.2	16.5
HISPANIC ORIGIN AND RACE														
Not Hispanic or Latino	5.5 ^a	5.7 ^a	6.2 ^a	5.9 ^a	5.7 ^a	5.8 ^a	6.1 ^a	7.6 ^a	8.2 ^a	8.9 ^a	9.2 ^a	9.2 ^a	10.1	11.0
White	5.8 ^a	5.9 ^a	6.5 ^a	6.1 ^a	6.2 ^a	5.9 ^a	6.5 ^a	7.8 ^a	8.9 ^a	9.7 ^a	10.0 ^a	9.8 ^a	10.7	11.4
Black or African American	3.9 ^a	5.1 ^a	4.6 ^a	5.7 ^a	4.5 ^a	5.4 ^a	4.8 ^a	6.2	6.4	5.9 ^a	6.0 ^a	7.1	6.9	8.0
AIAN	6.6	2.6 ^a	6.5	4.3 ^a	5.4	9.8	2.6 ^a	3.8 ^a	4.9	*	5.7	3.9 ^a	*	11.5
NHOPI	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Asian	5.3 ^a	3.9 ^a	4.7 ^a	5.0 ^a	4.3 ^a	5.0 ^a	2.6 ^a	8.1	6.6 ^a	5.5 ^a	9.3	7.9	9.2	11.3
Two or More Races	8.0 ^a	7.8 ^a	10.2	6.0 ^a	5.9 ^a	8.1 ^a	9.0 ^a	8.4 ^a	8.9 ^a	12.6	10.9	10.9	14.2	14.9
Hispanic or Latino	5.4 ^a	5.1 ^a	5.1 ^a	5.4 ^a	5.4 ^a	5.2 ^a	7.3 ^a	8.2 ^a	8.2 ^a	8.5 ^a	8.2 ^a	9.9	10.0	11.7

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.
NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section 3.3.5 in Chapter 3 of the 2019 National Survey on Drug Use and Health: Methodological Summary and Definitions).
NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.
NOTE: Respondents with unknown impairment data were excluded.

^a The difference between this estimate and the 2019 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2019.

Table 11.5A Receipt of Treatment for Depression in Past Year among Persons Aged 12 to 17 with Major Depressive Episode with Severe Impairment in Past Year, by Demographic Characteristics: Numbers in Thousands, 2006-2019

Demographic Characteristic	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
TOTAL	627 ^a	601 ^a	622 ^a	542 ^a	554 ^a	602 ^a	633 ^a	832 ^a	882 ^a	945 ^a	1,001 ^a	1,074 ^a	1,133 ^a	1,321
AGE GROUP														
12-13	92 ^a	85 ^a	91 ^a	78 ^a	66 ^a	91 ^a	91 ^a	150 ^a	143 ^a	142 ^a	131 ^a	153 ^a	177	218
14-15	232 ^a	216 ^a	194 ^a	184 ^a	215 ^a	200 ^a	239 ^a	320 ^a	307 ^a	374 ^a	369 ^a	364 ^a	406	492
16-17	304 ^a	300 ^a	336 ^a	280 ^a	273 ^a	311 ^a	302 ^a	363 ^a	433 ^a	429 ^a	501 ^a	558	550	612
GENDER														
Male	134 ^a	154 ^a	132 ^a	132 ^a	132 ^a	147 ^a	127 ^a	157 ^a	189 ^a	206 ^a	208 ^a	208 ^a	274	300
Female	493 ^a	447 ^a	489 ^a	410 ^a	422 ^a	455 ^a	506 ^a	675 ^a	693 ^a	739 ^a	793 ^a	866 ^a	859 ^a	1,021
HISPANIC ORIGIN AND RACE														
Not Hispanic or Latino	535 ^a	534 ^a	550 ^a	443 ^a	448 ^a	502 ^a	506 ^a	645 ^a	725 ^a	755 ^a	827 ^a	868 ^a	882 ^a	1,018
White	432 ^a	439 ^a	461 ^a	352 ^a	380 ^a	379 ^a	403 ^a	504 ^a	572 ^a	563 ^a	668 ^a	679 ^a	695	800
Black or African American	53 ^a	72	65 ^a	58 ^a	42 ^a	82	56 ^a	72	99	97	71	97	90	108
AIAN	*	*	*	*	*	*	*	*	*	*	*	*	*	*
NHOPI	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Asian	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Two or More Races	*	*	*	*	*	*	*	*	*	*	*	*	*	63
Hispanic or Latino	92 ^a	67 ^a	72 ^a	99 ^a	106 ^a	100 ^a	127 ^a	187 ^a	158 ^a	190 ^a	174 ^a	206 ^a	251	304

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.
NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section 3.3.5 in Chapter 3 of the *2019 National Survey on Drug Use and Health: Methodological Summary and Definitions*).
NOTE: Respondents with unknown past year treatment for depression data were excluded.
NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.
NOTE: Respondents with unknown impairment data were excluded.

^a The difference between this estimate and the 2019 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2019.

Table 11.5B Receipt of Treatment for Depression in Past Year among Persons Aged 12 to 17 with Major Depressive Episode with Severe Impairment in Past Year, by Demographic Characteristics: Percentages, 2006-2019

Demographic Characteristic	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
TOTAL	46.5	43.9 ^a	42.6 ^a	38.8 ^a	41.1 ^a	43.5 ^a	41.0 ^a	45.0 ^a	44.7 ^a	44.6 ^a	46.7	47.5	46.9	49.7
AGE GROUP														
12-13	44.1	42.3	38.3	33.8	28.7 ^a	42.3	32.0 ^a	48.5	38.7	37.0	37.6	46.2	47.2	41.9
14-15	44.8	43.2	38.6 ^a	35.6 ^a	45.0	41.0 ^a	40.5 ^a	43.2 ^a	43.4	45.4	47.5	42.3 ^a	45.6	50.8
16-17	48.8	44.9 ^a	46.9	43.2 ^a	42.7 ^a	45.5	45.4 ^a	45.3 ^a	48.1	47.1	49.2	52.0	47.8	52.3
GENDER														
Male	40.1	40.2	36.9	34.0	33.4	37.3	34.4	36.4	41.2	43.1	39.0	35.8	43.7	41.2
Female	48.6	45.4 ^a	44.5 ^a	40.7 ^a	44.3 ^a	45.9 ^a	43.2 ^a	47.6 ^a	45.7 ^a	45.1 ^a	49.3	51.5	48.1	52.9
HISPANIC ORIGIN AND RACE														
Not Hispanic or Latino	48.0	46.8	44.9 ^a	38.7 ^a	41.1 ^a	45.2 ^a	43.9 ^a	45.6 ^a	47.3	46.0 ^a	48.9	51.3	48.1	51.7
White	49.8 ^a	50.4 ^a	48.9 ^a	41.1 ^a	44.5 ^a	47.4 ^a	45.7 ^a	48.6 ^a	49.2 ^a	45.0 ^a	52.1	54.1	51.6	57.0
Black or African American	35.5	37.4	38.0	28.9	26.9 ^a	44.6	34.4	34.9	46.9	48.8	37.1	41.6	40.1	42.3
AIAN	*	*	*	*	*	*	*	*	*	*	*	*	*	*
NHOPI	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Asian	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Two or More Races	*	*	*	*	*	*	*	*	*	*	*	*	*	47.3
Hispanic or Latino	39.4	29.4 ^a	30.8 ^a	39.3	41.3	36.4	32.6 ^a	43.1	35.6	40.0	38.4	36.1	43.2	44.0

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.
NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section 3.3.5 in Chapter 3 of the *2019 National Survey on Drug Use and Health: Methodological Summary and Definitions*).
NOTE: Respondents with unknown past year treatment for depression data were excluded.
NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.
NOTE: Respondents with unknown impairment data were excluded.

^a The difference between this estimate and the 2019 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2019.