

**Table 7.17D Average Number of Days Used Specific Substances in Past Year among Past Year Users of Specific Substances Aged 12 or Older: Standard Errors, 2002-2019**

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
Marijuana	1.90	1.92	2.04	2.13	1.96	2.21	2.19	2.11	2.20	2.33	2.17	2.36	1.90	1.93	1.90	1.95	1.80	1.81
Cocaine	2.84	2.77	2.91	3.19	3.70	3.01	2.73	2.60	3.13	2.99	3.54	4.32	2.30	2.83	2.74	2.85	2.92	2.83
Crack	5.52	7.66	8.60	8.34	6.20	7.83	7.24	7.51	8.94	10.56	9.99	13.24	9.26	10.74	9.42	11.22	11.01	9.97
Heroin	20.23	20.82	16.36	9.23	23.92	16.36	14.82	13.04	10.54	14.77	15.69	12.86	9.59	10.75	11.84	11.36	11.84	15.37
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	1.41	1.08	1.55	1.26	1.94
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	3.95	2.35	2.81	3.17	3.29
Methamphetamine	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	7.07	8.69	8.54	7.58	8.38
Alcohol	1.02	0.98	1.03	0.99	1.00	0.97	1.02	1.05	0.91	0.94	0.98	0.91	0.77	0.80	0.82	0.80	0.80	0.82

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.  
NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section 3.3.5 in Chapter 3 of the *2019 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2019.

**Table 7.17P   Average Number of Days Used Specific Substances in Past Year among Past Year Users of Specific Substances Aged 12 or Older: *P* Values from Tests of Differences, 2019 Versus 2002-2018**

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
Marijuana	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0013	0.0006	0.0056	0.0779	0.0998	
Cocaine	0.0928	0.5397	0.0590	0.0019	0.0038	0.0545	0.1080	0.7916	0.8993	0.5178	0.1629	0.4702	0.5772	0.8176	0.8571	0.6934	0.7627	
Crack	0.4869	0.3907	0.9836	0.9078	0.3827	0.7928	0.3395	0.6107	0.6950	0.5744	0.7423	0.2069	0.9573	0.9237	0.5329	0.6719	0.6480	
Heroin	0.3291	0.2303	0.3276	0.0023	0.2194	0.2699	0.6993	0.0247	0.0686	0.6597	0.3612	0.7932	0.9739	0.7092	0.8066	0.5636	0.7204	
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.8047	0.4262	0.9930	0.5431	
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.3111	0.3762	0.4458	0.4777	
Methamphetamine	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.0279	0.3543	0.4067	0.3746	
Alcohol	0.0004	0.0003	0.0043	0.0002	0.0007	0.0035	0.0677	0.1188	0.0208	0.0078	0.0148	0.1022	0.9322	0.6039	0.7619	0.5122	0.2542	

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.  
NOTE:   Some 2006 to 2010 estimates that are used in the comparisons may differ from previously published estimates due to updates (see Section 3.3.5 in Chapter 3 of the *2019 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

Definitions:   Measures and terms are defined in Appendix A.

Source:   SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2019.

**Table 7.18D    Average Number of Days Used Specific Substances in Past Month among Past Month Users of Specific Substances Aged 12 or Older: Standard Errors, 2002-2019**

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
Marijuana	0.22	0.24	0.24	0.24	0.24	0.26	0.25	0.24	0.27	0.26	0.25	0.27	0.20	0.20	0.19	0.20	0.20	0.19
Cocaine	0.33	0.34	0.48	0.43	0.64	0.48	0.50	0.54	0.60	0.48	0.66	0.82	0.44	0.49	0.47	0.48	0.40	0.47
Crack	0.76	1.06	1.13	0.85	1.64	1.13	1.27	1.22	1.22	1.64	1.25	1.55	1.31	1.42	1.53	1.06	1.14	1.31
Heroin	2.74	2.38	2.66	1.62	2.38	2.59	2.16	2.23	1.88	1.90	1.74	1.63	1.22	1.61	1.67	1.32	1.39	2.39
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.34	0.18	0.26	0.27	0.19
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.48	0.52	0.54	0.69	0.50
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	0.91	1.05	1.07	1.05	0.98
Misuse of Pain Relievers <sup>1</sup>	--	--	--	--	--	--	--	--	--	--	--	--	--	0.35	0.38	0.38	0.48	0.48
Misuse of Stimulants	--	--	--	--	--	--	--	--	--	--	--	--	--	0.40	0.40	0.40	0.35	0.37
Misuse of Tranquilizers	--	--	--	--	--	--	--	--	--	--	--	--	--	0.35	0.41	0.33	0.48	0.44
Misuse of Sedatives	--	--	--	--	--	--	--	--	--	--	--	--	--	0.93	1.38	1.09	0.72	1.03
Cigarettes	0.14	0.15	0.14	0.14	0.14	0.15	0.15	0.14	0.16	0.15	0.15	0.16	0.13	0.14	0.14	0.15	0.16	0.16
Smokeless Tobacco	--	--	--	--	--	--	--	--	--	--	--	--	--	0.33	0.34	0.33	0.35	0.35
Cigars	0.22	0.27	0.25	0.25	0.26	0.25	0.27	0.27	0.28	0.28	0.32	0.29	0.23	0.27	0.27	0.26	0.28	0.29
Alcohol	0.10	0.10	0.10	0.09	0.10	0.09	0.10	0.10	0.09	0.10	0.09	0.09	0.08	0.08	0.08	0.07	0.08	0.08
Binge Alcohol Use <sup>2</sup>	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.06	0.07	0.06	0.07	0.07

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.  
NOTE:   Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section 3.3.5 in Chapter 3 of the *2019 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

<sup>1</sup> Prescription psychotherapeutic subtypes were revised in 2016; one effect was the comparability of codeine products between 2015 and 2016.

<sup>2</sup> Estimates of binge alcohol use include use by those who were heavy alcohol users.

Definitions:   Measures and terms are defined in Appendix A.

Source:   SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2019.

**Table 7.18P    Average Number of Days Used Specific Substances in Past Month among Past Month Users of Specific Substances Aged 12 or Older: *P* Values from Tests of Differences, 2019 Versus 2002-2018**

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
Marijuana	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0008	0.0000	0.0024	0.0113	0.0204	0.0144	0.0034	0.0377	0.6748	
Cocaine	0.8833	0.3561	0.0485	0.4583	0.1524	0.4330	0.5222	0.8278	0.9063	0.0913	0.2972	0.5414	0.8858	0.9864	0.6362	0.5189	0.2752	
Crack	0.1507	0.2140	0.4825	0.1731	0.5237	0.5007	0.9037	0.3009	0.6628	0.2028	0.8923	0.9316	0.7680	0.5494	0.8760	0.3941	0.3158	
Heroin	0.1478	0.6767	0.4094	0.3418	0.5347	0.6828	0.9834	0.3652	0.3642	0.1302	0.5090	0.6601	0.9617	0.8062	0.7153	0.9873	0.7358	
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.3286	0.6880	0.2749	0.1118	
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.8540	0.7589	0.9475	0.8675	
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	0.0552	0.8471	0.6670	0.7459	
Misuse of Pain Relievers <sup>1</sup>	--	--	--	--	--	--	--	--	--	--	--	--	--	0.9404	0.8505	0.7699	0.9673	
Misuse of Stimulants	--	--	--	--	--	--	--	--	--	--	--	--	--	0.5952	0.5824	0.5769	0.9605	
Misuse of Tranquilizers	--	--	--	--	--	--	--	--	--	--	--	--	--	0.5419	0.4251	0.3483	0.8388	
Misuse of Sedatives	--	--	--	--	--	--	--	--	--	--	--	--	--	0.1891	0.8466	0.6839	0.0949	
Cigarettes	0.0000	0.0000	0.0000	0.0000	0.0000	0.0007	0.0006	0.0009	0.0410	0.0038	0.0022	0.0245	0.3803	0.5628	0.8207	0.3733	0.8774	
Smokeless Tobacco	--	--	--	--	--	--	--	--	--	--	--	--	--	0.1710	0.3500	0.2639	0.0842	
Cigars	0.0000	0.0000	0.0000	0.0000	0.0001	0.0000	0.0006	0.0012	0.0413	0.0025	0.0939	0.0032	0.0005	0.3440	0.9123	0.9760	0.3556	
Alcohol	0.1968	0.1750	0.1134	0.3947	0.3248	0.5037	0.0256	0.0132	0.3867	0.4431	0.2068	0.6472	0.0144	0.8334	0.6918	0.8455	0.9242	
Binge Alcohol Use <sup>2</sup>	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.3803	0.5574	0.4257	0.5391	

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE:   Some 2006 to 2010 estimates that are used in the comparisons may differ from previously published estimates due to updates (see Section 3.3.5 in Chapter 3 of the *2019 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

<sup>1</sup> Prescription psychotherapeutic subtypes were revised in 2016; one effect was the comparability of codeine products between 2015 and 2016.

<sup>2</sup> Estimates of binge alcohol use that are used in the comparisons include use by those who were heavy alcohol users.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2019.