

Table 7.19A Daily or Almost Daily Use of Specific Substances in Past Year among All Persons Aged 12 or Older and among Past Year Users of Specific Substances Aged 12 or Older: Numbers in Thousands, 2002-2019

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
DAILY/ALMOST DAILY USE IN PAST YEAR AMONG TOTAL POPULATION																		
Marijuana	3,129 <sup>a</sup>	3,090 <sup>a</sup>	3,240 <sup>a</sup>	3,376 <sup>a</sup>	3,116 <sup>a</sup>	3,556 <sup>a</sup>	3,871 <sup>a</sup>	4,070 <sup>a</sup>	4,603 <sup>a</sup>	4,979 <sup>a</sup>	5,354 <sup>a</sup>	5,729 <sup>a</sup>	6,508 <sup>a</sup>	6,532 <sup>a</sup>	6,971 <sup>a</sup>	8,123 <sup>a</sup>	8,657 <sup>a</sup>	10,079
Cocaine	121	84	139	192	186	154	81	51	69	44	113	151	66	91	74	126	146	108
Crack	48	19	91	90	35	77	19	25	26	19	21	67	46	60	24	68	40	30
Heroin	60 <sup>a</sup>	30 <sup>a</sup>	55 <sup>a</sup>	20 <sup>a</sup>	151	40 <sup>a</sup>	71	48 <sup>a</sup>	62 <sup>a</sup>	101	150	92	131	110	186	158	132	150
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	27	22	47	29	57
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	29	14	35	25	22
Methamphetamine	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	135 <sup>a</sup>	161 <sup>a</sup>	197	229	310
Alcohol	9,678 <sup>a</sup>	9,638 <sup>a</sup>	9,745 <sup>a</sup>	10,165 <sup>a</sup>	10,322 <sup>a</sup>	10,081 <sup>a</sup>	10,651	11,090	10,028 <sup>a</sup>	10,793	10,831	10,587	11,239	11,434	11,563	11,893	10,978	11,603
DAILY/ALMOST DAILY USE IN PAST YEAR AMONG PAST YEAR USERS																		
Marijuana	3,129 <sup>a</sup>	3,090 <sup>a</sup>	3,240 <sup>a</sup>	3,376 <sup>a</sup>	3,116 <sup>a</sup>	3,556 <sup>a</sup>	3,871 <sup>a</sup>	4,070 <sup>a</sup>	4,603 <sup>a</sup>	4,979 <sup>a</sup>	5,354 <sup>a</sup>	5,729 <sup>a</sup>	6,508 <sup>a</sup>	6,532 <sup>a</sup>	6,971 <sup>a</sup>	8,123 <sup>a</sup>	8,657 <sup>a</sup>	10,079
Cocaine	121	84	139	192	186	154	81	51	69	44	113	151	66	91	74	126	146	108
Crack	48	19	91	90	35	77	19	25	*	*	21	*	46	60	24	*	40	30
Heroin	*	*	*	20 <sup>a</sup>	*	*	*	48 <sup>a</sup>	62 <sup>a</sup>	101	*	92	131	110	186	158	132	150
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	27	22	47	29	57
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	29	14	35	25	22
Methamphetamine	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	135 <sup>a</sup>	161 <sup>a</sup>	197	229	310
Alcohol	9,678 <sup>a</sup>	9,638 <sup>a</sup>	9,745 <sup>a</sup>	10,165 <sup>a</sup>	10,322 <sup>a</sup>	10,081 <sup>a</sup>	10,651	11,090	10,028 <sup>a</sup>	10,793	10,831	10,587	11,239	11,434	11,563	11,893	10,978	11,603

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.  
NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section 3.3.5 in Chapter 3 of the 2019 National Survey on Drug Use and Health: Methodological Summary and Definitions).  
<sup>a</sup> The difference between this estimate and the 2019 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

Definitions: Measures and terms are defined in Appendix A.  
Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2019.

Table 7.19B Daily or Almost Daily Use of Specific Substances in Past Year among All Persons Aged 12 or Older and among Past Year Users of Specific Substances Aged 12 or Older: Percentages, 2002-2019

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
<b>DAILY/ALMOST DAILY USE IN PAST YEAR AMONG TOTAL POPULATION</b>																		
Marijuana	1.3 <sup>a</sup>	1.3 <sup>a</sup>	1.3 <sup>a</sup>	1.4 <sup>a</sup>	1.3 <sup>a</sup>	1.4 <sup>a</sup>	1.5 <sup>a</sup>	1.6 <sup>a</sup>	1.8 <sup>a</sup>	1.9 <sup>a</sup>	2.1 <sup>a</sup>	2.2 <sup>a</sup>	2.5 <sup>a</sup>	2.4 <sup>a</sup>	2.6 <sup>a</sup>	3.0 <sup>a</sup>	3.2 <sup>a</sup>	3.7
Cocaine	0.1	0.0	0.1	0.1	0.1	0.1	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.1	0.0
Crack	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Heroin	0.0	0.0 <sup>a</sup>	0.0 <sup>a</sup>	0.0 <sup>a</sup>	0.1	0.0 <sup>a</sup>	0.0	0.0 <sup>a</sup>	0.0 <sup>a</sup>	0.0	0.1	0.0	0.0	0.0	0.1	0.1	0.0	0.1
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.0	0.0	0.0	0.0	0.0
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.0	0.0	0.0	0.0	0.0
Methamphetamine	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.1 <sup>a</sup>	0.1 <sup>a</sup>	0.1	0.1	0.1
Alcohol	4.1	4.1	4.1	4.2	4.2	4.1	4.3	4.4	4.0	4.2	4.2	4.0	4.2	4.3	4.3	4.4	4.0	4.2
<b>DAILY/ALMOST DAILY USE IN PAST YEAR AMONG PAST YEAR USERS</b>																		
Marijuana	12.2 <sup>a</sup>	12.2 <sup>a</sup>	12.7 <sup>a</sup>	13.3 <sup>a</sup>	12.3 <sup>a</sup>	14.1 <sup>a</sup>	15.0 <sup>a</sup>	14.2 <sup>a</sup>	15.7 <sup>a</sup>	16.7 <sup>a</sup>	17.0 <sup>a</sup>	17.4 <sup>a</sup>	18.5 <sup>a</sup>	18.1 <sup>a</sup>	18.6 <sup>a</sup>	19.8	19.9	20.9
Cocaine	2.0	1.4	2.5	3.5	3.1	2.7	1.5	1.1	1.5	1.1	2.4	3.6	1.5	1.9	1.5	2.1	2.6	2.0
Crack	3.1	1.4	7.0	6.5	2.4	5.3	1.7	2.4	*	*	2.3	*	5.9	7.2	2.7	*	5.2	3.8
Heroin	*	*	*	5.2 <sup>a</sup>	*	*	*	8.2 <sup>a</sup>	10.0	16.3	*	13.5	14.3	13.3	19.6	17.9	16.3	20.1
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.6	0.5	0.9	0.5	0.9
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	1.7	0.9	2.0	1.2	1.1
Methamphetamine	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	7.9 <sup>a</sup>	11.6	12.1	12.3	15.5
Alcohol	6.2	6.2	6.2	6.3	6.4	6.2	6.5	6.6	6.0	6.3	6.2	6.1	6.4	6.5	6.6	6.7	6.1	6.5

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.  
NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section 3.3.5 in Chapter 3 of the *2019 National Survey on Drug Use and Health: Methodological Summary and Definitions*).  
<sup>a</sup> The difference between this estimate and the 2019 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

Definitions: Measures and terms are defined in Appendix A.  
Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2019.

Table 7.20A Daily or Almost Daily Use of Specific Substances in Past Year among All Persons Aged 12 to 17 and among Past Year Users of Specific Substances Aged 12 to 17: Numbers in Thousands, 2002-2019

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
DAILY/ALMOST DAILY USE IN PAST YEAR AMONG TOTAL POPULATION																		
Marijuana	358 <sup>a</sup>	282	342	268	288	260	276	291	337	301	277	299	257	258	253	250	244	283
Cocaine	10	10	4	7	3	7	5	3	2	2	*	*	*	*	2	*	2	*
Crack	4	*	*	3	*	4	*	*	*	2	*	*	*	*	*	*	*	*
Heroin	2	2	3	*	*	*	1	3	*	*	*	*	3	*	2	*	*	*
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	3	*	3	*
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	5	11	19	5	5
Methamphetamine	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	2	3	2	*
Alcohol	102 <sup>a</sup>	89 <sup>a</sup>	91 <sup>a</sup>	76 <sup>a</sup>	87 <sup>a</sup>	84 <sup>a</sup>	61 <sup>a</sup>	52 <sup>a</sup>	50	60 <sup>a</sup>	44	43	61 <sup>a</sup>	58 <sup>a</sup>	36	47	24	29
DAILY/ALMOST DAILY USE IN PAST YEAR AMONG PAST YEAR USERS																		
Marijuana	358 <sup>a</sup>	282	342	268	288	260	276	291	337	301	277	299	257	258	253	250	244	283
Cocaine	10	10	4	7	3	7	5	3	2	*	0	*	*	*	*	*	*	*
Crack	4	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Heroin	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	1	3	0	3	1
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	5	11	19	5	5
Methamphetamine	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	*	*	*	*
Alcohol	102 <sup>a</sup>	89 <sup>a</sup>	91 <sup>a</sup>	76 <sup>a</sup>	87 <sup>a</sup>	84 <sup>a</sup>	61 <sup>a</sup>	52 <sup>a</sup>	50	60 <sup>a</sup>	44	43	61 <sup>a</sup>	58 <sup>a</sup>	36	47	24	29

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.  
NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section 3.3.5 in Chapter 3 of the 2019 National Survey on Drug Use and Health: Methodological Summary and Definitions).  
<sup>a</sup> The difference between this estimate and the 2019 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

Definitions: Measures and terms are defined in Appendix A.  
Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2019.

Table 7.20B Daily or Almost Daily Use of Specific Substances in Past Year among All Persons Aged 12 to 17 and among Past Year Users of Specific Substances Aged 12 to 17: Percentages, 2002-2019

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
DAILY/ALMOST DAILY USE IN PAST YEAR AMONG TOTAL POPULATION																		
Marijuana	1.4 <sup>a</sup>	1.1	1.4	1.1	1.1	1.0	1.1	1.2	1.4	1.2	1.1	1.2	1.0	1.0	1.0	1.0	1.0	1.1
Cocaine	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	*	0.0	*	0.0	*
Crack	0.0	*	*	0.0	*	0.0	*	*	*	0.0	*	*	*	*	*	*	*	*
Heroin	0.0	0.0	0.0	*	*	*	0.0	0.0	*	*	*	*	0.0	*	0.0	*	*	*
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	0.0	*	0.0	*
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.0	0.0	0.1	0.0	0.0
Methamphetamine	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	0.0	0.0	0.0	*
Alcohol	0.4 <sup>a</sup>	0.4 <sup>a</sup>	0.4 <sup>a</sup>	0.3 <sup>a</sup>	0.3 <sup>a</sup>	0.3 <sup>a</sup>	0.2 <sup>a</sup>	0.2 <sup>a</sup>	0.2	0.2 <sup>a</sup>	0.2	0.2	0.2 <sup>a</sup>	0.2 <sup>a</sup>	0.1	0.2	0.1	0.1
DAILY/ALMOST DAILY USE IN PAST YEAR AMONG PAST YEAR USERS																		
Marijuana	9.2	7.5	9.4	7.9	8.6	8.2	8.5	8.7	9.9	8.5	8.2	8.9	7.9	8.2	8.5	8.1	7.8	8.6
Cocaine	1.9	2.2	1.1	1.6	0.8	1.8	1.5	1.1	0.7	*	0.1	*	*	*	*	*	*	*
Crack	4.4	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Heroin	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.2	0.6	0.1	0.8	0.1
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.7	2.1	3.4	0.8	0.7
Methamphetamine	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	*	*	*	*
Alcohol	1.2 <sup>a</sup>	1.0 <sup>a</sup>	1.1 <sup>a</sup>	0.9	1.0 <sup>a</sup>	1.0 <sup>a</sup>	0.8	0.7	0.7	0.9	0.7	0.7	1.0	1.0	0.7	0.9	0.5	0.5

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.  
NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section 3.3.5 in Chapter 3 of the 2019 National Survey on Drug Use and Health: Methodological Summary and Definitions).  
<sup>a</sup> The difference between this estimate and the 2019 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

Definitions: Measures and terms are defined in Appendix A.  
Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2019.

Table 7.21A Daily or Almost Daily Use of Specific Substances in Past Year among All Persons Aged 18 or Older and among Past Year Users of Specific Substances Aged 18 or Older: Numbers in Thousands, 2002-2019

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
DAILY/ALMOST DAILY USE IN PAST YEAR AMONG TOTAL POPULATION																		
Marijuana	2,772 <sup>a</sup>	2,808 <sup>a</sup>	2,898 <sup>a</sup>	3,108 <sup>a</sup>	2,828 <sup>a</sup>	3,296 <sup>a</sup>	3,595 <sup>a</sup>	3,779 <sup>a</sup>	4,265 <sup>a</sup>	4,678 <sup>a</sup>	5,077 <sup>a</sup>	5,430 <sup>a</sup>	6,251 <sup>a</sup>	6,274 <sup>a</sup>	6,718 <sup>a</sup>	7,873 <sup>a</sup>	8,414 <sup>a</sup>	9,796
Cocaine	111	74	134	185	182	147	77	49	67	42	112	150	65	91	72	126	144	108
Crack	44	18	91	87	34	73	19	25	26	18	21	67	46	60	24	68	40	30
Heroin	58 <sup>a</sup>	28 <sup>a</sup>	53 <sup>a</sup>	19 <sup>a</sup>	151	40 <sup>a</sup>	69	44 <sup>a</sup>	62 <sup>a</sup>	101	150	91	127	110	183	158	132	150
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	26	19	47	25	56
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	25	*	16	20	17
Methamphetamine	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	135 <sup>a</sup>	159 <sup>a</sup>	194	228	310
Alcohol	9,577 <sup>a</sup>	9,549 <sup>a</sup>	9,654 <sup>a</sup>	10,089 <sup>a</sup>	10,235 <sup>a</sup>	9,997 <sup>a</sup>	10,590	11,039	9,978 <sup>a</sup>	10,733	10,787	10,544	11,178	11,375	11,527	11,846	10,954	11,574
DAILY/ALMOST DAILY USE IN PAST YEAR AMONG PAST YEAR USERS																		
Marijuana	2,772 <sup>a</sup>	2,808 <sup>a</sup>	2,898 <sup>a</sup>	3,108 <sup>a</sup>	2,828 <sup>a</sup>	3,296 <sup>a</sup>	3,595 <sup>a</sup>	3,779 <sup>a</sup>	4,265 <sup>a</sup>	4,678 <sup>a</sup>	5,077 <sup>a</sup>	5,430 <sup>a</sup>	6,251 <sup>a</sup>	6,274 <sup>a</sup>	6,718 <sup>a</sup>	7,873 <sup>a</sup>	8,414 <sup>a</sup>	9,796
Cocaine	111	74	134	185	182	147	77	49	67	42	112	150	65	91	72	126	144	108
Crack	44	18	91	87	34	73	19	25	*	*	21	*	46	60	24	*	40	30
Heroin	*	*	*	19 <sup>a</sup>	*	*	*	44 <sup>a</sup>	62 <sup>a</sup>	101	*	91	127	110	183	158	132	150
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	26	19	47	25	56
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	3	16	*	17
Methamphetamine	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	135 <sup>a</sup>	159 <sup>a</sup>	194	228	310
Alcohol	9,577 <sup>a</sup>	9,549 <sup>a</sup>	9,654 <sup>a</sup>	10,089 <sup>a</sup>	10,235 <sup>a</sup>	9,997 <sup>a</sup>	10,590	11,039	9,978 <sup>a</sup>	10,733	10,787	10,544	11,178	11,375	11,527	11,846	10,954	11,574

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.  
NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section 3.3.5 in Chapter 3 of the 2019 National Survey on Drug Use and Health: Methodological Summary and Definitions).  
<sup>a</sup> The difference between this estimate and the 2019 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

Definitions: Measures and terms are defined in Appendix A.  
Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2019.

Table 7.21B Daily or Almost Daily Use of Specific Substances in Past Year among All Persons Aged 18 or Older and among Past Year Users of Specific Substances Aged 18 or Older: Percentages, 2002-2019

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
<b>DAILY/ALMOST DAILY USE IN PAST YEAR AMONG TOTAL POPULATION</b>																		
Marijuana	1.3 <sup>a</sup>	1.3 <sup>a</sup>	1.3 <sup>a</sup>	1.4 <sup>a</sup>	1.3 <sup>a</sup>	1.5 <sup>a</sup>	1.6 <sup>a</sup>	1.7 <sup>a</sup>	1.9 <sup>a</sup>	2.0 <sup>a</sup>	2.2 <sup>a</sup>	2.3 <sup>a</sup>	2.6 <sup>a</sup>	2.6 <sup>a</sup>	2.7 <sup>a</sup>	3.2 <sup>a</sup>	3.4 <sup>a</sup>	3.9
Cocaine	0.1	0.0	0.1	0.1	0.1	0.1	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.1	0.1	0.0
Crack	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Heroin	0.0	0.0 <sup>a</sup>	0.0 <sup>a</sup>	0.0 <sup>a</sup>	0.1	0.0 <sup>a</sup>	0.0	0.0 <sup>a</sup>	0.0 <sup>a</sup>	0.0	0.1	0.0	0.1	0.0	0.1	0.1	0.1	0.1
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.0	0.0	0.0	0.0	0.0
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.0	*	0.0	0.0	0.0
Methamphetamine	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.1 <sup>a</sup>	0.1 <sup>a</sup>	0.1	0.1	0.1
Alcohol	4.6	4.5	4.5	4.6	4.6	4.5	4.7	4.9	4.4	4.6	4.6	4.4	4.7	4.7	4.7	4.8	4.4	4.6
<b>DAILY/ALMOST DAILY USE IN PAST YEAR AMONG PAST YEAR USERS</b>																		
Marijuana	12.7 <sup>a</sup>	13.1 <sup>a</sup>	13.3 <sup>a</sup>	14.1 <sup>a</sup>	12.8 <sup>a</sup>	15.0 <sup>a</sup>	15.9 <sup>a</sup>	14.9 <sup>a</sup>	16.5 <sup>a</sup>	17.9 <sup>a</sup>	18.0 <sup>a</sup>	18.3 <sup>a</sup>	19.6 <sup>a</sup>	19.1 <sup>a</sup>	19.4 <sup>a</sup>	20.8	20.8	21.8
Cocaine	2.1	1.4	2.6	3.6	3.2	2.7	1.5	1.1	1.6	1.2	2.5	3.7	1.5	2.0	1.5	2.2	2.7	2.0
Crack	3.0	1.4	7.4	6.5	2.4	5.3	1.7	2.5	*	*	2.4	*	6.1	7.3	2.7	*	5.3	3.9
Heroin	*	*	*	5.6 <sup>a</sup>	*	*	*	8.1 <sup>a</sup>	10.4	17.8	*	14.1	14.4	13.7	19.6	18.1	16.5	20.1
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.6	0.4	1.0	0.5	1.0
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	0.3	1.3	*	1.2
Methamphetamine	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	8.1 <sup>a</sup>	11.7	12.2	12.5	15.8
Alcohol	6.5	6.5	6.5	6.6	6.6	6.5	6.7	6.9	6.2	6.6	6.5	6.3	6.6	6.7	6.8	6.8	6.3	6.7

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.  
NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section 3.3.5 in Chapter 3 of the *2019 National Survey on Drug Use and Health: Methodological Summary and Definitions*).  
<sup>a</sup> The difference between this estimate and the 2019 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

Definitions: Measures and terms are defined in Appendix A.  
Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2019.

Table 7.22A Daily or Almost Daily Use of Specific Substances in Past Year among All Persons Aged 18 to 25 and among Past Year Users of Specific Substances Aged 18 to 25: Numbers in Thousands, 2002-2019

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
<b>DAILY/ALMOST DAILY USE IN PAST YEAR AMONG TOTAL POPULATION</b>																		
Marijuana	1,336 <sup>a</sup>	1,375 <sup>a</sup>	1,379 <sup>a</sup>	1,519 <sup>a</sup>	1,487 <sup>a</sup>	1,479 <sup>a</sup>	1,656 <sup>a</sup>	1,691 <sup>a</sup>	1,863 <sup>a</sup>	1,825 <sup>a</sup>	2,157 <sup>a</sup>	2,166 <sup>a</sup>	2,224 <sup>a</sup>	2,353	2,218 <sup>a</sup>	2,618	2,483	2,540
Cocaine	17	24	20	31	28	22	14	23	7	9	20	6	8	19	10	28	5	22
Crack	3	4	9	5	7	8	5	3	3	*	4	*	3	7	*	*	*	11
Heroin	6 <sup>a</sup>	8 <sup>a</sup>	8 <sup>a</sup>	11	8	16	24	17	33	28	43	25	17	29	38	41	24	26
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	14	5	16	6	6
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	2	7	2	*
Methamphetamine	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	15	28	25	25	30
Alcohol	681	733	681	668	678	669	710	672	642	604	636	599	645	726	721	783	659	689
<b>DAILY/ALMOST DAILY USE IN PAST YEAR AMONG PAST YEAR USERS</b>																		
Marijuana	1,336 <sup>a</sup>	1,375 <sup>a</sup>	1,379 <sup>a</sup>	1,519 <sup>a</sup>	1,487 <sup>a</sup>	1,479 <sup>a</sup>	1,656 <sup>a</sup>	1,691 <sup>a</sup>	1,863 <sup>a</sup>	1,825 <sup>a</sup>	2,157 <sup>a</sup>	2,166 <sup>a</sup>	2,224 <sup>a</sup>	2,353	2,218 <sup>a</sup>	2,618	2,483	2,540
Cocaine	17	24	20	31	28	22	14	23	7	9	20	6	8	19	10	28	5	22
Crack	3	4	9	5	7	8	5	3	3	*	*	*	*	*	*	*	*	*
Heroin	*	*	*	11	*	*	24	*	33	28	*	25	17	*	38	*	*	*
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	14	5	16	6	6
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	*	*	*	0
Methamphetamine	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	15	28	25	25	*
Alcohol	681	733	681	668	678	669	710	672	642	604	636	599	645	726	721	783	659	689

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.  
NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section 3.3.5 in Chapter 3 of the 2019 National Survey on Drug Use and Health: Methodological Summary and Definitions).  
<sup>a</sup> The difference between this estimate and the 2019 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

Definitions: Measures and terms are defined in Appendix A.  
Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2019.

Table 7.22B Daily or Almost Daily Use of Specific Substances in Past Year among All Persons Aged 18 to 25 and among Past Year Users of Specific Substances Aged 18 to 25: Percentages, 2002-2019

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
DAILY/ALMOST DAILY USE IN PAST YEAR AMONG TOTAL POPULATION																		
Marijuana	4.3 <sup>a</sup>	4.3 <sup>a</sup>	4.3 <sup>a</sup>	4.7 <sup>a</sup>	4.5 <sup>a</sup>	4.5 <sup>a</sup>	5.0 <sup>a</sup>	5.0 <sup>a</sup>	5.5 <sup>a</sup>	5.3 <sup>a</sup>	6.2 <sup>a</sup>	6.2 <sup>a</sup>	6.4 <sup>a</sup>	6.7 <sup>a</sup>	6.4 <sup>a</sup>	7.6	7.3	7.5
Cocaine	0.1	0.1	0.1	0.1	0.1	0.1	0.0	0.1	0.0	0.0	0.1	0.0	0.0	0.1	0.0	0.1	0.0	0.1
Crack	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	*	0.0	0.0	*	*	*	0.0
Heroin	0.0 <sup>a</sup>	0.0	0.0 <sup>a</sup>	0.0	0.0	0.0	0.1	0.1	0.1	0.1	0.1	0.1	0.0	0.1	0.1	0.1	0.1	0.1
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.0	0.0	0.0	0.0	0.0
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	0.0	0.0	0.0	*
Methamphetamine	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.0	0.1	0.1	0.1	0.1
Alcohol	2.2	2.3	2.1	2.1	2.1	2.0	2.2	2.0	1.9	1.8	1.8	1.7	1.8	2.1	2.1	2.3	1.9	2.0
DAILY/ALMOST DAILY USE IN PAST YEAR AMONG PAST YEAR USERS																		
Marijuana	14.5 <sup>a</sup>	15.2 <sup>a</sup>	15.4 <sup>a</sup>	16.7 <sup>a</sup>	16.2 <sup>a</sup>	16.4 <sup>a</sup>	18.1 <sup>a</sup>	16.3 <sup>a</sup>	18.2 <sup>a</sup>	17.3 <sup>a</sup>	19.8	19.7	19.9	20.9	19.5	21.9	21.0	21.3
Cocaine	0.8	1.1	0.9	1.4	1.2	1.0	0.8	1.3	0.4	0.6	1.3	0.4	0.5	1.0	0.5	1.3	0.3	1.2
Crack	1.2	1.4	3.3	1.5	2.5	3.4	2.3	1.6	1.4	*	*	*	*	*	*	*	*	*
Heroin	*	*	*	6.9	*	*	16.2	*	15.6	12.2	*	10.3	6.4	*	16.8	*	*	*
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.6	0.2	0.7	0.3	0.3
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	*	*	*	0.1
Methamphetamine	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	4.5	10.7	6.7	9.2	*
Alcohol	2.8	3.0	2.7	2.6	2.6	2.6	2.8	2.5	2.4	2.3 <sup>a</sup>	2.4	2.2 <sup>a</sup>	2.4	2.8	2.8	3.1	2.6	2.8

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.  
NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section 3.3.5 in Chapter 3 of the *2019 National Survey on Drug Use and Health: Methodological Summary and Definitions*).  
<sup>a</sup> The difference between this estimate and the 2019 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

Definitions: Measures and terms are defined in Appendix A.  
Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2019.



Table 7.23A Daily or Almost Daily Use of Specific Substances in Past Year among All Persons Aged 26 or Older and among Past Year Users of Specific Substances Aged 26 or Older: Numbers in Thousands, 2002-2019

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
DAILY/ALMOST DAILY USE IN PAST YEAR AMONG TOTAL POPULATION																		
Marijuana	1,435 <sup>a</sup>	1,433 <sup>a</sup>	1,519 <sup>a</sup>	1,589 <sup>a</sup>	1,341 <sup>a</sup>	1,817 <sup>a</sup>	1,939 <sup>a</sup>	2,088 <sup>a</sup>	2,402 <sup>a</sup>	2,853 <sup>a</sup>	2,920 <sup>a</sup>	3,264 <sup>a</sup>	4,027 <sup>a</sup>	3,921 <sup>a</sup>	4,500 <sup>a</sup>	5,255 <sup>a</sup>	5,930 <sup>a</sup>	7,256
Cocaine	94	51	115	154	154	125	62	26	60	33	92	145	57	72	62	98	138	86
Crack	40	14	83	82	27	65	14	22	24	16	17	66	43	53	24	68	39	19
Heroin	52	20 <sup>a</sup>	45	*	143	24 <sup>a</sup>	45	27 <sup>a</sup>	29 <sup>a</sup>	73	107	66	110	81	145	117	107	123
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	13	14	31	19	50
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	25	*	*	18	17
Methamphetamine	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	120 <sup>a</sup>	130 <sup>a</sup>	169	202	280
Alcohol	8,896 <sup>a</sup>	8,816 <sup>a</sup>	8,973 <sup>a</sup>	9,421 <sup>a</sup>	9,557 <sup>a</sup>	9,328 <sup>a</sup>	9,880	10,367	9,336 <sup>a</sup>	10,129	10,150	9,945	10,533	10,649	10,806	11,064	10,295	10,885
DAILY/ALMOST DAILY USE IN PAST YEAR AMONG PAST YEAR USERS																		
Marijuana	1,435 <sup>a</sup>	1,433 <sup>a</sup>	1,519 <sup>a</sup>	1,589 <sup>a</sup>	1,341 <sup>a</sup>	1,817 <sup>a</sup>	1,939 <sup>a</sup>	2,088 <sup>a</sup>	2,402 <sup>a</sup>	2,853 <sup>a</sup>	2,920 <sup>a</sup>	3,264 <sup>a</sup>	4,027 <sup>a</sup>	3,921 <sup>a</sup>	4,500 <sup>a</sup>	5,255 <sup>a</sup>	5,930 <sup>a</sup>	7,256
Cocaine	94	51	115	154	154	125	62	26	60	33	92	145	57	72	62	98	138	86
Crack	40	14	83	82	27	65	14	*	*	*	*	*	43	*	24	*	39	19
Heroin	*	*	*	*	*	*	*	*	*	*	*	*	110	81	145	117	107	*
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	14	31	19	50
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	1	*	*	*
Methamphetamine	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	120 <sup>a</sup>	130 <sup>a</sup>	169	202	280
Alcohol	8,896 <sup>a</sup>	8,816 <sup>a</sup>	8,973 <sup>a</sup>	9,421 <sup>a</sup>	9,557 <sup>a</sup>	9,328 <sup>a</sup>	9,880	10,367	9,336 <sup>a</sup>	10,129	10,150	9,945	10,533	10,649	10,806	11,064	10,295	10,885

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.  
NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section 3.3.5 in Chapter 3 of the 2019 National Survey on Drug Use and Health: Methodological Summary and Definitions).  
<sup>a</sup> The difference between this estimate and the 2019 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

Definitions: Measures and terms are defined in Appendix A.  
Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2019.

Table 7.23B Daily or Almost Daily Use of Specific Substances in Past Year among All Persons Aged 26 or Older and among Past Year Users of Specific Substances Aged 26 or Older: Percentages, 2002-2019

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
DAILY/ALMOST DAILY USE IN PAST YEAR AMONG TOTAL POPULATION																		
Marijuana	0.8 <sup>a</sup>	0.8 <sup>a</sup>	0.8 <sup>a</sup>	0.9 <sup>a</sup>	0.7 <sup>a</sup>	1.0 <sup>a</sup>	1.0 <sup>a</sup>	1.1 <sup>a</sup>	1.2 <sup>a</sup>	1.4 <sup>a</sup>	1.5 <sup>a</sup>	1.6 <sup>a</sup>	2.0 <sup>a</sup>	1.9 <sup>a</sup>	2.1 <sup>a</sup>	2.5 <sup>a</sup>	2.8 <sup>a</sup>	3.4
Cocaine	0.1	0.0	0.1	0.1	0.1	0.1	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.1	0.0
Crack	0.0	0.0	0.0 <sup>a</sup>	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Heroin	0.0	0.0 <sup>a</sup>	0.0	*	0.1	0.0 <sup>a</sup>	0.0	0.0 <sup>a</sup>	0.0 <sup>a</sup>	0.0	0.1	0.0	0.1	0.0	0.1	0.1	0.1	0.1
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.0	0.0	0.0	0.0	0.0
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.0	*	*	0.0	0.0
Methamphetamine	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.1 <sup>a</sup>	0.1 <sup>a</sup>	0.1	0.1	0.1
Alcohol	5.0	4.9	4.9	5.1	5.1	4.9	5.1	5.4	4.8	5.1	5.1	4.9	5.1	5.1	5.1	5.2	4.8	5.0
DAILY/ALMOST DAILY USE IN PAST YEAR AMONG PAST YEAR USERS																		
Marijuana	11.4 <sup>a</sup>	11.5 <sup>a</sup>	11.8 <sup>a</sup>	12.3 <sup>a</sup>	10.4 <sup>a</sup>	14.0 <sup>a</sup>	14.4 <sup>a</sup>	13.9 <sup>a</sup>	15.3 <sup>a</sup>	18.3 <sup>a</sup>	16.9 <sup>a</sup>	17.5 <sup>a</sup>	19.4 <sup>a</sup>	18.1 <sup>a</sup>	19.4 <sup>a</sup>	20.3	20.8	22.0
Cocaine	2.8	1.5	3.7	5.4	4.5	3.8	2.0	0.9	2.2	1.6	3.2	5.8	2.1	2.6	2.0	2.6	4.0	2.4
Crack	3.4	1.4	8.5	8.2	2.4	5.7	1.6	*	*	*	*	*	6.9	*	3.0	*	5.9	2.7
Heroin	*	*	*	*	*	*	*	*	*	*	*	*	17.8	13.7	20.5	17.8	16.8	*
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	0.7	1.4	0.7	1.6
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	0.2	*	*	*
Methamphetamine	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	8.9 <sup>a</sup>	11.9	13.9	13.1	16.7
Alcohol	7.2	7.3	7.3	7.4	7.5	7.2	7.5	7.7	6.9	7.4	7.2	7.0	7.3	7.4	7.5	7.5	6.9	7.3

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.  
NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section 3.3.5 in Chapter 3 of the *2019 National Survey on Drug Use and Health: Methodological Summary and Definitions*).  
<sup>a</sup> The difference between this estimate and the 2019 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

Definitions: Measures and terms are defined in Appendix A.  
Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2019.

Table 7.24A Daily or Almost Daily Use of Specific Substances in Past Year among All Persons Aged 12 to 20 and among Past Year Users of Specific Substances Aged 12 to 20: Numbers in Thousands, 2002-2019

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
DAILY/ALMOST DAILY USE IN PAST YEAR AMONG TOTAL POPULATION																		
Marijuana	976	900 <sup>a</sup>	886 <sup>a</sup>	971	949	858 <sup>a</sup>	1,052	1,011	1,036	1,050	1,161	1,108	1,077	1,100	964	1,131	1,078	1,059
Cocaine	19	20	10	14	14	14	9	13	4	11	15	3	4	10	8	8	3	6
Crack	5	4	2	4	2	4	3	*	3	3	3	*	*	4	*	*	*	4
Heroin	3	9	3	6	*	5	6	4	16	12	13	4	13	8	5	4	5	4
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	4	3	3	5	*
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	5	13	26	5	5
Methamphetamine	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	10	8	13	7	7
Alcohol	298 <sup>a</sup>	318 <sup>a</sup>	231 <sup>a</sup>	234 <sup>a</sup>	244 <sup>a</sup>	267 <sup>a</sup>	242 <sup>a</sup>	207 <sup>a</sup>	215 <sup>a</sup>	193 <sup>a</sup>	143	173	185	226 <sup>a</sup>	126	191	162	128
DAILY/ALMOST DAILY USE IN PAST YEAR AMONG PAST YEAR USERS																		
Marijuana	976	900 <sup>a</sup>	886 <sup>a</sup>	971	949	858 <sup>a</sup>	1,052	1,011	1,036	1,050	1,161	1,108	1,077	1,100	964	1,131	1,078	1,059
Cocaine	19	20	10	14	14	14	9	13	4	11	15	3	4	10	8	8	3	6
Crack	5	*	*	*	2	*	*	*	*	*	*	*	*	*	*	*	*	*
Heroin	3	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	4	3	3	5	1
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	5	13	26	5	5
Methamphetamine	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	*	*	*	*
Alcohol	298 <sup>a</sup>	318 <sup>a</sup>	231 <sup>a</sup>	234 <sup>a</sup>	244 <sup>a</sup>	267 <sup>a</sup>	242 <sup>a</sup>	207 <sup>a</sup>	215 <sup>a</sup>	193 <sup>a</sup>	143	173	185	226 <sup>a</sup>	126	191	162	128

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.  
NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section 3.3.5 in Chapter 3 of the *2019 National Survey on Drug Use and Health: Methodological Summary and Definitions*).  
<sup>a</sup> The difference between this estimate and the 2019 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

Definitions: Measures and terms are defined in Appendix A.  
Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2019.

Table 7.24B Daily or Almost Daily Use of Specific Substances in Past Year among All Persons Aged 12 to 20 and among Past Year Users of Specific Substances Aged 12 to 20: Percentages, 2002-2019

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
DAILY/ALMOST DAILY USE IN PAST YEAR AMONG TOTAL POPULATION																		
Marijuana	2.6	2.4	2.3 <sup>a</sup>	2.5	2.5	2.2 <sup>a</sup>	2.8	2.6	2.7	2.7	3.0	2.9	2.8	2.9	2.6	3.0	2.8	2.8
Cocaine	0.0 <sup>a</sup>	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Crack	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	0.0	0.0	*	*	0.0	*	*	*	0.0
Heroin	0.0	0.0	0.0	0.0	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.0	0.0	0.0	0.0	*
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.0	0.0	0.1	0.0	0.0
Methamphetamine	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.0	0.0	0.0	0.0	0.0
Alcohol	0.8 <sup>a</sup>	0.8 <sup>a</sup>	0.6 <sup>a</sup>	0.6 <sup>a</sup>	0.6 <sup>a</sup>	0.7 <sup>a</sup>	0.6 <sup>a</sup>	0.5 <sup>a</sup>	0.6 <sup>a</sup>	0.5 <sup>a</sup>	0.4	0.5	0.5	0.6 <sup>a</sup>	0.3	0.5 <sup>a</sup>	0.4	0.3
DAILY/ALMOST DAILY USE IN PAST YEAR AMONG PAST YEAR USERS																		
Marijuana	12.1	11.4 <sup>a</sup>	11.5 <sup>a</sup>	13.0	12.8	12.0 <sup>a</sup>	14.3	12.8	13.1	12.7	14.4	14.1	13.9	14.6	13.0	15.1	14.0	13.8
Cocaine	1.4	1.6	0.8	1.1	1.0	1.2	0.9	1.5	0.5	1.3	1.9	0.6	0.7	1.3	1.0	1.1	0.4	1.1
Crack	2.7	*	*	*	0.9	*	*	*	*	*	*	*	*	*	*	*	*	*
Heroin	2.3	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.3	0.2	0.3	0.4	0.1
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.5	1.7	3.4 <sup>a</sup>	0.6	0.5
Methamphetamine	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	*	*	*	*
Alcohol	1.7 <sup>a</sup>	1.8 <sup>a</sup>	1.3	1.3	1.4	1.5 <sup>a</sup>	1.4 <sup>a</sup>	1.2	1.3	1.2	0.9	1.2	1.3	1.7 <sup>a</sup>	1.0	1.5	1.3	1.0

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.  
NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section 3.3.5 in Chapter 3 of the 2019 National Survey on Drug Use and Health: Methodological Summary and Definitions).  
<sup>a</sup> The difference between this estimate and the 2019 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

Definitions: Measures and terms are defined in Appendix A.  
Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2019.

Table 7.25A Daily or Almost Daily Use of Specific Substances in Past Year among All Persons Aged 21 or Older and among Past Year Users of Specific Substances Aged 21 or Older: Numbers in Thousands, 2002-2019

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
DAILY/ALMOST DAILY USE IN PAST YEAR AMONG TOTAL POPULATION																		
Marijuana	2,153 <sup>a</sup>	2,191 <sup>a</sup>	2,354 <sup>a</sup>	2,405 <sup>a</sup>	2,168 <sup>a</sup>	2,698 <sup>a</sup>	2,819 <sup>a</sup>	3,059 <sup>a</sup>	3,566 <sup>a</sup>	3,929 <sup>a</sup>	4,193 <sup>a</sup>	4,621 <sup>a</sup>	5,431 <sup>a</sup>	5,431 <sup>a</sup>	6,008 <sup>a</sup>	6,991 <sup>a</sup>	7,579 <sup>a</sup>	9,020
Cocaine	102	64	129	178	171	139	72	38	64	33	98	148	62	82	67	118	143	103
Crack	42	15	89	86	33	73	17	24	24	16	19	67	45	56	24	68	40	26
Heroin	57 <sup>a</sup>	22 <sup>a</sup>	53 <sup>a</sup>	14 <sup>a</sup>	149	35 <sup>a</sup>	65	43 <sup>a</sup>	46 <sup>a</sup>	89	138	89	117	103	181	154	127	146
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	23	19	44	23	56
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	25	*	*	20	17
Methamphetamine	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	124 <sup>a</sup>	152 <sup>a</sup>	184	222	303
Alcohol	9,380 <sup>a</sup>	9,320 <sup>a</sup>	9,514 <sup>a</sup>	9,932 <sup>a</sup>	10,078 <sup>a</sup>	9,814 <sup>a</sup>	10,409	10,883	9,813 <sup>a</sup>	10,600	10,687	10,414	11,054	11,208	11,437	11,702	10,816	11,475
DAILY/ALMOST DAILY USE IN PAST YEAR AMONG PAST YEAR USERS																		
Marijuana	2,153 <sup>a</sup>	2,191 <sup>a</sup>	2,354 <sup>a</sup>	2,405 <sup>a</sup>	2,168 <sup>a</sup>	2,698 <sup>a</sup>	2,819 <sup>a</sup>	3,059 <sup>a</sup>	3,566 <sup>a</sup>	3,929 <sup>a</sup>	4,193 <sup>a</sup>	4,621 <sup>a</sup>	5,431 <sup>a</sup>	5,431 <sup>a</sup>	6,008 <sup>a</sup>	6,991 <sup>a</sup>	7,579 <sup>a</sup>	9,020
Cocaine	102	64	129	178	171	139	72	38	64	33	98	148	62	82	67	118	143	103
Crack	42	15	89	86	33	73	17	24	*	*	*	*	45	56	24	*	40	26
Heroin	*	*	*	14 <sup>a</sup>	*	*	*	*	46 <sup>a</sup>	*	*	89	117	103	181	154	127	146
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	23	19	44	23	56
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	1	9	*	17
Methamphetamine	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	124 <sup>a</sup>	152 <sup>a</sup>	184	222	303
Alcohol	9,380 <sup>a</sup>	9,320 <sup>a</sup>	9,514 <sup>a</sup>	9,932 <sup>a</sup>	10,078 <sup>a</sup>	9,814 <sup>a</sup>	10,409	10,883	9,813 <sup>a</sup>	10,600	10,687	10,414	11,054	11,208	11,437	11,702	10,816	11,475

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.  
NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section 3.3.5 in Chapter 3 of the 2019 National Survey on Drug Use and Health: Methodological Summary and Definitions).  
<sup>a</sup> The difference between this estimate and the 2019 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

Definitions: Measures and terms are defined in Appendix A.  
Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2019.

Table 7.25B Daily or Almost Daily Use of Specific Substances in Past Year among All Persons Aged 21 or Older and among Past Year Users of Specific Substances Aged 21 or Older: Percentages, 2002-2019

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
DAILY/ALMOST DAILY USE IN PAST YEAR AMONG TOTAL POPULATION																		
Marijuana	1.1 <sup>a</sup>	1.1 <sup>a</sup>	1.2 <sup>a</sup>	1.2 <sup>a</sup>	1.0 <sup>a</sup>	1.3 <sup>a</sup>	1.3 <sup>a</sup>	1.4 <sup>a</sup>	1.7 <sup>a</sup>	1.8 <sup>a</sup>	1.9 <sup>a</sup>	2.1 <sup>a</sup>	2.4 <sup>a</sup>	2.4 <sup>a</sup>	2.6 <sup>a</sup>	3.0 <sup>a</sup>	3.2 <sup>a</sup>	3.8
Cocaine	0.1	0.0	0.1	0.1	0.1	0.1	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.1	0.1	0.0
Crack	0.0	0.0	0.0 <sup>a</sup>	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Heroin	0.0	0.0 <sup>a</sup>	0.0	0.0 <sup>a</sup>	0.1	0.0 <sup>a</sup>	0.0	0.0 <sup>a</sup>	0.0 <sup>a</sup>	0.0	0.1	0.0	0.1	0.0	0.1	0.1	0.1	0.1
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.0	0.0	0.0	0.0	0.0
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.0	*	*	0.0	0.0
Methamphetamine	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.1 <sup>a</sup>	0.1 <sup>a</sup>	0.1	0.1	0.1
Alcohol	4.7	4.7	4.7	4.8	4.8	4.7	4.9	5.1	4.6	4.8	4.8	4.6	4.9	4.9	4.9	5.0	4.6	4.8
DAILY/ALMOST DAILY USE IN PAST YEAR AMONG PAST YEAR USERS																		
Marijuana	12.2 <sup>a</sup>	12.6 <sup>a</sup>	13.3 <sup>a</sup>	13.5 <sup>a</sup>	12.0 <sup>a</sup>	15.0 <sup>a</sup>	15.2 <sup>a</sup>	14.7 <sup>a</sup>	16.7 <sup>a</sup>	18.3 <sup>a</sup>	17.9 <sup>a</sup>	18.4 <sup>a</sup>	19.8 <sup>a</sup>	19.0 <sup>a</sup>	19.9 <sup>a</sup>	20.9	21.2	22.2
Cocaine	2.2	1.4	2.9	4.2	3.7	3.1	1.7	1.0	1.7	1.1	2.5	4.1	1.6	2.0	1.5	2.3	3.0	2.1
Crack	3.1	1.3	7.9	7.3	2.6	5.7	1.7	2.6	*	*	*	*	6.3	7.0	2.8	*	5.5	3.5
Heroin	*	*	*	5.2 <sup>a</sup>	*	*	*	*	9.3	*	*	14.9	14.4	13.6	20.5	18.4	16.7	20.0
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.7	0.6	1.2	0.5	1.2
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	0.1	0.9	*	1.5
Methamphetamine	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	7.9 <sup>a</sup>	12.0	12.4	12.9	16.1
Alcohol	6.8	6.8	6.8	6.9	7.0	6.7	7.0	7.2	6.5	6.9	6.8	6.5	6.8	6.9	7.1	7.1	6.5	6.9

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.  
NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section 3.3.5 in Chapter 3 of the *2019 National Survey on Drug Use and Health: Methodological Summary and Definitions*).  
<sup>a</sup> The difference between this estimate and the 2019 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

Definitions: Measures and terms are defined in Appendix A.  
Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2019.

**Table 7.26A    Daily or Almost Daily Use of Specific Substances in Past Month among All Persons Aged 12 or Older and among Past Month Users of Specific Substances Aged 12 or Older: Numbers in Thousands, 2002-2019**

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
<b>DAILY/ALMOST DAILY USE IN PAST MONTH AMONG TOTAL POPULATION</b>																		
Marijuana	4,756 <sup>a</sup>	4,894 <sup>a</sup>	4,880 <sup>a</sup>	5,089 <sup>a</sup>	5,104 <sup>a</sup>	5,099 <sup>a</sup>	5,454 <sup>a</sup>	6,181 <sup>a</sup>	6,935 <sup>a</sup>	7,074 <sup>a</sup>	7,601 <sup>a</sup>	8,144 <sup>a</sup>	9,240 <sup>a</sup>	9,240 <sup>a</sup>	9,995 <sup>a</sup>	10,829 <sup>a</sup>	11,966 <sup>a</sup>	13,830
Cocaine	113	133	263	209	343 <sup>a</sup>	176	168	157	178	69 <sup>a</sup>	129	185	116	113	187	137	125	175
Crack	42	76	118	75	121	99	66	90	55	20 <sup>a</sup>	73	95	58	52	96	59	53	95
Heroin	98	57 <sup>a</sup>	78 <sup>a</sup>	46 <sup>a</sup>	188	81 <sup>a</sup>	93	48 <sup>a</sup>	79 <sup>a</sup>	183	180	145	178	141	220	203	132	181
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	32	15	32	48 <sup>a</sup>	15
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	26	31	34	42	52
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	246 <sup>a</sup>	249 <sup>a</sup>	308	398	439
Misuse of Pain Relievers <sup>1</sup>	--	--	--	--	--	--	--	--	--	--	--	--	--	431	348	362	369	346
Misuse of Stimulants	--	--	--	--	--	--	--	--	--	--	--	--	--	89	91	122	76	104
Misuse of Tranquilizers	--	--	--	--	--	--	--	--	--	--	--	--	--	114	120	91	104	127
Misuse of Sedatives	--	--	--	--	--	--	--	--	--	--	--	--	--	29	39	39	*	26
Cigarettes (Daily Only) <sup>2</sup>	38,738 <sup>a</sup>	37,989 <sup>a</sup>	37,343 <sup>a</sup>	38,131 <sup>a</sup>	38,331 <sup>a</sup>	36,901 <sup>a</sup>	36,788 <sup>a</sup>	35,834 <sup>a</sup>	34,718 <sup>a</sup>	34,488 <sup>a</sup>	34,936 <sup>a</sup>	33,247 <sup>a</sup>	32,471 <sup>a</sup>	30,190 <sup>a</sup>	29,701 <sup>a</sup>	27,787	27,328	26,796
Smokeless Tobacco	--	--	--	--	--	--	--	--	--	--	--	--	--	5,399	5,336	5,149	4,890	4,975
Cigars	1,339 <sup>a</sup>	1,732 <sup>a</sup>	1,648 <sup>a</sup>	1,623 <sup>a</sup>	1,868 <sup>a</sup>	1,699 <sup>a</sup>	1,874 <sup>a</sup>	1,972	2,147	1,811 <sup>a</sup>	2,153	1,898	1,725 <sup>a</sup>	2,259	2,401	2,337	2,410	2,288
Alcohol	19,745 <sup>a</sup>	19,053 <sup>a</sup>	19,487 <sup>a</sup>	20,200	19,893	20,051	21,173	21,896	20,357	20,632	21,544	21,262	22,593	21,422	20,962	21,852	21,678	21,376
Binge Alcohol <sup>3</sup>	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	2,856	2,666	2,918	2,911	2,662
<b>DAILY/ALMOST DAILY USE IN PAST MONTH AMONG PAST MONTH USERS</b>																		
Marijuana	4,756 <sup>a</sup>	4,894 <sup>a</sup>	4,880 <sup>a</sup>	5,089 <sup>a</sup>	5,104 <sup>a</sup>	5,099 <sup>a</sup>	5,454 <sup>a</sup>	6,181 <sup>a</sup>	6,935 <sup>a</sup>	7,074 <sup>a</sup>	7,601 <sup>a</sup>	8,144 <sup>a</sup>	9,240 <sup>a</sup>	9,240 <sup>a</sup>	9,995 <sup>a</sup>	10,829 <sup>a</sup>	11,966 <sup>a</sup>	13,830
Cocaine	113	133	263	209	343 <sup>a</sup>	176	168	157	178	69 <sup>a</sup>	129	185	116	113	187	137	125	175
Crack	42	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Heroin	*	*	*	*	*	*	*	*	*	*	*	*	178	*	*	203	*	*
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	32	15	32	48 <sup>a</sup>	15
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	26	31	34	42	52
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	246 <sup>a</sup>	249 <sup>a</sup>	308	398	439
Misuse of Pain Relievers <sup>1</sup>	--	--	--	--	--	--	--	--	--	--	--	--	--	431	348	362	369	346
Misuse of Stimulants	--	--	--	--	--	--	--	--	--	--	--	--	--	89	91	122	76	104
Misuse of Tranquilizers	--	--	--	--	--	--	--	--	--	--	--	--	--	114	120	91	104	127
Misuse of Sedatives	--	--	--	--	--	--	--	--	--	--	--	--	--	*	*	*	*	*
Cigarettes (Daily Only) <sup>2</sup>	38,738 <sup>a</sup>	37,989 <sup>a</sup>	37,343 <sup>a</sup>	38,131 <sup>a</sup>	38,331 <sup>a</sup>	36,901 <sup>a</sup>	36,788 <sup>a</sup>	35,834 <sup>a</sup>	34,718 <sup>a</sup>	34,488 <sup>a</sup>	34,936 <sup>a</sup>	33,247 <sup>a</sup>	32,471 <sup>a</sup>	30,190 <sup>a</sup>	29,701 <sup>a</sup>	27,787	27,328	26,796
Smokeless Tobacco	--	--	--	--	--	--	--	--	--	--	--	--	--	5,399	5,336	5,149	4,890	4,975
Cigars	1,339 <sup>a</sup>	1,732 <sup>a</sup>	1,648 <sup>a</sup>	1,623 <sup>a</sup>	1,868 <sup>a</sup>	1,699 <sup>a</sup>	1,874 <sup>a</sup>	1,972	2,147	1,811 <sup>a</sup>	2,153	1,898	1,725 <sup>a</sup>	2,259	2,401	2,337	2,410	2,288
Alcohol	19,745	19,053 <sup>a</sup>	19,487 <sup>a</sup>	20,200	19,893	20,051	21,173	21,896	20,357	20,632	21,544	21,262	22,593	21,422	20,962	21,852	21,678	21,376
Binge Alcohol <sup>3</sup>	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	2,856	2,666	2,918	2,911	2,662
<b>SMOKED 1+ PACKS OF CIGARETTES PER DAY AMONG DAILY CIGARETTE USERS<sup>4</sup></b>	20,520 <sup>a</sup>	20,284 <sup>a</sup>	20,160 <sup>a</sup>	19,573 <sup>a</sup>	19,394 <sup>a</sup>	18,752 <sup>a</sup>	18,066 <sup>a</sup>	16,433 <sup>a</sup>	15,619 <sup>a</sup>	15,093 <sup>a</sup>	14,650 <sup>a</sup>	13,721 <sup>a</sup>	13,058 <sup>a</sup>	12,380 <sup>a</sup>	12,184 <sup>a</sup>	11,414	10,799	10,849

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.  
NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section 3.3.5 in Chapter 3 of the *2019 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

<sup>a</sup> The difference between this estimate and the 2019 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

<sup>1</sup> Prescription psychotherapeutic subtypes were revised in 2016; one effect was the comparability of codeine products between 2015 and 2016.

<sup>2</sup> Includes only respondents who smoked cigarettes on each of the past 30 days.

<sup>3</sup> Estimates of binge alcohol use include use by those who were heavy alcohol users.

<sup>4</sup> Respondents with missing data for number of cigarettes smoked per day were excluded from the analysis.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2019.

**Table 7.26B    Daily or Almost Daily Use of Specific Substances in Past Month among All Persons Aged 12 or Older and among Past Month Users of Specific Substances Aged 12 or Older: Percentages, 2002-2019**

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
<b>DAILY/ALMOST DAILY USE IN PAST MONTH AMONG TOTAL POPULATION</b>																		
Marijuana	2.0 <sup>a</sup>	2.1 <sup>a</sup>	2.0 <sup>a</sup>	2.1 <sup>a</sup>	2.1 <sup>a</sup>	2.1 <sup>a</sup>	2.2 <sup>a</sup>	2.5 <sup>a</sup>	2.7 <sup>a</sup>	2.7 <sup>a</sup>	2.9 <sup>a</sup>	3.1 <sup>a</sup>	3.5 <sup>a</sup>	3.5 <sup>a</sup>	3.7 <sup>a</sup>	4.0 <sup>a</sup>	4.4 <sup>a</sup>	5.0
Cocaine	0.0	0.1	0.1	0.1	0.1 <sup>a</sup>	0.1	0.1	0.1	0.1	0.0 <sup>a</sup>	0.0	0.1	0.0	0.0	0.1	0.1	0.0	0.1
Crack	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0 <sup>a</sup>	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Heroin	0.0	0.0 <sup>a</sup>	0.0	0.0 <sup>a</sup>	0.1	0.0	0.0	0.0 <sup>a</sup>	0.0 <sup>a</sup>	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.0	0.1
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.0	0.0	0.0	0.0 <sup>a</sup>	0.0
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.0	0.0	0.0	0.0	0.0
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	0.1 <sup>a</sup>	0.1 <sup>a</sup>	0.1	0.1	0.2
Misuse of Pain Relievers <sup>1</sup>	--	--	--	--	--	--	--	--	--	--	--	--	--	0.2	0.1	0.1	0.1	0.1
Misuse of Stimulants	--	--	--	--	--	--	--	--	--	--	--	--	--	0.0	0.0	0.0	0.0	0.0
Misuse of Tranquilizers	--	--	--	--	--	--	--	--	--	--	--	--	--	0.0	0.0	0.0	0.0	0.0
Misuse of Sedatives	--	--	--	--	--	--	--	--	--	--	--	--	--	0.0	0.0	0.0	*	0.0
Cigarettes (Daily Only) <sup>2</sup>	16.5 <sup>a</sup>	16.0 <sup>a</sup>	15.5 <sup>a</sup>	15.7 <sup>a</sup>	15.6 <sup>a</sup>	14.9 <sup>a</sup>	14.7 <sup>a</sup>	14.2 <sup>a</sup>	13.7 <sup>a</sup>	13.4 <sup>a</sup>	13.4 <sup>a</sup>	12.7 <sup>a</sup>	12.2 <sup>a</sup>	11.3 <sup>a</sup>	11.0 <sup>a</sup>	10.2	10.0	9.7
Smokeless Tobacco	--	--	--	--	--	--	--	--	--	--	--	--	--	2.0	2.0	1.9	1.8	1.8
Cigars	0.6 <sup>a</sup>	0.7	0.7	0.7 <sup>a</sup>	0.8	0.7	0.7	0.8	0.8	0.7	0.8	0.7	0.7 <sup>a</sup>	0.8	0.9	0.9	0.9	0.8
Alcohol	8.4 <sup>a</sup>	8.0	8.1	8.3	8.1	8.1	8.5 <sup>a</sup>	8.7 <sup>a</sup>	8.0	8.0	8.3	8.1	8.5 <sup>a</sup>	8.0	7.8	8.0	7.9	7.8
Binge Alcohol <sup>3</sup>	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	1.1	1.0	1.1	1.1	1.0
<b>DAILY/ALMOST DAILY USE IN PAST MONTH AMONG PAST MONTH USERS</b>																		
Marijuana	32.6 <sup>a</sup>	33.4 <sup>a</sup>	33.5 <sup>a</sup>	34.8 <sup>a</sup>	34.4 <sup>a</sup>	35.2 <sup>a</sup>	35.7 <sup>a</sup>	36.7 <sup>a</sup>	39.8 <sup>a</sup>	39.1 <sup>a</sup>	40.3 <sup>a</sup>	41.1	41.6	41.6	41.7	41.7	43.3	43.8
Cocaine	5.6	5.8	13.0	8.7	14.2	8.5	9.0	9.6	12.1	5.1	7.8	11.9	7.6	6.0	10.0	6.3	6.4	8.8
Crack	7.4	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Heroin	*	*	*	*	*	*	*	*	*	*	*	*	40.9	*	*	41.1	*	*
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	2.6	1.1	2.2	3.0 <sup>a</sup>	0.8
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	4.9	5.2	6.1	6.9	6.4
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	27.4	37.3	39.7	39.8	37.4
Misuse of Pain Relievers <sup>1</sup>	--	--	--	--	--	--	--	--	--	--	--	--	--	11.4	10.4	11.2	12.9	12.3
Misuse of Stimulants	--	--	--	--	--	--	--	--	--	--	--	--	--	5.4	5.2	6.6	4.5	6.7
Misuse of Tranquilizers	--	--	--	--	--	--	--	--	--	--	--	--	--	6.1	6.2	5.3	6.4	7.1
Misuse of Sedatives	--	--	--	--	--	--	--	--	--	--	--	--	--	*	*	*	*	*
Cigarettes (Daily Only) <sup>2</sup>	63.4 <sup>a</sup>	62.9 <sup>a</sup>	62.3 <sup>a</sup>	63.0 <sup>a</sup>	62.3 <sup>a</sup>	61.3 <sup>a</sup>	61.5 <sup>a</sup>	61.0 <sup>a</sup>	59.5	60.7 <sup>a</sup>	60.7 <sup>a</sup>	59.6	58.8	58.1	57.9	57.1	58.2	58.4
Smokeless Tobacco	--	--	--	--	--	--	--	--	--	--	--	--	--	60.0	60.9	60.0	61.3	58.4
Cigars	10.5 <sup>a</sup>	13.5 <sup>a</sup>	12.0 <sup>a</sup>	11.9 <sup>a</sup>	13.6 <sup>a</sup>	12.8 <sup>a</sup>	14.2 <sup>a</sup>	14.8 <sup>a</sup>	16.2 <sup>a</sup>	14.1 <sup>a</sup>	16.0 <sup>a</sup>	15.3 <sup>a</sup>	14.4 <sup>a</sup>	18.1	19.6	18.6	19.7	19.6
Alcohol	16.5 <sup>a</sup>	16.0	16.1	16.0	15.9	15.8	16.4 <sup>a</sup>	16.8 <sup>a</sup>	15.5	15.5	15.9	15.5	16.2	15.5	15.3	15.5	15.5	15.3
Binge Alcohol <sup>3</sup>	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	4.3	4.1	4.4	4.3	4.0
<b>SMOKED 1+ PACKS OF CIGARETTES PER DAY AMONG DAILY CIGARETTE USERS<sup>4</sup></b>																		
	53.1 <sup>a</sup>	53.5 <sup>a</sup>	54.0 <sup>a</sup>	51.4 <sup>a</sup>	50.6 <sup>a</sup>	50.9 <sup>a</sup>	49.2 <sup>a</sup>	45.9 <sup>a</sup>	45.1 <sup>a</sup>	43.8 <sup>a</sup>	42.0	41.3	40.3	41.1	41.1	41.2	39.6	40.5

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.  
NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section 3.3.5 in Chapter 3 of the *2019 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

<sup>a</sup> The difference between this estimate and the 2019 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

<sup>1</sup> Prescription psychotherapeutic subtypes were revised in 2016; one effect was the comparability of codeine products between 2015 and 2016.

<sup>2</sup> Includes only respondents who smoked cigarettes on each of the past 30 days.

<sup>3</sup> Estimates of binge alcohol use include use by those who were heavy alcohol users.

<sup>4</sup> Respondents with missing data for number of cigarettes smoked per day were excluded from the analysis.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2019.



**Table 7.27A    Daily or Almost Daily Use of Specific Substances in Past Month among All Persons Aged 12 to 17 and among Past Month Users of Specific Substances Aged 12 to 17: Numbers in Thousands, 2002-2019**

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
<b>DAILY/ALMOST DAILY USE IN PAST MONTH AMONG TOTAL POPULATION</b>																		
Marijuana	603 <sup>a</sup>	482	536 <sup>a</sup>	400	434	427	421	452	439	466	425	451	400	421	367	405	367	447
Cocaine	7	8	11	10	*	10	11	6	2	*	2	*	8	2	3	*	2	*
Crack	6	6	3	*	*	4	*	2	*	*	*	*	*	*	*	*	*	*
Heroin	4	3	6	*	*	*	2	2	*	12	*	*	5	*	*	*	4	*
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	2	3	2	3	*
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	6	12	10	4 <sup>a</sup>	22
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	*	2	5	4	2
Misuse of Pain Relievers <sup>1</sup>	--	--	--	--	--	--	--	--	--	--	--	--	--	7	8	10	*	3
Misuse of Stimulants	--	--	--	--	--	--	--	--	--	--	--	--	--	5	8	6	*	9
Misuse of Tranquilizers	--	--	--	--	--	--	--	--	--	--	--	--	--	12	6	7	10	2
Misuse of Sedatives	--	--	--	--	--	--	--	--	--	--	--	--	--	*	*	*	*	*
Cigarettes (Daily Only) <sup>2</sup>	1,021 <sup>a</sup>	904 <sup>a</sup>	828 <sup>a</sup>	704 <sup>a</sup>	701 <sup>a</sup>	659 <sup>a</sup>	510 <sup>a</sup>	511 <sup>a</sup>	459 <sup>a</sup>	440 <sup>a</sup>	360 <sup>a</sup>	271 <sup>a</sup>	292 <sup>a</sup>	208 <sup>a</sup>	129 <sup>a</sup>	96	99	75
Smokeless Tobacco	--	--	--	--	--	--	--	--	--	--	--	--	--	114 <sup>a</sup>	112 <sup>a</sup>	86	79	66
Cigars	78	59	76	57	58	44	52	62	48	45	31	27	36	32	49	36	35	47
Alcohol	165 <sup>a</sup>	137 <sup>a</sup>	163 <sup>a</sup>	136 <sup>a</sup>	104 <sup>a</sup>	121 <sup>a</sup>	83 <sup>a</sup>	108 <sup>a</sup>	87 <sup>a</sup>	77	69	55	55	55	34	37	27	48
Binge Alcohol <sup>3</sup>	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	12	5	14	12	16
<b>DAILY/ALMOST DAILY USE IN PAST MONTH AMONG PAST MONTH USERS</b>																		
Marijuana	603 <sup>a</sup>	482	536	400	434	427	421	452	439	466	425	451	400	421	367	405	367	447
Cocaine	7	8	*	10	1	*	*	*	*	*	*	*	*	*	*	*	*	*
Crack	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Heroin	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	*	*	*	*
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	*	*	4 <sup>a</sup>	22
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	*	*	*	*	*
Misuse of Pain Relievers <sup>1</sup>	--	--	--	--	--	--	--	--	--	--	--	--	--	7	8	10	1	*
Misuse of Stimulants	--	--	--	--	--	--	--	--	--	--	--	--	--	*	*	*	*	*
Misuse of Tranquilizers	--	--	--	--	--	--	--	--	--	--	--	--	--	*	*	*	*	*
Misuse of Sedatives	--	--	--	--	--	--	--	--	--	--	--	--	--	*	*	*	*	*
Cigarettes (Daily Only) <sup>2</sup>	1,021 <sup>a</sup>	904 <sup>a</sup>	828 <sup>a</sup>	704 <sup>a</sup>	701 <sup>a</sup>	659 <sup>a</sup>	510 <sup>a</sup>	511 <sup>a</sup>	459 <sup>a</sup>	440 <sup>a</sup>	360 <sup>a</sup>	271 <sup>a</sup>	292 <sup>a</sup>	208 <sup>a</sup>	129 <sup>a</sup>	96	99	75
Smokeless Tobacco	--	--	--	--	--	--	--	--	--	--	--	--	--	114 <sup>a</sup>	112 <sup>a</sup>	86	79	66
Cigars	78	59	76	57	58	44	52	62	48	45	31	27	36	32	49	36	35	47
Alcohol	165 <sup>a</sup>	137 <sup>a</sup>	163 <sup>a</sup>	136 <sup>a</sup>	104 <sup>a</sup>	121 <sup>a</sup>	83 <sup>a</sup>	108 <sup>a</sup>	87 <sup>a</sup>	77	69	55	55	55	34	37	27	48
Binge Alcohol <sup>3</sup>	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	12	5	14	12	16
<b>SMOKED 1+ PACKS OF CIGARETTES PER DAY AMONG DAILY CIGARETTE USERS<sup>4</sup></b>																		
	221	199	160	142	125	123	93	91	76	65	38	32	35	16	*	*	*	*

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.  
NOTE:    Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section 3.3.5 in Chapter 3 of the *2019 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

<sup>a</sup> The difference between this estimate and the 2019 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

<sup>1</sup> Prescription psychotherapeutic subtypes were revised in 2016; one effect was the comparability of codeine products between 2015 and 2016.

<sup>2</sup> Includes only respondents who smoked cigarettes on each of the past 30 days.

<sup>3</sup> Estimates of binge alcohol use include use by those who were heavy alcohol users.

<sup>4</sup> Respondents with missing data for number of cigarettes smoked per day were excluded from the analysis.

Definitions:    Measures and terms are defined in Appendix A.

Source:    SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2019.

Table 7.27B    Daily or Almost Daily Use of Specific Substances in Past Month among All Persons Aged 12 to 17 and among Past Month Users of Specific Substances Aged 12 to 17: Percentages, 2002-2019

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
<b>DAILY/ALMOST DAILY USE IN PAST MONTH AMONG TOTAL POPULATION</b>																		
Marijuana	2.4 <sup>a</sup>	1.9	2.1	1.6	1.7	1.7	1.7	1.8	1.8	1.9	1.7	1.8	1.6	1.7	1.5	1.6	1.5	1.8
Cocaine	0.0	0.0	0.0	0.0	*	0.0	0.0	0.0	0.0	*	0.0	*	0.0	0.0	0.0	*	0.0	*
Crack	0.0	0.0	0.0	*	*	0.0	*	0.0	*	*	*	*	*	*	*	*	*	*
Heroin	0.0	0.0	0.0	*	*	*	0.0	0.0	*	0.0	*	*	0.0	*	*	*	0.0	*
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.0	0.0	0.0	0.0	*
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.0	0.0	0.0	0.0 <sup>a</sup>	0.1
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	*	0.0	0.0	0.0	0.0
Misuse of Pain Relievers <sup>1</sup>	--	--	--	--	--	--	--	--	--	--	--	--	--	0.0	0.0	0.0	*	0.0
Misuse of Stimulants	--	--	--	--	--	--	--	--	--	--	--	--	--	0.0	0.0	0.0	*	0.0
Misuse of Tranquilizers	--	--	--	--	--	--	--	--	--	--	--	--	--	0.0	0.0	0.0	0.0	0.0
Misuse of Sedatives	--	--	--	--	--	--	--	--	--	--	--	--	--	*	*	*	*	*
Cigarettes (Daily Only) <sup>2</sup>	4.1 <sup>a</sup>	3.6 <sup>a</sup>	3.3 <sup>a</sup>	2.8 <sup>a</sup>	2.8 <sup>a</sup>	2.6 <sup>a</sup>	2.0 <sup>a</sup>	2.1 <sup>a</sup>	1.9 <sup>a</sup>	1.8 <sup>a</sup>	1.4 <sup>a</sup>	1.1 <sup>a</sup>	1.2 <sup>a</sup>	0.8 <sup>a</sup>	0.5 <sup>a</sup>	0.4	0.4	0.3
Smokeless Tobacco	--	--	--	--	--	--	--	--	--	--	--	--	--	0.5 <sup>a</sup>	0.5 <sup>a</sup>	0.3	0.3	0.3
Cigars	0.3	0.2	0.3	0.2	0.2	0.2	0.2	0.3	0.2	0.2	0.1	0.1	0.1	0.1	0.2	0.1	0.1	0.2
Alcohol	0.7 <sup>a</sup>	0.5 <sup>a</sup>	0.6 <sup>a</sup>	0.5 <sup>a</sup>	0.4 <sup>a</sup>	0.5 <sup>a</sup>	0.3 <sup>a</sup>	0.4 <sup>a</sup>	0.4 <sup>a</sup>	0.3	0.3	0.2	0.2	0.2	0.1	0.1	0.1	0.2
Binge Alcohol <sup>3</sup>	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.0	0.0	0.1	0.0	0.1
<b>DAILY/ALMOST DAILY USE IN PAST MONTH AMONG PAST MONTH USERS</b>																		
Marijuana	29.8 <sup>a</sup>	24.4	28.1	23.1	25.6	25.2	25.3	24.9	24.4	23.7	23.6	25.6	21.8	24.0	22.8	25.1	22.1	24.3
Cocaine	4.6	5.3	*	6.9	0.9	*	*	*	*	*	*	*	*	*	*	*	*	*
Crack	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Heroin	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	*	*	*	*
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	*	*	2.3 <sup>a</sup>	9.9
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	*	*	*	*	*
Misuse of Pain Relievers <sup>1</sup>	--	--	--	--	--	--	--	--	--	--	--	--	--	2.7	3.2	4.4	0.6	*
Misuse of Stimulants	--	--	--	--	--	--	--	--	--	--	--	--	--	*	*	*	*	*
Misuse of Tranquilizers	--	--	--	--	--	--	--	--	--	--	--	--	--	*	*	*	*	*
Misuse of Sedatives	--	--	--	--	--	--	--	--	--	--	--	--	--	*	*	*	*	*
Cigarettes (Daily Only) <sup>2</sup>	31.8 <sup>a</sup>	29.7 <sup>a</sup>	27.6 <sup>a</sup>	25.8 <sup>a</sup>	26.5 <sup>a</sup>	26.4 <sup>a</sup>	22.3 <sup>a</sup>	23.0 <sup>a</sup>	22.5 <sup>a</sup>	22.7 <sup>a</sup>	22.0 <sup>a</sup>	19.4 <sup>a</sup>	24.1 <sup>a</sup>	20.0 <sup>a</sup>	15.0	12.2	14.7	13.2
Smokeless Tobacco	--	--	--	--	--	--	--	--	--	--	--	--	--	31.0	31.8	25.6	28.4	26.5
Cigars	7.0	5.3 <sup>a</sup>	6.3 <sup>a</sup>	5.4 <sup>a</sup>	5.6 <sup>a</sup>	4.1 <sup>a</sup>	5.5 <sup>a</sup>	6.3 <sup>a</sup>	6.1 <sup>a</sup>	5.3 <sup>a</sup>	4.8 <sup>a</sup>	4.8 <sup>a</sup>	6.9	6.3 <sup>a</sup>	10.9	7.5	8.5	13.5
Alcohol	3.8 <sup>a</sup>	3.1	3.7 <sup>a</sup>	3.3	2.5	3.0	2.3	3.0	2.6	2.3	2.2	1.9	1.9	2.3	1.5	1.5	1.2	2.0
Binge Alcohol <sup>3</sup>	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.8	0.4	1.1	1.0	1.4
<b>SMOKED 1+ PACKS OF CIGARETTES PER DAY AMONG DAILY CIGARETTE USERS<sup>4</sup></b>																		
	21.8	22.0	19.4	20.1	17.9	18.7	18.4	17.9	16.7	14.8	10.8	11.9	11.9	7.8	*	*	*	*

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.  
NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section 3.3.5 in Chapter 3 of the *2019 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

<sup>a</sup> The difference between this estimate and the 2019 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

<sup>1</sup> Prescription psychotherapeutic subtypes were revised in 2016; one effect was the comparability of codeine products between 2015 and 2016.

<sup>2</sup> Includes only respondents who smoked cigarettes on each of the past 30 days.

<sup>3</sup> Estimates of binge alcohol use include use by those who were heavy alcohol users.

<sup>4</sup> Respondents with missing data for number of cigarettes smoked per day were excluded from the analysis.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2019.

**Table 7.28A    Daily or Almost Daily Use of Specific Substances in Past Month among All Persons Aged 18 or Older and among Past Month Users of Specific Substances Aged 18 or Older: Numbers in Thousands, 2002-2019**

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
<b>DAILY/ALMOST DAILY USE IN PAST MONTH AMONG TOTAL POPULATION</b>																		
Marijuana	4,153 <sup>a</sup>	4,412 <sup>a</sup>	4,344 <sup>a</sup>	4,689 <sup>a</sup>	4,670 <sup>a</sup>	4,672 <sup>a</sup>	5,033 <sup>a</sup>	5,729 <sup>a</sup>	6,496 <sup>a</sup>	6,608 <sup>a</sup>	7,176 <sup>a</sup>	7,693 <sup>a</sup>	8,840 <sup>a</sup>	8,820 <sup>a</sup>	9,628 <sup>a</sup>	10,424 <sup>a</sup>	11,599 <sup>a</sup>	13,383
Cocaine	106	124	251	200	342 <sup>a</sup>	166	157	151	176	69 <sup>a</sup>	127	184	108	111	184	137	123	175
Crack	36	70	114	75	120	95	66	88	55	20 <sup>a</sup>	73	94	57	52	96	59	53	95
Heroin	94	54 <sup>a</sup>	72 <sup>a</sup>	46 <sup>a</sup>	187	80 <sup>a</sup>	91	46 <sup>a</sup>	79 <sup>a</sup>	172	180	144	173	141	219	203	128	181
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	30	*	30	45	15
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	20	20	23	38	30
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	246 <sup>a</sup>	247 <sup>a</sup>	302	394	437
Misuse of Pain Relievers <sup>1</sup>	--	--	--	--	--	--	--	--	--	--	--	--	--	423	341	352	368	343
Misuse of Stimulants	--	--	--	--	--	--	--	--	--	--	--	--	--	84	83	116	75	95
Misuse of Tranquilizers	--	--	--	--	--	--	--	--	--	--	--	--	--	102	114	85	94	125
Misuse of Sedatives	--	--	--	--	--	--	--	--	--	--	--	--	--	29	39	39	*	26
Cigarettes (Daily Only) <sup>2</sup>	37,717 <sup>a</sup>	37,085 <sup>a</sup>	36,516 <sup>a</sup>	37,428 <sup>a</sup>	37,631 <sup>a</sup>	36,242 <sup>a</sup>	36,278 <sup>a</sup>	35,323 <sup>a</sup>	34,259 <sup>a</sup>	34,049 <sup>a</sup>	34,577 <sup>a</sup>	32,976 <sup>a</sup>	32,179 <sup>a</sup>	29,982 <sup>a</sup>	29,573 <sup>a</sup>	27,691	27,229	26,721
Smokeless Tobacco	--	--	--	--	--	--	--	--	--	--	--	--	--	5,285	5,224	5,063	4,811	4,910
Cigars	1,260 <sup>a</sup>	1,672 <sup>a</sup>	1,572 <sup>a</sup>	1,566 <sup>a</sup>	1,810 <sup>a</sup>	1,655 <sup>a</sup>	1,821 <sup>a</sup>	1,910	2,098	1,765 <sup>a</sup>	2,121	1,870	1,689 <sup>a</sup>	2,226	2,352	2,301	2,376	2,241
Alcohol	19,580 <sup>a</sup>	18,916 <sup>a</sup>	19,324 <sup>a</sup>	20,063	19,789 <sup>a</sup>	19,930	21,089	21,788	20,270	20,555	21,475	21,207	22,538	21,367	20,927	21,815	21,651	21,329
Binge Alcohol <sup>3</sup>	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	2,844	2,662	2,904	2,899	2,645
<b>DAILY/ALMOST DAILY USE IN PAST MONTH AMONG PAST MONTH USERS</b>																		
Marijuana	4,153 <sup>a</sup>	4,412 <sup>a</sup>	4,344 <sup>a</sup>	4,689 <sup>a</sup>	4,670 <sup>a</sup>	4,672 <sup>a</sup>	5,033 <sup>a</sup>	5,729 <sup>a</sup>	6,496 <sup>a</sup>	6,608 <sup>a</sup>	7,176 <sup>a</sup>	7,693 <sup>a</sup>	8,840 <sup>a</sup>	8,820 <sup>a</sup>	9,628 <sup>a</sup>	10,424 <sup>a</sup>	11,599 <sup>a</sup>	13,383
Cocaine	106	124	251	200	342 <sup>a</sup>	166	157	151	176	69 <sup>a</sup>	127	184	108	111	184	137	123	175
Crack	36	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Heroin	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	203	*	*
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	30	12	30	45	15
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	20	23	*	30
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	246 <sup>a</sup>	*	302	394	437
Misuse of Pain Relievers <sup>1</sup>	--	--	--	--	--	--	--	--	--	--	--	--	--	423	341	352	368	343
Misuse of Stimulants	--	--	--	--	--	--	--	--	--	--	--	--	--	84	83	116	75	95
Misuse of Tranquilizers	--	--	--	--	--	--	--	--	--	--	--	--	--	102	114	85	94	125
Misuse of Sedatives	--	--	--	--	--	--	--	--	--	--	--	--	--	*	*	*	*	*
Cigarettes (Daily Only) <sup>2</sup>	37,717 <sup>a</sup>	37,085 <sup>a</sup>	36,516 <sup>a</sup>	37,428 <sup>a</sup>	37,631 <sup>a</sup>	36,242 <sup>a</sup>	36,278 <sup>a</sup>	35,323 <sup>a</sup>	34,259 <sup>a</sup>	34,049 <sup>a</sup>	34,577 <sup>a</sup>	32,976 <sup>a</sup>	32,179 <sup>a</sup>	29,982 <sup>a</sup>	29,573 <sup>a</sup>	27,691	27,229	26,721
Smokeless Tobacco	--	--	--	--	--	--	--	--	--	--	--	--	--	5,285	5,224	5,063	4,811	4,910
Cigars	1,260 <sup>a</sup>	1,672 <sup>a</sup>	1,572 <sup>a</sup>	1,566 <sup>a</sup>	1,810 <sup>a</sup>	1,655 <sup>a</sup>	1,821 <sup>a</sup>	1,910	2,098	1,765 <sup>a</sup>	2,121	1,870	1,689 <sup>a</sup>	2,226	2,352	2,301	2,376	2,241
Alcohol	19,580 <sup>a</sup>	18,916 <sup>a</sup>	19,324 <sup>a</sup>	20,063	19,789	19,930	21,089	21,788	20,270	20,555	21,475	21,207	22,538	21,367	20,927	21,815	21,651	21,329
Binge Alcohol <sup>3</sup>	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	2,844	2,662	2,904	2,899	2,645
<b>SMOKED 1+ PACKS OF CIGARETTES PER DAY AMONG DAILY CIGARETTE USERS<sup>4</sup></b>	20,299 <sup>a</sup>	20,085 <sup>a</sup>	20,000 <sup>a</sup>	19,431 <sup>a</sup>	19,269 <sup>a</sup>	18,630 <sup>a</sup>	17,973 <sup>a</sup>	16,342 <sup>a</sup>	15,543 <sup>a</sup>	15,029 <sup>a</sup>	14,612 <sup>a</sup>	13,689 <sup>a</sup>	13,024 <sup>a</sup>	12,364 <sup>a</sup>	12,171 <sup>a</sup>	11,402	10,790	10,840

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.  
NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section 3.3.5 in Chapter 3 of the *2019 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

<sup>a</sup> The difference between this estimate and the 2019 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

<sup>1</sup> Prescription psychotherapeutic subtypes were revised in 2016; one effect was the comparability of codeine products between 2015 and 2016.

<sup>2</sup> Includes only respondents who smoked cigarettes on each of the past 30 days.

<sup>3</sup> Estimates of binge alcohol use include use by those who were heavy alcohol users.

<sup>4</sup> Respondents with missing data for number of cigarettes smoked per day were excluded from the analysis.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2019.

Table 7.28B Daily or Almost Daily Use of Specific Substances in Past Month among All Persons Aged 18 or Older and among Past Month Users of Specific Substances Aged 18 or Older: Percentages, 2002-2019

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
<b>DAILY/ALMOST DAILY USE IN PAST MONTH AMONG TOTAL POPULATION</b>																		
Marijuana	2.0 <sup>a</sup>	2.1 <sup>a</sup>	2.0 <sup>a</sup>	2.2 <sup>a</sup>	2.1 <sup>a</sup>	2.1 <sup>a</sup>	2.2 <sup>a</sup>	2.5 <sup>a</sup>	2.8 <sup>a</sup>	2.8 <sup>a</sup>	3.1 <sup>a</sup>	3.2 <sup>a</sup>	3.7 <sup>a</sup>	3.6 <sup>a</sup>	3.9 <sup>a</sup>	4.2 <sup>a</sup>	4.7 <sup>a</sup>	5.3
Cocaine	0.1	0.1	0.1	0.1	0.2 <sup>a</sup>	0.1	0.1	0.1	0.1	0.0 <sup>a</sup>	0.1	0.1	0.0	0.0	0.1	0.1	0.0	0.1
Crack	0.0	0.0	0.1	0.0	0.1	0.0	0.0	0.0	0.0	0.0 <sup>a</sup>	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Heroin	0.0	0.0 <sup>a</sup>	0.0	0.0 <sup>a</sup>	0.1	0.0	0.0	0.0 <sup>a</sup>	0.0 <sup>a</sup>	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.0	*	0.0	0.0	0.0
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.0	0.0	0.0	0.0	0.0
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	0.1 <sup>a</sup>	0.1 <sup>a</sup>	0.1	0.2	0.2
Misuse of Pain Relievers <sup>1</sup>	--	--	--	--	--	--	--	--	--	--	--	--	--	0.2	0.1	0.1	0.1	0.1
Misuse of Stimulants	--	--	--	--	--	--	--	--	--	--	--	--	--	0.0	0.0	0.0	0.0	0.0
Misuse of Tranquilizers	--	--	--	--	--	--	--	--	--	--	--	--	--	0.0	0.0	0.0	0.0	0.0
Misuse of Sedatives	--	--	--	--	--	--	--	--	--	--	--	--	--	0.0	0.0	0.0	*	0.0
Cigarettes (Daily Only) <sup>2</sup>	17.9 <sup>a</sup>	17.4 <sup>a</sup>	17.0 <sup>a</sup>	17.2 <sup>a</sup>	17.1 <sup>a</sup>	16.3 <sup>a</sup>	16.1 <sup>a</sup>	15.5 <sup>a</sup>	14.9 <sup>a</sup>	14.6 <sup>a</sup>	14.7 <sup>a</sup>	13.9 <sup>a</sup>	13.4 <sup>a</sup>	12.3 <sup>a</sup>	12.1 <sup>a</sup>	11.2	10.9	10.7
Smokeless Tobacco	--	--	--	--	--	--	--	--	--	--	--	--	--	2.2	2.1	2.0	1.9	2.0
Cigars	0.6 <sup>a</sup>	0.8	0.7 <sup>a</sup>	0.7 <sup>a</sup>	0.8	0.7	0.8	0.8	0.9	0.8	0.9	0.8	0.7 <sup>a</sup>	0.9	1.0	0.9	1.0	0.9
Alcohol	9.3 <sup>a</sup>	8.9	9.0	9.2 <sup>a</sup>	9.0	9.0	9.4 <sup>a</sup>	9.6 <sup>a</sup>	8.8	8.8	9.1	8.9	9.4 <sup>a</sup>	8.8	8.6	8.8	8.7	8.5
Binge Alcohol <sup>3</sup>	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	1.2	1.1	1.2	1.2	1.1
<b>DAILY/ALMOST DAILY USE IN PAST MONTH AMONG PAST MONTH USERS</b>																		
Marijuana	33.1 <sup>a</sup>	34.8 <sup>a</sup>	34.3 <sup>a</sup>	36.4 <sup>a</sup>	35.5 <sup>a</sup>	36.6 <sup>a</sup>	37.0 <sup>a</sup>	38.2 <sup>a</sup>	41.6 <sup>a</sup>	41.0 <sup>a</sup>	42.1 <sup>a</sup>	42.6	43.4	43.1	43.0	42.7	44.6	45.0
Cocaine	5.7	5.9	13.3	8.8	14.8	8.4	8.9	9.6	12.5	5.3	7.8	12.2	7.3	6.1	10.0	6.4	6.4	8.9
Crack	6.7	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Heroin	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	41.3	*	*
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	2.7	0.9	2.3	3.0 <sup>a</sup>	0.9
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	4.4	5.8	*	5.2
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	27.8	*	39.9	40.1	37.6
Misuse of Pain Relievers <sup>1</sup>	--	--	--	--	--	--	--	--	--	--	--	--	--	12.1	11.0	11.6	13.7	12.9
Misuse of Stimulants	--	--	--	--	--	--	--	--	--	--	--	--	--	5.5	5.1	6.8	4.9	6.6
Misuse of Tranquilizers	--	--	--	--	--	--	--	--	--	--	--	--	--	6.0	6.2	5.3	6.0	7.5
Misuse of Sedatives	--	--	--	--	--	--	--	--	--	--	--	--	--	*	*	*	*	*
Cigarettes (Daily Only) <sup>2</sup>	65.1 <sup>a</sup>	64.6 <sup>a</sup>	64.2 <sup>a</sup>	64.8 <sup>a</sup>	63.9 <sup>a</sup>	62.8 <sup>a</sup>	63.0 <sup>a</sup>	62.5 <sup>a</sup>	60.9	62.0 <sup>a</sup>	61.9 <sup>a</sup>	60.6	59.6	58.9	58.6	57.8	58.8	59.0
Smokeless Tobacco	--	--	--	--	--	--	--	--	--	--	--	--	--	61.3	62.1	61.4	62.5	59.4
Cigars	10.8 <sup>a</sup>	14.3 <sup>a</sup>	12.6 <sup>a</sup>	12.4 <sup>a</sup>	14.3 <sup>a</sup>	13.5 <sup>a</sup>	14.8 <sup>a</sup>	15.5 <sup>a</sup>	16.8	14.7 <sup>a</sup>	16.6	15.8 <sup>a</sup>	14.8 <sup>a</sup>	18.6	19.9	19.1	20.1	19.8
Alcohol	17.0 <sup>a</sup>	16.5	16.6	16.5	16.3	16.2	16.8 <sup>a</sup>	17.2 <sup>a</sup>	15.8	15.8	16.2	15.8	16.5	15.7	15.6	15.8	15.7	15.5
Binge Alcohol <sup>3</sup>	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	4.4	4.2	4.4	4.4	4.1
<b>SMOKED 1+ PACKS OF CIGARETTES PER DAY AMONG DAILY CIGARETTE USERS<sup>4</sup></b>	54.0 <sup>a</sup>	54.3 <sup>a</sup>	54.8 <sup>a</sup>	52.0 <sup>a</sup>	51.2 <sup>a</sup>	51.5 <sup>a</sup>	49.6 <sup>a</sup>	46.4 <sup>a</sup>	45.4 <sup>a</sup>	44.1 <sup>a</sup>	42.3	41.5	40.6	41.3	41.2	41.3	39.7	40.6

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.  
NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section 3.3.5 in Chapter 3 of the *2019 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

<sup>a</sup> The difference between this estimate and the 2019 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

<sup>1</sup> Prescription psychotherapeutic subtypes were revised in 2016; one effect was the comparability of codeine products between 2015 and 2016.

<sup>2</sup> Includes only respondents who smoked cigarettes on each of the past 30 days.

<sup>3</sup> Estimates of binge alcohol use include use by those who were heavy alcohol users.

<sup>4</sup> Respondents with missing data for number of cigarettes smoked per day were excluded from the analysis.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2019.

Table 7.29A Daily or Almost Daily Use of Specific Substances in Past Month among All Persons Aged 18 to 25 and among Past Month Users of Specific Substances Aged 18 to 25: Numbers in Thousands, 2002-2019

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
<b>DAILY/ALMOST DAILY USE IN PAST MONTH AMONG TOTAL POPULATION</b>																		
Marijuana	1,997 <sup>a</sup>	2,079 <sup>a</sup>	1,995 <sup>a</sup>	2,096 <sup>a</sup>	2,133 <sup>a</sup>	2,071 <sup>a</sup>	2,178 <sup>a</sup>	2,476 <sup>a</sup>	2,658 <sup>a</sup>	2,676 <sup>a</sup>	2,882 <sup>a</sup>	2,817 <sup>a</sup>	3,020 <sup>a</sup>	3,076	3,056 <sup>a</sup>	3,367	3,190	3,334
Cocaine	23	29	49	25	44	23	26	30	15	10	20	8 <sup>a</sup>	8 <sup>a</sup>	19	16	16	14	29
Crack	5	4	11	*	13	16	5	20	8	*	4	*	3	4	*	6	3	4
Heroin	10	9	8	26	9	18	24	14	29	54	52	48	33	41	40	55	25	25
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	13	3	9	7	9
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	6	13	7	*
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	34	20	38	34	46
Misuse of Pain Relievers <sup>1</sup>	--	--	--	--	--	--	--	--	--	--	--	--	--	67 <sup>a</sup>	34	66 <sup>a</sup>	36	28
Misuse of Stimulants	--	--	--	--	--	--	--	--	--	--	--	--	--	23	11	24	12	17
Misuse of Tranquilizers	--	--	--	--	--	--	--	--	--	--	--	--	--	30	20	22	19	24
Misuse of Sedatives	--	--	--	--	--	--	--	--	--	--	--	--	--	*	5	*	2	*
Cigarettes (Daily Only) <sup>2</sup>	6,549 <sup>a</sup>	6,726 <sup>a</sup>	6,574 <sup>a</sup>	6,349 <sup>a</sup>	6,149 <sup>a</sup>	5,829 <sup>a</sup>	5,623 <sup>a</sup>	5,458 <sup>a</sup>	5,358 <sup>a</sup>	5,207 <sup>a</sup>	4,967 <sup>a</sup>	4,591 <sup>a</sup>	4,263 <sup>a</sup>	3,923 <sup>a</sup>	3,237 <sup>a</sup>	2,939 <sup>a</sup>	2,438 <sup>a</sup>	2,097
Smokeless Tobacco	--	--	--	--	--	--	--	--	--	--	--	--	--	923 <sup>a</sup>	883 <sup>a</sup>	761	673	695
Cigars	280 <sup>a</sup>	320 <sup>a</sup>	355	363	371	371	360	379	423	391	389	341 <sup>a</sup>	352	417	392	463	430	435
Alcohol	1,741 <sup>a</sup>	1,820 <sup>a</sup>	1,881 <sup>a</sup>	1,923 <sup>a</sup>	1,941 <sup>a</sup>	1,890 <sup>a</sup>	1,923 <sup>a</sup>	1,847 <sup>a</sup>	1,798 <sup>a</sup>	1,645 <sup>a</sup>	1,717 <sup>a</sup>	1,660 <sup>a</sup>	1,649 <sup>a</sup>	1,532 <sup>a</sup>	1,447	1,425	1,389	1,314
Binge Alcohol <sup>3</sup>	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	294	249	226	237	232
<b>DAILY/ALMOST DAILY USE IN PAST MONTH AMONG PAST MONTH USERS</b>																		
Marijuana	1,997 <sup>a</sup>	2,079 <sup>a</sup>	1,995 <sup>a</sup>	2,096 <sup>a</sup>	2,133 <sup>a</sup>	2,071 <sup>a</sup>	2,178 <sup>a</sup>	2,476 <sup>a</sup>	2,658 <sup>a</sup>	2,676 <sup>a</sup>	2,882 <sup>a</sup>	2,817 <sup>a</sup>	3,020	3,076	3,056	3,367	3,190	3,334
Cocaine	23	29	49	25	44	23	26	30	15	10	20	8 <sup>a</sup>	8 <sup>a</sup>	19	16	16	14	29
Crack	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Heroin	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	13	3	9	7	9
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	*	*	*	*
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	*	*	*	*	*
Misuse of Pain Relievers <sup>1</sup>	--	--	--	--	--	--	--	--	--	--	--	--	--	67 <sup>a</sup>	34	66 <sup>a</sup>	36	28
Misuse of Stimulants	--	--	--	--	--	--	--	--	--	--	--	--	--	23	11	24	12	17
Misuse of Tranquilizers	--	--	--	--	--	--	--	--	--	--	--	--	--	30	20	22	19	24
Misuse of Sedatives	--	--	--	--	--	--	--	--	--	--	--	--	--	*	*	*	*	*
Cigarettes (Daily Only) <sup>2</sup>	6,549 <sup>a</sup>	6,726 <sup>a</sup>	6,574 <sup>a</sup>	6,349 <sup>a</sup>	6,149 <sup>a</sup>	5,829 <sup>a</sup>	5,623 <sup>a</sup>	5,458 <sup>a</sup>	5,358 <sup>a</sup>	5,207 <sup>a</sup>	4,967 <sup>a</sup>	4,591 <sup>a</sup>	4,263 <sup>a</sup>	3,923 <sup>a</sup>	3,237 <sup>a</sup>	2,939 <sup>a</sup>	2,438 <sup>a</sup>	2,097
Smokeless Tobacco	--	--	--	--	--	--	--	--	--	--	--	--	--	923 <sup>a</sup>	883 <sup>a</sup>	761	673	695
Cigars	280 <sup>a</sup>	320 <sup>a</sup>	355	363	371	371	360	379	423	391	389	341 <sup>a</sup>	352	417	392	463	430	435
Alcohol	1,741 <sup>a</sup>	1,820 <sup>a</sup>	1,881 <sup>a</sup>	1,923 <sup>a</sup>	1,941 <sup>a</sup>	1,890 <sup>a</sup>	1,923 <sup>a</sup>	1,847 <sup>a</sup>	1,798 <sup>a</sup>	1,645 <sup>a</sup>	1,717 <sup>a</sup>	1,660 <sup>a</sup>	1,649 <sup>a</sup>	1,532 <sup>a</sup>	1,447	1,425	1,389	1,314
Binge Alcohol <sup>3</sup>	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	294	249	226	237	232
<b>SMOKED 1+ PACKS OF CIGARETTES PER DAY AMONG DAILY CIGARETTE USERS<sup>4</sup></b>	2,556 <sup>a</sup>	2,494 <sup>a</sup>	2,296 <sup>a</sup>	2,342 <sup>a</sup>	2,116 <sup>a</sup>	1,915 <sup>a</sup>	1,774 <sup>a</sup>	1,609 <sup>a</sup>	1,461 <sup>a</sup>	1,361 <sup>a</sup>	1,247 <sup>a</sup>	1,021 <sup>a</sup>	959 <sup>a</sup>	880 <sup>a</sup>	848 <sup>a</sup>	736 <sup>a</sup>	569	524

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.  
NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section 3.3.5 in Chapter 3 of the 2019 National Survey on Drug Use and Health: Methodological Summary and Definitions).

<sup>a</sup> The difference between this estimate and the 2019 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

<sup>1</sup> Prescription psychotherapeutic subtypes were revised in 2016; one effect was the comparability of codeine products between 2015 and 2016.

<sup>2</sup> Includes only respondents who smoked cigarettes on each of the past 30 days.

<sup>3</sup> Estimates of binge alcohol use include use by those who were heavy alcohol users.

<sup>4</sup> Respondents with missing data for number of cigarettes smoked per day were excluded from the analysis.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2019.

**Table 7.29B    Daily or Almost Daily Use of Specific Substances in Past Month among All Persons Aged 18 to 25 and among Past Month Users of Specific Substances Aged 18 to 25: Percentages, 2002-2019**

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
<b>DAILY/ALMOST DAILY USE IN PAST MONTH AMONG TOTAL POPULATION</b>																		
Marijuana	6.4 <sup>a</sup>	6.6 <sup>a</sup>	6.2 <sup>a</sup>	6.5 <sup>a</sup>	6.5 <sup>a</sup>	6.3 <sup>a</sup>	6.6 <sup>a</sup>	7.4 <sup>a</sup>	7.8 <sup>a</sup>	7.8 <sup>a</sup>	8.3 <sup>a</sup>	8.1 <sup>a</sup>	8.6 <sup>a</sup>	8.8 <sup>a</sup>	8.8 <sup>a</sup>	9.8	9.4	9.9
Cocaine	0.1	0.1	0.2	0.1	0.1	0.1	0.1	0.1	0.0	0.0 <sup>a</sup>	0.1	0.0 <sup>a</sup>	0.0 <sup>a</sup>	0.1	0.0	0.0	0.0	0.1
Crack	0.0	0.0	0.0	*	0.0	0.0	0.0	0.1	0.0	*	0.0	*	0.0	0.0	*	0.0	0.0	0.0
Heroin	0.0	0.0	0.0	0.1	0.0	0.1	0.1	0.0	0.1	0.2	0.1	0.1	0.1	0.1	0.1	0.2	0.1	0.1
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.0	0.0	0.0	0.0	0.0
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	0.0	0.0	0.0	*
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	0.1	0.1	0.1	0.1	0.1
Misuse of Pain Relievers <sup>1</sup>	--	--	--	--	--	--	--	--	--	--	--	--	--	0.2 <sup>a</sup>	0.1	0.2 <sup>a</sup>	0.1	0.1
Misuse of Stimulants	--	--	--	--	--	--	--	--	--	--	--	--	--	0.1	0.0	0.1	0.0	0.1
Misuse of Tranquilizers	--	--	--	--	--	--	--	--	--	--	--	--	--	0.1	0.1	0.1	0.1	0.1
Misuse of Sedatives	--	--	--	--	--	--	--	--	--	--	--	--	--	*	0.0	*	0.0	*
Cigarettes (Daily Only) <sup>2</sup>	21.1 <sup>a</sup>	21.2 <sup>a</sup>	20.4 <sup>a</sup>	19.5 <sup>a</sup>	18.8 <sup>a</sup>	17.8 <sup>a</sup>	17.1 <sup>a</sup>	16.3 <sup>a</sup>	15.7 <sup>a</sup>	15.2 <sup>a</sup>	14.4 <sup>a</sup>	13.2 <sup>a</sup>	12.2 <sup>a</sup>	11.2 <sup>a</sup>	9.4 <sup>a</sup>	8.6 <sup>a</sup>	7.2 <sup>a</sup>	6.2
Smokeless Tobacco	--	--	--	--	--	--	--	--	--	--	--	--	--	2.6 <sup>a</sup>	2.6 <sup>a</sup>	2.2	2.0	2.1
Cigars	0.9 <sup>a</sup>	1.0 <sup>a</sup>	1.1	1.1	1.1	1.1	1.1	1.1	1.2	1.1	1.1	1.0 <sup>a</sup>	1.0	1.2	1.1	1.4	1.3	1.3
Alcohol	5.6 <sup>a</sup>	5.7 <sup>a</sup>	5.8 <sup>a</sup>	5.9 <sup>a</sup>	5.9 <sup>a</sup>	5.8 <sup>a</sup>	5.8 <sup>a</sup>	5.5 <sup>a</sup>	5.3 <sup>a</sup>	4.8 <sup>a</sup>	5.0 <sup>a</sup>	4.8 <sup>a</sup>	4.7 <sup>a</sup>	4.4	4.2	4.2	4.1	3.9
Binge Alcohol <sup>3</sup>	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.8	0.7	0.7	0.7	0.7
<b>DAILY/ALMOST DAILY USE IN PAST MONTH AMONG PAST MONTH USERS</b>																		
Marijuana	37.2 <sup>a</sup>	38.5 <sup>a</sup>	38.4 <sup>a</sup>	38.9 <sup>a</sup>	40.0 <sup>a</sup>	38.4 <sup>a</sup>	39.9 <sup>a</sup>	40.4	42.1	41.0	44.4	42.5	44.2	44.4	42.5	44.3	42.4	43.0
Cocaine	3.6	4.1	7.0	3.0	6.0	4.0	5.1	6.4	3.1	2.0	5.0	2.1	1.7 <sup>a</sup>	3.2	3.0	2.4	2.6	5.4
Crack	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Heroin	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	2.1	0.4	1.6	1.2	1.5
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	*	*	*	*
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	*	*	*	*	*
Misuse of Pain Relievers <sup>1</sup>	--	--	--	--	--	--	--	--	--	--	--	--	--	8.1	5.4	10.4	7.5	6.6
Misuse of Stimulants	--	--	--	--	--	--	--	--	--	--	--	--	--	3.0	1.5	3.3	2.1	2.9
Misuse of Tranquilizers	--	--	--	--	--	--	--	--	--	--	--	--	--	5.2	3.7	4.0	4.9	7.1
Misuse of Sedatives	--	--	--	--	--	--	--	--	--	--	--	--	--	*	*	*	*	*
Cigarettes (Daily Only) <sup>2</sup>	51.8 <sup>a</sup>	52.7 <sup>a</sup>	51.6 <sup>a</sup>	50.1 <sup>a</sup>	48.8 <sup>a</sup>	49.2 <sup>a</sup>	47.8 <sup>a</sup>	45.3 <sup>a</sup>	45.8 <sup>a</sup>	45.3 <sup>a</sup>	45.1 <sup>a</sup>	43.1 <sup>a</sup>	43.0 <sup>a</sup>	42.0 <sup>a</sup>	39.9 <sup>a</sup>	38.4	37.4	35.4
Smokeless Tobacco	--	--	--	--	--	--	--	--	--	--	--	--	--	48.7 <sup>a</sup>	48.9 <sup>a</sup>	46.7	45.0	41.9
Cigars	8.2 <sup>a</sup>	8.8 <sup>a</sup>	8.7 <sup>a</sup>	9.3 <sup>a</sup>	9.3 <sup>a</sup>	9.5 <sup>a</sup>	9.6 <sup>a</sup>	9.8 <sup>a</sup>	11.0 <sup>a</sup>	10.5 <sup>a</sup>	10.5 <sup>a</sup>	9.8 <sup>a</sup>	10.4 <sup>a</sup>	13.4	12.8 <sup>a</sup>	14.8	14.7	16.7
Alcohol	9.3 <sup>a</sup>	9.4 <sup>a</sup>	9.7 <sup>a</sup>	9.7 <sup>a</sup>	9.6 <sup>a</sup>	9.4 <sup>a</sup>	9.6 <sup>a</sup>	8.9 <sup>a</sup>	8.6 <sup>a</sup>	7.9	8.2 <sup>a</sup>	8.0	7.9	7.5	7.3	7.4	7.4	7.2
Binge Alcohol <sup>3</sup>	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	2.2	1.9	1.8	2.0	2.0
<b>SMOKED 1+ PACKS OF CIGARETTES PER DAY AMONG DAILY CIGARETTE USERS<sup>4</sup></b>																		
	39.1 <sup>a</sup>	37.1 <sup>a</sup>	34.9 <sup>a</sup>	36.9 <sup>a</sup>	34.4 <sup>a</sup>	32.9 <sup>a</sup>	31.6 <sup>a</sup>	29.5 <sup>a</sup>	27.3	26.1	25.1	22.3	22.5	22.5	26.2	25.0	23.4	25.0

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.  
NOTE:    Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section 3.3.5 in Chapter 3 of the *2019 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

<sup>a</sup> The difference between this estimate and the 2019 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

<sup>1</sup> Prescription psychotherapeutic subtypes were revised in 2016; one effect was the comparability of codeine products between 2015 and 2016.

<sup>2</sup> Includes only respondents who smoked cigarettes on each of the past 30 days.

<sup>3</sup> Estimates of binge alcohol use include use by those who were heavy alcohol users.

<sup>4</sup> Respondents with missing data for number of cigarettes smoked per day were excluded from the analysis.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2019.

**Table 7.30A Daily or Almost Daily Use of Specific Substances in Past Month among All Persons Aged 26 or Older and among Past Month Users of Specific Substances Aged 26 or Older: Numbers in Thousands, 2002-2019**

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
<b>DAILY/ALMOST DAILY USE IN PAST MONTH AMONG TOTAL POPULATION</b>																		
Marijuana	2,156 <sup>a</sup>	2,332 <sup>a</sup>	2,349 <sup>a</sup>	2,593 <sup>a</sup>	2,537 <sup>a</sup>	2,601 <sup>a</sup>	2,855 <sup>a</sup>	3,253 <sup>a</sup>	3,838 <sup>a</sup>	3,931 <sup>a</sup>	4,293 <sup>a</sup>	4,875 <sup>a</sup>	5,820 <sup>a</sup>	5,744 <sup>a</sup>	6,573 <sup>a</sup>	7,057 <sup>a</sup>	8,409 <sup>a</sup>	10,049
Cocaine	83	96	203	174	299	144	130	121	161	59	107	175	100	92	167	121	110	146
Crack	31	66	103	74	107	79	61	68	47	19 <sup>a</sup>	69	93	55	48	95	54	50	91
Heroin	84	45 <sup>a</sup>	64	20 <sup>a</sup>	178	61 <sup>a</sup>	67	32 <sup>a</sup>	49 <sup>a</sup>	118	128	96	140	100	179	149	103	156
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	17	*	21	38	*
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	18	14	11	30	30
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	212 <sup>a</sup>	227 <sup>a</sup>	264	360	391
Misuse of Pain Relievers <sup>1</sup>	--	--	--	--	--	--	--	--	--	--	--	--	--	356	307	286	332	315
Misuse of Stimulants	--	--	--	--	--	--	--	--	--	--	--	--	--	61	72	93	63	78
Misuse of Tranquilizers	--	--	--	--	--	--	--	--	--	--	--	--	--	72	95	63	75	101
Misuse of Sedatives	--	--	--	--	--	--	--	--	--	--	--	--	--	28	34	39	*	26
Cigarettes (Daily Only) <sup>2</sup>	31,168 <sup>a</sup>	30,359 <sup>a</sup>	29,942 <sup>a</sup>	31,079 <sup>a</sup>	31,481 <sup>a</sup>	30,414 <sup>a</sup>	30,655 <sup>a</sup>	29,865 <sup>a</sup>	28,901 <sup>a</sup>	28,841 <sup>a</sup>	29,610 <sup>a</sup>	28,385 <sup>a</sup>	27,916 <sup>a</sup>	26,059 <sup>a</sup>	26,336 <sup>a</sup>	24,751	24,792	24,624
Smokeless Tobacco	--	--	--	--	--	--	--	--	--	--	--	--	--	4,363	4,340	4,302	4,138	4,215
Cigars	980 <sup>a</sup>	1,353 <sup>a</sup>	1,217 <sup>a</sup>	1,203 <sup>a</sup>	1,439	1,284 <sup>a</sup>	1,461	1,531	1,676	1,374 <sup>a</sup>	1,733	1,530	1,337 <sup>a</sup>	1,809	1,960	1,838	1,945	1,806
Alcohol	17,839 <sup>a</sup>	17,095 <sup>a</sup>	17,443 <sup>a</sup>	18,140 <sup>a</sup>	17,848 <sup>a</sup>	18,041 <sup>a</sup>	19,167	19,941	18,472 <sup>a</sup>	18,911	19,757	19,547	20,889	19,835	19,481	20,389	20,262	20,015
Binge Alcohol <sup>3</sup>	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	2,550	2,413	2,678	2,663	2,413
<b>DAILY/ALMOST DAILY USE IN PAST MONTH AMONG PAST MONTH USERS</b>																		
Marijuana	2,156 <sup>a</sup>	2,332 <sup>a</sup>	2,349 <sup>a</sup>	2,593 <sup>a</sup>	2,537 <sup>a</sup>	2,601 <sup>a</sup>	2,855 <sup>a</sup>	3,253 <sup>a</sup>	3,838 <sup>a</sup>	3,931 <sup>a</sup>	4,293 <sup>a</sup>	4,875 <sup>a</sup>	5,820 <sup>a</sup>	5,744 <sup>a</sup>	6,573 <sup>a</sup>	7,057 <sup>a</sup>	8,409 <sup>a</sup>	10,049
Cocaine	83	96	203	174	299	144	130	121	161	59	107	*	100	92	167	121	110	146
Crack	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Heroin	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	9	21	38 <sup>a</sup>	6
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	*	*	*	*
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	212 <sup>a</sup>	*	264	360	391
Misuse of Pain Relievers <sup>1</sup>	--	--	--	--	--	--	--	--	--	--	--	--	--	356	307	286	332	315
Misuse of Stimulants	--	--	--	--	--	--	--	--	--	--	--	--	--	61	72	93	63	78
Misuse of Tranquilizers	--	--	--	--	--	--	--	--	--	--	--	--	--	72	95	63	75	101
Misuse of Sedatives	--	--	--	--	--	--	--	--	--	--	--	--	--	*	*	*	*	*
Cigarettes (Daily Only) <sup>2</sup>	31,168 <sup>a</sup>	30,359 <sup>a</sup>	29,942 <sup>a</sup>	31,079 <sup>a</sup>	31,481 <sup>a</sup>	30,414 <sup>a</sup>	30,655 <sup>a</sup>	29,865 <sup>a</sup>	28,901 <sup>a</sup>	28,841 <sup>a</sup>	29,610 <sup>a</sup>	28,385 <sup>a</sup>	27,916 <sup>a</sup>	26,059	26,336 <sup>a</sup>	24,751	24,792	24,624
Smokeless Tobacco	--	--	--	--	--	--	--	--	--	--	--	--	--	4,363	4,340	4,302	4,138	4,215
Cigars	980 <sup>a</sup>	1,353 <sup>a</sup>	1,217 <sup>a</sup>	1,203 <sup>a</sup>	1,439	1,284 <sup>a</sup>	1,461	1,531	1,676	1,374 <sup>a</sup>	1,733	1,530	1,337 <sup>a</sup>	1,809	1,960	1,838	1,945	1,806
Alcohol	17,839 <sup>a</sup>	17,095 <sup>a</sup>	17,443 <sup>a</sup>	18,140 <sup>a</sup>	17,848 <sup>a</sup>	18,041 <sup>a</sup>	19,167	19,941	18,472	18,911	19,757	19,547	20,889	19,835	19,481	20,389	20,262	20,015
Binge Alcohol <sup>3</sup>	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	2,550	2,413	2,678	2,663	2,413
<b>SMOKED 1+ PACKS OF CIGARETTES PER DAY AMONG DAILY CIGARETTE USERS<sup>4</sup></b>	17,743 <sup>a</sup>	17,591 <sup>a</sup>	17,704 <sup>a</sup>	17,089 <sup>a</sup>	17,153 <sup>a</sup>	16,715 <sup>a</sup>	16,199 <sup>a</sup>	14,733 <sup>a</sup>	14,082 <sup>a</sup>	13,668 <sup>a</sup>	13,365 <sup>a</sup>	12,667 <sup>a</sup>	12,065 <sup>a</sup>	11,484 <sup>a</sup>	11,324 <sup>a</sup>	10,666	10,221	10,315

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.  
NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section 3.3.5 in Chapter 3 of the *2019 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

<sup>a</sup> The difference between this estimate and the 2019 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

<sup>1</sup> Prescription psychotherapeutic subtypes were revised in 2016; one effect was the comparability of codeine products between 2015 and 2016.

<sup>2</sup> Includes only respondents who smoked cigarettes on each of the past 30 days.

<sup>3</sup> Estimates of binge alcohol use include use by those who were heavy alcohol users.

<sup>4</sup> Respondents with missing data for number of cigarettes smoked per day were excluded from the analysis.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2019.

Table 7.30B Daily or Almost Daily Use of Specific Substances in Past Month among All Persons Aged 26 or Older and among Past Month Users of Specific Substances Aged 26 or Older: Percentages, 2002-2019

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
<b>DAILY/ALMOST DAILY USE IN PAST MONTH AMONG TOTAL POPULATION</b>																		
Marijuana	1.2 <sup>a</sup>	1.3 <sup>a</sup>	1.3 <sup>a</sup>	1.4 <sup>a</sup>	1.4 <sup>a</sup>	1.4 <sup>a</sup>	1.5 <sup>a</sup>	1.7 <sup>a</sup>	2.0 <sup>a</sup>	2.0 <sup>a</sup>	2.1 <sup>a</sup>	2.4 <sup>a</sup>	2.8 <sup>a</sup>	2.8 <sup>a</sup>	3.1 <sup>a</sup>	3.3 <sup>a</sup>	3.9 <sup>a</sup>	4.6
Cocaine	0.0	0.1	0.1	0.1	0.2 <sup>a</sup>	0.1	0.1	0.1	0.1	0.0	0.1	0.1	0.0	0.0	0.1	0.1	0.1	0.1
Crack	0.0	0.0	0.1	0.0	0.1	0.0	0.0	0.0	0.0	0.0 <sup>a</sup>	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Heroin	0.0	0.0 <sup>a</sup>	0.0	0.0 <sup>a</sup>	0.1	0.0	0.0	0.0 <sup>a</sup>	0.0 <sup>a</sup>	0.1	0.0	0.0	0.1	0.0	0.1	0.1	0.0	0.1
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.0	*	0.0	0.0	*
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.0	0.0	0.0	0.0	0.0
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	0.1 <sup>a</sup>	0.1 <sup>a</sup>	0.1	0.2	0.2
Misuse of Pain Relievers <sup>1</sup>	--	--	--	--	--	--	--	--	--	--	--	--	--	0.2	0.1	0.1	0.2	0.1
Misuse of Stimulants	--	--	--	--	--	--	--	--	--	--	--	--	--	0.0	0.0	0.0	0.0	0.0
Misuse of Tranquilizers	--	--	--	--	--	--	--	--	--	--	--	--	--	0.0	0.0	0.0	0.0	0.0
Misuse of Sedatives	--	--	--	--	--	--	--	--	--	--	--	--	--	0.0	0.0	0.0	*	0.0
Cigarettes (Daily Only) <sup>2</sup>	17.4 <sup>a</sup>	16.8 <sup>a</sup>	16.4 <sup>a</sup>	16.8 <sup>a</sup>	16.8 <sup>a</sup>	16.0 <sup>a</sup>	16.0 <sup>a</sup>	15.4 <sup>a</sup>	14.8 <sup>a</sup>	14.5 <sup>a</sup>	14.8 <sup>a</sup>	14.0 <sup>a</sup>	13.6 <sup>a</sup>	12.5 <sup>a</sup>	12.5 <sup>a</sup>	11.6	11.5	11.4
Smokeless Tobacco	--	--	--	--	--	--	--	--	--	--	--	--	--	2.1	2.1	2.0	1.9	1.9
Cigars	0.5 <sup>a</sup>	0.7	0.7	0.6	0.8	0.7	0.8	0.8	0.9	0.7	0.9	0.8	0.7 <sup>a</sup>	0.9	0.9	0.9	0.9	0.8
Alcohol	9.9	9.4	9.5	9.8	9.5	9.5	10.0	10.3 <sup>a</sup>	9.5	9.5	9.9	9.6	10.2 <sup>a</sup>	9.5	9.3	9.6	9.4	9.2
Binge Alcohol <sup>3</sup>	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	1.2	1.1	1.3	1.2	1.1
<b>DAILY/ALMOST DAILY USE IN PAST MONTH AMONG PAST MONTH USERS</b>																		
Marijuana	30.0 <sup>a</sup>	32.1 <sup>a</sup>	31.4 <sup>a</sup>	34.5 <sup>a</sup>	32.5 <sup>a</sup>	35.2 <sup>a</sup>	35.1 <sup>a</sup>	36.6 <sup>a</sup>	41.3	41.0 <sup>a</sup>	40.6 <sup>a</sup>	42.7	43.0	42.4	43.3	42.0 <sup>a</sup>	45.5	45.6
Cocaine	6.7	6.7	16.8	12.2	18.8	10.3	10.4	10.9	17.7	7.1	8.8	*	9.8	7.4	12.9	8.2	7.8	10.2
Crack	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Heroin	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	1.6	3.0	4.2 <sup>a</sup>	0.5
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	*	*	*	*
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	28.0	*	43.5	40.3	37.2
Misuse of Pain Relievers <sup>1</sup>	--	--	--	--	--	--	--	--	--	--	--	--	--	13.3	12.4	12.0	15.0	14.1
Misuse of Stimulants	--	--	--	--	--	--	--	--	--	--	--	--	--	7.9	8.2	9.3	6.5	9.1
Misuse of Tranquilizers	--	--	--	--	--	--	--	--	--	--	--	--	--	6.3	7.3	6.0	6.4	7.6
Misuse of Sedatives	--	--	--	--	--	--	--	--	--	--	--	--	--	*	*	*	*	*
Cigarettes (Daily Only) <sup>2</sup>	68.8 <sup>a</sup>	68.0 <sup>a</sup>	67.8 <sup>a</sup>	68.9 <sup>a</sup>	67.9 <sup>a</sup>	66.3 <sup>a</sup>	67.0 <sup>a</sup>	67.2 <sup>a</sup>	64.8 <sup>a</sup>	66.5 <sup>a</sup>	66.0 <sup>a</sup>	64.9 <sup>a</sup>	63.3	62.7	62.2	61.5	62.3	62.5
Smokeless Tobacco	--	--	--	--	--	--	--	--	--	--	--	--	--	64.8	65.7	65.1	66.7	63.8
Cigars	11.9 <sup>a</sup>	16.7	14.4 <sup>a</sup>	13.9 <sup>a</sup>	16.6 <sup>a</sup>	15.4 <sup>a</sup>	17.2	18.0	19.4	16.6	19.1	18.2	16.6 <sup>a</sup>	20.4	22.4	20.6	21.9	20.7
Alcohol	18.4 <sup>a</sup>	18.0	18.0	17.8	17.7	17.6	18.2 <sup>a</sup>	18.8 <sup>a</sup>	17.2	17.3	17.7	17.3	18.0 <sup>a</sup>	17.2	17.0	17.2	17.0	16.8
Binge Alcohol <sup>3</sup>	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	4.9	4.7	5.1	4.9	4.5
<b>SMOKED 1+ PACKS OF CIGARETTES PER DAY AMONG DAILY CIGARETTE USERS<sup>4</sup></b>																		
	57.1 <sup>a</sup>	58.0 <sup>a</sup>	59.2 <sup>a</sup>	55.1 <sup>a</sup>	54.5 <sup>a</sup>	55.1 <sup>a</sup>	53.0 <sup>a</sup>	49.4 <sup>a</sup>	48.8 <sup>a</sup>	47.4 <sup>a</sup>	45.2 <sup>a</sup>	44.7	43.3	44.1	43.1	43.2	41.4	42.0

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.  
NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section 3.3.5 in Chapter 3 of the 2019 National Survey on Drug Use and Health: Methodological Summary and Definitions).

<sup>a</sup> The difference between this estimate and the 2019 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

<sup>1</sup> Prescription psychotherapeutic subtypes were revised in 2016; one effect was the comparability of codeine products between 2015 and 2016.

<sup>2</sup> Includes only respondents who smoked cigarettes on each of the past 30 days.

<sup>3</sup> Estimates of binge alcohol use include use by those who were heavy alcohol users.

<sup>4</sup> Respondents with missing data for number of cigarettes smoked per day were excluded from the analysis.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2019.



Table 7.31A Daily or Almost Daily Use of Specific Substances in Past Month among All Persons Aged 12 to 20 and among Past Month Users of Specific Substances Aged 12 to 20: Numbers in Thousands, 2002-2019

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
<b>DAILY/ALMOST DAILY USE IN PAST MONTH AMONG TOTAL POPULATION</b>																		
Marijuana	1,530	1,450	1,323 <sup>a</sup>	1,333 <sup>a</sup>	1,435	1,278 <sup>a</sup>	1,433	1,522	1,488	1,578	1,652	1,536	1,524	1,485	1,434	1,565	1,469	1,547
Cocaine	17	21	31 <sup>a</sup>	20	12	18	23 <sup>a</sup>	18	12	7	8	5	12	11	11	4	4	7
Crack	7	8	5	*	*	8	*	8	8	8	3	*	*	4	*	4	*	4
Heroin	7	9	6	8	4	7	9	5	8	33 <sup>a</sup>	17	11	18	7	4	*	7	4
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	10	3	5	6	3
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	6	14	19	4 <sup>a</sup>	22
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	19	*	23	15	9
Misuse of Pain Relievers <sup>1</sup>	--	--	--	--	--	--	--	--	--	--	--	--	--	15	18 <sup>a</sup>	25 <sup>a</sup>	11	5
Misuse of Stimulants	--	--	--	--	--	--	--	--	--	--	--	--	--	14	11	19	3	12
Misuse of Tranquilizers	--	--	--	--	--	--	--	--	--	--	--	--	--	18	17	10	23	6
Misuse of Sedatives	--	--	--	--	--	--	--	--	--	--	--	--	--	*	*	*	*	*
Cigarettes (Daily Only) <sup>2</sup>	3,374 <sup>a</sup>	3,209 <sup>a</sup>	3,028 <sup>a</sup>	2,819 <sup>a</sup>	2,769 <sup>a</sup>	2,550 <sup>a</sup>	2,430 <sup>a</sup>	2,330 <sup>a</sup>	2,221 <sup>a</sup>	2,146 <sup>a</sup>	1,756 <sup>a</sup>	1,559 <sup>a</sup>	1,511 <sup>a</sup>	1,190 <sup>a</sup>	900 <sup>a</sup>	771 <sup>a</sup>	663 <sup>a</sup>	535
Smokeless Tobacco	--	--	--	--	--	--	--	--	--	--	--	--	--	426 <sup>a</sup>	404 <sup>a</sup>	306	293	234
Cigars	207	205	220	232	199	190	194	237	219	217	169	163	159	195	191	219	195	208
Alcohol	632 <sup>a</sup>	624 <sup>a</sup>	647 <sup>a</sup>	625 <sup>a</sup>	587 <sup>a</sup>	553 <sup>a</sup>	533 <sup>a</sup>	549 <sup>a</sup>	467 <sup>a</sup>	386 <sup>a</sup>	354 <sup>a</sup>	328 <sup>a</sup>	337 <sup>a</sup>	278 <sup>a</sup>	233	274 <sup>a</sup>	230	189
Binge Alcohol <sup>3</sup>	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	69	43	66	51	38
<b>DAILY/ALMOST DAILY USE IN PAST MONTH AMONG PAST MONTH USERS</b>																		
Marijuana	1,530	1,450	1,323 <sup>a</sup>	1,333 <sup>a</sup>	1,435	1,278 <sup>a</sup>	1,433	1,522	1,488	1,578	1,652	1,536	1,524	1,485	1,434	1,565	1,469	1,547
Cocaine	17	21	31	20	12	18	23	18	12	7	8	*	*	11	*	4	*	*
Crack	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Heroin	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	10	*	5	6	3
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	6	14	19	4 <sup>a</sup>	22
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	*	*	*	*	*
Misuse of Pain Relievers <sup>1</sup>	--	--	--	--	--	--	--	--	--	--	--	--	--	15	18 <sup>a</sup>	25 <sup>a</sup>	11	5
Misuse of Stimulants	--	--	--	--	--	--	--	--	--	--	--	--	--	14	11	19	*	12
Misuse of Tranquilizers	--	--	--	--	--	--	--	--	--	--	--	--	--	18	17	10	23	6
Misuse of Sedatives	--	--	--	--	--	--	--	--	--	--	--	--	--	*	*	*	*	*
Cigarettes (Daily Only) <sup>2</sup>	3,374 <sup>a</sup>	3,209 <sup>a</sup>	3,028 <sup>a</sup>	2,819 <sup>a</sup>	2,769 <sup>a</sup>	2,550 <sup>a</sup>	2,430 <sup>a</sup>	2,330 <sup>a</sup>	2,221 <sup>a</sup>	2,146 <sup>a</sup>	1,756 <sup>a</sup>	1,559 <sup>a</sup>	1,511 <sup>a</sup>	1,190 <sup>a</sup>	900 <sup>a</sup>	771 <sup>a</sup>	663 <sup>a</sup>	535
Smokeless Tobacco	--	--	--	--	--	--	--	--	--	--	--	--	--	426 <sup>a</sup>	404 <sup>a</sup>	306	293	234
Cigars	207	205	220	232	199	190	194	237	219	217	169	163	159	195	191	219	195	208
Alcohol	632 <sup>a</sup>	624 <sup>a</sup>	647 <sup>a</sup>	625 <sup>a</sup>	587 <sup>a</sup>	553 <sup>a</sup>	533 <sup>a</sup>	549 <sup>a</sup>	467 <sup>a</sup>	386 <sup>a</sup>	354 <sup>a</sup>	328 <sup>a</sup>	337 <sup>a</sup>	278 <sup>a</sup>	233	274 <sup>a</sup>	230	189
Binge Alcohol <sup>3</sup>	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	69	43	66	51	38
<b>SMOKED 1+ PACKS OF CIGARETTES PER DAY AMONG DAILY CIGARETTE USERS<sup>4</sup></b>	1,013 <sup>a</sup>	925 <sup>a</sup>	834 <sup>a</sup>	856 <sup>a</sup>	734 <sup>a</sup>	621 <sup>a</sup>	604 <sup>a</sup>	547 <sup>a</sup>	467 <sup>a</sup>	464 <sup>a</sup>	313 <sup>a</sup>	280 <sup>a</sup>	273 <sup>a</sup>	165 <sup>a</sup>	221 <sup>a</sup>	174 <sup>a</sup>	131	85

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.  
NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section 3.3.5 in Chapter 3 of the *2019 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

<sup>a</sup> The difference between this estimate and the 2019 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

<sup>1</sup> Prescription psychotherapeutic subtypes were revised in 2016; one effect was the comparability of codeine products between 2015 and 2016.

<sup>2</sup> Includes only respondents who smoked cigarettes on each of the past 30 days.

<sup>3</sup> Estimates of binge alcohol use include use by those who were heavy alcohol users.

<sup>4</sup> Respondents with missing data for number of cigarettes smoked per day were excluded from the analysis.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2019.

Table 7.31B Daily or Almost Daily Use of Specific Substances in Past Month among All Persons Aged 12 to 20 and among Past Month Users of Specific Substances Aged 12 to 20: Percentages, 2002-2019

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
<b>DAILY/ALMOST DAILY USE IN PAST MONTH AMONG TOTAL POPULATION</b>																		
Marijuana	4.1	3.9	3.5 <sup>a</sup>	3.5 <sup>a</sup>	3.8	3.3 <sup>a</sup>	3.8	4.0	3.9	4.1	4.3	4.0	4.0	3.9	3.8	4.1	3.9	4.1
Cocaine	0.0	0.1	0.1 <sup>a</sup>	0.1	0.0	0.0	0.1 <sup>a</sup>	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Crack	0.0	0.0	0.0	*	*	0.0	*	0.0	0.0	*	0.0	*	*	0.0	*	0.0	*	0.0
Heroin	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1 <sup>a</sup>	0.0	0.0	0.0	0.0	0.0	*	0.0	0.0
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.0	0.0	0.0	0.0	0.0
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.0	0.0	0.0	0.0 <sup>a</sup>	0.1
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	0.0	*	0.1	0.0	0.0
Misuse of Pain Relievers <sup>1</sup>	--	--	--	--	--	--	--	--	--	--	--	--	--	0.0	0.0 <sup>a</sup>	0.1 <sup>a</sup>	0.0	0.0
Misuse of Stimulants	--	--	--	--	--	--	--	--	--	--	--	--	--	0.0	0.0	0.1	0.0	0.0
Misuse of Tranquilizers	--	--	--	--	--	--	--	--	--	--	--	--	--	0.0	0.0	0.0	0.1	0.0
Misuse of Sedatives	--	--	--	--	--	--	--	--	--	--	--	--	--	*	*	*	*	*
Cigarettes (Daily Only) <sup>2</sup>	9.1 <sup>a</sup>	8.6 <sup>a</sup>	8.0 <sup>a</sup>	7.4 <sup>a</sup>	7.3 <sup>a</sup>	6.6 <sup>a</sup>	6.4 <sup>a</sup>	6.1 <sup>a</sup>	5.8 <sup>a</sup>	5.6 <sup>a</sup>	4.6 <sup>a</sup>	4.1 <sup>a</sup>	4.0 <sup>a</sup>	3.1 <sup>a</sup>	2.4 <sup>a</sup>	2.0 <sup>a</sup>	1.7 <sup>a</sup>	1.4
Smokeless Tobacco	--	--	--	--	--	--	--	--	--	--	--	--	--	1.1 <sup>a</sup>	1.1 <sup>a</sup>	0.8	0.8	0.6
Cigars	0.6	0.5	0.6	0.6	0.5	0.5	0.5	0.6	0.6	0.6	0.4	0.4	0.4	0.5	0.5	0.6	0.5	0.5
Alcohol	1.7 <sup>a</sup>	1.7 <sup>a</sup>	1.7 <sup>a</sup>	1.6 <sup>a</sup>	1.5 <sup>a</sup>	1.4 <sup>a</sup>	1.4 <sup>a</sup>	1.4 <sup>a</sup>	1.2 <sup>a</sup>	1.0 <sup>a</sup>	0.9 <sup>a</sup>	0.9 <sup>a</sup>	0.9 <sup>a</sup>	0.7 <sup>a</sup>	0.6	0.7 <sup>a</sup>	0.6	0.5
Binge Alcohol <sup>3</sup>	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.2	0.1	0.2	0.1	0.1
<b>DAILY/ALMOST DAILY USE IN PAST MONTH AMONG PAST MONTH USERS</b>																		
Marijuana	34.3	32.5	31.6	31.8	35.2	31.0	34.5	34.0	32.5	32.2	35.6	34.5	33.4	33.6	32.8	35.5	32.1	33.4
Cocaine	4.2	5.0	8.1	4.5	2.6	5.2	7.3	8.2	5.3	2.6	4.5	*	*	5.0	*	2.1	*	*
Crack	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Heroin	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	2.7	*	1.3	1.8	0.7
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	2.7	6.7	8.6	1.8 <sup>a</sup>	8.3
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	*	*	*	*	*
Misuse of Pain Relievers <sup>1</sup>	--	--	--	--	--	--	--	--	--	--	--	--	--	2.8	4.1	6.3 <sup>a</sup>	3.4	1.6
Misuse of Stimulants	--	--	--	--	--	--	--	--	--	--	--	--	--	3.4	2.6	5.3	*	4.0
Misuse of Tranquilizers	--	--	--	--	--	--	--	--	--	--	--	--	--	4.8	5.3	3.2	9.6	2.8
Misuse of Sedatives	--	--	--	--	--	--	--	--	--	--	--	--	--	*	*	*	*	*
Cigarettes (Daily Only) <sup>2</sup>	42.1 <sup>a</sup>	41.1 <sup>a</sup>	39.5 <sup>a</sup>	38.3 <sup>a</sup>	38.4 <sup>a</sup>	37.4 <sup>a</sup>	36.2 <sup>a</sup>	34.6 <sup>a</sup>	34.6 <sup>a</sup>	34.5 <sup>a</sup>	32.6 <sup>a</sup>	31.4 <sup>a</sup>	34.7 <sup>a</sup>	30.9 <sup>a</sup>	27.8 <sup>a</sup>	25.6	25.6	23.4
Smokeless Tobacco	--	--	--	--	--	--	--	--	--	--	--	--	--	40.3 <sup>a</sup>	41.3 <sup>a</sup>	34.1	35.2	29.4
Cigars	7.5 <sup>a</sup>	7.4 <sup>a</sup>	7.1 <sup>a</sup>	7.9 <sup>a</sup>	7.0 <sup>a</sup>	6.4 <sup>a</sup>	7.1 <sup>a</sup>	8.5 <sup>a</sup>	8.6 <sup>a</sup>	8.5 <sup>a</sup>	7.6 <sup>a</sup>	8.0 <sup>a</sup>	8.4 <sup>a</sup>	11.0 <sup>a</sup>	11.5 <sup>a</sup>	13.1	12.7	16.5
Alcohol	5.9 <sup>a</sup>	5.7 <sup>a</sup>	6.0 <sup>a</sup>	5.8 <sup>a</sup>	5.4 <sup>a</sup>	5.1 <sup>a</sup>	5.3 <sup>a</sup>	5.3 <sup>a</sup>	4.7 <sup>a</sup>	4.0 <sup>a</sup>	3.8 <sup>a</sup>	3.8 <sup>a</sup>	3.9 <sup>a</sup>	3.6	3.2	3.7	3.2	2.7
Binge Alcohol <sup>3</sup>	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	1.4	0.9	1.5	1.2	0.9
<b>SMOKED 1+ PACKS OF CIGARETTES PER DAY AMONG DAILY CIGARETTE USERS<sup>4</sup></b>	30.1 <sup>a</sup>	28.9 <sup>a</sup>	27.6 <sup>a</sup>	30.4 <sup>a</sup>	26.5 <sup>a</sup>	24.4 <sup>a</sup>	24.9 <sup>a</sup>	23.5 <sup>a</sup>	21.1	21.6 <sup>a</sup>	17.9	18.0	18.1	14.0	24.6 <sup>a</sup>	22.6	19.8	16.0

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.  
NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section 3.3.5 in Chapter 3 of the *2019 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

<sup>a</sup> The difference between this estimate and the 2019 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

<sup>1</sup> Prescription psychotherapeutic subtypes were revised in 2016; one effect was the comparability of codeine products between 2015 and 2016.

<sup>2</sup> Includes only respondents who smoked cigarettes on each of the past 30 days.

<sup>3</sup> Estimates of binge alcohol use include use by those who were heavy alcohol users.

<sup>4</sup> Respondents with missing data for number of cigarettes smoked per day were excluded from the analysis.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2019.

**Table 7.32A Daily or Almost Daily Use of Specific Substances in Past Month among All Persons Aged 21 or Older and among Past Month Users of Specific Substances Aged 21 or Older: Numbers in Thousands, 2002-2019**

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
<b>DAILY/ALMOST DAILY USE IN PAST MONTH AMONG TOTAL POPULATION</b>																		
Marijuana	3,226 <sup>a</sup>	3,444 <sup>a</sup>	3,557 <sup>a</sup>	3,756 <sup>a</sup>	3,669 <sup>a</sup>	3,820 <sup>a</sup>	4,022 <sup>a</sup>	4,659 <sup>a</sup>	5,448 <sup>a</sup>	5,496 <sup>a</sup>	5,949 <sup>a</sup>	6,608 <sup>a</sup>	7,715 <sup>a</sup>	7,756 <sup>a</sup>	8,561 <sup>a</sup>	9,264 <sup>a</sup>	10,497 <sup>a</sup>	12,283
Cocaine	96	111	232	189	332	159	145	139	166	62 <sup>a</sup>	122	180	105	102	176	133	121	168
Crack	35	67	112	75	119	91	64	82	47	19 <sup>a</sup>	70	94	57	48	96	55	53	91
Heroin	90	48 <sup>a</sup>	72 <sup>a</sup>	38 <sup>a</sup>	184	73 <sup>a</sup>	84 <sup>a</sup>	43 <sup>a</sup>	71 <sup>a</sup>	151	163	134	160	134	216	203	125	177
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	22	12	27	42	13
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	20	18	15	38	30
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	227 <sup>a</sup>	247 <sup>a</sup>	284 <sup>a</sup>	384	429
Misuse of Pain Relievers <sup>1</sup>	--	--	--	--	--	--	--	--	--	--	--	--	--	416	330	336	358	341
Misuse of Stimulants	--	--	--	--	--	--	--	--	--	--	--	--	--	75	80	103	73	92
Misuse of Tranquilizers	--	--	--	--	--	--	--	--	--	--	--	--	--	96	103	81	81	121
Misuse of Sedatives	--	--	--	--	--	--	--	--	--	--	--	--	--	28	39	39	*	26
Cigarettes (Daily Only) <sup>2</sup>	35,364 <sup>a</sup>	34,781 <sup>a</sup>	34,315 <sup>a</sup>	35,312 <sup>a</sup>	35,562 <sup>a</sup>	34,351 <sup>a</sup>	34,358 <sup>a</sup>	33,504 <sup>a</sup>	32,497 <sup>a</sup>	32,343 <sup>a</sup>	33,180 <sup>a</sup>	31,689 <sup>a</sup>	30,961 <sup>a</sup>	29,000 <sup>a</sup>	28,801 <sup>a</sup>	27,016	26,665	26,262
Smokeless Tobacco	--	--	--	--	--	--	--	--	--	--	--	--	--	4,974	4,932	4,843	4,597	4,741
Cigars	1,132 <sup>a</sup>	1,526 <sup>a</sup>	1,428 <sup>a</sup>	1,391 <sup>a</sup>	1,669 <sup>a</sup>	1,509 <sup>a</sup>	1,679	1,734	1,927	1,594 <sup>a</sup>	1,984	1,735	1,566 <sup>a</sup>	2,063	2,210	2,118	2,216	2,080
Alcohol	19,113 <sup>a</sup>	18,429 <sup>a</sup>	18,841 <sup>a</sup>	19,575 <sup>a</sup>	19,306 <sup>a</sup>	19,498 <sup>a</sup>	20,639	21,346	19,890	20,247	21,190	20,934	22,256	21,143	20,729	21,577	21,448	21,187
Binge Alcohol <sup>3</sup>	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	2,787	2,623	2,852	2,860	2,623
<b>DAILY/ALMOST DAILY USE IN PAST MONTH AMONG PAST MONTH USERS</b>																		
Marijuana	3,226 <sup>a</sup>	3,444 <sup>a</sup>	3,557 <sup>a</sup>	3,756 <sup>a</sup>	3,669 <sup>a</sup>	3,820 <sup>a</sup>	4,022 <sup>a</sup>	4,659 <sup>a</sup>	5,448 <sup>a</sup>	5,496 <sup>a</sup>	5,949 <sup>a</sup>	6,608 <sup>a</sup>	7,715 <sup>a</sup>	7,756 <sup>a</sup>	8,561 <sup>a</sup>	9,264 <sup>a</sup>	10,497 <sup>a</sup>	12,283
Cocaine	96	111	232	189	332	159	145	139	166	62 <sup>a</sup>	122	180	105	102	176	133	121	168
Crack	35	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Heroin	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	203	*	*
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	22	12	27	42	13
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	*	*	*	*
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	227 <sup>a</sup>	*	284 <sup>a</sup>	384	429
Misuse of Pain Relievers <sup>1</sup>	--	--	--	--	--	--	--	--	--	--	--	--	--	416	330	336	358	341
Misuse of Stimulants	--	--	--	--	--	--	--	--	--	--	--	--	--	75	80	103	73	92
Misuse of Tranquilizers	--	--	--	--	--	--	--	--	--	--	--	--	--	96	103	81	81	121
Misuse of Sedatives	--	--	--	--	--	--	--	--	--	--	--	--	--	*	*	*	*	*
Cigarettes (Daily Only) <sup>2</sup>	35,364 <sup>a</sup>	34,781 <sup>a</sup>	34,315 <sup>a</sup>	35,312 <sup>a</sup>	35,562 <sup>a</sup>	34,351 <sup>a</sup>	34,358 <sup>a</sup>	33,504 <sup>a</sup>	32,497 <sup>a</sup>	32,343 <sup>a</sup>	33,180 <sup>a</sup>	31,689 <sup>a</sup>	30,961 <sup>a</sup>	29,000 <sup>a</sup>	28,801 <sup>a</sup>	27,016	26,665	26,262
Smokeless Tobacco	--	--	--	--	--	--	--	--	--	--	--	--	--	4,974	4,932	4,843	4,597	4,741
Cigars	1,132 <sup>a</sup>	1,526 <sup>a</sup>	1,428 <sup>a</sup>	1,391 <sup>a</sup>	1,669 <sup>a</sup>	1,509 <sup>a</sup>	1,679	1,734	1,927	1,594 <sup>a</sup>	1,984	1,735	1,566 <sup>a</sup>	2,063	2,210	2,118	2,216	2,080
Alcohol	19,113 <sup>a</sup>	18,429 <sup>a</sup>	18,841 <sup>a</sup>	19,575 <sup>a</sup>	19,306 <sup>a</sup>	19,498 <sup>a</sup>	20,639	21,346	19,890	20,247	21,190	20,934	22,256	21,143	20,729	21,577	21,448	21,187
Binge Alcohol <sup>3</sup>	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	2,787	2,623	2,852	2,860	2,623
<b>SMOKED 1+ PACKS OF CIGARETTES PER DAY AMONG DAILY CIGARETTE USERS<sup>4</sup></b>	19,507 <sup>a</sup>	19,359 <sup>a</sup>	19,326 <sup>a</sup>	18,717 <sup>a</sup>	18,660 <sup>a</sup>	18,131 <sup>a</sup>	17,462 <sup>a</sup>	15,886 <sup>a</sup>	15,152 <sup>a</sup>	14,629 <sup>a</sup>	14,337 <sup>a</sup>	13,441 <sup>a</sup>	12,786 <sup>a</sup>	12,215 <sup>a</sup>	11,963 <sup>a</sup>	11,239	10,668	10,763

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.  
NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section 3.3.5 in Chapter 3 of the *2019 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

<sup>a</sup> The difference between this estimate and the 2019 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

<sup>1</sup> Prescription psychotherapeutic subtypes were revised in 2016; one effect was the comparability of codeine products between 2015 and 2016.

<sup>2</sup> Includes only respondents who smoked cigarettes on each of the past 30 days.

<sup>3</sup> Estimates of binge alcohol use include use by those who were heavy alcohol users.

<sup>4</sup> Respondents with missing data for number of cigarettes smoked per day were excluded from the analysis.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2019.

Table 7.32B Daily or Almost Daily Use of Specific Substances in Past Month among All Persons Aged 21 or Older and among Past Month Users of Specific Substances Aged 21 or Older: Percentages, 2002-2019

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
<b>DAILY/ALMOST DAILY USE IN PAST MONTH AMONG TOTAL POPULATION</b>																		
Marijuana	1.6 <sup>a</sup>	1.7 <sup>a</sup>	1.8 <sup>a</sup>	1.8 <sup>a</sup>	1.8 <sup>a</sup>	1.8 <sup>a</sup>	1.9 <sup>a</sup>	2.2 <sup>a</sup>	2.5 <sup>a</sup>	2.5 <sup>a</sup>	2.7 <sup>a</sup>	2.9 <sup>a</sup>	3.4 <sup>a</sup>	3.4 <sup>a</sup>	3.7 <sup>a</sup>	4.0 <sup>a</sup>	4.5 <sup>a</sup>	5.2
Cocaine	0.0	0.1	0.1	0.1	0.2 <sup>a</sup>	0.1	0.1	0.1	0.1	0.0 <sup>a</sup>	0.1	0.1	0.0	0.0	0.1	0.1	0.1	0.1
Crack	0.0	0.0	0.1	0.0	0.1	0.0	0.0	0.0	0.0	0.0 <sup>a</sup>	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Heroin	0.0	0.0 <sup>a</sup>	0.0	0.0 <sup>a</sup>	0.1	0.0	0.0	0.0 <sup>a</sup>	0.0 <sup>a</sup>	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.0	0.0	0.0	0.0	0.0
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.0	0.0	0.0	0.0	0.0
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	0.1 <sup>a</sup>	0.1 <sup>a</sup>	0.1	0.2	0.2
Misuse of Pain Relievers <sup>1</sup>	--	--	--	--	--	--	--	--	--	--	--	--	--	0.2	0.1	0.1	0.2	0.1
Misuse of Stimulants	--	--	--	--	--	--	--	--	--	--	--	--	--	0.0	0.0	0.0	0.0	0.0
Misuse of Tranquilizers	--	--	--	--	--	--	--	--	--	--	--	--	--	0.0	0.0	0.0	0.0	0.1
Misuse of Sedatives	--	--	--	--	--	--	--	--	--	--	--	--	--	0.0	0.0	0.0	*	0.0
Cigarettes (Daily Only) <sup>2</sup>	17.9 <sup>a</sup>	17.4 <sup>a</sup>	16.9 <sup>a</sup>	17.2 <sup>a</sup>	17.1 <sup>a</sup>	16.4 <sup>a</sup>	16.2 <sup>a</sup>	15.7 <sup>a</sup>	15.1 <sup>a</sup>	14.8 <sup>a</sup>	15.0 <sup>a</sup>	14.1 <sup>a</sup>	13.6 <sup>a</sup>	12.6 <sup>a</sup>	12.4 <sup>a</sup>	11.5	11.3	11.1
Smokeless Tobacco	--	--	--	--	--	--	--	--	--	--	--	--	--	2.2	2.1	2.1	1.9	2.0
Cigars	0.6 <sup>a</sup>	0.8	0.7 <sup>a</sup>	0.7 <sup>a</sup>	0.8	0.7	0.8	0.8	0.9	0.7	0.9	0.8	0.7 <sup>a</sup>	0.9	1.0	0.9	0.9	0.9
Alcohol	9.7 <sup>a</sup>	9.2	9.3	9.6	9.3	9.3	9.7 <sup>a</sup>	10.0 <sup>a</sup>	9.2	9.2	9.6	9.3	9.8 <sup>a</sup>	9.2	8.9	9.2	9.1	8.9
Binge Alcohol <sup>3</sup>	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	1.2	1.1	1.2	1.2	1.1
<b>DAILY/ALMOST DAILY USE IN PAST MONTH AMONG PAST MONTH USERS</b>																		
Marijuana	31.9 <sup>a</sup>	33.8 <sup>a</sup>	34.2 <sup>a</sup>	36.0 <sup>a</sup>	34.1 <sup>a</sup>	36.9 <sup>a</sup>	36.2 <sup>a</sup>	37.7 <sup>a</sup>	42.5	41.7 <sup>a</sup>	41.9 <sup>a</sup>	43.0	43.8	43.5	43.7	42.9	45.5	45.5
Cocaine	5.9	6.0	14.2	9.7	16.7	9.1	9.4	9.8	13.3	5.7	8.2	13.0	7.8	6.2	10.4	6.8	6.8	9.2
Crack	6.7	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Heroin	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	41.7	*	*
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	2.5	1.2	2.6	3.3 <sup>a</sup>	0.8
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	*	*	*	*
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	26.9	*	40.3	40.1	37.7
Misuse of Pain Relievers <sup>1</sup>	--	--	--	--	--	--	--	--	--	--	--	--	--	12.8	11.4	11.9	14.2	13.6
Misuse of Stimulants	--	--	--	--	--	--	--	--	--	--	--	--	--	6.0	6.0	7.0	5.2	7.3
Misuse of Tranquilizers	--	--	--	--	--	--	--	--	--	--	--	--	--	6.4	6.3	5.7	5.8	7.7
Misuse of Sedatives	--	--	--	--	--	--	--	--	--	--	--	--	--	*	*	*	*	*
Cigarettes (Daily Only) <sup>2</sup>	66.6 <sup>a</sup>	66.1 <sup>a</sup>	65.7 <sup>a</sup>	66.4 <sup>a</sup>	65.4 <sup>a</sup>	64.4 <sup>a</sup>	64.7 <sup>a</sup>	64.4 <sup>a</sup>	62.6 <sup>a</sup>	63.9 <sup>a</sup>	63.6 <sup>a</sup>	62.4 <sup>a</sup>	60.8	60.3	59.9	59.1	60.1	60.2
Smokeless Tobacco	--	--	--	--	--	--	--	--	--	--	--	--	--	62.7	63.3	63.1	64.4	61.4
Cigars	11.3 <sup>a</sup>	15.2 <sup>a</sup>	13.4 <sup>a</sup>	13.0 <sup>a</sup>	15.4 <sup>a</sup>	14.6 <sup>a</sup>	16.0 <sup>a</sup>	16.4 <sup>a</sup>	18.0	15.4 <sup>a</sup>	17.7	16.7	15.6 <sup>a</sup>	19.2	20.8	19.5	20.8	19.9
Alcohol	17.5 <sup>a</sup>	17.0	17.1 <sup>a</sup>	17.0	16.9	16.8	17.4 <sup>a</sup>	17.8 <sup>a</sup>	16.4	16.4	16.8	16.3	17.0 <sup>a</sup>	16.2	16.0	16.2	16.2	16.0
Binge Alcohol <sup>3</sup>	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	4.5	4.3	4.6	4.6	4.3
<b>SMOKED 1+ PACKS OF CIGARETTES PER DAY AMONG DAILY CIGARETTE USERS<sup>4</sup></b>	55.3 <sup>a</sup>	55.7 <sup>a</sup>	56.4 <sup>a</sup>	53.1 <sup>a</sup>	52.5 <sup>a</sup>	52.9 <sup>a</sup>	50.9 <sup>a</sup>	47.5 <sup>a</sup>	46.7 <sup>a</sup>	45.2 <sup>a</sup>	43.3	42.5	41.4	42.2	41.6	41.7	40.1	41.0

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.  
NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section 3.3.5 in Chapter 3 of the *2019 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

<sup>a</sup> The difference between this estimate and the 2019 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

<sup>1</sup> Prescription psychotherapeutic subtypes were revised in 2016; one effect was the comparability of codeine products between 2015 and 2016.

<sup>2</sup> Includes only respondents who smoked cigarettes on each of the past 30 days.

<sup>3</sup> Estimates of binge alcohol use include use by those who were heavy alcohol users.

<sup>4</sup> Respondents with missing data for number of cigarettes smoked per day were excluded from the analysis.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2019.