

**Table 8.38A Major Depressive Episode (MDE) or MDE with Severe Impairment in Past Year among Persons Aged 18 or Older, and Receipt of Treatment for Depression in Past Year among Persons Aged 18 or Older with MDE or MDE with Severe Impairment in Past Year, by Gender and Age Group: Numbers in Thousands, 2018 and 2019**

Gender/Age Group	Had MDE (2018)	Had MDE (2019)	Had MDE with Severe Impairment <sup>1</sup> (2018)	Had MDE with Severe Impairment <sup>1</sup> (2019)	Received Treatment for Depression in Past Year among Persons with MDE <sup>2</sup> (2018)	Received Treatment for Depression in Past Year among Persons with MDE <sup>2</sup> (2019)	Received Treatment for Depression in Past Year among Persons with MDE with Severe Impairment <sup>1,2</sup> (2018)	Received Treatment for Depression in Past Year among Persons with MDE with Severe Impairment <sup>1,2</sup> (2019)
<b>TOTAL</b>	17,720 <sup>a</sup>	19,357	11,489 <sup>a</sup>	13,112	11,462 <sup>a</sup>	12,824	7,878 <sup>a</sup>	9,467
18-25	4,609 <sup>a</sup>	5,045	2,968 <sup>a</sup>	3,407	2,286 <sup>a</sup>	2,565	1,594 <sup>a</sup>	1,921
26 or Older	13,111 <sup>a</sup>	14,312	8,520 <sup>a</sup>	9,705	9,176 <sup>a</sup>	10,258	6,283 <sup>a</sup>	7,546
26-49	7,990 <sup>a</sup>	8,914	5,284 <sup>a</sup>	6,101	5,142 <sup>a</sup>	6,135	3,593 <sup>a</sup>	4,532
50 or Older	5,121	5,399	3,236	3,604	4,034	4,124	2,691	3,014
<b>MALE</b>	6,272 <sup>a</sup>	7,125	4,013 <sup>a</sup>	4,646	3,548	4,041	2,464	2,868
18-25	1,665	1,841	1,087	1,236	674	793	496	599
26 or Older	4,607 <sup>a</sup>	5,284	2,926	3,409	2,875	3,248	1,969	2,268
26-49	2,956	3,271	1,980	2,169	1,660	1,927	1,220	1,369
50 or Older	1,651	2,013	946	1,240	1,215	1,321	*	899
<b>FEMALE</b>	11,447	12,233	7,476 <sup>a</sup>	8,466	7,914 <sup>a</sup>	8,783	5,414 <sup>a</sup>	6,599
18-25	2,943 <sup>a</sup>	3,204	1,881 <sup>a</sup>	2,170	1,612	1,772	1,099 <sup>a</sup>	1,321
26 or Older	8,504	9,028	5,595 <sup>a</sup>	6,296	6,302 <sup>a</sup>	7,011	4,315 <sup>a</sup>	5,278
26-49	5,034 <sup>a</sup>	5,643	3,304 <sup>a</sup>	3,932	3,482 <sup>a</sup>	4,208	2,372 <sup>a</sup>	3,163
50 or Older	3,470	3,386	2,291	2,364	2,819	2,802	1,942	2,115

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

<sup>a</sup> The difference between this estimate and the 2019 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

<sup>1</sup> Respondents with unknown impairment data were excluded.

<sup>2</sup> Respondents with unknown past year treatment for depression data were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

**Table 8.38B Major Depressive Episode (MDE) or MDE with Severe Impairment in Past Year among Persons Aged 18 or Older, and Receipt of Treatment for Depression in Past Year among Persons Aged 18 or Older with MDE or MDE with Severe Impairment in Past Year, by Gender and Age Group: Percentages, 2018 and 2019**

Gender/Age Group	Had MDE (2018)	Had MDE (2019)	Had MDE with Severe Impairment <sup>1</sup> (2018)	Had MDE with Severe Impairment <sup>1</sup> (2019)	Received Treatment for Depression in Past Year among Persons with MDE <sup>2</sup> (2018)	Received Treatment for Depression in Past Year among Persons with MDE <sup>2</sup> (2019)	Received Treatment for Depression in Past Year among Persons with MDE with Severe Impairment <sup>1,2</sup> (2018)	Received Treatment for Depression in Past Year among Persons with MDE with Severe Impairment <sup>1,2</sup> (2019)
<b>TOTAL</b>	7.2 <sup>a</sup>	7.8	4.7 <sup>a</sup>	5.3	64.8	66.3	68.6 <sup>a</sup>	72.2
18-25	13.8 <sup>a</sup>	15.2	8.9 <sup>a</sup>	10.3	49.6	50.9	53.7	56.4
26 or Older	6.2 <sup>a</sup>	6.7	4.0 <sup>a</sup>	4.5	70.1	71.8	73.8 <sup>a</sup>	77.8
26-49	8.0 <sup>a</sup>	8.9	5.3 <sup>a</sup>	6.1	64.4 <sup>a</sup>	68.9	68.0 <sup>a</sup>	74.4
50 or Older	4.5	4.7	2.9	3.2	78.9	76.5	83.1	83.6
<b>MALE</b>	5.3 <sup>a</sup>	6.0	3.4 <sup>a</sup>	3.9	56.7	56.8	61.4	61.8
18-25	9.9	11.1	6.5 <sup>a</sup>	7.4	40.5	43.1	45.6	48.5
26 or Older	4.5 <sup>a</sup>	5.1	2.9	3.3	62.6	61.6	67.3	66.7
26-49	6.0	6.6	4.0	4.4	56.3	59.2	61.7	63.3
50 or Older	3.1	3.8	1.8	2.3	73.9	65.6	*	72.5
<b>FEMALE</b>	9.0	9.6	5.9 <sup>a</sup>	6.6	69.2	71.9	72.4 <sup>a</sup>	78.0
18-25	17.6 <sup>a</sup>	19.4	11.3 <sup>a</sup>	13.1	54.8	55.3	58.4	60.9
26 or Older	7.7	8.1	5.1	5.7	74.1	77.7	77.1 <sup>a</sup>	83.8
26-49	10.0 <sup>a</sup>	11.1	6.5 <sup>a</sup>	7.8	69.2 <sup>a</sup>	74.6	71.8 <sup>a</sup>	80.4
50 or Older	5.8	5.6	3.8	3.9	81.3	83.0	84.8	89.5

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

<sup>a</sup> The difference between this estimate and the 2019 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

<sup>1</sup> Respondents with unknown impairment data were excluded.

<sup>2</sup> Respondents with unknown past year treatment for depression data were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.