

Table 9.5A Major Depressive Episode (MDE) or MDE with Severe Impairment in Past Year among Persons Aged 12 to 17, and Receipt of Treatment for Depression in Past Year among Persons Aged 12 to 17 with MDE or MDE with Severe Impairment in Past Year, by Demographic Characteristics: Numbers in Thousands, 2018 and 2019

Demographic Characteristic	Had MDE (2018)	Had MDE (2019)	Had MDE with Severe Impairment ¹ (2018)	Had MDE with Severe Impairment ¹ (2019)	Received Treatment for Depression in Past Year among Persons with MDE ² (2018)	Received Treatment for Depression in Past Year among Persons with MDE ² (2019)	Received Treatment for Depression in Past Year among Persons with MDE with Severe Impairment ^{1,2} (2018)	Received Treatment for Depression in Past Year among Persons with MDE with Severe Impairment ^{1,2} (2019)
TOTAL	3,482 ^a	3,783	2,423 ^a	2,671	1,432 ^a	1,627	1,133 ^a	1,321
AGE GROUP								
12-13	632 ^a	816	382 ^a	530	251	302	177	218
14-15	1,250	1,341	891	970	514	585	406	492
16-17	1,600	1,626	1,150	1,171	667	740	550	612
GENDER								
Male	946 ^a	1,076	628 ^a	736	351	391	274	300
Female	2,537	2,708	1,795	1,935	1,081 ^a	1,236	859 ^a	1,021
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	2,601	2,755	1,841	1,981	1,100 ^a	1,251	882 ^a	1,018
White	1,905	1,975	1,354	1,415	870	985	695	800
Black or African American	334	367	225	256	115	130	90	108
AIAN	23	20	*	19	*	*	*	*
NHOPI	*	*	*	*	*	*	*	*
Asian	189	190	127	142	*	*	*	*
Two or More Races	137 ^a	189	109	134	48 ^a	83	*	63
Hispanic or Latino	882 ^a	1,028	582	690	332	376	251	304

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

^a The difference between this estimate and the 2019 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

¹ Respondents with unknown impairment data were excluded.

² Respondents with unknown past year treatment for depression data were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

Table 9.5B Major Depressive Episode (MDE) or MDE with Severe Impairment in Past Year among Persons Aged 12 to 17, and Receipt of Treatment for Depression in Past Year among Persons Aged 12 to 17 with MDE or MDE with Severe Impairment in Past Year, by Demographic Characteristics: Percentages, 2018 and 2019

Demographic Characteristic	Had MDE (2018)	Had MDE (2019)	Had MDE with Severe Impairment ¹ (2018)	Had MDE with Severe Impairment ¹ (2019)	Received Treatment for Depression in Past Year among Persons with MDE ² (2018)	Received Treatment for Depression in Past Year among Persons with MDE ² (2019)	Received Treatment for Depression in Past Year among Persons with MDE with Severe Impairment ^{1,2} (2018)	Received Treatment for Depression in Past Year among Persons with MDE with Severe Impairment ^{1,2} (2019)
TOTAL	14.4 ^a	15.7	10.0 ^a	11.1	41.4	43.3	46.9	49.7
AGE GROUP								
12-13	8.4 ^a	10.5	5.1 ^a	6.8	40.7	37.5	47.2	41.9
14-15	15.3	16.4	10.9	11.9	41.2	43.9	45.6	50.8
16-17	19.0	20.1	13.7	14.5	41.8	45.6	47.8	52.3
GENDER								
Male	7.7 ^a	8.8	5.1 ^a	6.0	37.5	36.8	43.7	41.2
Female	21.5	23.0	15.2	16.5	42.9	45.8	48.1	52.9
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	14.2	15.2	10.1	11.0	42.6	45.7	48.1	51.7
White	15.1	15.9	10.7	11.4	46.1	50.3	51.6	57.0
Black or African American	10.3	11.4	6.9	8.0	34.6	35.6	40.1	42.3
AIAN	15.2	12.2	*	11.5	*	*	*	*
NHOPI	*	*	*	*	*	*	*	*
Asian	13.6	15.1	9.2	11.3	*	*	*	*
Two or More Races	17.7	20.9	14.2	14.9	34.8	44.4	*	47.3
Hispanic or Latino	15.1	17.3	10.0	11.7	37.9	36.8	43.2	44.0

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

^a The difference between this estimate and the 2019 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

¹ Respondents with unknown impairment data were excluded.

² Respondents with unknown past year treatment for depression data were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.