

Mental Health Annual Report 2015–2020

Use of Mental Health Services: National Client-Level Data

DEPARTMENT OF HEALTH AND HUMAN SERVICES
Substance Abuse and Mental Health Services Administration

Acknowledgments

This report was prepared for the Substance Abuse and Mental Health Services Administration (SAMHSA), U.S. Department of Health and Human Services (HHS) under Contract No. 75S20320C00001. The Contracting Officer's Representative (COR) at SAMHSA's Center for Behavioral Health Statistics and Quality (CBHSQ) was Heydy Juarez.

SAMHSA complies with applicable federal civil rights laws and does not discriminate on the basis of race, color, national origin, age, disability, or sex. *SAMHSA cumple con las leyes federales de derechos civiles aplicables y no discrimina por motivos de raza, color, nacionalidad, edad, discapacidad o sexo.*

Public Domain Notice

All material appearing in this report is in the public domain and may be reproduced or copied without permission from SAMHSA. Citation of the source is appreciated. However, this publication may not be reproduced or distributed for a fee without the specific, written authorization of the Office of Communications, SAMHSA, HHS.

Recommended Citation

Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality. *Mental Health Annual Report: 2015–2020. Use of Mental Health Services: National Client-Level Data*. Rockville, MD: Substance Abuse and Mental Health Services Administration, 2022.

Electronic Access and Copies of Publication

This publication may be downloaded at
<https://www.samhsa.gov/data/data-we-collect/mh-cld-mental-health-client-level-data>

Originating Office

Center for Behavioral Health Statistics and Quality
Substance Abuse and Mental Health Services Administration
5600 Fishers Lane, Room 15SEH03
Rockville, Maryland 20857

October 2022

Table of Contents

List of Tables.....	ix
List of Figures.....	xxxv
Executive Summary.....	1
Background	5
Introduction	9
Chapter 1. Summary of Annual Data in State Client-Level Data Reporting: 2015–2020 ...	13
Individuals Served	13
Regional Distributions	16
Any-Mention Mental Health Diagnoses.....	17
Chapter 2. Characteristics of Children and Adolescents Aged 17 Years and Younger Receiving Mental Health Services.....	19
Individuals Served	19
Gender.....	19
Race	19
Ethnicity.....	20
Living Arrangements	20
SED Status	20
Client Treatment Status at the Start of the Reporting Period.....	20
SED Status, by Service Setting.....	20
Chapter 3. Characteristics of Young Adults Aged 18 to 24 Years Receiving Mental Health Services.....	21
Individuals Served	21
Gender.....	21
Race and Ethnicity	21
Education	22
Marital Status.....	22
Living Arrangements	22
SMI Status.....	22
Veteran Status	22
Employment Status	22
Client Treatment Status at the Start of the Reporting Period.....	22
Service Setting.....	23

SMI Status, by Service Setting	23
Chapter 4. Characteristics of Adults Aged 25 to 54 Years Receiving Mental Health Services	25
Individuals Served	25
Gender	25
Race	25
Ethnicity	26
Education	26
Marital Status	26
Living Arrangements	26
SMI Status	26
Veteran Status	26
Employment Status	27
Client Treatment Status at the Start of the Reporting Period	27
Service Setting	27
SMI Status, by Service Setting	27
Chapter 5. Characteristics of Older Adults Aged 55 Years and Older Receiving Mental Health Services	29
Individuals Served	29
Gender	29
Race	29
Ethnicity	30
Education	30
Marital Status	30
Living Arrangements	30
SMI Status	30
Veteran Status	30
Employment Status	31
Client Treatment Status at the Start of the Reporting Period	31
Service Setting	31
SMI Status, by Service Setting	31
Chapter 6. Characteristics of Individuals With Co-occurring Mental Health and Substance Use Disorders Receiving Mental Health Services	33
Individuals Served	33
Gender	33
Age	33
Race	34
Ethnicity	34
Substance Use Diagnosis	34
Living Arrangements	34
SMI/SED Status	34

Veteran Status	35
Employment Status	35
Service Setting.....	35
Client Treatment Status at the Start of the Reporting Period.....	35
Chapter 7. Client Outcomes Among Individuals Receiving Mental Health Services: 2015–2020	37
Employment Status	37
Transitions in Living Arrangements	38
Transitions in Employment Status	41
Chapter 8. Geographic and Clinical Trends: 2015–2020	45
Chapter 9. Client Characteristics by State or Jurisdiction: 2015–2020	47
Chapter 10. Client Characteristics by State or Jurisdiction: 2020	49
Tables.....	51
Appendix A. MH-TEDS/MH-CLD Crosswalk Tables.....	789
Appendix B. MH-TEDS and MH-CLD 2011–2020 reporting methods and 2020 reporting period, by state or jurisdiction.....	799
Appendix C. MH-TEDS and MH-CLD Data Dictionary of Included Variables	801
Appendix D. SMI/SED Federal Definitions.....	813
Serious Mental Illness (SMI).....	813
Serious Emotional Disturbance (SED).....	813
Appendix E. Mental Health and Substance Use Diagnosis Codes.....	815
Mental Health Diagnostic Groups With Corresponding ICD-9 Diagnostic Codes.....	815
Mental Health Diagnostic Groups with Corresponding ICD-10 Diagnostic Codes.....	820
TEDS Substance Use Diagnosis Codes	827
Appendix F. Terms and Acronyms	829
Appendix G. List of Contributors	831

List of Tables

Summary of Annual Data in State Client-Level Data Reporting: 2015–2020

1.1a	Individuals, by mental health diagnosis: Number, 2015–2020	52
1.1b	Individuals, by mental health diagnosis: Column percent distribution, 2015–2020	53
1.2a	Individuals, by gender and age: Number, 2015–2020	54
1.2b	Individuals, by gender and age: Column percent distribution, 2015–2020	55
1.3a	Individuals, by race and ethnicity: Number, 2015–2020	56
1.3b	Individuals, by race and ethnicity: Column percent distribution, 2015–2020	57
1.4a	Individuals aged 17 years and younger, by mental health diagnosis: Number, 2015–2020	58
1.4b	Individuals aged 17 years and younger, by mental health diagnosis: Column percent distribution, 2015–2020	59
1.5a	Individuals aged 18 to 24 years, by mental health diagnosis: Number, 2015–2020	60
1.5b	Individuals aged 18 to 24 years, by mental health diagnosis: Column percent distribution, 2015–2020	61
1.6a	Individuals aged 25 to 54 years, by mental health diagnosis: Number, 2015–2020	62
1.6b	Individuals aged 25 to 54 years, by mental health diagnosis: Column percent distribution, 2015–2020	63
1.7a	Individuals aged 55 years and older, by mental health diagnosis: Number, 2015–2020	64
1.7b	Individuals aged 55 years and older, by mental health diagnosis: Column percent distribution, 2015–2020	65
1.8a	Individuals, by marital status and living arrangement: Number, 2015–2020	66
1.8b	Individuals, by marital status and living arrangement: Column percent distribution, 2015–2020	67
1.9a	Individuals, by SMI/SED status and co-occurring mental health and substance use disorder status: Number, 2015–2020	68
1.9b	Individuals, by SMI/SED status and co-occurring mental health and substance use disorder status: Column percent distribution, 2015–2020	69
1.10a	Individuals aged 18 years and older, by employment status, ‘detailed not in labor force,’ education, and veteran status: Number, 2015–2020	70

List of Tables (Continued)

1.10b	Individuals aged 18 years and older, by employment status, ‘detailed not in labor force,’ education, and veteran status: Column percent distribution, 2015–2020	71
1.11a	Individuals, by timing of admission and mental health service setting: Number, 2015–2020	72
1.11b	Individuals, by timing of admission and mental health service setting: Column percent distribution, 2015–2020	73
1.12a	Individuals, by number of mental health diagnosis and any mention of mental health diagnosis: Number, 2015–2020	74
1.12b	Individuals, by number of mental health diagnosis and any mention of mental health diagnosis: Column percent distribution, 2015–2020	75
1.13a	Individuals, by Census region and division: Number, 2015–2020	76
1.13b	Individuals, by Census region and division: Column percent distribution, 2015–2020 ..	77
1.14a	Individuals with diagnosis of trauma- and stressor-related disorders, by gender, age, race, and ethnicity: Number, 2015–2020	78
1.14b	Individuals with diagnosis of trauma- and stressor-related disorders, by gender, age, race, and ethnicity: Column percent distribution, 2015–2020	80
1.15a	Individuals with diagnosis of anxiety disorders, by gender, age, race, and ethnicity: Number, 2015–2020	82
1.15b	Individuals with diagnosis of anxiety disorders, by gender, age, race, and ethnicity: Column percent distribution, 2015–2020	84
1.16a	Individuals with diagnosis of attention deficit/hyperactivity disorder, by gender, age, race, and ethnicity: Number, 2015–2020	86
1.16b	Individuals with diagnosis of attention deficit/hyperactivity disorder, by gender, age, race, and ethnicity: Column percent distribution, 2015–2020	88
1.17a	Individuals with diagnosis of bipolar disorders, by gender, age, race, and ethnicity: Number, 2015–2020	90
1.17b	Individuals with diagnosis of bipolar disorders, by gender, age, race, and ethnicity: Column percent distribution, 2015–2020	92
1.18a	Individuals with diagnosis of depressive disorders, by gender, age, race, and ethnicity: Number, 2015–2020	94
1.18b	Individuals with diagnosis of depressive disorders, by gender, age, race, and ethnicity: Column percent distribution, 2015–2020	96
1.19a	Individuals with diagnosis of schizophrenia and other psychotic disorders, by gender, age, race, and ethnicity: Number, 2015–2020	98

List of Tables (Continued)

1.19b	Individuals with diagnosis of schizophrenia and other psychotic disorders, by gender, age, race, and ethnicity: Column percent distribution, 2015–2020	100
1.20a	Individuals with co-occurring mental health and substance use disorders, by gender, age, race, and ethnicity: Number, 2015–2020	102
1.20b	Individuals with co-occurring mental health and substance use disorders, by gender, age, race, and ethnicity: Column percent distribution, 2015–2020	104
1.21a	Individuals without co-occurring mental health and substance use disorders, by gender, age, race, and ethnicity: Number, 2015–2020	106
1.21b	Individuals without co-occurring mental health and substance use disorders, by gender, age, race, and ethnicity: Column percent distribution, 2015–2020	108
1.22a	Individuals who received mental health services and U.S. population, by gender, marital status, and age group: Number, 2020	110
1.22b	Individuals who received mental health services and U.S. population, by gender, marital status, and age group: Column percent distribution, 2020	111
1.22c	Individuals who received mental health services and U.S. population, by gender, marital status, and age group: Row percent distribution, 2020	112
1.23a	Individuals who received mental health services and U.S. population, by race, ethnicity, and age group: Number, 2020	113
1.23b	Individuals who received mental health services and U.S. population, by race, ethnicity, and age group: Column percent distribution, 2020	114
1.23c	Individuals who received mental health services and U.S. population, by race, ethnicity, and age group: Row percent distribution, 2020	115
Characteristics of Children and Adolescents Aged 17 Years and Younger Receiving Mental Health Services		
2.1a	Individuals aged 17 years and younger, by gender, age, race, ethnicity, and SED status: Number, 2015–2020	116
2.1b	Individuals aged 17 years and younger, by gender, age, race, ethnicity, and SED status: Column percent distribution, 2015–2020	118
2.2a	Individuals aged 17 years and younger with SED, by gender, age, race, and ethnicity: Number, 2015–2020	120
2.2b	Individuals aged 17 years and younger with SED, by gender, age, race, and ethnicity: Column percent distribution, 2015–2020	122
2.3a	Individuals aged 17 years and younger, by mental health diagnosis, gender, race, and ethnicity: Number, 2020	124

List of Tables (Continued)

2.3b	Individuals aged 17 years and younger, by mental health diagnosis, gender, race, and ethnicity: Column percent distribution, 2020	125
2.3c	Individuals aged 17 years and younger, by mental health diagnosis, gender, race, and ethnicity: Row percent distribution, 2020	126
2.4a	Individuals aged 17 years and younger with SED, by mental health diagnosis, gender, race, and ethnicity: Number, 2020.....	127
2.4b	Individuals aged 17 years and younger with SED, by mental health diagnosis, gender, race, and ethnicity: Column percent distribution, 2020	128
2.4c	Individuals aged 17 years and younger with SED, by mental health diagnosis, gender, race, and ethnicity: Row percent distribution, 2020	129
2.5a	Individuals aged 17 years and younger, by mental health diagnosis, living arrangement, and SED status: Number, 2020	130
2.5b	Individuals aged 17 years and younger, by mental health diagnosis, living arrangement, and SED status: Column percent distribution, 2020	131
2.5c	Individuals aged 17 years and younger, by mental health diagnosis, living arrangement, and SED status: Row percent distribution, 2020	132
2.6a	Individuals aged 17 years and younger with SED, by mental health diagnosis and living arrangement: Number, 2020	133
2.6b	Individuals aged 17 years and younger with SED, by mental health diagnosis and living arrangement: Column percent distribution, 2020	134
2.6c	Individuals aged 17 years and younger with SED, by mental health diagnosis and living arrangement: Row percent distribution, 2020	135
2.7a	Individuals aged 17 years and younger, by mental health diagnosis, timing of admission, and mental health service setting: Number, 2020	136
2.7b	Individuals aged 17 years and younger, by mental health diagnosis, timing of admission, and mental health service setting: Column percent, 2020	137
2.7c	Individuals aged 17 years and younger, by mental health diagnosis, timing of admission, and mental health service setting: Row percent distribution, 2020	138
2.8a	Individuals aged 17 years and younger with SED, by mental health diagnosis, timing of admission, and mental health service setting: Number, 2020	139
2.8b	Individuals aged 17 years and younger with SED, by mental health diagnosis, timing of admission, and mental health service setting: Column percent, 2020	140
2.8c	Individuals aged 17 years and younger with SED, by mental health diagnosis, timing of admission, and mental health service setting: Row percent distribution, 2020	141

List of Tables (Continued)

2.9	Individuals aged 17 years and younger, by mental health service setting and SED status: Number and percent distribution, 2020	142
Characteristics of Young Adults Aged 18 to 24 Years Receiving Mental Health Services		
3.1a	Individuals aged 18 to 24 years, by gender, age, race, ethnicity, education, veteran status, and SMI status: Number, 2015–2020	143
3.1b	Individuals aged 18 to 24 years, by gender, age, race, ethnicity, education, veteran status, and SMI status: Column percent distribution, 2015–2020	145
3.2a	Individuals aged 18 to 24 years with SMI, by gender, age, race, ethnicity, education, and veteran status: Number, 2015–2020	147
3.2b	Individuals aged 18 to 24 years with SMI, by gender, age, race, ethnicity, education, and veteran status: Column percent distribution, 2015–2020	149
3.3a	Individuals aged 18 to 24 years, by mental health diagnosis, gender, race, and ethnicity: Number, 2020	151
3.3b	Individuals aged 18 to 24 years, by mental health diagnosis, gender, race, and ethnicity: Column percent distribution, 2020	152
3.3c	Individuals aged 18 to 24 years, by mental health diagnosis, gender, race, and ethnicity: Row percent distribution, 2020	153
3.4a	Individuals aged 18 to 24 years with SMI, by mental health diagnosis, gender, race, and ethnicity: Number, 2020	154
3.4b	Individuals aged 18 to 24 years with SMI, by mental health diagnosis, gender, race, and ethnicity: Column percent distribution, 2020	155
3.4c	Individuals aged 18 to 24 years with SMI, by mental health diagnosis, gender, race, and ethnicity: Row percent distribution, 2020	156
3.5a	Individuals aged 18 to 24 years, by mental health diagnosis, education, and marital status: Number, 2020	157
3.5b	Individuals aged 18 to 24 years, by mental health diagnosis, education, and marital status: Column percent distribution, 2020	158
3.5c	Individuals aged 18 to 24 years, by mental health diagnosis, education, and marital status: Row percent distribution, 2020	159
3.6a	Individuals aged 18 to 24 years with SMI, by mental health diagnosis, education, and marital status: Number, 2020	160
3.6b	Individuals aged 18 to 24 years with SMI, by mental health diagnosis, education, and marital status: Column percent distribution, 2020	161
3.6c	Individuals aged 18 to 24 years with SMI, by mental health diagnosis, education, marital status: Row percent distribution, 2020	162

List of Tables (Continued)

3.7a	Individuals aged 18 to 24 years, by mental health diagnosis, living arrangement, SMI status, and veteran status: Number, 2020	163
3.7b	Individuals aged 18 to 24 years, by mental health diagnosis, living arrangement, SMI status, and veteran status: Column percent distribution, 2020	164
3.7c	Individuals aged 18 to 24 years, by mental health diagnosis, living arrangement, SMI status, and veteran status: Row percent distribution, 2020	165
3.8a	Individuals aged 18 to 24 years with SMI, by mental health diagnosis, living arrangement and veteran status: Number, 2020	166
3.8b	Individuals aged 18 to 24 years with SMI, by mental health diagnosis, living arrangement and veteran status: Column percent distribution, 2020	167
3.8c	Individuals aged 18 to 24 years with SMI, by mental health diagnosis, living arrangement and veteran status: Row percent distribution, 2020	168
3.9a	Individuals aged 18 to 24 years, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Number, 2020	169
3.9b	Individuals aged 18 to 24 years, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Column percent distribution, 2020	170
3.9c	Individuals aged 18 to 24 years, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Row percent distribution, 2020	171
3.10a	Individuals aged 18 to 24 years with SMI, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Number, 2020	172
3.10b	Individuals aged 18 to 24 years with SMI, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Column percent distribution, 2020	173
3.10c	Individuals aged 18 to 24 years with SMI, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Row percent distribution, 2020	174
3.11a	Individuals aged 18 to 24 years, by mental health diagnosis, timing of admission, and mental health service setting: Number, 2020	175
3.11b	Individuals aged 18 to 24 years, by mental health diagnosis, timing of admission, and mental health service setting: Column percent distribution, 2020	176
3.11c	Individuals aged 18 to 24 years, by mental health diagnosis, timing of admission, and mental health service setting: Row percent distribution, 2020	177
3.12a	Individuals aged 18 to 24 years with SMI, by mental health diagnosis, timing of admission, and mental health service setting: Number, 2020	178
3.12b	Individuals aged 18 to 24 years with SMI, by mental health diagnosis, timing of admission, and mental health service setting: Column percent distribution, 2020	179

List of Tables (Continued)

3.12c	Individuals aged 18 to 24 years with SMI, by mental health diagnosis, timing of admission, and mental health service setting: Row percent distribution, 2020	180
3.13	Individuals aged 18 to 24 years, by mental health service setting and SMI status: Number and percent distribution, 2020	181

Characteristics of Adults Aged 25 to 54 Years Receiving Mental Health Services

4.1a	Individuals aged 25 to 54 years, by gender, age, race, ethnicity, education, veteran status, and SMI status: Number, 2015–2020	182
4.1b	Individuals aged 25 to 54 years, by gender, age, race, ethnicity, education, veteran status, and SMI status: Column percent distribution, 2015–2020	184
4.2a	Individuals aged 25 to 54 years with SMI, by gender, age, race, ethnicity, education, and veteran status: Number, 2015–2020	186
4.2b	Individuals aged 25 to 54 years with SMI, by gender, age, race, ethnicity, education, and veteran status: Column percent distribution, 2015–2020	188
4.3a	Individuals aged 25 to 54 years, by mental health diagnosis, gender, race, and ethnicity: Number, 2020	190
4.3b	Individuals aged 25 to 54 years, by mental health diagnosis, gender, race, and ethnicity: Column percent distribution, 2020	191
4.3c	Individuals aged 25 to 54 years, by mental health diagnosis, gender, race, and ethnicity: Row percent distribution, 2020	192
4.4a	Individuals aged 25 to 54 years with SMI, gender, race, and ethnicity: Number, 2020	193
4.4b	Individuals aged 25 to 54 years with SMI, by mental health diagnosis, gender, race, and ethnicity: Column percent distribution, 2020	194
4.4c	Individuals aged 25 to 54 years with SMI, by mental health diagnosis, gender, race, and ethnicity: Row percent distribution, 2020	195
4.5a	Individuals aged 25 to 54 years, by mental health diagnosis, education, and marital status: Number, 2020	196
4.5b	Individuals aged 25 to 54 years, by mental health diagnosis, education, and marital status: Column percent distribution, 2020	197
4.5c	Individuals aged 25 to 54 years, by mental health diagnosis, education, and marital status: Row percent distribution, 2020	198
4.6a	Individuals aged 25 to 54 years with SMI, by mental health diagnosis, education, and marital status: Number, 2020	199
4.6b	Individuals aged 25 to 54 years with SMI, by mental health diagnosis, education, and marital status: Column percent distribution, 2020	200

List of Tables (Continued)

4.6c	Individuals aged 25 to 54 years with SMI, by mental health diagnosis, education, and marital status: Row percent distribution, 2020	201
4.7a	Individuals aged 25 to 54 years, by mental health diagnosis, living arrangement, SMI status, and veteran status: Number, 2020	202
4.7b	Individuals aged 25 to 54 years, by mental health diagnosis, living arrangement, SMI status, and veteran status: Column percent distribution, 2020	203
4.7c	Individuals aged 25 to 54 years, by mental health diagnosis, living arrangement, SMI status, and veteran status: Row percent distribution, 2020	204
4.8a	Individuals aged 25 to 54 years with SMI, by mental health diagnosis, living arrangement, and veteran status: Number, 2020	205
4.8b	Individuals aged 25 to 54 years with SMI, by mental health diagnosis, living arrangement, and veteran status: Column percent distribution, 2020	206
4.8c	Individuals aged 25 to 54 years with SMI, by mental health diagnosis, living arrangement, and veteran status: Row percent distribution, 2020	207
4.9a	Individuals aged 25 to 54 years, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Number, 2020	208
4.9b	Individuals aged 25 to 54 years, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Column percent distribution, 2020	209
4.9c	Individuals aged 25 to 54 years, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Row percent distribution, 2020	210
4.10a	Individuals aged 25 to 54 years with SMI, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Number, 2020	211
4.10b	Individuals aged 25 to 54 years with SMI, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Column percent distribution, 2020	212
4.10c	Individuals aged 25 to 54 years with SMI, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Row percent distribution, 2020	213
4.11a	Individuals aged 25 to 54 years, by mental health diagnosis, timing of admission, and mental health service setting: Number, 2020	214
4.11b	Individuals aged 25 to 54 years, by mental health diagnosis, timing of admission, and mental health service setting: Column percent distribution, 2020	215
4.11c	Individuals aged 25 to 54 years, by mental health diagnosis, timing of admission, and mental health service setting: Row percent distribution, 2020	216
4.12a	Individuals aged 25 to 54 years with SMI, by mental health diagnosis, timing of admission, and mental health service setting: Number, 2020	217

List of Tables (Continued)

4.12b	Individuals aged 25 to 54 years with SMI, by mental health diagnosis, timing of admission, and mental health service setting: Column percent, 2020	218
4.12c	Individuals aged 25 to 54 years with SMI, by mental health diagnosis, timing of admission, and mental health service setting: Row percent distribution, 2020	219
4.13	Individuals aged 25 to 54 years, by mental health service setting and SMI status: Number and percent distribution, 2020	220

Characteristics of Older Adults Aged 55 Years and Older Receiving Mental Health Services

5.1a	Individuals aged 55 years and older, by gender, age, race, ethnicity, education, veteran status, and SMI status: Number, 2015–2020	221
5.1b	Individuals aged 55 years and older, by gender, age, race, ethnicity, education, veteran status, and SMI status: Column percent distribution, 2015–2020 ...	223
5.2a	Individuals aged 55 years and older with SMI, by gender, age, race, ethnicity, education, and veteran status: Number, 2015–2020	225
5.2b	Individuals aged 55 years and older with SMI, by gender, age, race, ethnicity, education, and veteran status: Column percent distribution, 2015–2020	227
5.3a	Individuals aged 55 years and older, by mental health diagnosis, gender, race, and ethnicity: Number, 2020	229
5.3b	Individuals aged 55 years and older, by mental health diagnosis, gender, race, and ethnicity: Column percent distribution, 2020	230
5.3c	Individuals aged 55 years and older, by mental health diagnosis, gender, race, and ethnicity: Row percent distribution, 2020	231
5.4a	Individuals aged 55 years and older with SMI, by mental health diagnosis, gender, race, and ethnicity: Number, 2020	232
5.4b	Individuals aged 55 years and older with SMI, by mental health diagnosis, gender, race, and ethnicity: Column percent distribution, 2020	233
5.4c	Individuals aged 55 years and older with SMI, by mental health diagnosis, gender, race, and ethnicity: Row percent distribution, 2020	234
5.5a	Individuals aged 55 years and older, by mental health diagnosis, education, and marital status: Number, 2020	235
5.5b	Individuals aged 55 years and older, by mental health diagnosis, education, and marital status: Column percent distribution, 2020	236
5.5c	Individuals aged 55 years and older, by mental health diagnosis, education, and marital status: Row percent distribution, 2020	237
5.6a	Individuals aged 55 years and older with SMI, by mental health diagnosis, education, and marital status: Number, 2020	238

List of Tables (Continued)

5.6b	Individuals aged 55 years and older with SMI, by mental health diagnosis, education, and marital status: Column percent distribution, 2020	239
5.6c	Individuals aged 55 years and older with SMI, by mental health diagnosis, education, and marital status: Row percent distribution, 2020	240
5.7a	Individuals aged 55 years and older, by mental health diagnosis, living arrangement, SMI status, and veteran status: Number, 2020	241
5.7b	Individuals aged 55 years and older, by mental health diagnosis, living arrangement, SMI status, and veteran status: Column percent distribution, 2020	242
5.7c	Individuals aged 55 years and older, by mental health diagnosis, living arrangement, SMI status, and veteran status: Row percent distribution, 2020	243
5.8a	Individuals aged 55 years and older with SMI, by mental health diagnosis, living arrangement, and veteran status: Number, 2020	244
5.8b	Individuals aged 55 years and older with SMI, by mental health diagnosis, living arrangement, and veteran status: Column percent distribution, 2020	245
5.8c	Individuals aged 55 years and older with SMI, by mental health diagnosis, living arrangement, and veteran status: Row percent distribution, 2020	246
5.9a	Individuals aged 55 years and older, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Number, 2020	247
5.9b	Individuals aged 55 years and older, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Column percent distribution, 2020	248
5.9c	Individuals aged 55 years and older, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Row percent distribution, 2020	249
5.10a	Individuals aged 55 years and older with SMI, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Number, 2020	250
5.10b	Individuals aged 55 years and older with SMI, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Column percent distribution, 2020	251
5.10c	Individuals aged 55 years and older with SMI, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Row percent distribution, 2020	252
5.11a	Individuals aged 55 years and older, by mental health diagnosis, timing of admission, and mental health service setting: Number, 2020	253
5.11b	Individuals aged 55 years and older, by mental health diagnosis, timing of admission, and mental health service setting: Column percent, 2020	254
5.11c	Individuals aged 55 years and older, by mental health diagnosis, timing of admission, and mental health service setting: Row percent distribution, 2020	255

List of Tables (Continued)

5.12a	Individuals aged 55 years and older with SMI, by mental health diagnosis, timing of admission, and mental health service setting: Number, 2020	256
5.12b	Individuals aged 55 years and older with SMI, by mental health diagnosis, timing of admission, and mental health service setting: Column percent, 2020	257
5.12c	Individuals aged 55 years and older with SMI, by mental health diagnosis, timing of admission, and mental health service setting: Row percent distribution, 2020	258
5.13	Individuals aged 55 years and older, by mental health service setting and SMI status: Number and percent distribution, 2020	259

Characteristics of Individuals With Co-occurring Mental Health and Substance Use Disorders Receiving Mental Health Services

6.1a	Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by gender, age, race, ethnicity, education, veteran status, and SMI/SED status: Number, 2015–2020	260
6.1b	Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by gender, age, race, ethnicity, education, veteran status, and SMI/SED status: Column percent distribution, 2015–2020	262
6.2a	Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, gender, and age: Number, 2020	264
6.2b	Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, gender, and age: Column percent distribution, 2020	265
6.2c	Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, gender, and age: Row percent distribution, 2020	266
6.3a	Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, race, and ethnicity: Number, 2020	267
6.3b	Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, race, and ethnicity: Column percent distribution, 2020	268
6.3c	Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, race, and ethnicity: Row percent distribution, 2020	269
6.4a	Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis and substance use diagnosis: Number, 2020	270

List of Tables (Continued)

6.4b	Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis and substance use diagnosis: Column percent distribution, 2020	271
6.4c	Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis and substance use diagnosis: Row percent distribution, 2020	272
6.5a	Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, education, and marital status: Number, 2020	273
6.5b	Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, education, and marital status: Column percent distribution, 2020	274
6.5c	Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, education, and marital status: Row percent distribution, 2020	275
6.6a	Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, living arrangement, SMI/SED status, and veteran status: Number, 2020	276
6.6b	Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, living arrangement, SMI/SED status, and veteran status: Column percent distribution, 2020	277
6.6c	Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, living arrangement, SMI/SED status, and veteran status: Row percent distribution, 2020	278
6.7a	Individuals aged 18 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Number, 2020	279
6.7b	Individuals aged 18 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Column percent distribution, 2020	280
6.7c	Individuals aged 18 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Row percent distribution, 2020	281
6.8a	Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, timing of admission, and mental health service setting: Number, 2020	282
6.8b	Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, timing of admission, and mental health service setting: Column percent, 2020	283

List of Tables (Continued)

6.8c	Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, timing of admission, and mental health service setting: Row percent distribution, 2020	284

Client Outcomes Among Individuals Receiving Mental Health Services: 2015–2020

7.1a	Individuals served in both 2017 and 2020, by living arrangement and selected characteristics in 2017: Number, 2017	285
7.1b	Individuals served in both 2017 and 2020, by living arrangement and selected characteristics in 2017: Number, 2017	288
7.1c	Individuals served in both 2017 and 2020, by living arrangement and selected characteristics in 2017: Column percent distribution, 2017	291
7.2a	Individuals served in both 2017 and 2020, by living arrangement and selected characteristics in 2020: Number, 2020	294
7.2b	Individuals served in both 2017 and 2020, by living arrangement and selected characteristics in 2020: Column percent distribution, 2020	297
7.2c	Individuals served in both 2017 and 2020, by living arrangement and selected characteristics in 2020: Row percent distribution, 2020	300
7.3a	Individuals served in both 2017 and 2020, by employment status and selected characteristics in 2017: Number, 2017	303
7.3b	Individuals served in both 2017 and 2020, by employment status and selected characteristics in 2017: Column percent distribution, 2020	306
7.3c	Individuals served in both 2017 and 2020, by employment status and selected characteristics in 2017: Row percent distribution, 2020	309
7.4a	Individuals served in both 2017 and 2020, by employment status and selected characteristics in 2020: Number, 2020	312
7.4b	Individuals served in both 2017 and 2020, by employment status and selected characteristics in 2020: Column percent distribution, 2020	315
7.4c	Individuals served in both 2017 and 2020, by employment status and selected characteristics in 2020: Row percent distribution, 2020	318
7.5a	Individuals aged 18 years and older served in both 2017 and 2020, by mental health diagnosis in 2017 and living arrangement in 2017 and 2020: Number ...	321
7.5b	Individuals aged 18 years and older served in both 2017 and 2020, by mental health diagnosis in 2017 and living arrangement in 2017 and 2020: Row percent distribution, 2020.....	323
7.6a	Individuals aged 18 years and older served in both 2017 and 2020, by mental health diagnosis in 2017 and employment status in 2017 and 2020: Number ..	325

List of Tables (Continued)

7.6b	Individuals aged 18 years and older served in both 2017 and 2020, by mental health diagnosis in 2017 and employment status in 2017 and 2020: Row percent	326
7.7	Individuals aged 18 years and older served in consecutive reporting periods, by mental health diagnosis and living arrangement: Number, 2015–2020	327
7.8a	Individuals aged 18 years and older served in both 2015 and 2017, by mental health diagnosis in 2015 and living arrangement in 2015 and 2017: Number ..	329
7.8b	Individuals aged 18 years and older served in both 2015 and 2017, by mental health diagnosis in 2015 and living arrangement in 2015 and 2017: Row percent distribution	331
7.9a	Individuals aged 18 years and older served in both 2015 and 2018, by mental health diagnosis in 2015 and living arrangement in 2015 and 2018: Number ..	333
7.9b	Individuals aged 18 years and older served in both 2015 and 2018, by mental health diagnosis in 2015 and living arrangement in 2015 and 2018: Row percent distribution	335
7.10a	Individuals aged 18 years and older served in both 2015 and 2019, by mental health diagnosis in 2015 and living arrangement in 2015 and 2019: Number ..	337
7.10b	Individuals aged 18 years and older served in both 2015 and 2019, by mental health diagnosis in 2015 and living arrangement in 2015 and 2019: Row percent distribution	339
7.11a	Individuals aged 18 years and older served in both 2015 and 2020, by mental health diagnosis in 2015 and living arrangement in 2015 and 2020: Number ..	341
7.11b	Individuals aged 18 years and older served in both 2015 and 2020, by mental health diagnosis in 2015 and living arrangement in 2015 and 2020: Row percent distribution	343
7.12	Individuals aged 18 years and older served in consecutive reporting periods, by mental health diagnosis and employment status: Number, 2015–2020	345
7.13a	Individuals aged 18 years and older served in both 2015 and 2017, by mental health diagnosis in 2015 and employment status in 2015 and 2017: Number ..	346
7.13b	Individuals aged 18 years and older served in both 2015 and 2017, by mental health diagnosis in 2015 and employment status in 2015 and 2017: Row percent distribution	347
7.14a	Individuals aged 18 years and older served in both 2015 and 2018 by mental health diagnosis in 2015 and employment status in 2015 and 2018: Number ..	348
7.14b	Individuals aged 18 years and older served in both 2015 and 2018 by mental health diagnosis in 2015 and employment status in 2015 and 2018: Row percent distribution	349

List of Tables (Continued)

7.15a	Individuals aged 18 years and older served in both 2015 and 2019 by mental health diagnosis in 2015 and employment status in 2015 and 2019: Number ..	350
7.15b	Individuals aged 18 years and older served in both 2015 and 2019 by mental health diagnosis in 2015 and employment status in 2015 and 2019: Row percent distribution	351
7.16a	Individuals aged 18 years and older served in both 2015 and 2020, by mental health diagnosis in 2015 and employment status in 2015 and 2020: Number ...	352
7.16b	Individuals aged 18 years and older served in both 2015 and 2020, by mental health diagnosis in 2015 and employment status in 2015 and 2020: Row percent distribution	353

Geographic and Clinical Trends: 2015–2020

8.1	Individuals, by mental health diagnosis, Census region, Census division, and state or jurisdiction: Number, 2020	354
8.2	Individuals per 100,000 population (2020 estimate), by mental health diagnosis, Census region, Census division, and state or jurisdiction: Number, 2020 ...	357
8.3	Individuals per 100,000 population, by mental health diagnosis, Census region, Census division, and state or jurisdiction: Number, 2020, adjusted to 2010 population estimate for age, gender, and race	360
8.4	Individuals with diagnosis of trauma- and stressor-related disorders, by Census region, Census division, and state or jurisdiction: Number, 2015–2020	363
8.5	Individuals with diagnosis of anxiety disorders, by Census region, Census division, and state or jurisdiction: Number, 2015–2020	366
8.6	Individuals with diagnosis of attention deficit/hyperactivity disorder, by Census region, Census division, and state or jurisdiction: Number, 2015–2020	369
8.7	Individuals with diagnosis of bipolar disorders, by Census region, Census division, and state or jurisdiction: Number, 2015–2020	372
8.8	Individuals with diagnosis of primary depressive disorders, by Census region, Census division, and state or jurisdiction: Number, 2015–2020	375
8.9	Individuals with diagnosis of primary schizophrenia and other psychotic disorders, by Census region, Census division, and state or jurisdiction: Number, 2015–2020	378
8.10	Individuals with co-occurring mental health and substance use disorders, by Census region, Census division, and state or jurisdiction: Number, 2015–2020	381

Client Characteristics by State or Jurisdiction: 2015–2020

9.1a	Alabama – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020	384

List of Tables (Continued)

9.1b	Alabama – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020	386
9.2a	Alaska – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020	388
9.2b	Alaska – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020	390
9.3a	Arizona – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020	392
9.3b	Arizona – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020	394
9.4a	Arkansas – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020	396
9.4b	Arkansas – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020	398
9.5a	California – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020	400
9.5b	California – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020	402
9.6a	Colorado – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020	404
9.6b	Colorado – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020	406
9.7a	Connecticut – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020	408
9.7b	Connecticut – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020	410
9.8a	Delaware – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020	412
9.8b	Delaware – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020	414
9.9a	District of Columbia – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020	416
9.9b	District of Columbia – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020	418
9.10a	Florida – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020	420

List of Tables (Continued)

9.10b	Florida – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020	422
9.11a	Georgia – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020	424
9.11b	Georgia – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020	426
9.12a	Hawaii – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020	428
9.12b	Hawaii – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020	430
9.13a	Idaho – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020	432
9.13b	Idaho – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020	434
9.14a	Illinois – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020	436
9.14b	Illinois – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020	438
9.15a	Indiana – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020	440
9.15b	Indiana – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020	442
9.16a	Iowa – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020	444
9.16b	Iowa – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020	446
9.17a	Kansas – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020	448
9.17b	Kansas – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020	450
9.18a	Kentucky – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020	452
9.18b	Kentucky – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020	454
9.19a	Louisiana – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020	456

List of Tables (Continued)

9.19b	Louisiana – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020	458
9.20a	Maine – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020	460
9.20b	Maine – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020	462
9.21a	Maryland – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020	464
9.21b	Maryland – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020	466
9.22a	Massachusetts – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020	468
9.22b	Massachusetts – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020	470
9.23a	Michigan – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020	472
9.23b	Michigan – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020	474
9.24a	Minnesota – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020	476
9.24b	Minnesota – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020	478
9.25a	Mississippi – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020	480
9.25b	Mississippi – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020	482
9.26a	Missouri – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020	484
9.26b	Missouri – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020	486
9.27a	Montana – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020	488
9.27b	Montana – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020	490
9.28a	Nebraska – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020	492

List of Tables (Continued)

9.28b	Nebraska – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020	494
9.29a	Nevada – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020	496
9.29b	Nevada – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020	498
9.30a	New Jersey – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020	500
9.30b	New Jersey – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020	502
9.31a	New Mexico – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020	504
9.31b	New Mexico – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020	506
9.32a	New York – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020	508
9.32b	New York – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020	510
9.33a	North Carolina – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020	512
9.33b	North Carolina – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020	514
9.34a	North Dakota – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020	516
9.34b	North Dakota – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020	518
9.35a	Ohio – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020	520
9.35b	Ohio – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020	522
9.36a	Oklahoma – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020	524
9.36b	Oklahoma – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020	526
9.37a	Oregon – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020	528

List of Tables (Continued)

9.37b	Oregon – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020	530
9.38a	Pennsylvania – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020	532
9.38b	Pennsylvania – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020	534
9.39a	Puerto Rico – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020	536
9.39b	Puerto Rico – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020	538
9.40a	Rhode Island – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020	540
9.40b	Rhode Island – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020	542
9.41a	South Carolina – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020	544
9.41b	South Carolina – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020	546
9.42a	South Dakota – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020	548
9.42b	South Dakota – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020	550
9.43a	Tennessee – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020	552
9.43b	Tennessee – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020	554
9.44a	Texas – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020	556
9.44b	Texas – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020	558
9.45a	Utah – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020	560
9.45b	Utah – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020	562
9.46a	Vermont – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020	564

List of Tables (Continued)

9.46b	Vermont – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020	566
9.47a	Virginia – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020	568
9.47b	Virginia – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020	570
9.48a	Washington – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020	572
9.48b	Washington – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020	574
9.49a	West Virginia – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020	576
9.49b	West Virginia – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020	578
9.50a	Wisconsin – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020	580
9.50b	Wisconsin – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020	582
9.51a	Wyoming – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020	584
9.51b	Wyoming – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020	586
9.52a	Other jurisdictions – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020	588
9.52b	Other jurisdictions – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020	590

Client Characteristics by State or Other Jurisdiction: 2020

10.1a	Alabama – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020	592
10.1b	Alabama – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020	594
10.2a	Alaska – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020	596
10.2b	Alaska – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020	598

List of Tables (Continued)

10.3a	Arizona – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020	600
10.3b	Arizona – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020	602
10.4a	Arkansas – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020	604
10.4b	Arkansas – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020	606
10.5a	California – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020	608
10.5b	California – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020	610
10.6a	Colorado – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020	612
10.6b	Colorado – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020	614
10.7a	Connecticut – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020	616
10.7b	Connecticut – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020	618
10.8a	Delaware – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020	620
10.8b	Delaware – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020	622
10.9a	District of Columbia – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020	624
10.9b	District of Columbia – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020	626
10.10a	Florida – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020	628
10.10b	Florida – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020	630
10.11a	Georgia – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020	632
10.11b	Georgia – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020	634

List of Tables (Continued)

10.12a	Hawaii – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020	636
10.12b	Hawaii – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020	638
10.13a	Idaho – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020	640
10.13b	Idaho – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020	642
10.14a	Illinois – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020	644
10.14b	Illinois – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020	646
10.15a	Indiana – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020	648
10.15b	Indiana – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020	650
10.16a	Iowa – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020	652
10.16b	Iowa – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020	654
10.17a	Kansas – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020	656
10.17b	Kansas – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020	658
10.18a	Kentucky – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020	660
10.18b	Kentucky – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020	662
10.19a	Louisiana – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020	664
10.19b	Louisiana – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020	666
10.20a	Massachusetts – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020	668
10.20b	Massachusetts – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020	670

List of Tables (Continued)

10.21a	Michigan – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020	672
10.21b	Michigan – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020	674
10.22a	Minnesota – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020	676
10.22b	Minnesota – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020	678
10.23a	Mississippi – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020	680
10.23b	Mississippi – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020	682
10.24a	Missouri – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020	684
10.24b	Missouri – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020	686
10.25a	Montana – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020	688
10.25b	Montana – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020	690
10.26a	Nebraska – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020	692
10.26b	Nebraska – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020	694
10.27a	Nevada – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020	696
10.27b	Nevada – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020	698
10.28a	New Jersey – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020	700
10.28b	New Jersey – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020	702
10.29a	New Mexico – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020	704
10.29b	New Mexico – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020	706

List of Tables (Continued)

10.30a New York – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020	708
10.30b New York – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020	710
10.31a North Carolina – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020	712
10.31b North Carolina – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020	714
10.32a North Dakota – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020	716
10.32b North Dakota – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020	718
10.33a Ohio – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020	720
10.33b Ohio – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020	722
10.34a Oklahoma – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020	724
10.34b Oklahoma – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020	726
10.35a Oregon – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020	728
10.35b Oregon – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020	730
10.36a Pennsylvania – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020	732
10.36b Pennsylvania – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020	734
10.37a Puerto Rico – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020	736
10.37b Puerto Rico – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020	738
10.38a Rhode Island – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020	740
10.38b Rhode Island – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020	742

List of Tables (Continued)

10.39a	South Carolina – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020	744
10.39b	South Carolina – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020	746
10.40a	South Dakota – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020	748
10.40b	South Dakota – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020	750
10.41a	Tennessee – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020	752
10.41b	Tennessee – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020	754
10.42a	Texas – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020	756
10.42b	Texas – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020	758
10.43a	Utah – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020	760
10.43b	Utah – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020	762
10.44a	Vermont – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020	764
10.44b	Vermont – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020	766
10.45a	Virginia – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020	768
10.45b	Virginia – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020	770
10.46a	Washington – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020	772
10.46b	Washington – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020	774
10.47a	Wisconsin – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020	776
10.47b	Wisconsin – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020	778

List of Tables (Continued)

10.48a	Wyoming – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020	780
10.48b	Wyoming – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020	782
10.49a	Other jurisdictions – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020	784
10.49b	Other jurisdictions – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020	786

List of Figures

Client Outcomes Among Individuals Receiving Mental Health Services: 2015–2020

Figure 1. Census regions and Census divisions of the United States.....	16
Figure 2. Living arrangements in 2020 of individuals aged 18 years and older who reported living in a private residence in 2017, by 2017 mental health diagnosis	38
Figure 3. Living arrangements in 2020 of individuals aged 18 years and older who experienced homelessness in 2017, by 2017 mental health diagnosis.....	39
Figure 4. Living arrangements in 2020 of individuals aged 18 years and older who reported being in jail or a correctional facility in 2017, by 2017 mental health diagnosis.....	40
Figure 5. Employment status in 2020 of individuals aged 18 years and older who reported being employed in 2017, by 2017 mental health diagnosis.....	41
Figure 6. Employment status in 2020 of individuals aged 18 years and older who reported being unemployed in 2017, by 2017 mental health diagnosis.....	42
Figure 7. Employment status in 2020 of individuals aged 18 years and older who were not in the labor force in 2017, by 2017 mental health diagnosis	43

Executive Summary

This report presents results from the Mental Health Client-Level Data (MH-CLD) and Mental Health Treatment Episode Data Set (MH-TEDS) for individuals receiving mental health services from state mental health systems in 2020, as well as selected trends in data collected from such individuals between 2015 and 2020. The report provides information on mental health diagnoses, mental health treatment settings, and demographic and substance use characteristics of individuals in mental health treatment in facilities that reported to individual state administrative data systems.

In the 2020 reporting period¹, 50 states and other jurisdictions (hereafter, “states”) submitted 6,945,521 records for individuals receiving mental health services. American Samoa, the Federated States of Micronesia, Guam, Maine, Marshall Islands, Maryland, New Hampshire, U.S. Virgin Islands, and West Virginia did not report data for the 2020 reporting period, as of March 3, 2022 [Table 1.1a and Appendix Table B1].

Of the 6,945,521 eligible client-level records for the 2020 reporting period, 85.4 percent had valid mental health and/or substance use diagnosis codes. Such records form the basis for most tables in this report. However, individual records may not have data for all variables; therefore, the analysis of specific variables may be based on a smaller number of individuals [Tables 1.1a–b].

State Client-Level Data Reporting: 2015–2020

For the 2015 reporting period, 45 states reported that a total of 5,165,254 individuals received mental health services; for the 2016 reporting period, 47 states reported that a total of 5,656,750 individuals received mental health services; for the 2017 reporting period, 49 states reported that a total of 6,171,990 individuals received mental health services; for the 2018 reporting period, 49 states reported that a total of 6,418,247 individuals received mental health services; for the 2019 reporting period, 48 states reported that a total of 6,548,372 individuals received mental health services; and for the 2020 reporting period, 50 states reported that a total of 6,945,521 individuals received mental health services [Table 1.1a].

In all six reporting periods from 2015 through 2020, between 27.1 and 28.9 percent of individuals served were aged 17 years and younger; between 10.1 and 10.4 percent were aged 18 to 24 years; between 44.3 and 46.2 percent were aged 25 to 54 years; and between 15.2 and 17.0 percent were aged 55 years and older [Table 1.2b]. The estimates presented here were aggregated using data presented in Table 1.2b. For a detailed breakdown by age categories and years, please refer to table 1.2b.

¹ For most states, the 2020 reporting period was July 1, 2019, to June 30, 2020. See Appendix B, Table B1.

Children and Adolescents Aged 17 Years and Younger Receiving Mental Health Services: 2020

A total of 1,915,199 children and adolescents aged 17 years and younger were served in the 2020 reporting period [Table 2.1a]; 1,637,967 had a valid mental health diagnosis code [Table 2.3a].

Among males aged 17 years and younger served in the 2020 reporting period, the most frequently reported diagnoses were ADHD (38.1 percent), trauma- and stressor-related disorders (26.7 percent), and anxiety disorders (15.6 percent) [Table 2.3c].

Among females aged 17 years and younger served in the 2020 reporting period, the most frequently reported diagnoses were trauma- and stressor-related disorders (34.8 percent), depressive disorders (26.3 percent), and anxiety disorders (24.8 percent) [Table 2.3c].

Young Adults Aged 18 to 24 Years Receiving Mental Health Services: 2020

A total of 724,047 young adults aged 18 to 24 years were served in 2020 [Table 3.1a]; 608,665 had a valid mental health diagnosis code [Table 3.3a].

Depressive disorders were the most frequently reported diagnoses among individuals aged 18 to 24 years served in the 2020 reporting period, across all genders, races, ethnicities and marital statuses [Tables 3.3c and 3.5c].

Adults Aged 25 to 54 Years Receiving Mental Health Services: 2020

A total of 3,123,221 adults aged 25 to 54 years were served in 2020 [Table 4.1a]; 2,667,813 had a valid mental health diagnosis code [Table 4.3a].

Among males aged 25 to 54 years, the most frequently reported diagnoses were depressive disorders (29.0 percent), schizophrenia and other psychotic disorders (24.9 percent), and anxiety disorders (20.9 percent); among females, the most frequently reported diagnoses were depressive disorders (38.8 percent), anxiety disorders (30.6 percent), and trauma- and stressor- related disorders (23.5percent) [Table 4.3c].

Older Adults Aged 55 Years and Older Receiving Mental Health Services: 2020

A total of 1,178,713 adults aged 55 years and older were served in 2020 [Table 5.1a]; 1,010,080 had a valid mental health diagnosis code [Table 5.3a].

Among males aged 55 years and older served in the 2020 reporting period, the most frequently reported diagnoses were depressive disorders (33.5 percent), schizophrenia and other psychotic disorders (29.8 percent), and anxiety disorders (16.5 percent); among females served during this period, the most frequently reported diagnoses were depressive disorders (43.0 percent), anxiety disorders (24.3 percent), and schizophrenia and other psychotic disorders (18.9 percent) [Table 5.3c].

Individuals With Co-Occurring Mental Health and Substance Use Disorders Receiving Mental Health Services: 2020

A total of 1,663,657 individuals aged 12 and older with co-occurring mental and substance use disorders were served in 2020 [Table 6.1a].

For males with co-occurring mental and substance use disorders aged 12 years or older served during the 2020 reporting period, the most frequently reported diagnoses were depressive disorders (31.9 percent), schizophrenia and other psychotic disorders (24.1 percent), and anxiety disorders (19.8 percent); among females, the most frequently reported diagnoses were depressive disorders (40.8 percent), anxiety disorders (27.6 percent), and trauma- and stressor-related disorders (23.8 percent) [Table 6.2c].

Client Outcomes Among Individuals Receiving Mental Health Services in 2017 and 2020

Among individuals served in both the 2017 and 2020 reporting periods, 94.9 percent of those who lived in a private residence in 2017 also reported living in a private residence in 2020; 1.9 percent experienced homelessness; 1.4 percent transitioned to residential care; and 1.7 percent, combined, transitioned to institutional settings, jail, or other living arrangements [Table 7.5b].

Among individuals aged 18 years and older who were served in both the 2017 and 2020 reporting periods, 68.9 percent of those who were employed in 2017 also reported being employed in 2020; 11.4 percent transitioned to being unemployed, and 19.8 percent had left the labor force [Table 7.6b].

Individuals Who Received Mental Health Services per 100,000 Population, by State or Jurisdiction in 2020, Adjusted for Age, Gender, and Race

Tables 8.2 and 8.3 present, respectively, the number and adjusted number of individuals per 100,000 population served in the 2020 reporting period, by mental health diagnosis, Census region, Census division, and state or jurisdiction.

Among individuals served in the 2020 reporting period, the mental health diagnoses with the largest per 100,000 population ratios (adjusted for age, gender, and ethnicity) were depressive disorders (535 per 100,000 population), anxiety disorders (415 per 100,000 population), and trauma- and stressor-related disorders (381 per 100,000 population) [Table 8.3].

Across all Census divisions in the 2020 reporting period except the Mountain division, depressive disorders were the mental health diagnosis with the highest adjusted ratio per 100,000 population, ranging from 256 per 100,000 population for New England to 1,048 per 100,000 population for the West North Central division [Table 8.3].

Background

Mental health services data are currently collected for and reported to the Substance Abuse and Mental Health Services Administration's (SAMHSA) Center for Behavioral Health Statistics and Quality (CBHSQ). State mental health agencies (SMHAs) are the state entities with primary responsibility for reporting the data in accordance with the reporting terms and conditions of the Behavioral Health Services Information System (BHSIS) Agreements funded by SAMHSA. The BHSIS Agreement stipulates that states and jurisdictions submit Uniform Reporting System (URS) data and client-level data by December 1 each year, in compliance with mental health block grant reporting requirements (42 U.S.C. §300x–6).

URS data reporting started in 2002 in an effort spearheaded by SAMHSA's Center for Mental Health Services (CMHS) to collect both system performance measures and client outcomes at an aggregate level. The collection of client-level data came later.

In 2008, SAMHSA, through CMHS, tested the feasibility of transitioning from aggregate reporting of SAMHSA's mental health national outcome measures (NOMs)² to individual-level data reporting. This led to the development of the Mental Health Client-Level Data (MH-CLD) reporting method. The development was supported by data infrastructure grants (DIGs), which were a vehicle through which states and jurisdictions received financial and technical assistance from SAMHSA to assist in their capacity-building efforts toward meeting the block grant reporting requirements.

The general framework for the MH-CLD involves a compilation of the demographic, clinical, and outcome data of individuals served by the SMHA within a state-defined 12-month reporting period. States may choose the calendar year or the state fiscal year as their reporting period. Individuals served is defined as all enrolled individuals who received mental health and support services, including screening, assessment, crisis services, and telemedicine, from programs operated or funded by the SMHA during the reporting period.

Two data sets are submitted each reporting period: Basic Client Information (BCI) is due December 1, and the State Hospital Readmission (SHR) data set is due March 1 of the following year. These two data sets inform five mental health NOMs: access to services/capacity, stability in housing/residential status, 30-day and 180-day readmission to state hospital, adult employment or children's school attendance and education level, and criminal justice involvement. There are five additional NOMs—social connectedness, decreased mental illness symptomatology, perception of care, cost-effectiveness, and use of evidence-based practices—that have not been converted into client-level data reporting and continue to be captured in the URS. Data are reported in accordance with the Health Insurance Portability and Accountability Act (HIPAA) rules for non-protected health information (non-PHI).

In 2010, SAMHSA, through CBHSQ, began a parallel effort of testing the feasibility of building a national behavioral health database that integrates both mental health and substance use data using a uniform platform for data reporting. This effort led to the integration of mental health data into the well-established system of reporting for substance use client-level data, known as the Treatment

²National outcome measures can be found in the [Mental Health Client-Level Data State Instruction Manual](#).

Episode Data Set (TEDS).³ To enable reporting of mental health client data to TEDS, the system was modified to incorporate data elements that capture the characteristics unique to the mental health services system and the individuals receiving mental health services. These characteristics include, among others, the extended period of time that an individual may be engaged in treatment, the recovery process, and the simultaneous receipt of different types of services from different service providers. This method of reporting mental health data at the individual level is referred to as the Mental Health Treatment Episode Data Set (MH-TEDS).

Unlike MH-CLD, MH-TEDS is structured around treatment events, that is, admissions and discharges from specific service settings. Admission and discharge records can be linked to track treatment episodes and the treatment services received by individuals. Thus, with MH-TEDS, both the individual client and the treatment episode can serve as a unit of analysis, whereas with MH-CLD the client is the sole unit of analysis. The common TEDS platform in which both the substance use and mental health data are reported enhances the ability to report data for people with co-occurring mental and substance use disorders. It also offers optional data fields for individuals with mental illness that are not captured in MH-CLD, such as referral source, details on criminal justice referral, income sources, and health insurance.

MH-TEDS is a compilation of demographic, substance use, mental health, clinical, legal, and socioeconomic characteristics of individuals who are primarily receiving publicly funded mental health and/or substance use services. MH-TEDS data support program performance measurement and management goals. The same set of mental health NOMs enumerated under MH-CLD is also supported by MH-TEDS. Thus, while both MH-TEDS and MH-CLD collect similar client-level data, the collection methodology differs.

National mental health client-level data reporting was launched in 2011, when MH-CLD was the only reporting method available. The number of states reporting MH-CLD or MH-TEDS has increased from 19 in 2011 to 50 in 2020 (see Appendix B for details). It is important to note that the level and extent of reporting varies. Note also that New York and Tennessee submit data only for a subset of the total number of individuals that they served during a reporting period.

Beginning in the 2014 reporting period, states could choose which method—MH-CLD or MH-TEDS—to use for data reporting. In 2020, 33 states and two other jurisdictions (Commonwealth of the Northern Mariana Islands and the Republic of Palau) used MH-CLD; 14 states (Alaska, Connecticut, Delaware, Florida, Georgia, Kansas, Louisiana, Michigan, Mississippi, Nebraska, North Carolina, Oklahoma, Pennsylvania, and Vermont), the District of Columbia, and Puerto Rico used MH-TEDS [Appendix Table B1]. Connecticut reports data through both MH-CLD (for children and adolescents) and MH-TEDS (for adults).

In summary, this report presents the client-level data reported for the 2015 through 2020 reporting periods. Data from all six reporting periods are presented in Chapters 1 through 6. Chapter 7 presents outcomes for individuals served across two or more reporting periods. Chapter 8 presents, for each state or jurisdiction, the number of individuals served in the 2015–2020 reporting periods, according to select mental health diagnoses for age, gender, and race based on the U.S. state-level population.

³Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality. Treatment Episode Data Set (TEDS): 2019. *National Admissions to and Discharges From Publicly Funded Substance Use Treatment*. Rockville, MD: Substance Abuse and Mental Health Services Administration, 2021.

Chapter 9 presents state-level data for the 2015 through 2020 reporting periods, and Chapter 10 presents state-level data only for the 2020 reporting period.

See Appendix A for a crosswalk of data fields collected in MH-TEDS and MH-CLD.

Introduction

This report presents mental health client-level data submitted for the 2015 through 2020 reporting periods using the Mental Health Client-Level Data (MH-CLD) and the Mental Health Treatment Episode Data Set (MH-TEDS). This report provides information on the demographics, characteristics, and outcomes of individuals served by state mental health agencies (SMHAs) for these six reporting periods, with particular focus on the most recent reporting period. The Substance Abuse and Mental Health Services Administration (SAMHSA) has provided support to states to build capacity to collect and report client-level data.

Methodology

Reconciling data from MH-TEDS and MH-CLD

Both the MH-TEDS and MH-CLD include individuals served through SMHAs during the state-defined 12-month reporting period. Most states elect to use the state fiscal year, which, for the 2020 reporting period, usually ran from July 1, 2019, through June 30, 2020; however, some states define their fiscal year differently. See Appendix B for details.

MH-CLD presented in this report contains one record per individual served. Data items are populated by the states based on the status of the individual, either at the start and end of the state's reporting period, or the most recently available data. MH-TEDS contains one record for each admission to, and discharge from, a particular service type/setting (referred to as a treatment episode) within the client's treatment continuum during the reporting period. Data collected from MH-TEDS admission records and update/discharge records are stored as two separate data sets that are linked using a HIPAA-compliant, non-protected health information, unique client identifier located in both files. Linking the records allows for the creation of a single client record comparable to the MH-CLD format.

The process of creating an individual record using the MH-TEDS data allows for the reporting of these individuals in both the MH-TEDS and MH-CLD data sets. Tables in all chapters of this report use individuals as the unit of analysis for the combined data from MH-TEDS and MH-CLD.

States continually review and improve their data collection and processing. When systematic errors are identified, states may revise or replace historical MH-TEDS and MH-CLD data files. While this process represents an improvement in the data system, the historical statistics in this report will differ slightly from those in earlier reports.

Selection of reporting variables

MH-TEDS and MH-CLD collect a variety of data items, including demographics (e.g., age, gender, race, and ethnicity), national outcome measures (NOMs, e.g., living arrangements, employment status, criminal justice involvement, and school attendance), and other clinical measures (e.g., co-occurring mental health and substance use disorders). Throughout this report, data for living arrangements and employment are presented according to individuals' status at

the end of the given reporting period. Valid data on criminal justice involvement were reported for less than 41 percent of the individuals served for each reporting period, and arrest data were excluded from this report. Global Assessment of Functioning (GAF) and Children's Global Assessment (CGAS) scores were dropped from the fifth edition of the Diagnostic and Statistical Manual of Mental Disorders (DSM- 5) and thus are excluded from this report due to insufficient data. See Appendix C for a description of the variables; see Appendix D for the federal definitions of serious mental illness (SMI) and serious emotional disturbance (SED).

Mental health diagnoses are reported in the MH-TEDS and MH-CLD as codes from either the International Classification of Diseases, Clinical Modification (ICD-9-CM/ICD-10-CM), or from the DSM-IV/5. Up to three mental health diagnostic codes may be reported for each client. For this report, diagnostic codes were categorized based on a modified version of the Clinical Classification Software (CCS). The mental health section of the CCS has 15 diagnostic categories and 25 subcategories; in all, 12 mental health diagnoses were selected for tabulation, with all other mental health diagnoses grouped as other mental health diagnoses for a given table. See Appendix E for details.

The 12 mental health diagnostic groups used in this report are: trauma- and stressor-related disorders; anxiety disorders; attention deficit/hyperactivity disorder (ADHD); conduct disorder; delirium, dementia, and amnestic and other cognitive disorders (delirium/dementia); bipolar disorders; depressive disorders; oppositional defiant disorder; personality disorders; pervasive developmental disorders; schizophrenia and other psychotic disorders; and alcohol or substance-related disorders.

Records with partially complete data have been retained in this report. For example, an entire client record is not removed from the report because a single data item is missing. When records include missing or invalid data for a specific variable, those records are excluded from tabulations of that variable. The total number of records on which a percentage distribution is based is reported in each table.

Stratification of reporting population and subgroups

In general, this report presents national- and state-level data from the MH-CLD in the 2020 reporting period, as well as trend data from the 2015–2020 reporting periods. There are three major levels of data reporting: (1) the entire population within each age category, (2) the subgroup population of individuals with, or at risk for, serious emotional disturbance (SED) or serious mental illness (SMI), and (3) the entire population within each mental health diagnosis category.

Because different stages of life are associated with distinct forms of mental and behavioral disorders and with distinctive treatment service utilization patterns, this report examines the characteristics of individuals receiving mental health services by the following age groups: children and adolescents (aged 17 years and younger), young adults (aged 18 to 24 years), adults (aged 25 to 54 years), and

older adults (aged 55 years and older). Accordingly, Chapters 2 through 5, which focus on specific age groups, include age-specific sets of mental health diagnoses.⁴

Additionally, this report includes the subpopulation of individuals with co-occurring mental and substance use disorders. For this subpopulation, the report includes individuals aged 12 years and older to match the population used in the substance use TEDS reports. Determination of individuals with co-occurring mental and substance use disorders was based on the variable in the data set called co-occurring mental and substance use disorders.

Identification of individuals with a serious mental illness (SMI) or serious emotional disturbance (SED) is based on each state's definitions for SMI and SED. This is reported as a single variable: SMI/SED status. Individuals aged 17 years and younger who are indicated in the data set as having SED, or being at risk of SED, are included in the SED subpopulation, while individuals aged 18 years and older who are indicated as having an SMI are included in the SMI subpopulation. Exceptions may exist for a few states where young adults (aged 18 to 21 years) continue to receive mental health services from the state's child mental health system. They are included in the SED subpopulation as reported by the state.

Cohort individual analysis

Chapter 7 presents data on cohorts of individuals who received mental health services through SMHAs during two or more reporting periods. While the narrative of Chapter 7 focuses specifically on a cohort of individuals served in both the 2017 and 2020 reporting periods, additional tables for Chapter 7 encompass cohorts of individuals served in other reporting year combinations, such as in both 2015 and 2016, or in both 2015 and 2020. Individuals were selected for analysis if records with the same unique identification number, from the same state, indicated that they were served in both of the given reporting periods.

Limitations

For both MH-TEDS and MH-CLD, states report only individuals served through SMHAs during the state-defined 12-month reporting period. These data do not represent the total national demand for mental health treatment or describe the mental health status of the national population. In addition, New York and Tennessee report only a subset of the total individuals served through their SMHAs.

This report uses any mention of a mental health diagnosis (whether first-mention, second-mention, or third-mention) to group individuals. This means that individuals may be counted for up to three diagnoses within the same table. For this reason, caution must be used when comparing tables from this report with tables in previous years' reports that counted only primary (first mention) mental health diagnoses. About 85.4 percent of all individuals in the 2020 data have at least one mental health diagnosis, while 23.1 percent have exactly two mental health diagnoses, and 7.2 percent have exactly three mental health diagnoses [Tables 1.12a–b].

⁴For example, individuals with a mental health diagnosis of schizophrenia and other psychotic disorders are grouped with *other* for persons aged 17 years and younger in Chapter 2. Although the subject of recent interest, this diagnostic category is not reported separately for this age group. Less than 1 percent of all individuals served in this age group reported schizophrenia and other psychotic disorders as a diagnosis.

Chapter 1

Summary of Annual Data in State Client-Level Data Reporting: 2015–2020

This chapter provides a brief background on state reporting of mental health client-level data and a summary of the data submitted for the 2015 through 2020 reporting periods.⁵ Data on NOMs and clinical measures are presented by demographic characteristics and mental health diagnoses⁶ for all individuals.

Caution must be used when comparing data over time. The number of states submitting client-level data has varied for each reporting period: 45 states reported 2015 data, 47 states reported 2016 data, 49 states reported 2017 data, 49 states reported 2018 data, 48 states reported 2019 data, and 50 states reported 2020 data.

All data tables represent sums of individuals who received publicly funded mental health services—individuals served—through the auspices of state mental health agencies (SMHAs).

Individuals Served

Tables 1.1a–b. For the 2015 reporting period, 45 states reported that a total of 5,165,254 individuals received mental health services; for the 2016 reporting period, 47 states reported that a total of 5,656,750 individuals received mental health services; for the 2017 reporting period, 49 states reported that a total of 6,171,990 individuals received mental health services; for the 2018 reporting period, 49 states reported that a total of 6,418,247 individuals received mental health services; for the 2019 reporting period, 48 states reported that a total of 6,548,372 individuals received mental health services; and for the 2020 reporting period, 50 states reported that a total of 6,945,521 individuals received mental health services.⁷

Gender

Tables 1.2a–b. Females accounted for between 51.3 and 51.7 percent of all individuals served in every reporting period between 2015 and 2020.

Age group

Tables 1.2a–b. In all six reporting periods from 2015 through 2020, between 27.1 and 28.9 percent of individuals served were aged 17 years and younger; between 10.1 and 10.4 percent were aged 18 to 24 years; between 44.3 and 46.2 percent were aged 25 to 54 years; and between 15.2 and 17.0 percent were aged 55 years and older. The most frequently reported diagnoses among specific age

⁵Hereafter, “states” may refer to any of the 50 states, the District of Columbia, the Commonwealth of Puerto Rico, and the combined jurisdictions of the Republic of Palau and the Commonwealth of the Northern Mariana Islands.

⁶Except where otherwise noted, throughout this report, the term “diagnosis” refers to all reported mental health diagnoses, including first-mention, second-mention, and third-mention mental health diagnoses on the client-level data record.

⁷Data were not submitted or were incomplete for one or more years in some states or jurisdictions; see Appendix B.

groups across the six reporting periods are reported in Tables 1.4a–1.7b and are further discussed in Chapters 2 through 5.

Race and ethnicity⁸

Tables 1.3a–b. For each of the reporting periods from 2015 through 2020, Whites represented between 65.0 and 67.1 percent of the individuals served; Blacks or African Americans represented between 19.6 and 20.9 percent; and American Indians or Alaska Natives, Asians, Native Hawaiians or Other Pacific Islanders, and other groups, combined, represented between 12.0 and 14.4 percent.⁹

Hispanics accounted for between 16.2 and 19.8 percent of all individuals served in each of the six reporting periods from 2015 through 2020.

Table 1.23b. Blacks or African Americans accounted for 19.6 percent of individuals served in the 2020 reporting period but just 13.5 percent of the 2020 U.S. population.¹⁰

Marital status

Tables 1.8a–b. Among individuals aged 18 years and older in each reporting period from 2015 through 2020, more than half reported never being married (ranging between 55.5 and 65.7 percent); those who were married at the time of reporting ranged between 12.9 and 17.5 percent.

Tables 1.22a–c. Among individuals aged 25 to 54 years served in the 2020 reporting period, 15.1 percent were currently married. In contrast, 56.5 percent of the 2020 U.S. population in that age group were currently married.¹¹

Living arrangements

Tables 1.8a–b. Individuals living in a private residence accounted for between 83.6 and 86.1 percent of those served in the six reporting periods from 2015 through 2020; between 3.5 and 4.3 percent experienced homelessness in each of those reporting periods.

SMI/SED status

Tables 1.9a–b. Individuals with serious mental illness (SMI) accounted for between 72.3 and 77.9 percent of individuals aged 18 years and older who were served in any of the six reporting periods from 2015 through 2020. Children and adolescents aged 17 years and younger who had, or were at risk for, serious emotional disturbance (SED) accounted for between 70.5 and 78.2 percent of individuals aged 17 years and younger who were served in any of the six reporting periods from 2015 through 2020.

⁸Data on specific Hispanic or Latino origin (hereafter referred to as “Hispanic origin”) are presented for a subset of states that reported information on individuals’ specific Hispanic or Latino origin.

⁹Other includes individuals who reported two or more races or some other race.

¹⁰Source: U.S. Census Bureau, Population Estimates Program, SC-EST2020-alldata6: “Annual State Resident Population Estimates for 6 Race Groups (5 Race Alone Groups and Two or More Races) by Age, Sex, and Hispanic Origin: April 1, 2010, to July 1, 2019, April 1 2020 and July 1, 2020,” <https://www2.census.gov/programs-surveys/popest/datasets/2010-2020/state/asrh/SC-EST2020-ALLDATA6.csv>.

¹¹Source: U.S. Census Bureau, Current Population Survey, 2019 Annual Social and Economic Supplement (ASEC), November 2020, Table A1. <https://www.census.gov/data/tables/2020/demo/families/cps-2020.html>.

Employment

Tables 1.10a–b. Among individuals aged 18 years and older served in any of the six reporting periods from 2015 through 2020, the proportion employed ranged between 20.8 and 25.3 percent, the proportion unemployed ranged between 26.9 and 31.4 percent, and the proportion not in the labor force ranged between 43.3 and 51.1 percent.

Education

Tables 1.10a–b. Individuals who completed Grade 12 or a general equivalency diploma (GED) accounted for between 46.3 and 47.8 percent of individuals aged 18 years and older who were served in any of the six reporting periods from 2015–2020; between 20.3 and 24.8 percent completed at least a year of education beyond Grade 12.

Veteran status

Tables 1.10a–b. In the 2020 reporting period, veterans accounted for 4.9 percent of individuals aged 18 years and older served.

Client treatment status at the start of the reporting period

Tables 1.11a–b. Among individuals served in any of the six reporting periods from 2015 through 2020, both those newly admitted to treatment¹² ranged between 46.5 to 53.6 percent, and those continuing treatment¹³ ranged between 46.4 and 53.5 percent.

Service setting

Tables 1.11a–b. For the 2015 to 2020 reporting periods, the proportion of individuals who received mental health services from community-based programs ranged between 97.1 and 97.5 percent. Further analysis of the settings in which individuals received mental health services during the 2020 reporting period, specifically, is presented in Chapters 2 through 6.

Demographic characteristics of individuals with selected mental health diagnoses

Tables 1.14a–1.19b present the gender, age, race, and ethnicity of individuals with the following diagnoses: trauma- and stressor-related disorders [Tables 1.14a–b], anxiety disorders [Tables 1.15a–b], attention deficit/hyperactivity disorder [Tables 1.16a–b], bipolar disorders [Tables 1.17a–b], depressive disorders [Tables 1.18a–b], and schizophrenia and other psychotic disorders [Tables 1.19a–b].

¹²Individuals who were newly admitted to treatment either (a) had not previously received a service and started receiving services during the reporting year, or (b) had previously received a service, were discharged or inactive for an extended period of time, and resumed receiving services during the reporting period.

¹³Individuals continuing treatment consists of those who were admitted prior to the start of the reporting period and who received a service during the reporting period.


Demographic characteristics, by status for co-occurring mental health and substance use disorders

Tables 1.20a–1.21b present the gender, age, race, and ethnicity of individuals grouped by the presence or absence of co-occurring mental health and substance use disorders.

Regional Distributions

Tables 1.13a–b and Figure 1. About a third (between 27.6 and 36.6 percent) of all individuals who received mental health services in the 2015 through 2020 reporting periods were in the South. The Census regions with the next largest proportions of individuals who received mental health services in the 2015 through 2020 reporting periods were the West (between 24.3 and 28.8 percent), the Midwest (between 21.7 and 27.1 percent), and the Northeast (between 10.9 and 20.7 percent).

Figure 1. Census regions and Census divisions of the United States



SOURCE: U.S. Census Bureau.

Any-Mention Mental Health Diagnoses

Tables 1.12a–b present the number and percentage distribution of individuals who received any mention—whether first, second, or third—of the listed mental health diagnoses for the 2015, 2016, 2017, 2018, 2019 and 2020 reporting periods. Differences among reporting periods should be interpreted with great caution; variation may be more a function of the mix and number of reporting states than of actual differences.

CHAPTER 2

CHARACTERISTICS OF CHILDREN AND ADOLESCENTS AGED 17 YEARS AND YOUNGER RECEIVING MENTAL HEALTH SERVICES

This chapter presents data on children and adolescents aged 17 years and younger who received mental health services through SMHAs in the 2015–2020 reporting periods, with particular focus on the 2020 reporting period. Data on NOMs and clinical measures are presented by demographic characteristics and mental health diagnoses for all children and adolescents. Characteristics for the subpopulation of children and adolescents with, or at risk for, SED are presented where the percentages notably differ from the characteristics of all children and adolescents.

Individuals Served

Tables 2.1a–b present the total number of individuals aged 17 years and younger served in the 2015 through 2020 reporting periods, by demographic characteristics.

The number of individuals served who were aged 17 years and younger increased from 1,473,157 in the 2015 reporting period to 1,915,199 in the 2020 reporting period.

Tables 2.3a–2.4c. In the 2020 reporting period, 1,637,967 individuals aged 17 years and younger had a valid mental health diagnosis code, of which 1,155,930 were reported to have, or be at risk of, SED.

Only individuals with valid mental health diagnoses are included in these and all subsequent tabulations of mental health diagnoses in this chapter.

Gender

Tables 2.3a–2.4c. Tables 2.3a–c present the most frequently reported diagnoses, by gender, among individuals aged 17 years and younger, while Tables 2.4a–c present the most frequently reported diagnoses, by gender, for individuals aged 17 years and younger with SED. Males accounted for 55.4 percent and females accounted for 44.6 percent of individuals aged 17 years and younger who were served in the 2020 reporting period with a valid mental health diagnosis.

Among males aged 17 years and younger served in the 2020 reporting period, the most frequently reported diagnoses were ADHD (38.1 percent), trauma- and stressor-related disorders (26.7 percent), and anxiety disorders (15.6 percent). Among females aged 17 years and younger served in the 2020 reporting period, the most frequently reported diagnoses were trauma- and stressor-related disorders (34.8 percent), depressive disorders (26.3 percent), and anxiety disorders (24.8 percent)

Race

Tables 2.3a–2.4c. The most frequently reported diagnoses among individuals aged 17 years and younger served in the 2020 reporting period varied by race. Trauma- and stressor-related disorders were reported most frequently among American Indians or Alaska Natives (38.4 percent), Native

Hawaiian or Other Pacific Islanders (35.2 percent), Whites (30.9 percent), and Asians (27.3 percent); ADHD was the most frequently reported diagnosis among Blacks or African Americans (39.4 percent).

Ethnicity

Tables 2.3a–2.4c. Among Hispanic individuals aged 17 years and younger served in the 2020 reporting period, the most frequently reported diagnoses were trauma- and stressor-related disorders (28.9 percent); next were ADHD (23.4 percent) and depressive disorders (20.6 percent). Among non-Hispanics, the most frequently reported diagnoses were ADHD (32.8 percent), trauma- and stressor-related disorders (30.6 percent), and anxiety disorders (20.6 percent).

Living Arrangements

Tables 2.5a–2.6c. Among individuals aged 17 years and younger served in the 2020 reporting period, trauma- and stressor-related disorders were the most frequently reported diagnoses among those who were in foster care (56.6 percent), experienced homelessness (44.8 percent), were in residential care (34.3 percent), or were in other living arrangements¹⁴ (30.6 percent). During the same period, individuals aged 17 years and younger living in a private residence were most frequently reported with an ADHD diagnosis (32.1 percent).

SED Status

Tables 2.5a–2.6c. Among individuals aged 17 years and younger with SED served in the 2020 reporting period, the most frequently reported diagnoses were ADHD (29.4 percent) and trauma- and stressor-related disorders (28.7 percent); among those without SED, the most frequently reported diagnoses were trauma- and stressor-related disorders (34.4 percent).

Client Treatment Status at the Start of the Reporting Period

Tables 2.7a–2.8c. Among individuals aged 17 years and younger who were continuing treatment during the 2020 reporting period, the most frequently reported diagnosis was ADHD (35.2 percent); among those who were newly admitted to treatment in the 2020 reporting period, the most frequently reported diagnoses were trauma- and stressor-related disorders (33.8 percent).

SED Status, by Service Setting

Table 2.9. SED status varied by service setting among individuals aged 17 years and younger served in the 2020 reporting period varying from 97.0 percent in institutions under the justice system, to 90.1 percent in state psychiatric hospitals, 77.5 percent in residential treatment centers, 72.3 percent in community-based settings, and 74.4 percent in other psychiatric inpatient settings. It should be noted that individuals may have received services in more than one setting and therefore may be counted in multiple rows of Table 2.9.

¹⁴“Other living arrangements” includes crisis residences and other undefined residences.

Chapter 3

Characteristics of Young Adults Aged 18 to 24 Years Receiving Mental Health Services

This chapter presents data on young adults aged 18 to 24 years who received mental health services through SMHAs in the 2015–2020 reporting periods, with particular focus on the 2020 reporting period. Data on NOMs and clinical measures for all young adults are presented by demographic characteristics and mental health diagnoses. Characteristics for the subpopulation of young adults with SMI are presented where the percentages notably differ from all young adults who received mental health services.

Individuals Served

Tables 3.1a–b present the total number of individuals aged 18 to 24 years served in the 2015 through 2020 reporting periods, by demographic characteristics.

The number of individuals served who were aged 18 to 24 years increased from 521,586 in the 2015 reporting period to 724,047 in the 2020 reporting period.

Tables 3.3a–3.4c. In the 2020 reporting period, 608,665 individuals aged 18 to 24 years had a valid mental health diagnosis code, of which 414,898 were reported to have SMI.

Only individuals with valid mental health diagnoses are included in these and all subsequent tabulations of mental health diagnoses in this chapter.

Gender

Tables 3.3a–c. Females accounted for 54.0 percent of individuals aged 18 to 24 years who were served during the 2020 reporting period with known mental health diagnoses. The most frequently reported diagnoses among males were depressive disorders (29.5 percent), anxiety disorders (22.2 percent), trauma- and stressor-related disorders (16.1 percent), and schizophrenia and other psychotic disorders (14.6 percent). Among females aged 18 to 24 years, the most frequently reported diagnoses were depressive disorders (42.9 percent), anxiety disorders (33.5 percent), trauma- and stressor-related disorders (24.1 percent); schizophrenia and other psychotic disorders were reported in 5.1 percent of females aged 18 to 24 years.

Race and Ethnicity

Tables 3.3a–c. Depressive disorders were the most frequently reported diagnoses among individuals of every racial and ethnic group aged 18 to 24 years served during the 2020 reporting period, ranging from 32.5 percent among Blacks or African Americans to 40.1 percent among Native Hawaiian or Other Pacific Islander. There was no substantial difference of diagnosis frequencies among individuals of Hispanic and Non-Hispanic origin.

Education

Tables 3.5a–3.6c. Among individuals of every education level above 8th grade aged 18 to 24 years served during the 2020 reporting period, the most frequently reported diagnoses were depressive disorders, ranging from 38.8 percent of individuals who completed Grade 9 to 11 to 47.5 percent among individuals who completed at least a year of education beyond Grade 12. Among individuals aged 18 to 24 completing 8th grade educations or less, the most frequently reported diagnoses were anxiety disorders (29.3 percent) and depressive disorders (28.9 percent).

Marital Status

Tables 3.5a–3.6c. Among individuals across all marital status categories aged 18 to 24 years who were served during the 2020 reporting period, the most frequently reported diagnoses were depressive disorders, ranging from 35.3 percent among widowed individuals to 47.7 percent among divorced individuals.

Living Arrangements

Tables 3.7a–3.8c. Among individuals aged 18 to 24 years who were served during the 2020 reporting period, depressive disorders were the most frequently reported diagnoses among those who had other living arrangements (39.4 percent), those who lived in a private residence (39.1 percent), those who were in institutional settings (36.4 percent), those who experienced homelessness (34.2 percent), and those who were in residential care (26.7 percent). Trauma- and stressor-related disorders were the most frequently reported diagnoses for those who were in foster care (33.0 percent) and jail or correctional facilities (31.9 percent).

SMI Status

Tables 3.7a–3.8c. Among individuals aged 18 to 24 years served during the 2020 reporting period, depressive disorders were the most frequently reported diagnoses for those who had SMI (42.8 percent). Anxiety disorders were the most frequently reported diagnosis for those who did not have SMI (25.8 percent).

Veteran Status

Tables 3.7a–3.8c. Among individuals aged 18 to 24 years served during the 2020 reporting period, depressive disorders were the most frequently reported diagnoses for those who were veterans (40.9 percent) and those who were not veterans (35.7 percent).

Employment Status

Tables 3.9a–3.10c. Among individuals aged 18 to 24 years served in the 2020 reporting period, 28.6 percent were employed, 28.8 percent were unemployed, and 42.6 percent were not in the labor force; of those not in the labor force, 44.0 percent were students.

Client Treatment Status at the Start of the Reporting Period

Tables 3.11a–3.12c. Among individuals aged 18 to 24 years served during the 2020 reporting period, the most frequently reported diagnoses were depressive disorders, regardless of individuals' treatment

status at the start of the reporting period (39.5 percent among individuals who were newly admitted to treatment and 33.6 percent among individuals who were continuing treatment).

Service Setting

Tables 3.11a–3.12c. The most frequently reported diagnoses varied by mental health service setting. Individuals may have received services in more than one service setting during the reporting period and therefore may be counted in multiple rows of Tables 3.11a–c and Tables 3.12a–c.

Among individuals aged 18 to 24 years served in the 2020 reporting period, the most frequently reported diagnoses were schizophrenia and other psychotic disorders for those in state psychiatric hospitals (42.1 percent); depressive disorders were the most frequently reported diagnoses among those served in other psychiatric inpatient facilities (40.6 percent), community-based programs (36.9 percent), and residential treatment centers (26.8 percent). Trauma- and stressor-related disorders were the most frequently reported diagnoses among those in institutions under the justice system (35.3 percent).

SMI Status, by Service Setting

Table 3.13. Among individuals aged 18 to 24 years served during the 2020 reporting period, SMI status varied by service setting. It should be noted that individuals may have received services in more than one setting and therefore may be counted in multiple rows of Table 3.13.

Chapter 4

Characteristics of Adults Aged 25 to 54 Years Receiving Mental Health Services

This chapter presents data on adults aged 25 to 54 years who received mental health services through SMHAs in the 2015–2020 reporting periods, with particular focus on the 2020 reporting period. Data on NOMs and clinical measures for all adults aged 25 to 54 years old are presented by demographic characteristics and mental health diagnoses. Characteristics for the subpopulation of adults with SMI are presented where the percentages notably differ from the characteristics of all adults aged 25 to 54 years old.

Individuals Served

Tables 4.1a–b present the total of individuals aged 25 to 54 years reported in the 2015 through 2020 reporting periods, by demographic characteristics.

The number of individuals served who were aged 25 to 54 years increased from 2,377,733 in the 2015 reporting period to 3,123,221 in the 2020 reporting period.

Tables 4.3a–4.4b. In the 2020 reporting period, 2,667,813 individuals aged 25 to 54 years had a valid mental health diagnosis code, of which 1,947,307 were reported to have SMI.

Only individuals with valid mental health diagnoses are included in these and all subsequent tabulations of mental health diagnoses in this chapter.

Gender

Tables 4.3a–4.4c. Females accounted for 54.1 percent of individuals aged 25 to 54 years who were served during the 2020 reporting period. The most frequently reported diagnoses among females were depressive disorders (38.8 percent), anxiety disorders (30.6 percent), and trauma- and stressor-related disorders (23.5 percent). Among males, the most frequently reported diagnoses were depressive disorders (29.0 percent), schizophrenia and other psychotic disorders (24.9 percent), and anxiety disorders (20.9 percent).

Race

Tables 4.3a–4.4c. Among individuals of all races, who were aged 25 to 54 years and were served in the 2020 reporting period, depressive disorders were the most frequently reported diagnoses; depressive disorders were reported among 35.5 percent of Whites, 32.8 percent of Native Hawaiians or Other Pacific Islanders, 32.0 percent of Blacks or African Americans, 32.0 percent of Asians, and 32.4 percent of American Indian or Alaska Natives.

Ethnicity

Tables 4.3a–4.4c. Among individuals aged 25 to 54 years served in the 2020 reporting period, depressive disorders were the most frequently reported diagnoses for both those of non-Hispanic origin (35.1 percent) and Hispanic origin (32.3 percent).

Education

Tables 4.5a–4.6c. Among individuals aged 25 to 54 years served in the 2020 reporting period, depressive disorders were the most frequently reported diagnoses, ranging from 33.7 percent among those who completed at most Grade 8 to 39.4 percent among those who completed at least a year of education beyond Grade 12.

Marital Status

Tables 4.5a–4.6c. Depressive disorders were the most frequently reported diagnoses among individuals aged 25 to 54 years served during the 2020 reporting period across all marital status categories, ranging from 30.5 percent of individuals who never married to 40.1 percent of individuals who were separated. Among individuals in this age group with schizophrenia and other psychotic disorders, 77.4 percent were never married.

Living Arrangements

Tables 4.7a–4.8c. Among individuals aged 25 to 54 years served during the 2020 reporting period who experienced homelessness, lived in a private residence, or had other living arrangements, the most frequently reported diagnoses were depressive disorders, ranging from 32.6 percent of individuals experiencing homelessness to 36.3 percent of individuals living in a private residence. Schizophrenia and other psychotic disorders were the most frequently reported diagnoses among those living in residential care (42.9 percent), in an institutional setting (41.3 percent), or in foster care (27.7 percent). Trauma- and stressor-related disorders were the most frequently reported diagnoses among those living in jails or correctional facilities (28.5 percent).

SMI Status

Tables 4.7a–4.8c. Among individuals with SMI who were aged 25 to 54 years and were served in the 2020 reporting period, the most frequently reported diagnoses were depressive disorders (38.2 percent); the next most frequent diagnoses for those with SMI were anxiety disorders (26.6 percent) and schizophrenia and other psychotic disorders (21.3 percent). Among those without SMI, the most frequent diagnoses were and trauma- and stressor-related disorders (26.0 percent).

Among individuals with schizophrenia and other psychotic disorders who were aged 25 to 54 years and were served in the 2020 reporting period, 92.3 percent were reported to have SMI.

Veteran Status

Tables 4.7a–4.8c. Among individuals aged 25 to 54 years served during the 2020 reporting period, depressive disorders were the most frequently reported diagnoses both for those who were veterans (36.8 percent) and those who were not veterans (33.4 percent).

Veterans and non-veterans did not differ substantially in their frequency of trauma- and stressor-related disorders (19.6 and 20.3 percent, respectively).

Employment Status

Tables 4.9a–4.10c. Among individuals aged 25 to 54 years served in the 2020 reporting period, 28.5 percent were employed, 34.5 percent were unemployed, and 37.0 percent were not in the labor force. Among individuals in this age group, the most frequent diagnoses across all employment statuses were depressive disorders, ranging from 35.2 percent of individuals not in the labor force to 41.8 percent of individuals with full time employment.

Client Treatment Status at the Start of the Reporting Period

Tables 4.11a–4.12c. Among individuals aged 25 to 54 years served in the 2020 reporting period, depressive disorders were the most frequently reported diagnoses, regardless of individuals' treatment status at the start of the reporting period; depressive disorders were reported for 36.6 percent of individuals who were newly admitted to treatment and for 32.4 percent of individuals continuing treatment. The second most frequently reported diagnoses for individuals newly admitted to treatment and for individuals continuing treatment were anxiety disorders (27.8 percent and 24.8 percent, respectively).

Service Setting

Tables 4.11a–4.12c. It should be noted that individuals may have received services in more than one service setting during the reporting period and therefore may be counted in multiple rows of Tables 4.11a–c and 4.12a–c.

Community-based programs accounted for 96.7 percent of individuals aged 25 to 54 years served during the 2020 reporting period. Among individuals aged 25 to 54 years served in community-based programs during the 2020 reporting period, the most frequent diagnoses were depressive disorders (34.6 percent), anxiety disorders (26.8 percent), and trauma- and stressor-related disorders (20.3 percent). Depressive disorders were also the most frequently reported diagnoses among those aged 25 to 54 years who were served during the 2020 reporting period in other psychiatric inpatient settings (36.4 percent). Schizophrenia and other psychotic disorders were the most frequently reported diagnoses among individuals aged 25 to 54 years who were served during the 2020 reporting period in state psychiatric hospitals (56.7 percent) and residential treatment centers (41.2 percent). Among individuals aged 25 to 54 years served in institutions under the justice system during the 2020 reporting period, trauma- and stressor-related disorders were the most frequently reported diagnoses (31.5 percent).

SMI Status, by Service Setting

Table 4.13 presents the number and percentage of individuals aged 25 to 54 years served during the 2020 reporting period, by mental health service setting and SMI status. It should be noted that individuals may have received services in more than one setting and therefore may be counted in multiple rows of Table 4.13.

Chapter 5

Characteristics of Older Adults Aged 55 Years and Older Receiving Mental Health Services

This chapter presents data on individuals aged 55 years and older who received mental health services through SMHAs in the 2015–2020 reporting periods, with particular focus on the 2020 reporting period. Data on NOMs and clinical measures for older adults are presented by diagnostic categories and mental health diagnoses. Characteristics for the subpopulation of older adults with SMI are presented where the percentages notably differ from the characteristics of all individuals aged 55 and older.

Individuals Served

Tables 5.1a–b present the total number of individuals aged 55 years and older served in the 2015 through 2020 reporting periods, by demographic characteristics.

The number of individuals served who were aged 55 years and older increased from 781,018 in the 2015 reporting period to 1,178,713 in the 2020 reporting period.

Tables 5.3a–5.4b. In the 2020 reporting period, 1,010,080 individuals aged 55 years and older had a valid mental health diagnosis code, of which 744,318 were reported to have SMI.

Only individuals with valid mental health diagnoses are included in these and all subsequent tabulations of mental health diagnoses in this chapter.

Gender

Tables 5.3a–5.4c. Females accounted for 58.3 percent of individuals aged 55 years and older who were served in the 2020 reporting period. Among females, the most frequently reported diagnoses were depressive disorders (43.0 percent), anxiety disorders (24.3 percent), and schizophrenia and other psychotic disorders (18.9 percent). Among males, the most frequently reported diagnoses were depressive disorders (33.5 percent), schizophrenia and other psychotic disorders (29.8 percent), and anxiety disorders (16.5 percent).

Race

Tables 5.3a–5.4c. Among individuals of all races, except for Blacks or African Americans, who were aged 55 years and older and were served in the 2020 reporting period, the most frequently reported diagnoses were depressive disorders, ranging from 37.3 percent of American Indians or Alaska Natives to 46.0 percent of Asians; among Blacks or African Americans, the two most frequently reported diagnoses were schizophrenia and other psychotic disorders (38.3 percent) and depressive disorders (34.5 percent).

Ethnicity

Tables 5.3a–5.4c. Among individuals aged 55 years and older served during the 2020 reporting period, the most frequently reported diagnoses were depressive disorders both for those of Hispanic origin (41.9 percent) and non-Hispanic origin (39.0 percent). Schizophrenia and other psychotic disorders were the second most frequently reported diagnoses among non-Hispanics (24.2 percent), while anxiety disorders were the second most frequently reported diagnoses among Hispanics (21.7 percent).

Education

Tables 5.5a–5.6c. Among individuals across all levels of educational attainment who were aged 55 years or older and were served in the 2020 reporting period, the most frequently reported diagnoses were depressive disorders, ranging from 38.7 percent of individuals with a 9th to 11th grade education to 42.7 percent of individuals with less than a 9th grade education.

Marital Status

Tables 5.5a–5.6c. Among never-married individuals aged 55 years and older who were served in the 2020 reporting period, the most frequently reported diagnoses were schizophrenia and other psychotic disorders (35.9 percent). Depressive disorders were the most frequently reported diagnoses for individuals with any other marital status, ranging from 41.7 percent of those divorced to 45.3 percent of those widowed.

Living Arrangements

Tables 5.7a–5.8c. Among individuals aged 55 years and older served in the 2020 reporting period who were living in a private residence, in institutional settings, experienced homelessness, or had other living arrangements, the most frequently reported diagnoses were depressive disorders, ranging from 36.3 percent among those living in institutional settings to 41.6 percent among those living in a private residence. Schizophrenia and other psychotic disorders were the most frequently reported diagnoses among those living in residential care (57.6 percent), foster care (50.4 percent), or a jail or correctional facility (31.6 percent).

SMI Status

Tables 5.7a–5.8c. Among individuals aged 55 years and older served in the 2020 reporting period, the most frequently reported diagnoses were depressive disorders, both among those who had SMI (42.1 percent) and those who did not have SMI (30.1 percent). For those with SMI, the second most frequently reported diagnoses were schizophrenia and other psychotic disorders (28.1 percent); among those without SMI, the second most frequently reported diagnoses were anxiety disorders (21.7 percent).

Veteran Status

Tables 5.7a–5.8c. Among individuals aged 55 years and older served during the 2020 reporting period, depressive disorders were the most frequently reported diagnoses both for those who were veterans (37.3 percent) and those who were not veterans (37.8 percent).

Schizophrenia and other psychotic disorders were reported among 22.8 percent of veterans and 23.2 percent of non-veterans aged 55 years and older who were served in the 2020 reporting period.

Employment Status

Tables 5.9a–5.10c. Among individuals aged 55 years and older served in the 2020 reporting period, 13.1 percent were employed, 22.1 percent were unemployed, and 64.8 percent were not in the labor force (of which 56.2 percent were disabled and 13.4 percent were retired). Among individuals across all employment statuses who were aged 55 years and older and were served in the 2020 reporting period, the most frequently reported diagnoses were depressive disorders, ranging from 41.1 percent among those unemployed to 48.8 percent among those employed full time.

Client Treatment Status at the Start of the Reporting Period

Tables 5.11a–5.12c. Among individuals aged 55 years and older served in the 2020 reporting period, depressive disorders were the most frequently reported diagnoses, regardless of individuals' treatment status at the start of the reporting period; depressive disorders were reported for 41.8 percent of those newly admitted to treatment in the 2020 reporting period and for 37.8 percent of individuals who were continuing treatment. Schizophrenia and other psychotic disorders were the second most frequently reported diagnoses in the same period for individuals aged 55 years and older who were continuing treatment (26.7 percent). Among individuals aged 55 years and older who were newly admitted to treatment in the 2020 reporting period, the second most frequently reported diagnoses were anxiety disorders (23.5 percent).

Service Setting

Tables 5.11a–5.12c. Individuals may have received services in more than one service setting and therefore may be counted in multiple rows of Tables 5.11a–c and 5.12a–c.

Community-based programs accounted for 97.0 percent of individuals aged 55 years and older served during the 2020 reporting period. Depressive disorders were the most frequent diagnoses among individuals aged 55 years and older who were served during the 2020 reporting period in community-based programs (39.5 percent) and other psychiatric inpatient settings (40.2 percent). Schizophrenia and other psychotic disorders were the most frequently reported diagnoses among individuals aged 55 years and older served during the same period in state psychiatric hospitals (59.2 percent), residential treatment centers (51.5 percent), and institutions under the justice system (26.7 percent).

SMI Status, by Service Setting

Table 5.13 presents the number and percentage distribution of individuals aged 55 years and older who were served in the 2020 reporting period by mental health service setting and SMI status. It should be noted that individuals may have received services in more than one setting during the reporting period and therefore may be counted in multiple rows of Table 5.13. During the 2020 reporting period, community-based programs reported the lowest proportion among all service settings of individuals aged 55 years and older who had SMI (72.8 percent); other psychiatric inpatient settings had the highest proportions of individuals aged 55 years and older with SMI (90.8 percent).

Chapter 6

Characteristics of Individuals With Co-occurring Mental Health and Substance Use Disorders Receiving Mental Health Services

This chapter presents data on individuals aged 12 years and older reported to have co-occurring mental health and substance use disorders who received mental health services through SMHAs during the 2015–2020 reporting periods, with particular focus on the 2020 reporting period. Data on NOMs and clinical measures for all individuals are presented by demographic characteristics and mental health diagnoses. It should be noted that the definition for co-occurring mental health and substance use disorders includes those who have a reported, identified substance use disorder but may not have a formal substance use diagnosis in addition to a reported mental health diagnosis. It should also be noted that data reported in this chapter exclude individuals whose only mental health diagnoses were alcohol or substance-related disorders.

Individuals Served

Tables 1.2a and 6.1a. States reported a total of 1,663,657 individuals with co-occurring mental health and substance use disorders aged 12 years and older who were served in the 2020 reporting period, accounting for 24.0 percent of all individuals served during this period.

Mental health diagnoses for individuals with co-occurring mental health and substance use disorders aged 12 and older who were served in the 2020 reporting period differed somewhat by gender, age group, race, ethnicity, living arrangements, employment, not in labor force details, service setting, and timing of admission. Mental health diagnoses for this cohort were largely similar across SMI/SED status.

Gender

Tables 6.2a–c. For males with co-occurring mental health and substance use disorders aged 12 years or older served during the 2020 reporting period, the most frequently reported diagnoses were depressive disorders (31.9 percent), schizophrenia and other psychotic disorders (24.1 percent), and anxiety disorders (19.8 percent). Among their female counterparts, the most frequently reported diagnoses were depressive disorders (40.8 percent), anxiety disorders (27.6 percent), and trauma- and stressor-related disorders (23.8 percent).

Age

Tables 6.2a–c. Among individuals with co-occurring mental health and substance use disorders served in the 2020 reporting period, for all age categories except the 12 to 14 years age group, the most frequently reported diagnoses were depressive disorders, ranging between 34.5 percent in the 30 to 34 years age group and 40.8 percent in the 55 to 59 years age group. Among individuals with

co-occurring mental health and substance use disorders in the 12 to 14 years age group, the most frequently reported diagnoses were trauma- and stressor-related disorders (28.9 percent).

Race

Tables 6.3a–c. Among individuals with co-occurring mental health and substance use disorders served during the 2020 reporting period, the most frequently reported diagnoses were depressive disorders for American Indian or Alaska Natives (42.5 percent), Whites (37.9 percent), Native Hawaiians or Other Pacific Islanders (34.1 percent), Blacks or African Americans (32.1 percent), and Asians (31.6 percent).

Ethnicity

Tables 6.3a–c. Among individuals with co-occurring mental health and substance use disorders served in the 2020 reporting period, the most frequently reported diagnoses were depressive disorders for both those of Hispanic origin (37.3 percent) and non-Hispanic origin (36.6 percent).

Substance Use Diagnosis

Tables 6.4a–c. Among individuals with co-occurring mental health and substance use disorders served during the 2020 reporting period, mental health diagnoses showed no major differences by substance use diagnosis. Depressive disorders were the most frequently reported diagnoses for individuals who had co-occurring alcohol dependence (43.0 percent), opioid dependence (41.7 percent), cocaine dependence (37.0 percent), and marijuana dependence (34.3 percent). Depressive disorders were also the most frequently reported mental health diagnoses for those who had co-occurring non-dependent use of opioids (43.2 percent), alcohol (39.3 percent), marijuana (36.3 percent), and cocaine (35.6 percent). Depressive disorders were also the most frequently reported diagnoses for individuals with alcohol intoxication (45.5 percent), alcoholic psychoses (44.3 percent), and drug psychoses (35.0 percent).

Living Arrangements

Tables 6.6a–c. Among individuals with co-occurring mental health and substance use diagnoses served during the 2020 reporting period, depressive disorders were the most frequently reported diagnoses for those who experienced homelessness (38.2 percent), those living in private residences (38.0 percent), and among those in other living arrangements (36.9 percent). Schizophrenia and other psychotic disorders were the most frequently reported diagnoses among those in residential care (44.7 percent) and institutional settings (37.6 percent). Trauma- and stressor-related disorders were the most frequently reported diagnoses among those in jail/correctional facilities (30.8 percent) and foster care (30.7 percent).

SMI/SED Status

Tables 6.6a–c. Among individuals who had co-occurring mental health and substance use disorders and were served during the 2020 reporting period, depressive disorders were the most frequently reported diagnoses among those with SMI (39.3 percent) and those with SED or at risk of it (41.7 percent). Trauma- and stressor-related disorders were the most frequently reported diagnoses among those without SMI/SED (30.9 percent).

Veteran Status

Tables 6.6a–c. Among individuals who had co-occurring mental health and substance use disorders and who were served during the 2020 reporting period, depressive disorders were the most frequently reported diagnoses both for those who were veterans (39.5 percent) and those who were not (36.8 percent).

Among individuals who had co-occurring mental health and substance use disorders who were served during the 2020 reporting period, schizophrenia and other psychotic disorders were reported among 18.1 percent of veterans and 20.8 percent of non-veterans.

Employment Status

Tables 6.7a–c. Among individuals who had co-occurring mental health and substance use disorders, who were aged 18 years and older, and who were served during the 2020 reporting period, 24.4 percent were employed, 38.5 percent were unemployed, and 37.1 percent were not in the labor force; the most frequently reported diagnoses were depressive disorders across all employment statuses (43.5 percent among the employed, 39.5 percent among the unemployed, and 36.8 percent among those not in the labor force).

Service Setting

Tables 6.8a–c. Individuals may have received services in more than one service setting during the reporting period and therefore may be counted in multiple rows of Tables 6.8a–c.

Among individuals with co-occurring mental health and substance use disorders served during the 2020 reporting period, the most frequently reported diagnosis varied by service setting. Schizophrenia and other psychotic disorders were the most frequently reported diagnoses for those in state psychiatric hospitals (54.8 percent) and residential treatment centers (40.0 percent). Depressive disorders were the most frequently reported diagnoses for those in other psychiatric inpatient settings (39.3 percent) and community-based programs (36.7 percent). Trauma- and stressor-related disorders were the most frequently reported diagnoses for those in institutions under the justice system (34.9 percent).

Client Treatment Status at the Start of the Reporting Period

Tables 6.8a–c. Among individuals who had co-occurring mental health and substance use disorders and who were served during the 2020 reporting period, depressive disorders were the most frequently reported diagnoses, regardless of individuals' treatment status at the start of the reporting period; depressive disorders were reported for 38.0 percent of those newly admitted to treatment in the 2020 reporting period and for 34.5 percent of individuals who were continuing treatment.

Chapter 7

Client Outcomes Among Individuals Receiving Mental Health Services: 2015–2020

This chapter presents data on cohorts of individuals who received mental health services through SMHAs during two or more reporting periods. While the following narrative focuses specifically on a cohort of individuals served in both the 2017 and 2020 reporting periods [Tables 7.1a–7.6b], additional tables for Chapter 7 [Tables 7.7–7.16b] encompass cohorts of individuals served across various other combinations of reporting periods between 2015 and 2020. The length of treatment was not taken into consideration. NOMs are presented by demographic characteristics, mental health diagnoses, and clinical measures for all individuals, based on the last reported status in each period.

Individuals were selected for analysis by matching any individual records in 2017 with any in 2020 from the same state and with the same unique identification number; other individuals served in the 2017 and 2020 reporting periods were excluded from analysis in this chapter. The resulting data set contained 1,296,090 unique clients [Table 7.1a].

Living Arrangements

Tables 7.1a–7.2c. States reported a total of 1,296,090 individuals served in both the 2017 and 2020 reporting periods who had known living arrangements in both reporting years.

Among individuals served in both the 2017 and 2020 reporting periods, those living in a private residence accounted for 85.5 percent in 2017 and 85.3 percent in 2020; those living in residential care accounted for 5.5 and 5.7 percent respectively in 2017 and 2020; those who experienced homelessness accounted for 3.1 and 3.4 percent respectively in 2017 and 2020; those who were in institutional settings accounted for 1.7 percent in both 2017 and 2020; those in jails/correctional facilities accounted for 1.2 percent in both 2017 and 2020; and those in foster care accounted for 1.4 percent in both 2017 and 2020.

Employment Status


Tables 7.3a–7.4c. States reported a total of 665,210 individuals aged 18 years and older in 2020 who were served in both the 2017 and 2020 reporting periods and who had known employment status in both years. Among individuals aged 18 years and older in 2020 who were served in both the 2017 and 2020 reporting periods, those who were not in the labor force accounted for 58.3 and 58.5 percent respectively in 2017 and 2020; those who were unemployed accounted for 22.9 and 22.3 percent respectively in 2017 and 2020; and those who were employed accounted for 18.8 and 19.2 percent respectively in 2017 and 2020.

Transitions in Living Arrangements

Individuals living in a private residence in 2017

Tables 7.5a–b and Figure 2. Among individuals served in both the 2017 and 2020 reporting periods, 94.9 percent of those who lived in a private residence in 2017 also reported living in a private residence in 2020; 1.9 percent transitioned to experiencing homelessness; 1.4 percent transitioned to residential care; and 1.7 percent, combined, transitioned to institutional settings, jail, or other living arrangements.

Figure 2. Living arrangements in 2020 of individuals aged 18 years and older who reported living in a private residence in 2017, by 2017 mental health diagnosis




NOTES: Percentages may not sum to 100 percent due to rounding. Less than 1 percent of individuals served in 2017 and 2020 moved into foster homes or foster care in 2020.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2017 and 2020 reporting periods.

Individuals who experienced homelessness in 2017

Tables 7.5a–b and Figure 3. Among individuals aged 18 years and older served in both the 2017 and 2020 reporting periods, 54.7 percent who experienced homelessness in 2017 also reported experiencing homelessness in 2020; 35.9 percent transitioned to a private residence, 4.3 percent to residential care, 1.3 percent to institutional settings, 1.4 percent to jail or correctional facilities, and 2.2 percent to other living arrangements.

Figure 3. Living arrangements in 2020 of individuals aged 18 years and older who experienced homelessness in 2017, by 2017 mental health diagnosis




NOTES: Percentages may not sum to 100 percent due to rounding. Less than 1 percent of individuals served in 2017 and 2020 moved into foster homes or foster care in 2020.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2017 and 2020 reporting periods.

Individuals in jail or correctional facilities in 2017

Tables 7.5a–b and Figure 4. Among individuals aged 18 years and older served in both the 2017 and 2020 reporting periods, 61.4 percent of those who were in jail or correctional facilities in 2017 also reported being in jail or correctional facilities in 2020; 23.9 percent transitioned to private residences, 5.6 percent to experiencing homelessness, 4.6 percent to residential care, and 2.9 percent to institutional settings.

Figure 4. Living arrangements in 2020 of individuals aged 18 years and older who reported being in jail or a correctional facility in 2017, by 2017 mental health diagnosis



NOTES: Percentages may not sum to 100 percent due to rounding. Less than 1 percent of individuals served in 2017 and 2020 moved into foster homes or foster care in 2020.


SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2017 and 2020 reporting periods.

Transitions in Employment Status

Individuals who were employed in 2017

Tables 7.6a–b and Figure 5. Among individuals aged 18 years and older who were served in both the 2017 and 2020 reporting periods, 68.9 percent of those who were employed in 2017 also reported being employed in 2020; 11.4 percent transitioned to being unemployed, and 19.8 percent had left the labor force.

Figure 5. Employment status in 2020 of individuals aged 18 years and older who reported being employed in 2017, by 2017 mental health diagnosis




NOTES: Percentages may not sum to 100 percent due to rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2017 and 2020 reporting periods.

Individuals who were unemployed in 2017

Tables 7.6a–b and Figure 6. Among individuals served in both the 2017 and 2020 reporting periods, 69.2 percent of those who were unemployed in 2017 also reported being unemployed in 2020; 20.0 percent of those unemployed in 2017 reported in 2020 that they were not in the labor force, and 10.8 percent reported that they were employed.

Figure 6. Employment status in 2020 of individuals aged 18 years and older who reported being unemployed in 2017, by 2017 mental health diagnosis




NOTES: Percentages may not sum to 100 percent due to rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2017 and 2020 reporting periods.

Individuals who were not in the labor force in 2017

Tables 7.6a–b and Figure 7. Among individuals served in both the 2017 and 2020 reporting periods overall, 85.7 percent of those who were not in the labor force in 2017 also reported not being in the labor force in 2020; 7.7 percent transitioned to unemployment, and 6.6 percent transitioned to employment in 2020. The pattern of transition from not in the labor force to other employment statuses was similar among individuals served with bipolar and depressive disorders compared with individuals across all selected mental health diagnoses combined.

Figure 7. Employment status in 2020 of individuals aged 18 years and older who were not in the labor force in 2017, by 2017 mental health diagnosis



NOTES: Percentages may not sum to 100 percent due to rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2017 and 2020 reporting periods.

Chapter 8

Geographic and Clinical Trends: 2015–2020

It is important to note that differences among reporting periods should be interpreted with great caution; variation may be more a function of the mix and number of reporting states than of actual differences. Throughout this chapter, states/jurisdictions may refer to any of the 50 states, the District of Columbia, the Commonwealth of Puerto Rico, and the combined jurisdictions of the Republic of Palau and the Commonwealth of the Northern Mariana Islands.

Table 8.1 presents the number of individuals served in the 2020 reporting period, by mental health diagnosis, Census region, Census division, and state or jurisdiction.

Tables 8.2 and 8.3 present, respectively, the number and adjusted number of individuals per 100,000 population served in the 2020 reporting period, by mental health diagnosis, Census region, Census division, and state or jurisdiction.¹⁵

Among individuals served in the 2020 reporting period, the mental health diagnoses with the largest per 100,000 population ratios (adjusted for age, gender, and ethnicity) were depressive disorders (535 per 100,000 population), anxiety disorders (415 per 100,000 population), and trauma- and stressor-related disorders (381 per 100,000 population) [Table 8.3].

Across all Census divisions in the 2020 reporting period except the Mountain division, depressive disorders were the mental health diagnosis with the highest adjusted ratio per 100,000 population, ranging from 256 per 100,000 population for New England to 1,048 per 100,000 population for the West North Central division. Anxiety disorders were the mental health diagnosis with the highest adjusted ratio per 100,000 population in the Mountain division (676 per 100,000 population) [Table 8.3].

In 29 of the 47 reporting states and jurisdictions, depressive disorders were the diagnoses with the highest adjusted ratio of individuals per 100,000 population. In Alaska, Colorado, New Mexico, South Dakota, and Vermont, trauma and stressor-related disorders were the diagnoses with the highest adjusted ratio of individuals per 100,000 population. In Arizona, Iowa, Minnesota, Montana, North Dakota, Oregon, Utah, Washington, and Wisconsin, anxiety disorders were the diagnoses with the highest adjusted ratio of individuals per 100,000 population. In Hawaii, Idaho, Massachusetts, and New York, schizophrenia and other psychotic disorders were the diagnoses with the highest ratio of individuals per 100,000 population [Table 8.3].

Tables 8.4–8.9 present—by Census region, Census division, and state or jurisdiction—the number of individuals served in the 2015 through 2020 reporting periods, grouped by the following diagnoses:

¹⁵The adjusted number of individuals per 100,000 population served in the 2020 reporting period is calculated based on age, sex, and race of the mid-year 2010 population (according to vintage 2020 estimates). States that submitted insufficient data for inclusion in this report, as well as their reference populations, are excluded from these calculations. U.S. Census Bureau, Population Estimates Program, SC-EST2020-alldata6: “Annual State Resident Population Estimates for 6 Race Groups (5 Race Alone Groups and Two or More Races) by Age, Sex, and Hispanic Origin: April 1, 2010, to July 1, 2020.” <https://www2.census.gov/programs-surveys/popest/datasets/2010-2020/state/asrh/SC-EST2020-ALLDATA6.csv>.

trauma- and stressor-related disorders [Table 8.4], anxiety disorders [Table 8.5], attention deficit/hyperactivity disorder [Table 8.6], bipolar disorders [Table 8.7], depressive disorders [Table 8.8], and schizophrenia and other psychotic disorders [Table 8.9].

Table 8.10 presents the number of individuals with co-occurring mental health and substance use disorders who were served in the 2015 through 2020 reporting periods, by Census region, Census division, and state or jurisdiction. It should be noted that the definition for co-occurring mental health and substance use disorders includes those who have a reported, identified substance use disorder but may not have a formal substance use diagnosis in addition to a reported mental health diagnosis. It should also be noted that data reported in this chapter exclude individuals whose only mental health diagnoses were substance use disorders.

Chapter 9

Client Characteristics by State or Jurisdiction: 2015–2020

Tables 9.1a–9.52b present, for each state or jurisdiction, the number and percent distribution of individuals served in the 2015 through 2020 reporting periods, according to mental health diagnosis, gender, age, race, ethnicity, and veteran status.

Chapter 10

Client Characteristics by State or Jurisdiction: 2020

Tables 10.1a–10.49b present, for each state or jurisdiction, the number and percent distribution of individuals served in the 2020 reporting period, according to mental health diagnosis, gender, age, race, ethnicity, and veteran status.

States that submitted no or insufficient data for the 2020 reporting period—Maine, Maryland, New Hampshire, and West Virginia—do not appear in tables for this chapter.

Tables

Table 1.1a. Individuals, by mental health diagnosis: Number, 2015–2020

<i>Mental health diagnosis¹</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Number of states reporting	45	47	49	49	48	50
Total	5,165,254	5,656,750	6,171,990	6,418,247	6,548,372	6,945,521
Trauma- and stressor-related disorders	853,482	878,639	1,035,177	1,161,339	1,253,914	1,295,221
Anxiety disorders	755,391	837,961	1,052,774	1,239,939	1,339,890	1,404,997
Attention deficit/hyperactivity disorder+	510,737	560,910	621,999	696,888	672,592	654,495
Conduct disorder+	147,020	139,246	135,740	133,315	128,455	117,672
Delirium/dementia	33,432	26,246	26,082	29,837	29,147	31,576
Mood disorders	2,219,816	2,132,801	2,278,835	2,439,327	2,479,529	2,504,101
Bipolar disorders	855,430	747,805	724,193	734,667	704,177	713,775
Depressive disorders	1,419,968	1,429,859	1,597,103	1,756,277	1,824,785	1,834,097
Oppositional defiant disorder+	165,302	168,642	191,914	206,728	201,124	181,674
Pervasive developmental disorders+	69,481	94,969	98,479	109,585	96,230	110,316
Personality disorders	169,840	147,069	165,439	174,277	163,177	161,378
Schizophrenia and other psychotic disorders	640,161	622,833	728,180	770,554	742,888	765,575
Alcohol or substance-related disorders	269,224	234,950	244,899	273,451	296,773	384,246
Other mental disorders ²	1,186,202	849,710	713,449	858,184	860,407	885,016
No/deferred diagnosis ³	650,131	1,014,081	911,364	752,807	769,603	1,016,968

+ Indicates diagnoses typically identified in childhood (age < 18 years).

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ No/deferred diagnosis includes unknown, not collected, missing, invalid, or other non-mental health diagnoses.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 1.1b. Individuals, by mental health diagnosis: Column percent distribution, 2015–2020

<i>Mental health diagnosis¹</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Number of states reporting	45	47	49	49	48	50
Total	100.0	100.0	100.0	100.0	100.0	100.0
Trauma- and stressor-related disorders	16.5	15.5	16.8	18.1	19.1	18.6
Anxiety disorders	14.6	14.8	17.1	19.3	20.5	20.2
Attention deficit/hyperactivity disorder+	9.9	9.9	10.1	10.9	10.3	9.4
Conduct disorder+	2.8	2.5	2.2	2.1	2.0	1.7
Delirium/dementia	†	†	†	†	†	†
Mood disorders	43.0	37.7	36.9	38.0	37.9	36.1
Bipolar disorders	16.6	13.2	11.7	11.4	10.8	10.3
Depressive disorders	27.5	25.3	25.9	27.4	27.9	26.4
Oppositional defiant disorder+	3.2	3.0	3.1	3.2	3.1	2.6
Pervasive developmental disorders+	1.3	1.7	1.6	1.7	1.5	1.6
Personality disorders	3.3	2.6	2.7	2.7	2.5	2.3
Schizophrenia and other psychotic disorders	12.4	11.0	11.8	12.0	11.3	11.0
Alcohol or substance-related disorders	5.2	4.2	4.0	4.3	4.5	5.5
Other mental disorders ²	23.0	15.0	11.6	13.4	13.1	12.7
No/deferred diagnosis ³	12.6	17.9	14.8	11.7	11.8	14.6

+ Indicates diagnoses typically identified in childhood (age < 18 years); † Less than 1 percent of individuals.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ No/deferred diagnosis includes unknown, not collected, missing, invalid, or other non-mental health diagnoses.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 1.2a. Individuals, by gender and age: Number, 2015–2020

<i>Gender and age</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Number of states reporting	45	47	49	49	48	50
Total	5,165,254	5,656,750	6,171,990	6,418,247	6,548,372	6,945,521
Gender						
Male	2,501,184	2,726,456	2,989,848	3,117,023	3,153,702	3,349,353
Female	2,647,114	2,901,222	3,153,644	3,284,966	3,375,245	3,574,913
Total	5,148,298	5,627,678	6,143,492	6,401,989	6,528,947	6,924,266
Age¹						
0 to 11 years	710,211	789,831	814,625	899,746	918,686	922,028
12 to 14 years	365,574	390,461	405,592	446,037	478,309	493,952
15 to 17 years	397,372	430,652	450,187	476,732	494,610	499,219
18 to 20 years	225,984	250,346	273,807	293,478	310,263	325,599
21 to 24 years	295,602	323,282	353,177	359,746	366,599	398,448
25 to 29 years	417,692	471,105	536,765	557,084	568,472	614,608
30 to 34 years	424,129	467,115	521,379	535,903	547,828	607,046
35 to 39 years	376,740	422,607	480,949	500,964	509,507	559,204
40 to 44 years	359,482	377,450	404,802	411,461	416,636	462,221
45 to 49 years	381,214	405,443	434,660	431,728	418,937	434,063
50 to 54 years	418,476	442,656	465,607	453,612	434,856	446,079
55 to 59 years	342,046	377,066	418,358	429,411	429,053	451,653
60 to 64 years	215,213	242,842	278,831	297,124	307,640	336,524
65 years and older	223,759	252,091	322,359	320,124	341,885	390,536
Total	5,153,494	5,642,947	6,161,098	6,413,150	6,543,281	6,941,180

¹ Age is based on the individual's age at the midpoint of the reporting period.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 1.2b. Individuals, by gender and age: Column percent distribution, 2015–2020

<i>Gender and age</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Number of states reporting	45	47	49	49	48	50
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	48.6	48.4	48.7	48.7	48.3	48.4
Female	51.4	51.6	51.3	51.3	51.7	51.6
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age¹						
0 to 11 years	13.8	14.0	13.2	14.0	14.0	13.3
12 to 14 years	7.1	6.9	6.6	7.0	7.3	7.1
15 to 17 years	7.7	7.6	7.3	7.4	7.6	7.2
18 to 20 years	4.4	4.4	4.4	4.6	4.7	4.7
21 to 24 years	5.7	5.7	5.7	5.6	5.6	5.7
25 to 29 years	8.1	8.3	8.7	8.7	8.7	8.9
30 to 34 years	8.2	8.3	8.5	8.4	8.4	8.7
35 to 39 years	7.3	7.5	7.8	7.8	7.8	8.1
40 to 44 years	7.0	6.7	6.6	6.4	6.4	6.7
45 to 49 years	7.4	7.2	7.1	6.7	6.4	6.3
50 to 54 years	8.1	7.8	7.6	7.1	6.6	6.4
55 to 59 years	6.6	6.7	6.8	6.7	6.6	6.5
60 to 64 years	4.2	4.3	4.5	4.6	4.7	4.8
65 years and older	4.3	4.5	5.2	5.0	5.2	5.6
Total	100.0	100.0	100.0	100.0	100.0	100.0

¹ Age is based on the individual's age at the midpoint of the reporting period.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 1.3a. Individuals, by race and ethnicity: Number, 2015–2020

<i>Race and ethnicity</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Number of states reporting	45	47	49	49	48	50
Total	5,165,254	5,656,750	6,171,990	6,418,247	6,548,372	6,945,521
Race						
American Indian or Alaska Native	69,240	71,774	78,927	89,207	101,338	133,372
Asian ¹	63,803	74,829	85,680	94,950	90,574	91,731
Black or African American	981,718	1,058,487	1,157,760	1,197,807	1,242,263	1,230,018
Native Hawaiian or Other Pacific Islander	13,637	15,566	10,996	13,984	15,320	15,293
White	3,136,010	3,469,322	3,743,980	3,892,086	3,889,169	4,137,523
Other	432,966	480,702	558,153	609,153	642,858	682,949
Total	4,697,374	5,170,680	5,635,496	5,897,187	5,981,522	6,290,886
Ethnicity						
Not of Hispanic or Latino origin	3,470,405	3,781,453	4,639,457	4,746,793	4,802,670	5,015,183
Hispanic or Latino origin	700,094	771,838	899,234	945,157	977,132	1,241,591
Total	4,170,499	4,553,291	5,538,691	5,691,950	5,779,802	6,256,774
Specific Hispanic or Latino origin						
Mexican	55,759	43,840	52,764	52,613	54,828	52,863
Puerto Rican	51,894	44,178	51,200	27,842	27,155	28,561
Cuban	9,609	8,925	8,821	9,598	9,659	11,049
Other specific	52,113	51,422	59,103	60,670	50,729	54,642
Total	169,375	148,365	171,888	150,723	142,371	147,115

¹ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 1.3b. Individuals, by race and ethnicity: Column percent distribution, 2015–2020

<i>Race and ethnicity</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Number of states reporting	45	47	49	49	48	50
Total	100.0	100.0	100.0	100.0	100.0	100.0
Race						
American Indian or Alaska Native	1.5	1.4	1.4	1.5	1.7	2.1
Asian ¹	1.4	1.4	1.5	1.6	1.5	1.5
Black or African American	20.9	20.5	20.5	20.3	20.8	19.6
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	66.8	67.1	66.4	66.0	65.0	65.8
Other	9.2	9.3	9.9	10.3	10.7	10.9
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	83.2	83.0	83.8	83.4	83.1	80.2
Hispanic or Latino origin	16.8	17.0	16.2	16.6	16.9	19.8
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	32.9	29.5	30.7	34.9	38.5	35.9
Puerto Rican	30.6	29.8	29.8	18.5	19.1	19.4
Cuban	5.7	6.0	5.1	6.4	6.8	7.5
Other specific	30.8	34.7	34.4	40.3	35.6	37.1
Total	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

¹ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 1.4a. Individuals aged 17 years and younger, by mental health diagnosis: Number, 2015–2020

<i>Mental health diagnosis¹</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Number of states reporting	45	47	49	49	48	50
Total individuals aged 0 to 17 years	1,473,157	1,610,944	1,670,404	1,822,515	1,891,605	1,915,199
Trauma- and stressor-related disorders	342,828	369,110	401,584	466,743	512,144	496,671
Anxiety disorders	174,269	201,398	241,484	289,008	321,681	322,311
Attention deficit/hyperactivity disorder+	400,594	442,962	476,533	528,942	504,773	481,629
Conduct disorder	132,323	124,330	112,268	109,868	104,510	93,111
Mood disorders	346,026	308,921	280,054	310,595	339,839	326,670
Bipolar disorders	129,000	84,347	44,013	36,917	30,784	27,577
Depressive disorders	223,842	228,960	238,691	276,414	311,619	301,271
Oppositional defiant disorder	150,151	158,241	173,032	187,997	182,926	162,854
Pervasive developmental disorders	46,923	69,239	67,327	71,563	61,616	69,912
Alcohol or substance-related disorders	14,474	14,582	16,023	14,527	14,456	14,469
Other mental disorders ²	423,501	357,337	234,264	296,814	303,884	306,827
No/deferred diagnosis ³	177,170	198,670	200,834	192,940	211,591	277,232

+ Indicates diagnoses typically identified in childhood (age < 18 years).

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ No/deferred diagnosis includes unknown, not collected, missing, invalid, or other non-mental health diagnoses.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 1.4b. Individuals aged 17 years and younger, by mental health diagnosis: Column percent distribution, 2015–2020

<i>Mental health diagnosis¹</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Number of states reporting	45	47	49	49	48	50
Total individuals aged 0 to 17 years	100.0	100.0	100.0	100.0	100.0	100.0
Trauma- and stressor-related disorders	23.3	22.9	24.0	25.6	27.1	25.9
Anxiety disorders	11.8	12.5	14.5	15.9	17.0	16.8
Attention deficit/hyperactivity disorder+	27.2	27.5	28.5	29.0	26.7	25.1
Conduct disorder	9.0	7.7	6.7	6.0	5.5	4.9
Mood disorders	23.5	19.2	16.8	17.0	18.0	17.1
Bipolar disorders	8.8	5.2	2.6	2.0	1.6	1.4
Depressive disorders	15.2	14.2	14.3	15.2	16.5	15.7
Oppositional defiant disorder	10.2	9.8	10.4	10.3	9.7	8.5
Pervasive developmental disorders	3.2	4.3	4.0	3.9	3.3	3.7
Alcohol or substance-related disorders	†	†	†	†	†	†
Other mental disorders ²	28.7	22.2	14.0	16.3	16.1	16.0
No/deferred diagnosis ³	12.0	12.3	12.0	10.6	11.2	14.5

+ Indicates diagnoses typically identified in childhood (age < 18 years); † Less than 1 percent of individuals.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ No/deferred diagnosis includes unknown, not collected, missing, invalid, or other non-mental health diagnoses.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 1.5a. Individuals aged 18 to 24 years, by mental health diagnosis: Number, 2015–2020

<i>Mental health diagnosis¹</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Number of states reporting	45	47	49	49	48	50
Total individuals aged 18 to 24 years	521,586	573,628	626,984	653,224	676,862	724,047
Trauma- and stressor-related disorders	80,077	80,595	95,492	106,465	117,690	124,460
Anxiety disorders	82,194	92,974	117,460	142,779	160,820	172,077
Attention deficit/hyperactivity disorder+	45,238	46,266	55,491	61,571	59,923	59,814
Mood disorders	244,686	235,569	244,204	270,279	286,089	291,508
Bipolar disorders	102,853	84,978	75,266	75,828	73,865	73,317
Depressive disorders	148,514	155,944	173,671	200,372	218,018	223,326
Personality disorders	16,711	14,805	17,129	18,668	18,720	18,210
Schizophrenia and other psychotic disorders	52,130	51,073	56,553	59,182	57,181	57,417
Alcohol or substance-related disorders	31,750	30,211	27,825	30,160	31,355	38,462
Other mental disorders ²	142,876	115,230	109,894	125,055	122,313	126,760
No/deferred diagnosis ³	73,712	113,635	109,194	91,954	89,845	115,382

+ Indicates diagnoses typically identified in childhood (age < 18 years).

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ No/deferred diagnosis includes unknown, not collected, missing, invalid, or other non-mental health diagnoses.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 1.5b. Individuals aged 18 to 24 years, by mental health diagnosis: Column percent distribution, 2015–2020

<i>Mental health diagnosis¹</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Number of states reporting	45	47	49	49	48	50
Total individuals aged 18 to 24 years	100.0	100.0	100.0	100.0	100.0	100.0
Trauma- and stressor-related disorders	15.4	14.1	15.2	16.3	17.4	17.2
Anxiety disorders	15.8	16.2	18.7	21.9	23.8	23.8
Attention deficit/hyperactivity disorder+	8.7	8.1	8.9	9.4	8.9	8.3
Mood disorders	46.9	41.1	38.9	41.4	42.3	40.3
Bipolar disorders	19.7	14.8	12.0	11.6	10.9	10.1
Depressive disorders	28.5	27.2	27.7	30.7	32.2	30.8
Personality disorders	3.2	2.6	2.7	2.9	2.8	2.5
Schizophrenia and other psychotic disorders	10.0	8.9	9.0	9.1	8.4	7.9
Alcohol or substance-related disorders	6.1	5.3	4.4	4.6	4.6	5.3
Other mental disorders ²	27.4	20.1	17.5	19.1	18.1	17.5
No/deferred diagnosis ³	14.1	19.8	17.4	14.1	13.3	15.9

+ Indicates diagnoses typically identified in childhood (age < 18 years); † Less than 1 percent of individuals.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ No/deferred diagnosis includes unknown, not collected, missing, invalid, or other non-mental health diagnoses.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 1.6a. Individuals aged 25 to 54 years, by mental health diagnosis: Number, 2015–2020

<i>Mental health diagnosis¹</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Number of states reporting	45	47	49	49	48	50
Total individuals aged 25 to 54 years	2,377,733	2,586,376	2,844,162	2,890,752	2,896,236	3,123,221
Trauma- and stressor-related disorders	351,786	353,438	429,776	471,563	499,051	532,647
Anxiety disorders	394,114	429,853	540,116	627,453	663,300	697,741
Attention deficit/hyperactivity disorder+	59,056	64,605	80,561	96,101	97,427	101,545
Mood disorders	1,233,210	1,200,187	1,290,941	1,356,111	1,345,453	1,350,315
Bipolar disorders	503,196	462,356	472,147	480,124	460,666	463,761
Depressive disorders	764,501	766,130	846,568	909,837	916,864	914,629
Personality disorders	117,581	100,940	111,376	115,822	107,097	105,153
Schizophrenia and other psychotic disorders	394,841	384,495	439,456	464,306	444,316	460,519
Alcohol or substance-related disorders	181,536	158,782	159,940	181,216	197,999	262,739
Other mental disorders ²	501,004	332,670	320,011	372,099	363,045	376,509
No/deferred diagnosis ³	294,714	487,863	438,183	343,530	331,742	455,408

+ Indicates diagnoses typically identified in childhood (age < 18 years).

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ No/deferred diagnosis includes unknown, not collected, missing, invalid, or other non-mental health diagnoses.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 1.6b. Individuals aged 25 to 54 years, by mental health diagnosis: Column percent distribution, 2015–2020

<i>Mental health diagnosis¹</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Number of states reporting	45	47	49	49	48	50
Total individuals aged 25 to 54 years	100.0	100.0	100.0	100.0	100.0	100.0
Trauma- and stressor-related disorders	14.8	13.7	15.1	16.3	17.2	17.1
Anxiety disorders	16.6	16.6	19.0	21.7	22.9	22.3
Attention deficit/hyperactivity disorder+	2.5	2.5	2.8	3.3	3.4	3.3
Mood disorders	51.9	46.4	45.4	46.9	46.5	43.2
Bipolar disorders	21.2	17.9	16.6	16.6	15.9	14.8
Depressive disorders	32.2	29.6	29.8	31.5	31.7	29.3
Personality disorders	4.9	3.9	3.9	4.0	3.7	3.4
Schizophrenia and other psychotic disorders	16.6	14.9	15.5	16.1	15.3	14.7
Alcohol or substance-related disorders	7.6	6.1	5.6	6.3	6.8	8.4
Other mental disorders ²	21.1	12.9	11.3	12.9	12.5	12.1
No/deferred diagnosis ³	12.4	18.9	15.4	11.9	11.5	14.6

+ Indicates diagnoses typically identified in childhood (age < 18 years); † Less than 1 percent of individuals.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ No/deferred diagnosis includes unknown, not collected, missing, invalid, or other non-mental health diagnoses.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 1.7a. Individuals aged 55 years and older, by mental health diagnosis: Number, 2015–2020

<i>Mental health diagnosis¹</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Number of states reporting	45	47	49	49	48	50
Total individuals aged 55 years and older	781,018	871,999	1,019,548	1,046,659	1,078,578	1,178,713
Trauma- and stressor-related disorders	77,941	73,897	106,774	115,827	123,919	140,825
Anxiety disorders	104,155	112,470	152,455	180,092	193,124	212,400
Delirium/dementia	21,020	16,617	17,800	20,154	20,225	22,719
Mood disorders	391,062	382,483	460,118	500,920	506,456	534,843
Bipolar disorders	118,279	113,898	131,504	141,445	138,518	148,935
Depressive disorders	280,287	275,297	335,871	368,556	376,883	394,254
Personality disorders	31,889	27,406	33,611	36,395	34,412	35,333
Schizophrenia and other psychotic disorders	177,180	171,580	216,573	233,583	228,777	236,477
Alcohol or substance-related disorders	39,924	29,761	40,120	47,283	52,634	68,119
Other mental disorders ²	181,515	108,362	131,386	151,802	153,735	162,424
No/deferred diagnosis ³	103,441	212,879	161,566	123,070	135,791	168,633

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ No/deferred diagnosis includes unknown, not collected, missing, invalid, or other non-mental health diagnoses.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 1.7b. Individuals aged 55 years and older, by mental health diagnosis: Column percent distribution, 2015–2020

<i>Mental health diagnosis¹</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Number of states reporting	45	47	49	49	48	50
Total individuals aged 55 years and older	100.0	100.0	100.0	100.0	100.0	100.0
Trauma- and stressor-related disorders	10.0	8.5	10.5	11.1	11.5	11.9
Anxiety disorders	13.3	12.9	15.0	17.2	17.9	18.0
Delirium/dementia	2.7	1.9	1.7	1.9	1.9	1.9
Mood disorders	50.1	43.9	45.1	47.9	47.0	45.4
Bipolar disorders	15.1	13.1	12.9	13.5	12.8	12.6
Depressive disorders	35.9	31.6	32.9	35.2	34.9	33.4
Personality disorders	4.1	3.1	3.3	3.5	3.2	3.0
Schizophrenia and other psychotic disorders	22.7	19.7	21.2	22.3	21.2	20.1
Alcohol or substance-related disorders	5.1	3.4	3.9	4.5	4.9	5.8
Other mental disorders ²	23.2	12.4	12.9	14.5	14.3	13.8
No/deferred diagnosis ³	13.2	24.4	15.8	11.8	12.6	14.3

† Less than 1 percent of individuals.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ No/deferred diagnosis includes unknown, not collected, missing, invalid, or other non-mental health diagnoses.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 1.8a. Individuals, by marital status and living arrangement: Number, 2015–2020

<i>Marital status and living arrangement</i>		<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Number of states reporting		45	47	49	49	48	50
Total		5,165,254	5,656,750	6,171,990	6,418,247	6,548,372	6,945,521
Marital status							
Total individuals aged 18 years and older		3,680,337	4,032,003	4,490,694	4,590,635	4,651,676	5,025,981
Never married		1,079,592	1,294,949	1,598,751	1,770,697	1,622,157	1,735,105
Currently married		340,806	359,596	395,587	347,966	409,107	414,207
Separated		132,951	152,003	158,900	142,841	162,069	164,344
Divorced		330,081	350,710	394,449	358,704	397,090	376,627
Widowed		61,882	69,157	76,300	75,960	80,749	97,170
Total		1,945,312	2,226,415	2,623,987	2,696,168	2,671,172	2,787,453
Living arrangement							
Homeless		113,877	115,064	140,380	159,289	173,330	178,804
Foster home/foster care		47,826	41,547	51,837	52,895	54,095	57,536
Residential care		120,538	113,317	144,599	144,090	149,831	136,341
Institutional setting		53,520	48,984	56,109	62,006	72,775	82,933
Jail/correctional facility		61,764	58,322	64,690	67,413	69,632	64,375
Private residence		2,713,945	2,643,459	3,053,623	3,237,767	3,565,364	3,610,160
Other		120,073	140,242	142,893	84,964	82,993	63,346
Total		3,231,543	3,160,935	3,654,131	3,808,424	4,168,020	4,193,495

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 1.8b. Individuals, by marital status and living arrangement: Column percent distribution, 2015–2020

<i>Marital status and living arrangement</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Number of states reporting	45	47	49	49	48	50
Total	100.0	100.0	100.0	100.0	100.0	100.0
Marital status						
Total individuals aged 18 years and older	100.0	100.0	100.0	100.0	100.0	100.0
Never married	55.5	58.2	60.9	65.7	60.7	62.2
Currently married	17.5	16.2	15.1	12.9	15.3	14.9
Separated	6.8	6.8	6.1	5.3	6.1	5.9
Divorced	17.0	15.8	15.0	13.3	14.9	13.5
Widowed	3.2	3.1	2.9	2.8	3.0	3.5
Total	100.0	100.0	100.0	100.0	100.0	100.0
Living arrangement						
Homeless	3.5	3.6	3.8	4.2	4.2	4.3
Foster home/foster care	1.5	1.3	1.4	1.4	1.3	1.4
Residential care	3.7	3.6	4.0	3.8	3.6	3.3
Institutional setting	1.7	1.5	1.5	1.6	1.7	2.0
Jail/correctional facility	1.9	1.8	1.8	1.8	1.7	1.5
Private residence	84.0	83.6	83.6	85.0	85.5	86.1
Other	3.7	4.4	3.9	2.2	2.0	1.5
Total	100.0	100.0	100.0	100.0	100.0	100.0

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 1.9a. Individuals, by SMI/SED status and co-occurring mental health and substance use disorder status: Number, 2015–2020

<i>SMI/SED status and co-occurring mental health and substance use disorder status</i>		<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Number of states reporting		45	47	49	49	48	50
Total		5,165,254	5,656,750	6,171,990	6,418,247	6,548,372	6,945,521
SMI status (individuals aged 18 years and older)¹							
SMI		2,246,718	2,347,345	2,685,739	2,965,701	3,092,582	3,106,523
Not SMI		861,104	790,910	972,736	959,178	876,370	1,052,783
Total		3,107,822	3,138,255	3,658,475	3,924,879	3,968,952	4,159,306
SED status (individuals aged 17 years and younger)¹							
SED		928,981	970,333	1,046,722	1,188,099	1,290,470	1,155,930
Not SED		346,559	406,446	396,327	413,625	359,090	445,567
Total		1,275,540	1,376,779	1,443,049	1,601,724	1,649,560	1,601,497
Co-occurring mental health and substance use disorders (individuals aged 12 years and older)²							
Yes		838,701	858,569	1,150,251	1,247,319	1,513,156	2,071,949
No		3,047,552	3,265,186	3,461,762	3,625,123	3,439,066	3,187,722
Total³		3,886,253	4,123,755	4,612,013	4,872,442	4,952,222	5,259,671

SED = serious emotional disturbance; SMI = serious mental illness.

¹ The federal definitions for SED and SMI status can be found in Appendix D.

² Co-occurring mental health and substance use disorders are only presented for individuals aged 12 years and older.

³ Total number of individuals aged 12 years and older for whom co-occurring mental health and substance use disorders were reported.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 1.9b. Individuals, by SMI/SED status and co-occurring mental health and substance use disorder status: Column percent distribution, 2015–2020

<i>SMI/SED status and co-occurring mental health and substance use disorder status</i>							
		2015	2016	2017	2018	2019	2020
Number of states reporting		45	47	49	49	48	50
	Total	100.0	100.0	100.0	100.0	100.0	100.0
SMI status (individuals aged 18 years and older)¹							
SMI		72.3	74.8	73.4	75.6	77.9	74.7
Not SMI		27.7	25.2	26.6	24.4	22.1	25.3
	Total	100.0	100.0	100.0	100.0	100.0	100.0
SED status (individuals aged 17 years and younger)¹							
SED		72.8	70.5	72.5	74.2	78.2	72.2
Not SED		27.2	29.5	27.5	25.8	21.8	27.8
	Total	100.0	100.0	100.0	100.0	100.0	100.0
Co-occurring mental health and substance use disorders (individuals aged 12 years and older)²							
Yes		21.6	20.8	24.9	25.6	30.6	39.4
No		78.4	79.2	75.1	74.4	69.4	60.6
	Total³	100.0	100.0	100.0	100.0	100.0	100.0

SED = serious emotional disturbance; SMI = serious mental illness.

¹ The federal definitions for SED and SMI status can be found in Appendix D.

² Co-occurring mental health and substance use disorders are only presented for individuals aged 12 years and older.

³ Total percentage of individuals aged 12 years and older for whom co-occurring mental health and substance use disorders were reported.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 1.10a. Individuals aged 18 years and older, by employment status, 'detailed not in labor force,' education, and veteran status: Number, 2015–2020

<i>Employment status, 'detailed not in labor force,' education, and veteran status</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Number of states reporting	45	47	49	49	48	50
Total individuals aged 18 years and older	3,680,337	4,032,003	4,490,694	4,590,635	4,651,676	5,025,981
Employment status						
Employed	411,832	409,574	471,492	520,475	539,659	604,808
Full time	206,665	211,218	239,098	264,824	268,872	317,780
Part time	139,087	133,802	162,213	174,400	170,547	196,696
Unemployed	569,924	501,286	659,044	652,624	659,715	750,437
Not in labor force (see detail below)	1,002,269	953,235	1,044,335	1,077,094	1,018,793	1,037,008
Total	1,984,025	1,864,095	2,174,871	2,250,193	2,218,167	2,392,253
Detailed not in labor force						
Total individuals aged 18 years and older not in labor force	1,002,269	953,235	1,044,335	1,077,094	1,018,793	1,037,008
Disabled	441,233	440,335	495,590	527,718	459,635	445,941
Retired	38,501	39,203	45,794	47,648	47,320	48,150
Student	80,564	75,722	83,082	83,901	86,604	86,575
Homemaker	48,239	47,939	49,703	43,196	40,734	38,206
Sheltered/non-competitive employment	11,761	9,439	13,032	13,100	11,758	9,823
Other	381,971	340,597	357,134	361,531	372,742	408,313
Highest school grade completed						
0 to 8	218,709	221,065	232,542	223,527	233,933	236,111
9 to 11	330,094	329,490	386,717	399,119	395,551	429,713
12 (or GED)	794,920	826,314	946,667	1,058,086	1,052,288	1,142,344
More than 12	341,656	360,244	477,392	530,668	533,462	594,737
Total	1,685,379	1,737,113	2,043,318	2,211,400	2,215,234	2,402,905
Veteran status						
Veteran	<i>n/c</i>	<i>n/c</i>	40,983	44,310	78,675	100,847
Not a veteran	<i>n/c</i>	<i>n/c</i>	1,242,710	1,810,457	1,874,545	1,957,844
Total	<i>n/c</i>	<i>n/c</i>	1,283,693	1,854,767	1,953,220	2,058,691

n/c Not collected.

GED = general equivalency diploma.

NOTES: Total number of client characteristics may not equal the total number of clients due to missing values. *Employed* includes "full-time/part-time not differentiated," which is not presented separately.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 1.10b. Individuals aged 18 years and older, by employment status, ‘detailed not in labor force,’ education, and veteran status: Column percent distribution, 2015–2020

<i>Employment status, ‘detailed not in labor force,’ education, and veteran status</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Number of states reporting	45	47	49	49	48	50
Total individuals aged 18 years and older	100.0	100.0	100.0	100.0	100.0	100.0
Employment status						
Employed	20.8	22.0	21.7	23.1	24.3	25.3
Full time	10.4	11.3	11.0	11.8	12.1	13.3
Part time	7.0	7.2	7.5	7.8	7.7	8.2
Unemployed	28.7	26.9	30.3	29.0	29.7	31.4
Not in labor force (see detail below)	50.5	51.1	48.0	47.9	45.9	43.3
Total	100.0	100.0	100.0	100.0	100.0	100.0
Detailed not in labor force						
Total individuals aged 18 years and older not in labor force	100.0	100.0	100.0	100.0	100.0	100.0
Disabled	44.0	46.2	47.5	49.0	45.1	43.0
Retired	3.8	4.1	4.4	4.4	4.6	4.6
Student	8.0	7.9	8.0	7.8	8.5	8.3
Homemaker	4.8	5.0	4.8	4.0	4.0	3.7
Sheltered/non-competitive employment	1.2	†	1.2	1.2	1.2	†
Other	38.1	35.7	34.2	33.6	36.6	39.4
Highest school grade completed						
0 to 8	13.0	12.7	11.4	10.1	10.6	9.8
9 to 11	19.6	19.0	18.9	18.0	17.9	17.9
12 (or GED)	47.2	47.6	46.3	47.8	47.5	47.5
More than 12	20.3	20.7	23.4	24.0	24.1	24.8
Total	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status						
Veteran	<i>n/c</i>	<i>n/c</i>	3.2	2.4	4.0	4.9
Not a veteran	<i>n/c</i>	<i>n/c</i>	96.8	97.6	96.0	95.1
Total	<i>n/c</i>	<i>n/c</i>	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals; *n/c* Not collected.

GED = general equivalency diploma.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding. *Employed* includes “full-time/part-time not differentiated,” which is not presented separately.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 1.11a. Individuals, by timing of admission and mental health service setting: Number, 2015–2020

<i>Timing of admission and mental health service setting</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Number of states reporting	45	47	49	49	48	50
Timing of admission						
New individuals	2,492,447	3,033,794	3,213,221	3,247,301	3,170,931	3,231,365
Continuing individuals	2,672,807	2,622,956	2,958,769	3,170,946	3,377,441	3,714,156
Total	5,165,254	5,656,750	6,171,990	6,418,247	6,548,372	6,945,521
Mental health service setting^{1,2}						
State psychiatric hospital	121,456	126,645	124,240	122,461	125,261	122,227
Community-based program	5,027,153	5,500,239	5,994,811	6,258,772	6,379,100	6,764,338
Residential treatment center	43,296	56,502	67,567	68,517	75,501	75,583
Other psychiatric inpatient	177,533	236,651	264,310	275,311	264,961	264,304
Institutions under the justice system	70,512	69,043	72,495	71,582	72,583	64,723
Total	5,165,254	5,656,749	6,171,990	6,418,247	6,548,372	6,945,521

¹ Numbers may not sum to total, because individuals could receive services in more than one service setting during the reporting period.

² Mental health services setting category definitions can be found in Appendix C.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 1.11b. Individuals, by timing of admission and mental health service setting: Column percent, distribution 2015–2020

<i>Timing of admission and mental health service setting</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Number of states reporting	45	47	49	49	48	50
Timing of admission						
New individuals	48.3	53.6	52.1	50.6	48.4	46.5
Continuing individuals	51.7	46.4	47.9	49.4	51.6	53.5
Total	100.0	100.0	100.0	100.0	100.0	100.0
Mental health service setting^{1,2}						
State psychiatric hospital	2.4	2.2	2.0	1.9	1.9	1.8
Community-based program	97.3	97.2	97.1	97.5	97.4	97.4
Residential treatment center	†	†	1.1	1.1	1.2	1.1
Other psychiatric inpatient	3.4	4.2	4.3	4.3	4.0	3.8
Institutions under the justice system	1.4	1.2	1.2	1.1	1.1	†
Total	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

¹ Percentages may not sum to 100 percent, because individuals could receive services in more than one service setting during the reporting period.

² Mental health services setting category definitions can be found in Appendix C.

NOTE: Percentages for timing of admission may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

**Table 1.12a. Individuals, by number of mental health diagnosis and any mention of mental health diagnosis:
Number, 2015–2020**

<i>Number of diagnoses and any mention of mental health diagnosis</i>		<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Number of states reporting		45	47	49	49	48	50
	Total	5,165,254	5,656,750	6,171,990	6,418,247	6,548,372	6,945,521
Number of mental health diagnoses							
None		650,131	1,014,081	911,364	752,807	769,603	1,016,968
1		2,479,153	2,963,234	3,537,386	3,635,677	3,718,681	3,819,824
2		1,511,232	1,262,572	1,371,620	1,579,857	1,585,318	1,605,846
3		524,738	416,863	351,620	449,906	474,770	502,883
Any mention of mental health diagnosis^{1,2}							
Trauma- and stressor-related disorders		853,482	878,639	1,035,177	1,161,339	1,253,914	1,295,221
Anxiety disorders		755,391	837,961	1,052,774	1,239,939	1,339,890	1,404,997
Attention deficit/hyperactivity disorder		510,737	560,910	621,999	696,888	672,592	654,495
Conduct disorder		147,020	139,246	135,740	133,315	128,455	117,672
Mood disorders		2,219,816	2,132,801	2,278,835	2,439,327	2,479,529	2,504,101
Bipolar disorders		855,430	747,805	724,193	734,667	704,177	713,775
Depressive disorders		1,419,968	1,429,859	1,597,103	1,756,277	1,824,785	1,834,097
Oppositional defiant disorder		165,302	168,642	191,914	206,728	201,124	181,674
Pervasive developmental disorders		69,481	94,969	98,479	109,585	96,230	110,316
Schizophrenia and other psychotic disorders		640,161	622,833	728,180	770,554	742,888	765,575
Personality disorders		169,840	147,069	165,439	174,277	163,177	161,378
Alcohol or substance-related disorders		269,224	234,950	244,899	273,451	296,773	384,246
Delirium/dementia		33,432	26,246	26,082	29,837	29,147	31,576
Other mental disorders ³		1,186,202	849,710	713,449	858,184	860,407	885,016
No/deferred diagnosis		650,131	1,014,081	911,364	752,807	769,603	1,016,968

¹ Numbers may not sum to the total, because up to three diagnoses may be reported for an individual.

² Up to three diagnoses can be reported for each individual; “any mention” includes all diagnoses reported. For this reason, the sum of any mention of bipolar and depressive disorders may be larger than any mention of mood disorders. Mental health diagnostic category definitions can be found in Appendix E.

³ “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

**Table 1.12b. Individuals, by number of mental health diagnosis and any mention of mental health diagnosis:
Column percent distribution, 2015–2020**

<i>Number of diagnoses and any mention of mental health diagnosis</i>		<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Number of states reporting		45	47	49	49	48	50
	Total	100.0	100.0	100.0	100.0	100.0	100.0
Number of mental health diagnoses							
None		12.6	17.9	14.8	11.7	11.8	14.6
1		48.0	52.4	57.3	56.6	56.8	55.0
2		29.3	22.3	22.2	24.6	24.2	23.1
3		10.2	7.4	5.7	7.0	7.3	7.2
Any mention of mental health diagnosis^{1,2}							
Trauma- and stressor-related disorders		16.5	15.5	16.8	18.1	19.1	18.6
Anxiety disorders		14.6	14.8	17.1	19.3	20.5	20.2
Attention deficit/hyperactivity disorder		9.9	9.9	10.1	10.9	10.3	9.4
Conduct disorder		2.8	2.5	2.2	2.1	2.0	1.7
Mood disorders		43.0	37.7	36.9	38.0	37.9	36.1
Bipolar disorders		16.6	13.2	11.7	11.4	10.8	10.3
Depressive disorders		27.5	25.3	25.9	27.4	27.9	26.4
Oppositional defiant disorder		3.2	3.0	3.1	3.2	3.1	2.6
Pervasive developmental disorders		1.3	1.7	1.6	1.7	1.5	1.6
disorders		12.4	11.0	11.8	12.0	11.3	11.0
Personality disorders		3.3	2.6	2.7	2.7	2.5	2.3
Alcohol or substance-related disorders		5.2	4.2	4.0	4.3	4.5	5.5
Delirium/dementia		†	†	†	†	†	†
Other mental disorders ³		23.0	15.0	11.6	13.4	13.1	12.7
No/deferred diagnosis		12.6	17.9	14.8	11.7	11.8	14.6

† Less than 1 percent of individuals.

¹ Percentages may not sum to 100 percent, because up to three diagnoses may be reported for an individual.

² Up to three diagnoses can be reported for each individual; “any mention” includes all diagnoses reported. For this reason, the sum of any mention of bipolar and depressive disorders may be larger than any mention of mood disorders. Mental health diagnostic category definitions can be found in Appendix E.

³ “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 1.13a. Individuals, by Census region and division: Number, 2015–2020

<i>Census region and Census division</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Number of states reporting	45	47	49	49	48	50
Total	5,141,885	5,638,807	6,146,072	6,414,937	6,543,899	6,941,801
Region¹						
Northeast	560,347	1,115,327	1,274,412	1,226,718	1,256,271	1,247,225
Midwest	1,220,159	1,223,133	1,484,436	1,598,102	1,771,702	1,836,424
South	1,882,068	1,831,834	1,854,663	1,928,812	1,926,414	1,915,757
West	1,479,311	1,468,513	1,532,561	1,661,305	1,589,512	1,942,395
Division¹						
New England	180,721	191,349	240,464	181,976	184,370	171,766
Middle Atlantic	379,626	923,978	1,033,948	1,044,742	1,071,901	1,075,459
East North Central	752,146	721,577	982,894	989,257	1,082,519	993,750
West North Central	468,013	501,556	501,542	608,845	689,183	842,674
South Atlantic	878,731	800,461	812,765	849,366	821,994	823,628
East South Central	467,125	461,375	441,287	463,160	469,034	454,263
West South Central	536,212	569,998	600,611	616,286	635,386	637,866
Mountain	565,442	592,573	650,495	696,577	521,422	872,748
Pacific	913,869	875,940	882,066	964,728	1,068,090	1,069,647

¹ The Commonwealth of the Northern Mariana Islands, the Republic of Palau, and the Commonwealth of Puerto Rico are not included in any Census region or division.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 1.13b. Individuals, by Census region and division: Column percent distribution, 2015–2020

<i>Census region and Census division</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Number of states reporting	45	47	49	49	48	50
Total	100.0	100.0	100.0	100.0	100.0	100.0
Region¹						
Northeast	10.9	19.8	20.7	19.1	19.2	18.0
Midwest	23.7	21.7	24.2	24.9	27.1	26.5
South	36.6	32.5	30.2	30.1	29.4	27.6
West	28.8	26.0	24.9	25.9	24.3	28.0
Division¹						
New England	3.5	3.4	3.9	2.8	2.8	2.5
Middle Atlantic	7.4	16.4	16.8	16.3	16.4	15.5
East North Central	14.6	12.8	16.0	15.4	16.5	14.3
West North Central	9.1	8.9	8.2	9.5	10.5	12.1
South Atlantic	17.1	14.2	13.2	13.2	12.6	11.9
East South Central	9.1	8.2	7.2	7.2	7.2	6.5
West South Central	10.4	10.1	9.8	9.6	9.7	9.2
Mountain	11.0	10.5	10.6	10.9	8.0	12.6
Pacific	17.8	15.5	14.4	15.0	16.3	15.4

¹ The Commonwealth of the Northern Mariana Islands, the Republic of Palau, and the Commonwealth of Puerto Rico are not included in any Census region or division.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 1.14a. Individuals with diagnosis of trauma- and stressor-related disorders, by gender, age, race, and ethnicity: Number, 2015–2020

<i>Gender, age, race, and ethnicity</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Number of states reporting	45	47	49	49	48	50
Total individuals with diagnosis of trauma- and stressor-related disorders	853,482	878,639	1,035,177	1,161,339	1,253,914	1,295,221
Gender						
Male	340,514	354,225	422,131	478,010	517,556	530,123
Female	511,057	521,667	609,526	680,404	732,105	761,230
Total	851,571	875,892	1,031,657	1,158,414	1,249,661	1,291,353
Age¹						
0 to 11 years	190,427	207,478	223,391	261,172	283,663	269,732
12 to 14 years	79,120	83,028	90,758	106,229	120,397	120,196
15 to 17 years	73,281	78,604	87,435	99,342	108,084	106,743
18 to 20 years	35,735	36,936	43,098	50,033	55,778	58,020
21 to 24 years	44,342	43,659	52,394	56,432	61,912	66,440
25 to 29 years	64,741	66,624	82,813	90,997	98,393	103,368
30 to 34 years	66,475	67,549	82,437	90,957	98,498	105,831
35 to 39 years	59,678	61,603	76,761	85,948	91,831	98,512
40 to 44 years	54,599	53,168	62,865	69,386	73,931	80,815
45 to 49 years	53,561	53,089	63,601	68,879	70,216	73,926
50 to 54 years	52,732	51,405	61,299	65,396	66,182	70,195
55 to 59 years	37,702	37,934	48,323	54,117	56,889	62,154
60 to 64 years	21,199	20,897	28,083	32,401	34,854	39,736
65 years and older	19,040	15,066	30,368	29,309	32,176	38,935
Total	852,632	877,040	1,033,626	1,160,598	1,252,804	1,294,603

Continued. See notes at end of table.

Table 1.14a. Individuals with diagnosis of trauma- and stressor-related disorders, by gender, age, race, and ethnicity: Number, 2015–2020 (continued)

<i>Gender, age, race, and ethnicity</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	16,560	17,765	19,395	22,320	26,984	33,147
Asian ²	11,401	12,914	13,604	15,435	14,905	14,546
Black or African American	135,336	133,344	169,589	189,804	209,762	209,379
Native Hawaiian or Other Pacific Islander	2,470	2,646	2,123	3,042	3,268	3,147
White	529,438	554,685	643,157	710,199	760,857	783,816
Other	78,490	81,937	98,833	117,517	127,406	137,128
Total	773,695	803,291	946,701	1,058,317	1,143,182	1,181,163
Ethnicity						
Not of Hispanic or Latino origin	576,764	606,353	786,405	861,308	927,697	944,583
Hispanic or Latino origin	128,085	140,256	161,258	173,279	191,424	235,861
Total	704,849	746,609	947,663	1,034,587	1,119,121	1,180,444
Specific Hispanic or Latino origin						
Mexican	12,608	9,107	15,298	15,843	17,228	16,255
Puerto Rican	5,377	4,312	5,143	4,864	4,672	4,857
Cuban	874	835	1,075	1,057	1,157	1,217
Other specific	9,931	9,155	14,000	14,462	13,586	14,034
Total	28,790	23,409	35,516	36,226	36,643	36,363

¹ Age is based on the individual's age at the midpoint of the reporting period.

² For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTES: Total number of client characteristics may not equal the total number of clients due to missing values. This table refers to all mental health diagnoses, which is up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 1.14b. Individuals with diagnosis of trauma- and stressor-related disorders, by gender, age, race, and ethnicity: Column percent distribution, 2015–2020

<i>Gender, age, race, and ethnicity</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Number of states reporting	45	47	49	49	48	50
Total individuals with diagnosis of trauma- and stressor-related disorders	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	40.0	40.4	40.9	41.3	41.4	41.1
Female	60.0	59.6	59.1	58.7	58.6	58.9
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age¹						
0 to 11 years	22.3	23.7	21.6	22.5	22.6	20.8
12 to 14 years	9.3	9.5	8.8	9.2	9.6	9.3
15 to 17 years	8.6	9.0	8.5	8.6	8.6	8.2
18 to 20 years	4.2	4.2	4.2	4.3	4.5	4.5
21 to 24 years	5.2	5.0	5.1	4.9	4.9	5.1
25 to 29 years	7.6	7.6	8.0	7.8	7.9	8.0
30 to 34 years	7.8	7.7	8.0	7.8	7.9	8.2
35 to 39 years	7.0	7.0	7.4	7.4	7.3	7.6
40 to 44 years	6.4	6.1	6.1	6.0	5.9	6.2
45 to 49 years	6.3	6.1	6.2	5.9	5.6	5.7
50 to 54 years	6.2	5.9	5.9	5.6	5.3	5.4
55 to 59 years	4.4	4.3	4.7	4.7	4.5	4.8
60 to 64 years	2.5	2.4	2.7	2.8	2.8	3.1
65 years and older	2.2	1.7	2.9	2.5	2.6	3.0
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 1.14b. Individuals with diagnosis of trauma- and stressor-related disorders, by gender, age, race, and ethnicity: Column percent distribution, 2015–2020 (continued)

<i>Gender, age, race, and ethnicity</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	2.1	2.2	2.0	2.1	2.4	2.8
Asian ²	1.5	1.6	1.4	1.5	1.3	1.2
Black or African American	17.5	16.6	17.9	17.9	18.3	17.7
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	68.4	69.1	67.9	67.1	66.6	66.4
Other	10.1	10.2	10.4	11.1	11.1	11.6
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	81.8	81.2	83.0	83.3	82.9	80.0
Hispanic or Latino origin	18.2	18.8	17.0	16.7	17.1	20.0
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	43.8	38.9	43.1	43.7	47.0	44.7
Puerto Rican	18.7	18.4	14.5	13.4	12.8	13.4
Cuban	3.0	3.6	3.0	2.9	3.2	3.3
Other specific	34.5	39.1	39.4	39.9	37.1	38.6
Total	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

¹ Age is based on the individual's age at the midpoint of the reporting period.

² For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding. This table refers to all mental health diagnoses, which is up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 1.15a. Individuals with diagnosis of anxiety disorders, by gender, age, race, and ethnicity: Number, 2015–2020

<i>Gender, age, race, and ethnicity</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Number of states reporting	45	46	49	49	48	50
Total individuals with diagnosis of anxiety disorders	755,391	837,961	1,052,774	1,239,939	1,339,890	1,404,997
Gender						
Male	294,414	323,185	404,124	472,491	506,271	527,839
Female	459,936	505,409	646,155	765,248	830,360	874,137
Total	754,350	828,594	1,050,279	1,237,739	1,336,631	1,401,976
Age¹						
0 to 11 years	77,092	86,173	99,363	116,021	127,063	122,054
12 to 14 years	44,682	51,578	62,815	76,573	86,891	90,106
15 to 17 years	52,495	63,647	79,306	96,414	107,727	110,151
18 to 20 years	34,290	40,228	51,298	64,555	74,469	78,536
21 to 24 years	47,904	52,746	66,162	78,224	86,351	93,541
25 to 29 years	73,601	82,823	105,070	124,710	135,555	141,726
30 to 34 years	75,563	82,664	103,905	121,468	131,352	139,315
35 to 39 years	65,137	73,035	94,313	111,425	119,227	126,570
40 to 44 years	59,919	63,370	77,676	89,809	95,258	103,009
45 to 49 years	58,809	63,086	78,982	90,445	91,914	94,986
50 to 54 years	61,085	64,875	80,170	89,596	89,994	92,135
55 to 59 years	48,065	52,938	68,008	80,097	83,438	88,716
60 to 64 years	29,352	32,606	43,053	52,650	56,402	62,442
65 years and older	26,738	26,926	41,394	47,345	53,284	61,242
Total	754,732	836,695	1,051,515	1,239,332	1,338,925	1,404,529

Continued. See notes at end of table.

Table 1.15a. Individuals with diagnosis of anxiety disorders, by gender, age, race, and ethnicity: Number, 2015–2020 (continued)

<i>Gender, age, race, and ethnicity</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	11,169	11,909	15,619	19,466	23,743	29,027
Asian ²	7,523	9,269	12,343	15,019	15,232	15,418
Black or African American	78,421	85,392	112,096	131,804	147,607	150,861
Native Hawaiian or Other Pacific Islander	2,679	3,433	1,699	2,700	2,925	2,963
White	542,741	598,787	749,417	871,238	925,271	966,447
Other	56,409	64,472	85,609	107,994	118,047	128,138
Total	698,942	773,262	976,783	1,148,221	1,232,825	1,292,854
Ethnicity						
Not of Hispanic or Latino origin	552,427	610,738	831,861	932,081	994,041	1,053,876
Hispanic or Latino origin	95,551	112,339	140,447	164,132	177,425	240,877
Total	647,978	723,077	972,308	1,096,213	1,171,466	1,294,753
Specific Hispanic or Latino origin						
Mexican	9,167	6,097	9,992	10,466	11,862	12,064
Puerto Rican	3,525	3,285	4,067	4,461	4,224	4,147
Cuban	935	894	1,006	1,025	1,216	1,446
Other specific	7,227	7,221	9,594	10,608	10,680	11,901
Total	20,854	17,497	24,659	26,560	27,982	29,558

¹ Age is based on the individual's age at the midpoint of the reporting period.

² For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTES: Total number of client characteristics may not equal the total number of clients due to missing values. This table refers to all mental health diagnoses, which is up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 1.15b. Individuals with diagnosis of anxiety disorders, by gender, age, race, and ethnicity: Column percent distribution, 2015–2020

<i>Gender, age, race, and ethnicity</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Number of states reporting	45	46	49	49	48	50
Total individuals with diagnosis of anxiety disorders	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	39.0	39.0	38.5	38.2	37.9	37.6
Female	61.0	61.0	61.5	61.8	62.1	62.4
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age¹						
0 to 11 years	10.2	10.3	9.4	9.4	9.5	8.7
12 to 14 years	5.9	6.2	6.0	6.2	6.5	6.4
15 to 17 years	7.0	7.6	7.5	7.8	8.0	7.8
18 to 20 years	4.5	4.8	4.9	5.2	5.6	5.6
21 to 24 years	6.3	6.3	6.3	6.3	6.4	6.7
25 to 29 years	9.8	9.9	10.0	10.1	10.1	10.1
30 to 34 years	10.0	9.9	9.9	9.8	9.8	9.9
35 to 39 years	8.6	8.7	9.0	9.0	8.9	9.0
40 to 44 years	7.9	7.6	7.4	7.2	7.1	7.3
45 to 49 years	7.8	7.5	7.5	7.3	6.9	6.8
50 to 54 years	8.1	7.8	7.6	7.2	6.7	6.6
55 to 59 years	6.4	6.3	6.5	6.5	6.2	6.3
60 to 64 years	3.9	3.9	4.1	4.2	4.2	4.4
65 years and older	3.5	3.2	3.9	3.8	4.0	4.4
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 1.15b. Individuals with diagnosis of anxiety disorders, by gender, age, race, and ethnicity: Column percent distribution, 2015–2020
(continued)

<i>Gender, age, race, and ethnicity</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	1.6	1.5	1.6	1.7	1.9	2.2
Asian ²	1.1	1.2	1.3	1.3	1.2	1.2
Black or African American	11.2	11.0	11.5	11.5	12.0	11.7
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	77.7	77.4	76.7	75.9	75.1	74.8
Other	8.1	8.3	8.8	9.4	9.6	9.9
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	85.3	84.5	85.6	85.0	84.9	81.4
Hispanic or Latino origin	14.7	15.5	14.4	15.0	15.1	18.6
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	44.0	34.8	40.5	39.4	42.4	40.8
Puerto Rican	16.9	18.8	16.5	16.8	15.1	14.0
Cuban	4.5	5.1	4.1	3.9	4.3	4.9
Other specific	34.7	41.3	38.9	39.9	38.2	40.3
Total	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

¹ Age is based on the individual's age at the midpoint of the reporting period.

² For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding. This table refers to all mental health diagnoses, which is up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 1.16a. Individuals with diagnosis of attention deficit/hyperactivity disorder, by gender, age, race, and ethnicity: Number, 2015–2020

<i>Gender, age, race, and ethnicity</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Number of states reporting	45	46	49	49	48	50
Total individuals with diagnosis of attention deficit/hyperactivity disorder	510,737	560,910	621,999	696,888	672,592	654,495
Gender						
Male	348,956	380,266	421,616	472,402	452,699	439,432
Female	160,551	179,135	198,275	222,395	217,312	212,894
Total	509,507	559,401	619,891	694,797	670,011	652,326
Age¹						
0 to 11 years	226,682	253,565	271,139	301,556	288,030	270,758
12 to 14 years	100,998	109,987	118,267	132,315	127,879	125,510
15 to 17 years	72,914	79,410	87,127	95,071	88,864	85,361
18 to 20 years	27,181	28,694	33,155	37,061	36,061	35,499
21 to 24 years	18,057	17,572	22,336	24,510	23,862	24,315
25 to 29 years	18,367	19,087	24,848	29,330	29,177	29,583
30 to 34 years	14,037	15,002	19,017	22,812	23,583	24,866
35 to 39 years	9,470	10,944	13,730	16,948	17,750	18,674
40 to 44 years	6,909	7,811	9,149	10,902	11,366	12,189
45 to 49 years	5,517	6,434	7,621	9,141	8,721	9,239
50 to 54 years	4,756	5,327	6,196	6,968	6,830	6,994
55 to 59 years	3,113	3,656	4,563	5,447	5,484	5,806
60 to 64 years	1,736	2,070	2,508	3,140	3,159	3,531
65 years and older	866	1,027	2,014	1,617	1,674	2,045
Total	510,603	560,586	621,670	696,818	672,440	654,370

Continued. See notes at end of table.

Table 1.16a. Individuals with diagnosis of attention deficit/hyperactivity disorder, by gender, age, race, and ethnicity: Number, 2015–2020 (continued)

<i>Gender, age, race, and ethnicity</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	6,535	6,150	5,979	7,358	7,256	9,022
Asian ²	2,543	3,048	3,432	4,940	4,150	4,199
Black or African American	116,109	121,503	137,153	144,067	143,308	129,730
Native Hawaiian or Other Pacific Islander	1,338	1,470	731	1,144	1,071	1,043
White	303,410	338,343	373,242	417,880	395,412	390,624
Other	45,406	56,540	66,181	73,361	70,374	70,284
Total	475,341	527,054	586,718	648,750	621,571	604,902
Ethnicity						
Not of Hispanic or Latino origin	364,275	404,135	485,668	516,323	499,565	490,524
Hispanic or Latino origin	70,486	85,556	98,881	105,607	95,399	114,465
Total	434,761	489,691	584,549	621,930	594,964	604,989
Specific Hispanic or Latino origin						
Mexican	6,500	4,203	5,371	5,966	5,627	5,723
Puerto Rican	3,154	2,384	3,303	3,078	2,238	2,213
Cuban	502	440	473	524	487	539
Other specific	5,413	5,091	5,991	6,222	5,199	5,890
Total	15,569	12,118	15,138	15,790	13,551	14,365

¹ Age is based on the individual's age at the midpoint of the reporting period.

² For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTES: Total number of client characteristics may not equal the total number of clients due to missing values. This table refers to all mental health diagnoses, which is up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 1.16b. Individuals with diagnosis of attention deficit/hyperactivity disorder, by gender, age, race, and ethnicity: Column percent distribution, 2015–2020

<i>Gender, age, race, and ethnicity</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Number of states reporting	45	46	49	49	48	50
Total individuals with diagnosis of attention deficit/hyperactivity disorder	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	68.5	68.0	68.0	68.0	67.6	67.4
Female	31.5	32.0	32.0	32.0	32.4	32.6
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age¹						
0 to 11 years	44.4	45.2	43.6	43.3	42.8	41.4
12 to 14 years	19.8	19.6	19.0	19.0	19.0	19.2
15 to 17 years	14.3	14.2	14.0	13.6	13.2	13.0
18 to 20 years	5.3	5.1	5.3	5.3	5.4	5.4
21 to 24 years	3.5	3.1	3.6	3.5	3.5	3.7
25 to 29 years	3.6	3.4	4.0	4.2	4.3	4.5
30 to 34 years	2.7	2.7	3.1	3.3	3.5	3.8
35 to 39 years	1.9	2.0	2.2	2.4	2.6	2.9
40 to 44 years	1.4	1.4	1.5	1.6	1.7	1.9
45 to 49 years	1.1	1.1	1.2	1.3	1.3	1.4
50 to 54 years	†	†	†	†	1.0	1.1
55 to 59 years	†	†	†	†	†	†
60 to 64 years	†	†	†	†	†	†
65 years and older	†	†	†	†	†	†
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 1.16b. Individuals with diagnosis of attention deficit/hyperactivity disorder, by gender, age, race, and ethnicity: Column percent distribution, 2015–2020 (continued)

<i>Gender, age, race, and ethnicity</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	1.4	1.2	1.0	1.1	1.2	1.5
Asian ²	†	†	†	†	†	†
Black or African American	24.4	23.1	23.4	22.2	23.1	21.4
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	63.8	64.2	63.6	64.4	63.6	64.6
Other	9.6	10.7	11.3	11.3	11.3	11.6
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	83.8	82.5	83.1	83.0	84.0	81.1
Hispanic or Latino origin	16.2	17.5	16.9	17.0	16.0	18.9
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	41.7	34.7	35.5	37.8	41.5	39.8
Puerto Rican	20.3	19.7	21.8	19.5	16.5	15.4
Cuban	3.2	3.6	3.1	3.3	3.6	3.8
Other specific	34.8	42.0	39.6	39.4	38.4	41.0
Total	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

¹ Age is based on the individual's age at the midpoint of the reporting period.

² For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding. This table refers to all mental health diagnoses, which is up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 1.17a. Individuals with diagnosis of bipolar disorders, by gender, age, race, and ethnicity: Number, 2015–2020

<i>Gender, age, race, and ethnicity</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Number of states reporting	45	46	48	49	47	50
Total individuals with diagnosis of bipolar disorders	855,430	747,805	724,193	734,667	704,177	713,775
Gender						
Male	366,514	311,002	289,998	292,146	278,384	281,146
Female	485,658	433,484	431,946	441,048	424,033	431,079
Total	852,172	744,486	721,944	733,194	702,417	712,225
Age¹						
0 to 11 years	36,904	22,162	8,862	6,101	4,662	3,642
12 to 14 years	38,361	24,297	12,194	10,116	8,372	7,535
15 to 17 years	53,735	37,888	22,957	20,700	17,750	16,400
18 to 20 years	41,407	33,771	27,831	28,267	27,519	26,409
21 to 24 years	61,446	51,207	47,435	47,561	46,346	46,908
25 to 29 years	88,534	79,973	80,288	82,064	80,701	80,116
30 to 34 years	90,974	82,596	82,969	85,366	84,084	85,559
35 to 39 years	83,033	78,794	82,566	85,611	82,508	82,599
40 to 44 years	79,636	71,562	72,384	73,364	71,009	73,525
45 to 49 years	80,165	74,747	76,869	77,185	71,704	71,354
50 to 54 years	80,854	74,684	77,071	76,534	70,660	70,608
55 to 59 years	58,038	56,389	62,348	66,254	63,688	66,313
60 to 64 years	33,560	32,988	37,268	41,131	40,235	43,610
65 years and older	26,681	24,521	31,888	34,060	34,595	39,012
Total	853,328	745,579	722,930	734,314	703,833	713,590

Continued. See notes at end of table.

Table 1.17a. Individuals with diagnosis of bipolar disorders, by gender, age, race, and ethnicity: Number, 2015–2020 (continued)

<i>Gender, age, race, and ethnicity</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	10,031	7,924	7,511	7,846	7,814	8,828
Asian ²	6,522	6,173	6,402	7,371	5,990	5,937
Black or African American	142,733	120,330	118,355	121,064	121,353	116,699
Native Hawaiian or Other Pacific Islander	2,232	2,037	1,058	1,245	1,170	1,191
White	582,011	517,766	508,784	513,600	484,621	492,718
Other	51,817	49,135	42,483	48,859	48,518	50,810
Total	795,346	703,365	684,593	699,985	669,466	676,183
Ethnicity						
Not of Hispanic or Latino origin	642,245	566,104	589,760	599,707	577,208	573,885
Hispanic or Latino origin	82,807	78,609	67,445	69,743	67,737	89,965
Total	725,052	644,713	657,205	669,450	644,945	663,850
Specific Hispanic or Latino origin						
Mexican	5,697	2,969	3,753	3,646	3,596	3,436
Puerto Rican	4,880	3,499	3,724	3,585	3,326	3,359
Cuban	1,367	1,109	1,042	1,087	980	1,069
Other specific	7,124	5,086	5,520	5,867	4,636	5,128
Total	19,068	12,663	14,039	14,185	12,538	12,992

¹ Age is based on the individual's age at the midpoint of the reporting period.

² For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTES: Total number of client characteristics may not equal the total number of clients due to missing values. This table refers to all mental health diagnoses, which is up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 1.17b. Individuals with diagnosis of bipolar disorders, by gender, age, race, and ethnicity: Column percent distribution, 2015–2020

<i>Gender, age, race, and ethnicity</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Number of states reporting	45	46	48	49	47	50
Total individuals with diagnosis of bipolar disorders	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	43.0	41.8	40.2	39.8	39.6	39.5
Female	57.0	58.2	59.8	60.2	60.4	60.5
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age¹						
0 to 11 years	4.3	3.0	1.2	†	†	†
12 to 14 years	4.5	3.3	1.7	1.4	1.2	1.1
15 to 17 years	6.3	5.1	3.2	2.8	2.5	2.3
18 to 20 years	4.9	4.5	3.8	3.8	3.9	3.7
21 to 24 years	7.2	6.9	6.6	6.5	6.6	6.6
25 to 29 years	10.4	10.7	11.1	11.2	11.5	11.2
30 to 34 years	10.7	11.1	11.5	11.6	11.9	12.0
35 to 39 years	9.7	10.6	11.4	11.7	11.7	11.6
40 to 44 years	9.3	9.6	10.0	10.0	10.1	10.3
45 to 49 years	9.4	10.0	10.6	10.5	10.2	10.0
50 to 54 years	9.5	10.0	10.7	10.4	10.0	9.9
55 to 59 years	6.8	7.6	8.6	9.0	9.0	9.3
60 to 64 years	3.9	4.4	5.2	5.6	5.7	6.1
65 years and older	3.1	3.3	4.4	4.6	4.9	5.5
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 1.17b. Individuals with diagnosis of bipolar disorders, by gender, age, race, and ethnicity: Column percent distribution, 2015–2020
(continued)

<i>Gender, age, race, and ethnicity</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	1.3	1.1	1.1	1.1	1.2	1.3
Asian ²	†	†	†	1.1	†	†
Black or African American	17.9	17.1	17.3	17.3	18.1	17.3
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	73.2	73.6	74.3	73.4	72.4	72.9
Other	6.5	7.0	6.2	7.0	7.2	7.5
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	88.6	87.8	89.7	89.6	89.5	86.4
Hispanic or Latino origin	11.4	12.2	10.3	10.4	10.5	13.6
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	29.9	23.4	26.7	25.7	28.7	26.4
Puerto Rican	25.6	27.6	26.5	25.3	26.5	25.9
Cuban	7.2	8.8	7.4	7.7	7.8	8.2
Other specific	37.4	40.2	39.3	41.4	37.0	39.5
Total	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

¹ Age is based on the individual's age at the midpoint of the reporting period.

² For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding. This table refers to all mental health diagnoses, which is up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 1.18a. Individuals with diagnosis of depressive disorders, by gender, age, race, and ethnicity: Number, 2015–2020

<i>Gender, age, race, and ethnicity</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Number of states reporting	45	47	49	49	48	50
Total individuals with diagnosis of depressive disorders	1,419,968	1,429,859	1,597,103	1,756,277	1,824,785	1,834,097
Gender						
Male	520,408	525,321	591,495	654,794	682,420	686,701
Female	895,383	899,495	1,001,706	1,098,402	1,138,081	1,143,469
Total	1,415,791	1,424,816	1,593,201	1,753,196	1,820,501	1,830,170
Age¹						
0 to 11 years	37,922	36,984	38,110	43,636	53,453	47,352
12 to 14 years	74,271	74,380	76,795	90,310	104,387	104,148
15 to 17 years	111,649	117,596	123,786	142,468	153,779	149,771
18 to 20 years	65,131	69,616	77,983	92,445	102,672	103,649
21 to 24 years	83,383	86,328	95,688	107,927	115,346	119,677
25 to 29 years	121,021	126,925	145,249	162,535	171,130	173,133
30 to 34 years	124,944	127,308	142,782	156,064	162,628	166,299
35 to 39 years	116,078	119,569	136,665	149,661	153,362	154,273
40 to 44 years	118,225	114,303	122,530	130,889	132,085	134,925
45 to 49 years	132,427	129,770	140,787	147,236	141,907	136,417
50 to 54 years	151,806	148,255	158,555	163,452	155,752	149,582
55 to 59 years	127,236	128,194	144,511	157,014	155,537	154,816
60 to 64 years	79,968	80,857	95,049	107,619	110,495	115,094
65 years and older	73,083	66,246	96,311	103,923	110,851	124,344
Total	1,417,144	1,426,331	1,594,801	1,755,179	1,823,384	1,833,480

Continued. See notes at end of table.

Table 1.18a. Individuals with diagnosis of depressive disorders, by gender, age, race, and ethnicity: Number, 2015–2020 (continued)

<i>Gender, age, race, and ethnicity</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	20,909	20,100	22,764	25,501	29,092	35,768
Asian ²	21,808	22,336	24,275	27,622	26,807	26,378
Black or African American	240,532	234,512	266,590	289,820	311,301	293,485
Native Hawaiian or Other Pacific Islander	3,956	4,388	2,818	3,745	4,227	4,067
White	901,285	923,561	1,028,753	1,118,952	1,138,617	1,151,672
Other	113,140	123,462	135,703	162,752	176,243	187,781
Total	1,301,630	1,328,359	1,480,903	1,628,392	1,686,287	1,699,151
Ethnicity						
Not of Hispanic or Latino origin	1,000,550	1,011,761	1,227,149	1,320,859	1,363,355	1,367,174
Hispanic or Latino origin	181,034	202,504	224,986	245,853	260,903	320,270
Total	1,181,584	1,214,265	1,452,135	1,566,712	1,624,258	1,687,444
Specific Hispanic or Latino origin						
Mexican	16,697	10,457	15,523	16,048	17,148	16,586
Puerto Rican	8,406	6,972	8,145	8,539	8,154	8,041
Cuban	3,302	3,097	3,050	3,048	3,022	3,483
Other specific	15,403	13,814	17,094	18,004	16,190	17,514
Total	43,808	34,340	43,812	45,639	44,514	45,624

¹ Age is based on the individual's age at the midpoint of the reporting period.

² For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTES: Total number of client characteristics may not equal the total number of clients due to missing values. This table refers to all mental health diagnoses, which is up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022..

Table 1.18b. Individuals with diagnosis of depressive disorders, by gender, age, race, and ethnicity: Column percent distribution, 2015–2020

<i>Gender, age, race, and ethnicity</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Number of states reporting	45	47	49	49	48	50
Total individuals with diagnosis of depressive disorders	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	36.8	36.9	37.1	37.3	37.5	37.5
Female	63.2	63.1	62.9	62.7	62.5	62.5
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age¹						
0 to 11 years	2.7	2.6	2.4	2.5	2.9	2.6
12 to 14 years	5.2	5.2	4.8	5.1	5.7	5.7
15 to 17 years	7.9	8.2	7.8	8.1	8.4	8.2
18 to 20 years	4.6	4.9	4.9	5.3	5.6	5.7
21 to 24 years	5.9	6.1	6.0	6.1	6.3	6.5
25 to 29 years	8.5	8.9	9.1	9.3	9.4	9.4
30 to 34 years	8.8	8.9	9.0	8.9	8.9	9.1
35 to 39 years	8.2	8.4	8.6	8.5	8.4	8.4
40 to 44 years	8.3	8.0	7.7	7.5	7.2	7.4
45 to 49 years	9.3	9.1	8.8	8.4	7.8	7.4
50 to 54 years	10.7	10.4	9.9	9.3	8.5	8.2
55 to 59 years	9.0	9.0	9.1	8.9	8.5	8.4
60 to 64 years	5.6	5.7	6.0	6.1	6.1	6.3
65 years and older	5.2	4.6	6.0	5.9	6.1	6.8
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 1.18b. Individuals with diagnosis of depressive disorders, by gender, age, race, and ethnicity: Column percent distribution, 2015–2020 (continued)

<i>Gender, age, race, and ethnicity</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	1.6	1.5	1.5	1.6	1.7	2.1
Asian ²	1.7	1.7	1.6	1.7	1.6	1.6
Black or African American	18.5	17.7	18.0	17.8	18.5	17.3
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	69.2	69.5	69.5	68.7	67.5	67.8
Other	8.7	9.3	9.2	10.0	10.5	11.1
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	84.7	83.3	84.5	84.3	83.9	81.0
Hispanic or Latino origin	15.3	16.7	15.5	15.7	16.1	19.0
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	38.1	30.5	35.4	35.2	38.5	36.4
Puerto Rican	19.2	20.3	18.6	18.7	18.3	17.6
Cuban	7.5	9.0	7.0	6.7	6.8	7.6
Other specific	35.2	40.2	39.0	39.4	36.4	38.4
Total	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

¹ Age is based on the individual's age at the midpoint of the reporting period.

² For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding. This table refers to all mental health diagnoses, which is up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 1.19a. Individuals with diagnosis of schizophrenia and other psychotic disorders, by gender, age, race, and ethnicity: Number, 2015–2020

<i>Gender, age, race, and ethnicity</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Number of states reporting	45	46	48	49	48	50
Total individuals with diagnosis of schizophrenia and other psychotic disorders	640,161	622,833	728,180	770,554	742,888	765,575
Gender						
Male	391,560	383,477	449,489	477,761	461,361	476,145
Female	244,053	234,530	275,446	291,528	280,251	288,121
Total	635,613	618,007	724,935	769,289	741,612	764,266
Age¹						
0 to 11 years	2,530	2,258	2,708	2,349	2,122	1,797
12 to 14 years	3,064	2,912	3,201	3,238	3,002	2,753
15 to 17 years	6,487	6,146	6,748	6,925	6,592	6,245
18 to 20 years	15,330	14,725	16,251	16,865	16,831	16,670
21 to 24 years	36,800	36,348	40,302	42,317	40,350	40,747
25 to 29 years	56,776	57,460	67,591	73,823	73,032	75,835
30 to 34 years	64,009	63,212	72,443	77,346	75,433	80,370
35 to 39 years	59,885	61,092	72,864	79,856	78,021	82,122
40 to 44 years	59,504	56,797	64,659	69,298	67,260	72,456
45 to 49 years	69,614	65,842	73,930	76,081	70,781	71,172
50 to 54 years	85,053	80,092	87,969	87,902	79,789	78,564
55 to 59 years	76,415	74,810	88,493	92,977	87,388	87,454
60 to 64 years	50,696	50,531	62,836	69,052	68,148	70,419
65 years and older	50,069	46,239	65,244	71,554	73,241	78,604
Total	636,232	618,464	725,239	769,583	741,990	765,208

Continued. See notes at end of table.

Table 1.19a. Individuals with diagnosis of schizophrenia and other psychotic disorders, by gender, age, race, and ethnicity: Number, 2015–2020 (continued)

<i>Gender, age, race, and ethnicity</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	6,571	6,571	7,575	8,354	8,456	10,173
Asian ²	15,281	15,235	16,902	19,196	18,701	17,642
Black or African American	194,496	187,878	221,203	234,767	229,501	241,020
Native Hawaiian or Other Pacific Islander	2,914	3,127	1,892	2,161	2,286	2,145
White	318,606	319,285	372,455	388,398	366,278	372,795
Other	42,929	43,970	56,837	69,709	71,721	72,550
Total	580,797	576,066	676,864	722,585	696,943	716,325
Ethnicity						
Not of Hispanic or Latino origin	450,075	460,728	559,893	597,929	574,499	587,311
Hispanic or Latino origin	63,720	69,266	82,541	92,660	91,501	107,660
Total	513,795	529,994	642,434	690,589	666,000	694,971
Specific Hispanic or Latino origin						
Mexican	3,501	2,915	4,070	3,956	3,967	3,860
Puerto Rican	3,339	2,844	3,749	4,113	4,502	4,565
Cuban	1,422	1,375	1,579	1,557	1,482	1,516
Other specific	5,182	5,059	6,057	6,606	5,041	5,514
Total	13,444	12,193	15,455	16,232	14,992	15,455

¹ Age is based on the individual's age at the midpoint of the reporting period.

² For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTES: Total number of client characteristics may not equal the total number of clients due to missing values. This table refers to all mental health diagnoses, which is up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 1.19b. Individuals with diagnosis of schizophrenia and other psychotic disorders, by gender, age, race, and ethnicity: Column percent distribution, 2015–2020

<i>Gender, age, race, and ethnicity</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Number of states reporting	45	46	48	49	48	50
Total individuals with diagnosis of schizophrenia and other psychotic disorders	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	61.6	62.1	62.0	62.1	62.2	62.3
Female	38.4	37.9	38.0	37.9	37.8	37.7
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age¹						
0 to 11 years	†	†	†	†	†	†
12 to 14 years	†	†	†	†	†	†
15 to 17 years	1.0	†	†	†	†	†
18 to 20 years	2.4	2.4	2.2	2.2	2.3	2.2
21 to 24 years	5.8	5.9	5.6	5.5	5.4	5.3
25 to 29 years	8.9	9.3	9.3	9.6	9.8	9.9
30 to 34 years	10.1	10.2	10.0	10.1	10.2	10.5
35 to 39 years	9.4	9.9	10.0	10.4	10.5	10.7
40 to 44 years	9.4	9.2	8.9	9.0	9.1	9.5
45 to 49 years	10.9	10.6	10.2	9.9	9.5	9.3
50 to 54 years	13.4	13.0	12.1	11.4	10.8	10.3
55 to 59 years	12.0	12.1	12.2	12.1	11.8	11.4
60 to 64 years	8.0	8.2	8.7	9.0	9.2	9.2
65 years and older	7.9	7.5	9.0	9.3	9.9	10.3
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 1.19b. Individuals with diagnosis of schizophrenia and other psychotic disorders, by gender, age, race, and ethnicity: Column percent distribution, 2015–2020 (continued)

<i>Gender, age, race, and ethnicity</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	1.1	1.1	1.1	1.2	1.2	1.4
Asian ²	2.6	2.6	2.5	2.7	2.7	2.5
Black or African American	33.5	32.6	32.7	32.5	32.9	33.6
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	54.9	55.4	55.0	53.8	52.6	52.0
Other	7.4	7.6	8.4	9.6	10.3	10.1
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	87.6	86.9	87.2	86.6	86.3	84.5
Hispanic or Latino origin	12.4	13.1	12.8	13.4	13.7	15.5
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	26.0	23.9	26.3	24.4	26.5	25.0
Puerto Rican	24.8	23.3	24.3	25.3	30.0	29.5
Cuban	10.6	11.3	10.2	9.6	9.9	9.8
Other specific	38.5	41.5	39.2	40.7	33.6	35.7
Total	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

¹ Age is based on the individual's age at the midpoint of the reporting period.

² For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding. This table refers to all mental health diagnoses, which is up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 1.20a. Individuals with co-occurring mental health and substance use disorders, by gender, age, race, and ethnicity: Number, 2015–2020

<i>Gender, age, race, and ethnicity</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Number of states reporting	44	44	46	46	46	49
Total individuals with co-occurring mental health and substance use disorders¹	750,256	685,036	927,904	1,073,395	1,318,814	1,783,845
Gender						
Male	419,787	380,595	499,349	575,371	690,535	901,050
Female	330,024	303,909	425,157	496,040	625,338	879,767
Total	749,811	684,504	924,506	1,071,411	1,315,873	1,780,817
Age²						
0 to 11 years	2,025	2,006	22,998	23,657	53,591	119,695
12 to 14 years	7,915	7,468	19,190	18,617	36,682	72,138
15 to 17 years	35,092	33,046	45,987	48,803	64,160	93,684
18 to 20 years	30,768	28,262	36,644	42,613	56,194	77,477
21 to 24 years	57,227	50,825	64,267	73,144	89,569	117,078
25 to 29 years	90,205	84,149	111,447	132,700	157,854	197,270
30 to 34 years	95,197	87,726	114,073	134,794	159,794	201,555
35 to 39 years	80,659	77,483	104,408	125,281	147,478	185,646
40 to 44 years	75,035	65,677	82,694	96,492	114,562	149,637
45 to 49 years	81,331	71,122	88,735	99,947	111,954	137,622
50 to 54 years	85,803	75,755	93,412	103,115	113,457	138,263
55 to 59 years	60,064	56,406	76,167	90,658	104,980	134,590
60 to 64 years	30,343	29,050	41,409	51,217	63,255	88,164
65 years and older	17,761	15,173	24,938	31,327	43,792	70,533
Total	749,425	684,148	926,369	1,072,365	1,317,322	1,783,352

Continued. See notes at end of table.

Table 1.20a. Individuals with co-occurring mental health and substance use disorders, by gender, age, race, and ethnicity: Number, 2015–2020 (continued)

<i>Gender, age, race, and ethnicity</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	13,693	13,209	15,667	19,438	27,676	34,487
Asian ³	6,269	7,170	7,906	9,740	10,035	14,562
Black or African American	135,423	122,733	177,093	206,471	260,083	387,382
Native Hawaiian or Other Pacific Islander	2,143	1,680	1,781	2,243	2,983	3,330
White	478,797	440,180	593,135	679,019	830,679	1,110,251
Other	61,418	58,398	79,897	101,539	130,939	160,810
Total	697,743	643,370	875,479	1,018,450	1,262,395	1,710,822
Ethnicity						
Not of Hispanic or Latino origin	557,566	530,491	757,150	867,304	1,068,223	1,467,001
Hispanic or Latino origin	91,915	90,867	102,697	114,837	153,810	219,444
Total	649,481	621,358	859,847	982,141	1,222,033	1,686,445
Specific Hispanic or Latino origin						
Mexican	6,004	2,743	4,741	5,812	9,695	11,540
Puerto Rican	6,986	5,091	6,642	6,994	15,199	17,250
Cuban	852	638	887	973	6,922	8,461
Other specific	7,484	5,533	6,934	7,588	12,970	16,880
Total	21,326	14,005	19,204	21,367	44,786	54,131

¹ Total individuals with co-occurring mental health and substance use disorders who had at least one valid mental health diagnosis other than an alcohol- or substance-related disorder.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 1.20b. Individuals with co-occurring mental health and substance use disorders, by gender, age, race, and ethnicity: Column percent distribution, 2015–2020

<i>Gender, age, race, and ethnicity</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Number of states reporting	44	44	46	46	46	49
Total individuals with co-occurring mental health and substance use disorders¹	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	56.0	55.6	54.0	53.7	52.5	50.6
Female	44.0	44.4	46.0	46.3	47.5	49.4
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age²						
0 to 11 years	†	†	2.5	2.2	4.1	6.7
12 to 14 years	1.1	1.1	2.1	1.7	2.8	4.0
15 to 17 years	4.7	4.8	5.0	4.6	4.9	5.3
18 to 20 years	4.1	4.1	4.0	4.0	4.3	4.3
21 to 24 years	7.6	7.4	6.9	6.8	6.8	6.6
25 to 29 years	12.0	12.3	12.0	12.4	12.0	11.1
30 to 34 years	12.7	12.8	12.3	12.6	12.1	11.3
35 to 39 years	10.8	11.3	11.3	11.7	11.2	10.4
40 to 44 years	10.0	9.6	8.9	9.0	8.7	8.4
45 to 49 years	10.9	10.4	9.6	9.3	8.5	7.7
50 to 54 years	11.4	11.1	10.1	9.6	8.6	7.8
55 to 59 years	8.0	8.2	8.2	8.5	8.0	7.5
60 to 64 years	4.0	4.2	4.5	4.8	4.8	4.9
65 years and older	2.4	2.2	2.7	2.9	3.3	4.0
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 1.20b. Individuals with co-occurring mental health and substance use disorders, by gender, age, race, and ethnicity: Column percent distribution, 2015–2020 (continued)

<i>Gender, age, race, and ethnicity</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	2.0	2.1	1.8	1.9	2.2	2.0
Asian ³	†	1.1	†	†	†	†
Black or African American	19.4	19.1	20.2	20.3	20.6	22.6
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	68.6	68.4	67.7	66.7	65.8	64.9
Other	8.8	9.1	9.1	10.0	10.4	9.4
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	85.8	85.4	88.1	88.3	87.4	87.0
Hispanic or Latino origin	14.2	14.6	11.9	11.7	12.6	13.0
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	28.2	19.6	24.7	27.2	21.6	21.3
Puerto Rican	32.8	36.4	34.6	32.7	33.9	31.9
Cuban	4.0	4.6	4.6	4.6	15.5	15.6
Other specific	35.1	39.5	36.1	35.5	29.0	31.2
Total	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

¹ Total individuals with co-occurring mental health and substance use disorders who had at least one valid mental health diagnosis other than an alcohol- or substance-related disorder.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 1.21a. Individuals without co-occurring mental health and substance use disorders, by gender, age, race, and ethnicity: Number, 2015–2020

<i>Gender, age, race, and ethnicity</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Number of states reporting	44	44	46	46	46	48
Total individuals without co-occurring mental health and substance use disorders	3,343,624	3,437,247	3,746,707	4,013,406	3,856,296	3,541,912
Gender						
Male	1,540,792	1,594,358	1,723,559	1,845,739	1,764,454	1,615,109
Female	1,788,025	1,825,156	2,009,768	2,157,394	2,078,749	1,914,593
Total	3,328,817	3,419,514	3,733,327	4,003,133	3,843,203	3,529,702
Age¹						
0 to 11 years	589,681	645,141	629,626	710,115	689,814	615,153
12 to 14 years	288,945	305,563	306,094	343,968	345,987	319,552
15 to 17 years	280,651	302,307	306,989	333,818	329,777	295,582
18 to 20 years	144,611	153,252	165,117	183,075	181,609	164,078
21 to 24 years	171,805	172,696	190,166	200,886	192,433	177,223
25 to 29 years	236,046	244,369	277,092	294,805	281,612	255,677
30 to 34 years	237,467	241,933	267,814	279,879	267,580	247,426
35 to 39 years	215,831	222,753	252,418	267,312	252,816	231,916
40 to 44 years	208,506	204,084	220,838	229,374	215,228	200,960
45 to 49 years	219,763	217,686	239,559	244,804	222,223	198,529
50 to 54 years	244,128	238,661	261,105	261,912	235,691	211,708
55 to 59 years	208,493	208,659	242,080	256,088	239,183	220,654
60 to 64 years	138,018	138,493	169,695	188,108	181,738	175,523
65 years and older	150,266	130,154	210,492	216,684	217,781	224,618
Total	3,334,211	3,425,751	3,739,085	4,010,828	3,853,472	3,538,599

Continued. See notes at end of table.

Table 1.21a. Individuals without co-occurring mental health and substance use disorders, by gender, age, race, and ethnicity: Number, 2015–2020 (continued)

<i>Gender, age, race, and ethnicity</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	45,221	42,529	48,281	53,316	54,073	68,686
Asian ²	48,957	52,671	59,618	65,749	62,065	56,551
Black or African American	606,311	589,771	662,987	683,775	698,387	542,149
Native Hawaiian or Other Pacific Islander	10,047	12,303	7,027	9,605	9,538	8,282
White	2,001,375	2,117,111	2,307,099	2,435,513	2,280,974	2,081,146
Other	300,488	335,794	367,624	417,513	421,895	406,203
Total	3,012,399	3,150,179	3,452,636	3,665,471	3,526,932	3,163,017
Ethnicity						
Not of Hispanic or Latino origin	2,244,427	2,379,293	2,871,519	2,933,837	2,932,239	2,413,563
Hispanic or Latino origin	452,574	528,577	575,808	625,958	640,863	782,017
Total	2,697,001	2,907,870	3,447,327	3,559,795	3,573,102	3,195,580
Specific Hispanic or Latino origin						
Mexican	36,641	27,880	39,877	39,429	40,288	36,365
Puerto Rican	11,422	9,322	8,550	8,935	7,442	6,078
Cuban	1,624	1,424	1,760	1,634	1,820	1,629
Other specific	26,039	23,512	31,544	32,094	31,294	30,421
Total	75,726	62,138	81,731	82,092	80,844	74,493

¹ Age is based on the individual's age at the midpoint of the reporting period.

² For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 1.21b. Individuals without co-occurring mental health and substance use disorders, by gender, age, race, and ethnicity: Column percent distribution, 2015–2020

<i>Gender, age, race, and ethnicity</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Number of states reporting	44	44	46	46	46	48
Total individuals without co-occurring mental health and substance use disorders	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	46.3	46.6	46.2	46.1	45.9	45.8
Female	53.7	53.4	53.8	53.9	54.1	54.2
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age¹						
0 to 11 years	17.7	18.8	16.8	17.7	17.9	17.4
12 to 14 years	8.7	8.9	8.2	8.6	9.0	9.0
15 to 17 years	8.4	8.8	8.2	8.3	8.6	8.4
18 to 20 years	4.3	4.5	4.4	4.6	4.7	4.6
21 to 24 years	5.2	5.0	5.1	5.0	5.0	5.0
25 to 29 years	7.1	7.1	7.4	7.4	7.3	7.2
30 to 34 years	7.1	7.1	7.2	7.0	6.9	7.0
35 to 39 years	6.5	6.5	6.8	6.7	6.6	6.6
40 to 44 years	6.3	6.0	5.9	5.7	5.6	5.7
45 to 49 years	6.6	6.4	6.4	6.1	5.8	5.6
50 to 54 years	7.3	7.0	7.0	6.5	6.1	6.0
55 to 59 years	6.3	6.1	6.5	6.4	6.2	6.2
60 to 64 years	4.1	4.0	4.5	4.7	4.7	5.0
65 years and older	4.5	3.8	5.6	5.4	5.7	6.3
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 1.21b. Individuals without co-occurring mental health and substance use disorders, by gender, age, race, and ethnicity: Column percent distribution, 2015–2020 (continued)

<i>Gender, age, race, and ethnicity</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	1.5	1.4	1.4	1.5	1.5	2.2
Asian ²	1.6	1.7	1.7	1.8	1.8	1.8
Black or African American	20.1	18.7	19.2	18.7	19.8	17.1
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	66.4	67.2	66.8	66.4	64.7	65.8
Other	10.0	10.7	10.6	11.4	12.0	12.8
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	83.2	81.8	83.3	82.4	82.1	75.5
Hispanic or Latino origin	16.8	18.2	16.7	17.6	17.9	24.5
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	48.4	44.9	48.8	48.0	49.8	48.8
Puerto Rican	15.1	15.0	10.5	10.9	9.2	8.2
Cuban	2.1	2.3	2.2	2.0	2.3	2.2
Other specific	34.4	37.8	38.6	39.1	38.7	40.8
Total	100.0	100.0	100.0	100.0	100.0	100.0

¹ Age is based on the individual's age at the midpoint of the reporting period.

² For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 1.22a. Individuals who received mental health services and U.S. population, by gender, marital status, and age group: Number, 2020

Gender and marital status	Age group				
	0–17 years	18–24 years	25–54 years	55 years and older	Total
Individuals who received mental health services ¹					
Gender					
Male	1,059,019	335,403	1,455,295	499,199	3,348,916
Female	850,828	385,360	1,661,394	676,894	3,574,476
Total	1,909,847	720,763	3,116,689	1,176,093	6,923,392
Marital status					
Never married	1,254,530	353,845	1,100,734	280,526	2,989,635
Currently married	617	14,334	259,167	140,706	414,824
Separated	336	3,926	111,514	48,904	164,680
Divorced	817	3,551	206,244	166,832	377,444
Widowed	49	466	33,630	63,074	97,219
Total	1,256,349	376,122	1,711,289	700,042	4,043,802
U.S. population					
Gender					
Male	37,194,403	15,335,918	64,414,939	45,310,942	162,256,202
Female	35,627,710	14,691,079	64,157,032	52,752,100	167,227,921
Total	72,822,113	30,026,997	128,571,971	98,063,042	329,484,123
Marital status					
Never married	n/a	26,696,000	39,771,000	7,812,000	74,279,000
Currently married	n/a	1,985,000	71,329,000	59,855,000	133,169,000
Separated	n/a	170,000	2,730,000	1,544,000	4,444,000
Divorced	n/a	154,000	11,137,000	14,036,000	25,327,000
Widowed	n/a	47,000	1,209,000	13,475,000	14,731,000
Total	n/a	29,051,000	126,173,000	96,721,000	251,945,000

n/a Population detail required for marital status was not available.

¹ Age is based on the age of the individual who received mental health services at the midpoint of the reporting period.

SOURCES: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.; U.S. population: U.S. Census Bureau, Current Population Survey, 2020 Annual Social and Economic Supplement, December 2020, Table A1.

<https://www2.census.gov/programs-surveys/demo/tables/families/2020/cps-2020/ta01-all.xls> ; Population: U.S. Census Bureau, Population Estimates Program, SC-EST2020-alldata6: Annual State Resident Population Estimates for 6 Race Groups (5 Race Alone Groups and Two or More Races) by Age, Sex, and Hispanic Origin: April 1, 2010 to July 1, 2019; April 1, 2020; and July 1, 2020.

<https://www2.census.gov/programs-surveys/popest/datasets/2010-2020/state/asrh/SC-EST2020-ALLDATA6.csv>.

Table 1.22b. Individuals who received mental health services and U.S. population, by gender, marital status, and age group: Column percent distribution, 2020

Gender and marital status	Age group				
	0–17 years	18–24 years	25–54 years	55 years and older	Total
<i>Individuals who received mental health services[†]</i>					
Gender					
Male	55.5	46.5	46.7	42.4	48.4
Female	44.5	53.5	53.3	57.6	51.6
Total	100.0	100.0	100.0	100.0	100.0
Marital status					
Never married	99.9	94.1	64.3	40.1	73.9
Currently married	†	3.8	15.1	20.1	10.3
Separated	†	1.0	6.5	7.0	4.1
Divorced	†	†	12.1	23.8	9.3
Widowed	†	†	2.0	9.0	2.4
Total	100.0	100.0	100.0	100.0	100.0
<i>U.S. population</i>					
Gender					
Male	51.1	51.1	50.1	46.2	49.2
Female	48.9	48.9	49.9	53.8	50.8
Total	100.0	100.0	100.0	100.0	100.0
Marital status					
Never married	n/a	91.9	31.5	8.1	29.5
Currently married	n/a	6.8	56.5	61.9	52.9
Separated	n/a	†	2.2	1.6	1.8
Divorced	n/a	†	8.8	14.5	10.1
Widowed	n/a	†	†	13.9	5.8
Total	n/a	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals; n/a Population detail required for marital status was not available.

[†] Age is based on the age of the individual who received mental health services at the midpoint of the reporting period.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCES: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.; U.S. population: U.S. Census Bureau, Current Population Survey, 2020 Annual Social and Economic Supplement, December 2020, Table A1. <https://www2.census.gov/programs-surveys/demo/tables/families/2020/cps-2020/ta01a1-all.xls> ; Population: U.S. Census Bureau, Population Estimates Program, SC-EST2020-alldata6: Annual State Resident Population Estimates for 6 Race Groups (5 Race Alone Groups and Two or More Races) by Age, Sex, and Hispanic Origin: April 1, 2010 to July 1, 2019; April 1, 2020; and July 1, 2020. <https://www2.census.gov/programs-surveys/popest/datasets/2010-2020/state/asrh/SC-EST2020-ALLDATA6.csv>.

Table 1.22c. Individuals who received mental health services and U.S. population, by gender, marital status, and age group: Row percent distribution, 2020

Gender and marital status	Age group				
	0–17 years	18–24 years	25–54 years	55 years and older	Total
Individuals who received mental health services ¹					
Gender					
Male	31.6	10.0	43.5	14.9	100.0
Female	23.8	10.8	46.5	18.9	100.0
Total	27.6	10.4	45.0	17.0	100.0
Marital status					
Never married	42.0	11.8	36.8	9.4	100.0
Currently married	†	3.5	62.5	33.9	100.0
Separated	†	2.4	67.7	29.7	100.0
Divorced	†	†	54.6	44.2	100.0
Widowed	†	†	34.6	64.9	100.0
Total	31.1	9.3	42.3	17.3	100.0
U.S. population					
Gender					
Male	22.9	9.5	39.7	27.9	100.0
Female	21.3	8.8	38.4	31.5	100.0
Total	22.1	9.1	39.0	29.8	100.0
Marital status					
Never married	n/a	35.9	53.5	10.5	100.0
Currently married	n/a	1.5	53.6	44.9	100.0
Separated	n/a	3.8	61.4	34.7	100.0
Divorced	n/a	†	44.0	55.4	100.0
Widowed	n/a	†	8.2	91.5	100.0
Total	n/a	11.5	50.1	38.4	100.0

† Less than 1 percent of individuals; n/a Population detail required for marital status was not available.

¹ Age is based on the age of the individual who received mental health services at the midpoint of the reporting period.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCES: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.; U.S. population: U.S. Census Bureau, Current Population Survey, 2020 Annual Social and Economic Supplement, December 2020, Table A1.

<https://www2.census.gov/programs-surveys/demo/tables/families/2020/cps-2020/ta011-all.xls> ; Population: U.S. Census Bureau, Population Estimates Program, SC-EST2020-alldata6: Annual State Resident Population Estimates for 6 Race Groups (5 Race Alone Groups and Two or More Races) by Age, Sex, and Hispanic Origin: April 1, 2010 to July 1, 2019; April 1, 2020; and July 1, 2020.

<https://www2.census.gov/programs-surveys/popest/datasets/2010-2020/state/asrh/SC-EST2020-ALLDATA6.csv>.

Table 1.23a. Individuals who received mental health services and U.S. population, by race, ethnicity, and age group: Number, 2020

Race and ethnicity	Age group				
	0–17 years	18–24 years	25–54 years	55 years and older	Total
Individuals who received mental health services¹					
Race					
American Indian or Alaska Native	39,909	14,620	63,295	15,543	133,367
Asian²	20,287	10,093	41,416	19,923	91,719
Black or African American	312,263	126,633	570,602	220,420	1,229,918
Native Hawaiian or Other Pacific Islander	4,392	1,808	7,157	1,930	15,287
White	1,023,729	417,017	1,946,322	750,151	4,137,219
Other	277,290	84,694	251,296	69,464	682,744
Total	1,677,870	654,865	2,880,088	1,077,431	6,290,254
Ethnicity					
Not of Hispanic or Latino origin	1,227,720	505,877	2,364,429	916,771	5,014,797
Hispanic or Latino origin	458,587	144,063	486,305	152,404	1,241,359
Total	1,686,307	649,940	2,850,734	1,069,175	6,256,156
U.S. population					
Race					
American Indian or Alaska Native	1,229,661	452,967	1,733,118	877,244	4,292,990
Asian	4,172,530	1,758,735	9,227,549	4,853,464	20,012,278
Black or African American	11,116,083	4,550,480	18,268,568	10,595,981	44,531,112
Native Hawaiian or Other Pacific Islander	237,720	83,109	366,284	156,580	843,693
White	51,859,580	21,949,686	96,004,577	80,495,881	250,309,724
Other	4,206,539	1,232,020	2,971,875	1,083,892	9,494,326
Total	72,822,113	30,026,997	128,571,971	98,063,042	329,484,123
Ethnicity					
Not of Hispanic or Latino origin	54,190,278	23,104,146	102,951,526	87,925,294	268,171,244
Hispanic or Latino origin	18,631,835	6,922,851	25,620,445	10,137,748	61,312,879
Total	72,822,113	30,026,997	128,571,971	98,063,042	329,484,123

¹ Age is based on the age of the individual who received mental health services at the midpoint of the reporting period.

² For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

SOURCES: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.; Population: U.S. Census Bureau, Population Estimates Program, SC-EST2020-alldata6: Annual State Resident Population Estimates for 6 Race Groups (5 Race Alone Groups and Two or More Races) by Age, Sex, and Hispanic Origin: April 1, 2010 to July 1, 2019; April 1, 2020; and July 1, 2020. <https://www2.census.gov/programs-surveys/popest/datasets/2010-2020/state/asrh/SC-EST2020-ALLDATA6.csv>.

Table 1.23b. Individuals who received mental health services and U.S. population, by race, ethnicity, and age group: Column percent distribution, 2020

Race and ethnicity	Age group				
	0–17 years	18–24 years	25–54 years	55 years and older	Total
	Individuals who received mental health services ¹				
Race					
American Indian or Alaska Native	2.4	2.2	2.2	1.4	2.1
Asian ²	1.2	1.5	1.4	1.8	1.5
Black or African American	18.6	19.3	19.8	20.5	19.6
Native Hawaiian or Other Pacific Islander	†	†	†	†	†
White	61.0	63.7	67.6	69.6	65.8
Other	16.5	12.9	8.7	6.4	10.9
Total	100.0	100.0	100.0	100.0	100.0
Ethnicity					
Not of Hispanic or Latino origin	72.8	77.8	82.9	85.7	80.2
Hispanic or Latino origin	27.2	22.2	17.1	14.3	19.8
Total	100.0	100.0	100.0	100.0	100.0
U.S. population					
Race					
American Indian or Alaska Native	1.7	1.5	1.3	†	1.3
Asian	5.7	5.9	7.2	4.9	6.1
Black or African American	15.3	15.2	14.2	10.8	13.5
Native Hawaiian or Other Pacific Islander	†	†	†	†	†
White	71.2	73.1	74.7	82.1	76.0
Other	5.8	4.1	2.3	1.1	2.9
Total	100.0	100.0	100.0	100.0	100.0
Ethnicity					
Not of Hispanic or Latino origin	74.4	76.9	80.1	89.7	81.4
Hispanic or Latino origin	25.6	23.1	19.9	10.3	18.6
Total	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

¹ Age is based on the age of the individual who received mental health services at the midpoint of the reporting period.

² For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCES: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.; Population: U.S. Census Bureau, Population Estimates Program, SC-EST2020-alldata6: Annual State Resident Population Estimates for 6 Race Groups (5 Race Alone Groups and Two or More Races) by Age, Sex, and Hispanic Origin: April 1, 2010 to July 1, 2019; April 1, 2020; and July 1, 2020. <https://www2.census.gov/programs-surveys/popest/datasets/2010-2020/state/asrh/SC-EST2020-ALLDATA6.csv>.

Table 1.23c. Individuals who received mental health services and U.S. population, by race, ethnicity, and age group: Row percent distribution, 2020

Race and ethnicity	Age group				
	0–17 years	18–24 years	25–54 years	55 years and older	Total
Individuals who received mental health services ¹					
Race					
American Indian or Alaska Native	29.9	11.0	47.5	11.7	100.0
Asian ²	22.1	11.0	45.2	21.7	100.0
Black or African American	25.4	10.3	46.4	17.9	100.0
Native Hawaiian or Other Pacific Islander	28.7	11.8	46.8	12.6	100.0
White	24.7	10.1	47.0	18.1	100.0
Other	40.6	12.4	36.8	10.2	100.0
Total	26.7	10.4	45.8	17.1	100.0
Ethnicity					
Not of Hispanic or Latino origin	24.5	10.1	47.1	18.3	100.0
Hispanic or Latino origin	36.9	11.6	39.2	12.3	100.0
Total	27.0	10.4	45.6	17.1	100.0
U.S. population					
Race					
American Indian or Alaska Native	28.6	10.6	40.4	20.4	100.0
Asian	20.8	8.8	46.1	24.3	100.0
Black or African American	25.0	10.2	41.0	23.8	100.0
Native Hawaiian or Other Pacific Islander	28.2	9.9	43.4	18.6	100.0
White	20.7	8.8	38.4	32.2	100.0
Other	44.3	13.0	31.3	11.4	100.0
Total	22.1	9.1	39.0	29.8	100.0
Ethnicity					
Not of Hispanic or Latino origin	20.2	8.6	38.4	32.8	100.0
Hispanic or Latino origin	30.4	11.3	41.8	16.5	100.0
Total	22.1	9.1	39.0	29.8	100.0

¹ Age is based on the age of the individual who received mental health services at the midpoint of the reporting period.

² For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCES: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.; Population: U.S. Census Bureau, Population Estimates Program, SC-EST2020-alldata6: Annual State Resident Population Estimates for 6 Race Groups (5 Race Alone Groups and Two or More Races) by Age, Sex, and Hispanic Origin: April 1, 2010 to July 1, 2019; April 1, 2020; and July 1, 2020. <https://www2.census.gov/programs-surveys/popest/datasets/2010-2020/state/asrh/SC-EST2020-ALLDATA6.csv>.

Table 2.1a. Individuals aged 17 years and younger, by gender, age, race, ethnicity, and SED status: Number, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Number of states reporting	45	47	49	49	48	50
Total individuals aged 0 to 17 years	1,473,157	1,610,944	1,670,404	1,822,515	1,891,605	1,915,199
Gender						
Male	836,862	914,616	941,595	1,027,209	1,050,663	1,059,019
Female	633,671	690,267	724,057	790,216	834,398	850,828
Total	1,470,533	1,604,883	1,665,652	1,817,425	1,885,061	1,909,847
Age¹						
0 to 11 years	710,211	789,831	814,625	899,746	918,686	922,028
12 to 14 years	365,574	390,461	405,592	446,037	478,309	493,952
15 to 17 years	397,372	430,652	450,187	476,732	494,610	499,219
Total	1,473,157	1,610,944	1,670,404	1,822,515	1,891,605	1,915,199
Race						
American Indian or Alaska Native	22,179	22,280	23,717	27,261	30,968	39,909
Asian ²	11,749	14,737	17,522	19,498	19,958	20,287
Black or African American	292,895	303,691	320,898	331,362	348,443	312,263
Native Hawaiian or Other Pacific Islander	3,850	4,478	3,239	4,437	5,244	4,392
White	793,681	893,070	932,229	1,005,454	1,010,002	1,023,729
Other	205,282	220,149	242,625	258,128	272,356	277,290
Total	1,329,636	1,458,405	1,540,230	1,646,140	1,686,971	1,677,870

Continued. See notes at end of table.

Table 2.1a. Individuals aged 17 years and younger, by gender, age, race, ethnicity, and SED status: Number, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Ethnicity						
Not of Hispanic or Latino origin	912,344	1,002,179	1,171,583	1,206,929	1,241,793	1,227,720
Hispanic or Latino origin	309,023	332,201	368,810	396,574	408,639	458,587
Total	1,221,367	1,334,380	1,540,393	1,603,503	1,650,432	1,686,307
Specific Hispanic or Latino origin						
Mexican	28,027	20,592	25,404	25,761	26,539	25,103
Puerto Rican	12,942	10,455	9,614	6,422	5,198	5,984
Cuban	1,387	1,260	1,314	1,420	1,612	1,701
Other specific	19,289	17,917	20,610	21,197	18,605	19,266
Total	61,645	50,224	56,942	54,800	51,954	52,054
SED status³						
SED	1,007,979	1,042,065	1,140,786	1,275,291	1,420,637	1,333,084
Not SED	386,662	476,614	472,108	489,682	408,180	510,942
Total	1,394,641	1,518,679	1,612,894	1,764,973	1,828,817	1,844,026

SED = serious emotional disturbance.

¹ Age is based on the individual's age at the midpoint of the reporting period.

² For some states, "Asian" includes "Pacific Islanders." See Appendix C.

³ The federal definition for SED status can be found in Appendix D.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 2.1b. Individuals aged 17 years and younger, by gender, age, race, ethnicity, and SED status: Column percent distribution, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Number of states reporting	45	47	49	49	48	50
Total individuals aged 0 to 17 years	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	56.9	57.0	56.5	56.5	55.7	55.5
Female	43.1	43.0	43.5	43.5	44.3	44.5
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age¹						
0 to 11 years	48.2	49.0	48.8	49.4	48.6	48.1
12 to 14 years	24.8	24.2	24.3	24.5	25.3	25.8
15 to 17 years	27.0	26.7	27.0	26.2	26.1	26.1
Total	100.0	100.0	100.0	100.0	100.0	100.0
Race						
American Indian or Alaska Native	1.7	1.5	1.5	1.7	1.8	2.4
Asian ²	†	1.0	1.1	1.2	1.2	1.2
Black or African American	22.0	20.8	20.8	20.1	20.7	18.6
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	59.7	61.2	60.5	61.1	59.9	61.0
Other	15.4	15.1	15.8	15.7	16.1	16.5
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 2.1b. Individuals aged 17 years and younger, by gender, age, race, ethnicity, and SED status: Column percent distribution, 2015–2020
(continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Ethnicity						
Not of Hispanic or Latino origin	74.7	75.1	76.1	75.3	75.2	72.8
Hispanic or Latino origin	25.3	24.9	23.9	24.7	24.8	27.2
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	45.5	41.0	44.6	47.0	51.1	48.2
Puerto Rican	21.0	20.8	16.9	11.7	10.0	11.5
Cuban	2.2	2.5	2.3	2.6	3.1	3.3
Other specific	31.3	35.7	36.2	38.7	35.8	37.0
Total	100.0	100.0	100.0	100.0	100.0	100.0
SED status³						
SED	72.3	68.6	70.7	72.3	77.7	72.3
Not SED	27.7	31.4	29.3	27.7	22.3	27.7
Total	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

SED = serious emotional disturbance.

¹ Age is based on the individual's age at the midpoint of the reporting period.

² For some states, "Asian" includes "Pacific Islanders." See Appendix C.

³ The federal definition for SED status can be found in Appendix D.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 2.2a. Individuals aged 17 years and younger with SED, by gender, age, race, and ethnicity: Number, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Number of states reporting	45	47	48	49	48	50
Total individuals with SED aged 0 to 17 years¹	1,007,979	1,042,065	1,140,786	1,275,291	1,420,637	1,333,084
Gender						
Male	579,106	604,205	648,699	718,015	785,298	726,517
Female	427,000	435,738	489,191	554,565	630,668	602,592
Total	1,006,106	1,039,943	1,137,890	1,272,580	1,415,966	1,329,109
Age²						
0 to 11 years	475,798	492,813	543,760	608,733	685,794	632,751
12 to 14 years	255,186	260,770	283,252	320,749	362,381	349,172
15 to 17 years	276,995	288,482	313,774	345,809	372,462	351,161
Total	1,007,979	1,042,065	1,140,786	1,275,291	1,420,637	1,333,084
Race						
American Indian or Alaska Native	12,427	12,329	14,821	17,617	25,323	27,985
Asian ³	7,935	9,716	11,424	13,502	15,043	13,987
Black or African American	218,409	206,597	220,966	237,846	257,462	214,137
Native Hawaiian or Other Pacific Islander	2,862	3,245	2,449	3,609	4,457	3,097
White	528,472	567,741	604,681	655,865	728,317	692,445
Other	160,057	166,891	202,056	215,667	225,812	221,478
Total	930,162	966,519	1,056,397	1,144,106	1,256,414	1,173,129

Continued. See notes at end of table.

Table 2.2a. Individuals aged 17 years and younger with SED, by gender, age, race, and ethnicity: Number, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Ethnicity						
Not of Hispanic or Latino origin	633,760	664,713	765,238	802,168	874,874	841,847
Hispanic or Latino origin	221,982	239,442	279,790	295,611	339,505	333,885
Total	855,742	904,155	1,045,028	1,097,779	1,214,379	1,175,732
Specific Hispanic or Latino origin						
Mexican	16,895	13,445	17,789	18,134	18,244	16,791
Puerto Rican	8,003	7,131	5,728	5,093	2,841	3,306
Cuban	1,196	1,097	1,149	1,258	773	723
Other specific	14,289	13,256	15,197	15,842	12,462	12,959
Total	40,383	34,929	39,863	40,327	34,320	33,779

SED = serious emotional disturbance.

¹ The federal definition for SED status can be found in Appendix D.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 2.2b. Individuals aged 17 years and younger with SED, by gender, age, race, and ethnicity: Column percent distribution, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Number of states reporting	45	47	48	49	48	50
Total individuals with SED aged 0 to 17 years¹	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	57.6	58.1	57.0	56.4	55.5	54.7
Female	42.4	41.9	43.0	43.6	44.5	45.3
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age²						
0 to 11 years	47.2	47.3	47.7	47.7	48.3	47.5
12 to 14 years	25.3	25.0	24.8	25.2	25.5	26.2
15 to 17 years	27.5	27.7	27.5	27.1	26.2	26.3
Total	100.0	100.0	100.0	100.0	100.0	100.0
Race						
American Indian or Alaska Native	1.3	1.3	1.4	1.5	2.0	2.4
Asian ³	†	1.0	1.1	1.2	1.2	1.2
Black or African American	23.5	21.4	20.9	20.8	20.5	18.3
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	56.8	58.7	57.2	57.3	58.0	59.0
Other	17.2	17.3	19.1	18.9	18.0	18.9
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 2.2b. Individuals aged 17 years and younger with SED, by gender, age, race, and ethnicity: Column percent distribution, 2015–2020
(continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Ethnicity						
Not of Hispanic or Latino origin	74.1	73.5	73.2	73.1	72.0	71.6
Hispanic or Latino origin	25.9	26.5	26.8	26.9	28.0	28.4
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	41.8	38.5	44.6	45.0	53.2	49.7
Puerto Rican	19.8	20.4	14.4	12.6	8.3	9.8
Cuban	3.0	3.1	2.9	3.1	2.3	2.1
Other specific	35.4	38.0	38.1	39.3	36.3	38.4
Total	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

SED = serious emotional disturbance.

¹ The federal definition for SED status can be found in Appendix D.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 2.3a. Individuals aged 17 years and younger, by mental health diagnosis, gender, race, and ethnicity: Number, 2020

		Mental health diagnosis¹									
Gender, race, and ethnicity		Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Conduct disorder	Mood disorders		ODD	PDD	Other mental disorders¹
							Bipolar disorders	Depressive disorders			
Total individuals aged 0 to 17 years		1,637,967	496,671	322,311	481,629	93,111	27,577	301,271	162,854	69,912	319,355
Gender											
Male		905,484	241,832	141,181	345,420	66,913	13,317	109,628	112,133	55,697	187,281
Female		727,593	253,120	180,414	134,542	25,885	14,155	190,996	50,188	14,050	131,416
Total		1,633,077	494,952	321,595	479,962	92,798	27,472	300,624	162,321	69,747	318,697
Race											
American Indian or Alaska Native		36,333	13,960	7,297	6,508	1,300	351	7,261	1,999	1,103	10,050
Asian²		15,154	4,132	3,226	2,663	592	200	3,930	710	1,111	3,287
Black or African American		273,438	76,030	30,339	107,622	23,854	5,122	37,247	41,784	8,422	48,780
Native Hawaiian or Other Pacific Islander		3,558	1,252	755	734	201	48	810	243	165	672
White		891,148	275,278	199,330	269,564	45,944	16,816	167,419	84,847	41,487	172,208
Other		245,414	70,399	47,396	57,823	13,321	3,132	55,420	22,741	8,033	46,552
Total		1,465,045	441,051	288,343	444,914	85,212	25,669	272,087	152,324	60,321	281,549
Ethnicity											
Not of Hispanic or Latino origin		1,073,643	328,403	221,493	352,004	66,985	20,476	191,961	122,521	44,085	194,301
Hispanic or Latino origin		393,918	113,991	71,415	92,163	17,667	5,018	81,078	29,017	15,021	86,651
Total		1,467,561	442,394	292,908	444,167	84,652	25,494	273,039	151,538	59,106	280,952
Specific Hispanic or Latino origin											
Mexican		24,442	9,068	5,294	4,790	980	198	6,309	1,735	728	3,570
Puerto Rican		4,838	1,356	642	1,707	357	94	866	454	185	899
Cuban		1,616	435	232	398	88	19	288	74	47	436
Other specific		18,493	6,432	3,873	4,656	1,255	207	4,582	1,870	591	2,802
Total		49,389	17,291	10,041	11,551	2,680	518	12,045	4,133	1,551	7,707

ADHD = attention deficit/hyperactivity disorder; ODD = oppositional defiant disorder; PDD = pervasive developmental disorders.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes substance use diagnoses. See Appendix E. Mental health diagnostic category definitions can be found in Appendix E.

² For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 2.3b. Individuals aged 17 years and younger, by mental health diagnosis, gender, race, and ethnicity: Column percent distribution, 2020

Gender, race, and ethnicity	Mental health diagnosis ¹									
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Conduct disorder	Mood disorders		ODD	PDD	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Total individuals aged 0 to 17 years	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender										
Male	55.4	48.9	43.9	72.0	72.1	48.5	36.5	69.1	79.9	58.8
Female	44.6	51.1	56.1	28.0	27.9	51.5	63.5	30.9	20.1	41.2
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race										
American Indian or Alaska Native	2.5	3.2	2.5	1.5	1.5	1.4	2.7	1.3	1.8	3.6
Asian ²	1.0	†	1.1	†	†	†	1.4	†	1.8	1.2
Black or African American	18.7	17.2	10.5	24.2	28.0	20.0	13.7	27.4	14.0	17.3
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	†	†	†	†
White	60.8	62.4	69.1	60.6	53.9	65.5	61.5	55.7	68.8	61.2
Other	16.8	16.0	16.4	13.0	15.6	12.2	20.4	14.9	13.3	16.5
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity										
Not of Hispanic or Latino origin	73.2	74.2	75.6	79.3	79.1	80.3	70.3	80.9	74.6	69.2
Hispanic or Latino origin	26.8	25.8	24.4	20.7	20.9	19.7	29.7	19.1	25.4	30.8
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin										
Mexican	49.5	52.4	52.7	41.5	36.6	38.2	52.4	42.0	46.9	46.3
Puerto Rican	9.8	7.8	6.4	14.8	13.3	18.1	7.2	11.0	11.9	11.7
Cuban	3.3	2.5	2.3	3.4	3.3	3.7	2.4	1.8	3.0	5.7
Other specific	37.4	37.2	38.6	40.3	46.8	40.0	38.0	45.2	38.1	36.4
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; ODD = oppositional defiant disorder; PDD = pervasive developmental disorders.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes substance use diagnoses. See Appendix E. Mental health diagnostic category definitions can be found in Appendix E.

² For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 2.3c. Individuals aged 17 years and younger, by mental health diagnosis, gender, race, and ethnicity: Row percent distribution, 2020

Gender, race, and ethnicity	Total	Mental health diagnosis ¹								
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Conduct disorder	Mood disorders		ODD	PDD	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Total individuals aged 0 to 17 years	100.0	30.3	19.7	29.4	5.7	1.7	18.4	9.9	4.3	19.5
Gender										
Male	100.0	26.7	15.6	38.1	7.4	1.5	12.1	12.4	6.2	20.7
Female	100.0	34.8	24.8	18.5	3.6	1.9	26.3	6.9	1.9	18.1
Total	100.0	30.3	19.7	29.4	5.7	1.7	18.4	9.9	4.3	19.5
Race										
American Indian or Alaska Native	100.0	38.4	20.1	17.9	3.6	†	20.0	5.5	3.0	27.7
Asian ²	100.0	27.3	21.3	17.6	3.9	1.3	25.9	4.7	7.3	21.7
Black or African American	100.0	27.8	11.1	39.4	8.7	1.9	13.6	15.3	3.1	17.8
Native Hawaiian or Other Pacific Islander	100.0	35.2	21.2	20.6	5.6	1.3	22.8	6.8	4.6	18.9
White	100.0	30.9	22.4	30.2	5.2	1.9	18.8	9.5	4.7	19.3
Other	100.0	28.7	19.3	23.6	5.4	1.3	22.6	9.3	3.3	19.0
Total	100.0	30.1	19.7	30.4	5.8	1.8	18.6	10.4	4.1	19.2
Ethnicity										
Not of Hispanic or Latino origin	100.0	30.6	20.6	32.8	6.2	1.9	17.9	11.4	4.1	18.1
Hispanic or Latino origin	100.0	28.9	18.1	23.4	4.5	1.3	20.6	7.4	3.8	22.0
Total	100.0	30.1	20.0	30.3	5.8	1.7	18.6	10.3	4.0	19.1
Specific Hispanic or Latino origin										
Mexican	100.0	37.1	21.7	19.6	4.0	†	25.8	7.1	3.0	14.6
Puerto Rican	100.0	28.0	13.3	35.3	7.4	1.9	17.9	9.4	3.8	18.6
Cuban	100.0	26.9	14.4	24.6	5.4	1.2	17.8	4.6	2.9	27.0
Other specific	100.0	34.8	20.9	25.2	6.8	1.1	24.8	10.1	3.2	15.2
Total	100.0	35.0	20.3	23.4	5.4	1.0	24.4	8.4	3.1	15.6

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; ODD = oppositional defiant disorder; PDD = pervasive developmental disorders.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes substance use diagnoses. See Appendix E. Mental health diagnostic category definitions can be found in Appendix E.

² For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 2.4a. Individuals aged 17 years and younger with SED, by mental health diagnosis, gender, race, and ethnicity: Number, 2020

Gender, race, and ethnicity	Total	Mental health diagnosis ¹								
		Trauma- and stressor- related disorders	Anxiety disorders	ADHD	Conduct disorder	Mood disorders		ODD	PDD	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Total individuals with SED aged 0 to 17 years²	1,155,930	332,184	264,993	339,561	66,916	23,982	256,237	123,208	42,263	233,570
Gender										
Male	626,381	161,209	115,986	242,850	48,152	11,536	93,291	84,894	33,883	135,804
Female	525,671	169,608	148,449	95,302	18,513	12,357	162,416	37,852	8,244	97,274
Total	1,152,052	330,817	264,435	338,152	66,665	23,893	255,707	122,746	42,127	233,078
Race										
American Indian or Alaska Native	26,160	9,907	6,186	4,136	925	306	6,170	1,604	628	7,540
Asian ³	11,354	2,876	2,749	1,876	426	169	3,491	540	712	2,424
Black or African American	191,557	49,690	25,604	78,592	17,280	4,468	32,074	31,327	5,205	35,205
Native Hawaiian or Other Pacific Islander	2,824	1,014	659	567	176	43	734	207	90	502
White	608,807	178,545	159,778	187,497	31,656	14,533	138,196	62,751	27,111	123,934
Other	200,074	55,741	42,529	43,731	10,804	2,839	50,694	18,749	5,288	38,973
Total	1,040,776	297,773	237,505	316,399	61,267	22,358	231,359	115,178	39,034	208,578
Ethnicity										
Not of Hispanic or Latino origin	748,652	219,317	177,633	253,468	46,691	17,591	157,878	91,101	30,605	141,430
Hispanic or Latino origin	291,196	78,894	63,220	60,499	13,513	4,572	73,870	23,020	7,892	66,546
Total	1,039,848	298,211	240,853	313,967	60,204	22,163	231,748	114,121	38,497	207,976
Specific Hispanic or Latino origin										
Mexican	16,559	6,129	3,719	3,528	746	154	4,596	1,357	387	2,541
Puerto Rican	2,633	700	489	953	168	70	694	323	87	524
Cuban	708	283	158	180	30	11	185	40	18	127
Other specific	12,723	4,393	2,907	3,454	1,007	158	3,398	1,473	333	1,900
Total	32,623	11,505	7,273	8,115	1,951	393	8,873	3,193	825	5,092

ADHD = attention deficit/hyperactivity disorder; ODD = oppositional defiant disorder; PDD = pervasive developmental disorders; SED = serious emotional disturbance.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SED status can be found in Appendix D.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 2.4b. Individuals aged 17 years and younger with SED, by mental health diagnosis, gender, race, and ethnicity: Column percent distribution, 2020

Gender, race, and ethnicity	Total	Mental health diagnosis ¹								
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Conduct disorder	Mood disorders		ODD	PDD	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Total individuals with SED aged 0 to 17 years²	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender										
Male	54.4	48.7	43.9	71.8	72.2	48.3	36.5	69.2	80.4	58.3
Female	45.6	51.3	56.1	28.2	27.8	51.7	63.5	30.8	19.6	41.7
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race										
American Indian or Alaska Native	2.5	3.3	2.6	1.3	1.5	1.4	2.7	1.4	1.6	3.6
Asian ³	1.1	†	1.2	†	†	†	1.5	†	1.8	1.2
Black or African American	18.4	16.7	10.8	24.8	28.2	20.0	13.9	27.2	13.3	16.9
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	†	†	†	†
White	58.5	60.0	67.3	59.3	51.7	65.0	59.7	54.5	69.5	59.4
Other	19.2	18.7	17.9	13.8	17.6	12.7	21.9	16.3	13.5	18.7
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity										
Not of Hispanic or Latino origin	72.0	73.5	73.8	80.7	77.6	79.4	68.1	79.8	79.5	68.0
Hispanic or Latino origin	28.0	26.5	26.2	19.3	22.4	20.6	31.9	20.2	20.5	32.0
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin										
Mexican	50.8	53.3	51.1	43.5	38.2	39.2	51.8	42.5	46.9	49.9
Puerto Rican	8.1	6.1	6.7	11.7	8.6	17.8	7.8	10.1	10.5	10.3
Cuban	2.2	2.5	2.2	2.2	1.5	2.8	2.1	1.3	2.2	2.5
Other specific	39.0	38.2	40.0	42.6	51.6	40.2	38.3	46.1	40.4	37.3
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; ODD = oppositional defiant disorder; PDD = pervasive developmental disorders; SED = serious emotional disturbance.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SED status can be found in Appendix D.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 2.4c. Individuals aged 17 years and younger with SED, by mental health diagnosis, gender, race, and ethnicity: Row percent distribution, 2020

Gender, race, and ethnicity	Total	Mental health diagnosis ¹								
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Conduct disorder	Mood disorders		ODD	PDD	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Total individuals with SED aged 0 to 17 years²	100.0	28.7	22.9	29.4	5.8	2.1	22.2	10.7	3.7	20.2
Gender										
Male	100.0	25.7	18.5	38.8	7.7	1.8	14.9	13.6	5.4	21.7
Female	100.0	32.3	28.2	18.1	3.5	2.4	30.9	7.2	1.6	18.5
Total	100.0	28.7	23.0	29.4	5.8	2.1	22.2	10.7	3.7	20.2
Race										
American Indian or Alaska Native	100.0	37.9	23.6	15.8	3.5	1.2	23.6	6.1	2.4	28.8
Asian ³	100.0	25.3	24.2	16.5	3.8	1.5	30.7	4.8	6.3	21.3
Black or African American	100.0	25.9	13.4	41.0	9.0	2.3	16.7	16.4	2.7	18.4
Native Hawaiian or Other Pacific Islander	100.0	35.9	23.3	20.1	6.2	1.5	26.0	7.3	3.2	17.8
White	100.0	29.3	26.2	30.8	5.2	2.4	22.7	10.3	4.5	20.4
Other	100.0	27.9	21.3	21.9	5.4	1.4	25.3	9.4	2.6	19.5
Total	100.0	28.6	22.8	30.4	5.9	2.1	22.2	11.1	3.8	20.0
Ethnicity										
Not of Hispanic or Latino origin	100.0	29.3	23.7	33.9	6.2	2.3	21.1	12.2	4.1	18.9
Hispanic or Latino origin	100.0	27.1	21.7	20.8	4.6	1.6	25.4	7.9	2.7	22.9
Total	100.0	28.7	23.2	30.2	5.8	2.1	22.3	11.0	3.7	20.0
Specific Hispanic or Latino origin										
Mexican	100.0	37.0	22.5	21.3	4.5	†	27.8	8.2	2.3	15.3
Puerto Rican	100.0	26.6	18.6	36.2	6.4	2.7	26.4	12.3	3.3	19.9
Cuban	100.0	40.0	22.3	25.4	4.2	1.6	26.1	5.6	2.5	17.9
Other specific	100.0	34.5	22.8	27.1	7.9	1.2	26.7	11.6	2.6	14.9
Total	100.0	35.3	22.3	24.9	6.0	1.2	27.2	9.8	2.5	15.6

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; ODD = oppositional defiant disorder; PDD = pervasive developmental disorders; SED = serious emotional disturbance.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SED status can be found in Appendix D.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 2.5a. Individuals aged 17 years and younger, by mental health diagnosis, living arrangement, and SED status: Number, 2020

Living arrangement and SED status	Total	Mental health diagnosis ¹								
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Conduct disorder	Mood disorders		ODD	PDD	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Total individuals aged 0 to 17 years	1,637,967	496,671	322,311	481,629	93,111	27,577	301,271	162,854	69,912	319,355
Living arrangement										
Homeless	4,319	1,934	599	791	215	96	720	339	93	938
Foster home/foster care	35,862	20,289	5,245	7,844	1,930	554	4,378	2,989	736	7,170
Residential care	12,321	4,221	2,279	3,759	1,444	641	3,407	1,843	378	3,850
Private residence	861,422	257,205	188,353	276,808	52,825	15,880	176,906	97,249	35,595	187,152
Other	20,604	6,304	4,381	4,387	1,932	583	5,776	1,898	393	5,227
Total	934,528	289,953	200,857	293,589	58,346	17,754	191,187	104,318	37,195	204,337
SED status²										
SED	1,155,930	332,184	264,993	339,561	66,916	23,982	256,237	123,208	42,263	233,570
Not SED	445,567	153,092	50,349	131,081	24,422	2,951	38,423	35,367	24,109	78,207
Total	1,601,497	485,276	315,342	470,642	91,338	26,933	294,660	158,575	66,372	311,777

ADHD = attention deficit/hyperactivity disorder; ODD = oppositional defiant disorder; PDD = pervasive developmental disorders; SED = serious emotional disturbance.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SED status can be found in Appendix D.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 2.5b. Individuals aged 17 years and younger, by mental health diagnosis, living arrangement, and SED status: Column percent distribution, 2020

<i>Living arrangement and SED status</i>		<i>Mental health diagnosis¹</i>								
		<i>Trauma- and stressor-related disorders</i>				<i>Mood disorders</i>		<i>Other mental disorders¹</i>		
						<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
	<i>Total</i>							<i>ODD</i>	<i>PDD</i>	
Total individuals aged 0 to 17 years	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Living arrangement										
Homeless	†	†	†	†	†	†	†	†	†	†
Foster home/foster care	3.8	7.0	2.6	2.7	3.3	3.1	2.3	2.9	2.0	3.5
Residential care	1.3	1.5	1.1	1.3	2.5	3.6	1.8	1.8	1.0	1.9
Private residence	92.2	88.7	93.8	94.3	90.5	89.4	92.5	93.2	95.7	91.6
Other	2.2	2.2	2.2	1.5	3.3	3.3	3.0	1.8	1.1	2.6
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
SED status²										
SED	72.2	68.5	84.0	72.1	73.3	89.0	87.0	77.7	63.7	74.9
Not SED	27.8	31.5	16.0	27.9	26.7	11.0	13.0	22.3	36.3	25.1
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; ODD = oppositional defiant disorder; PDD = pervasive developmental disorders; SED = serious emotional disturbance.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SED status can be found in Appendix D.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 2.5c. Individuals aged 17 years and younger, by mental health diagnosis, living arrangement, and SED status: Row percent distribution, 2020

Living arrangement and SED status	Total	Mental health diagnosis ¹								
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Conduct disorder	Mood disorders		ODD	PDD	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Total individuals aged 0 to 17 years	100.0	30.3	19.7	29.4	5.7	1.7	18.4	9.9	4.3	19.5
Living arrangement										
Homeless	100.0	44.8	13.9	18.3	5.0	2.2	16.7	7.8	2.2	21.7
Foster home/foster care	100.0	56.6	14.6	21.9	5.4	1.5	12.2	8.3	2.1	20.0
Residential care	100.0	34.3	18.5	30.5	11.7	5.2	27.7	15.0	3.1	31.2
Private residence	100.0	29.9	21.9	32.1	6.1	1.8	20.5	11.3	4.1	21.7
Other	100.0	30.6	21.3	21.3	9.4	2.8	28.0	9.2	1.9	25.4
Total	100.0	31.0	21.5	31.4	6.2	1.9	20.5	11.2	4.0	21.9
SED status²										
SED	100.0	28.7	22.9	29.4	5.8	2.1	22.2	10.7	3.7	20.2
Not SED	100.0	34.4	11.3	29.4	5.5	†	8.6	7.9	5.4	17.6
Total	100.0	30.3	19.7	29.4	5.7	1.7	18.4	9.9	4.1	19.5

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; ODD = oppositional defiant disorder; PDD = pervasive developmental disorders; SED = serious emotional disturbance.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SED status can be found in Appendix D.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 2.6a. Individuals aged 17 years and younger with SED, by mental health diagnosis and living arrangement: Number, 2020

Mental health diagnosis ¹										
Living arrangement	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Conduct disorder	Mood disorders		ODD	PDD	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Total individuals with SED aged 0 to 17 years ²	1,155,930	332,184	264,993	339,561	66,916	23,982	256,237	123,208	42,263	233,570
Living arrangement										
Homeless	3,550	1,613	511	646	187	78	604	291	62	785
Foster home/foster care	30,216	16,992	4,783	6,946	1,596	489	3,956	2,597	595	6,214
Residential care	11,030	3,706	2,092	3,477	1,307	608	3,202	1,666	318	3,476
Private residence	660,036	192,126	151,948	223,946	41,126	13,782	146,046	79,868	23,067	149,810
Other	18,704	5,647	4,086	3,911	1,794	541	5,429	1,668	331	4,757
Total	723,536	220,084	163,420	238,926	46,010	15,498	159,237	86,090	24,373	165,042

ADHD = attention deficit/hyperactivity disorder; ODD = oppositional defiant disorder; PDD = pervasive developmental disorders; SED = serious emotional disturbance.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SED status can be found in Appendix D.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 2.6b. Individuals aged 17 years and younger with SED, by mental health diagnosis and living arrangement: Column percent distribution, 2020

Living arrangement	Total	Mental health diagnosis ¹								
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Conduct disorder	Mood disorders		ODD	PDD	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Total individuals with SED aged 0 to 17 years²	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Living arrangement										
Homeless	†	†	†	†	†	†	†	†	†	†
Foster home/foster care	4.2	7.7	2.9	2.9	3.5	3.2	2.5	3.0	2.4	3.8
Residential care	1.5	1.7	1.3	1.5	2.8	3.9	2.0	1.9	1.3	2.1
Private residence	91.2	87.3	93.0	93.7	89.4	88.9	91.7	92.8	94.6	90.8
Other	2.6	2.6	2.5	1.6	3.9	3.5	3.4	1.9	1.4	2.9
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; ODD = oppositional defiant disorder; PDD = pervasive developmental disorders; SED = serious emotional disturbance.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SED status can be found in Appendix D.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 2.6c. Individuals aged 17 years and younger with SED, by mental health diagnosis and living arrangement: Row percent distribution, 2020

Mental health diagnosis ¹										
Living arrangement		Trauma- and stressor-related disorders				Mood disorders		Other mental disorders ¹		
			Anxiety disorders	ADHD	Conduct disorder	Bipolar disorders	Depressive disorders			
	Total							ODD	PDD	
Total individuals with SED aged 0 to 17 years ²	100.0	28.7	22.9	29.4	5.8	2.1	22.2	10.7	3.7	20.2
Living arrangement										
Homeless	100.0	45.4	14.4	18.2	5.3	2.2	17.0	8.2	1.7	22.1
Foster home/foster care	100.0	56.2	15.8	23.0	5.3	1.6	13.1	8.6	2.0	20.6
Residential care	100.0	33.6	19.0	31.5	11.8	5.5	29.0	15.1	2.9	31.5
Private residence	100.0	29.1	23.0	33.9	6.2	2.1	22.1	12.1	3.5	22.7
Other	100.0	30.2	21.8	20.9	9.6	2.9	29.0	8.9	1.8	25.4
Total	100.0	30.4	22.6	33.0	6.4	2.1	22.0	11.9	3.4	22.8

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; ODD = oppositional defiant disorder; PDD = pervasive developmental disorders; SED = serious emotional disturbance.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SED status can be found in Appendix D.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 2.7a. Individuals aged 17 years and younger, by mental health diagnosis, timing of admission, and mental health service setting: Number, 2020

Timing of admission and mental health service setting	Total	Mental health diagnosis ¹								
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Conduct disorder	Mood disorders		ODD	PDD	Other mental disorders ¹
						Bipolar disorder	Depressive disorder			
Total individuals aged 0 to 17 years	1,637,967	496,671	322,311	481,629	93,111	27,577	301,271	162,854	69,912	319,355
Timing of admission										
New individuals	785,139	265,416	154,633	181,785	42,661	9,725	157,969	65,513	22,579	139,706
Continuing individuals	852,828	231,255	167,678	299,844	50,450	17,852	143,302	97,341	47,333	179,649
Mental health service setting^{2,3}										
State psychiatric hospital	8,516	2,155	1,432	2,918	809	651	2,835	1,231	352	3,148
Community-based program	1,624,251	493,455	320,859	478,928	91,731	26,957	296,531	161,385	69,630	315,088
Residential treatment center	12,653	3,569	1,865	3,334	1,304	901	3,836	2,062	343	4,498
Other psychiatric inpatient	30,078	6,818	6,610	6,599	1,574	1,620	12,851	2,814	426	9,732
Institutions under the justice system	3,787	1,043	443	593	989	136	767	488	27	1,436

ADHD = attention deficit/hyperactivity disorder; ODD = oppositional defiant disorder; PDD = pervasive developmental disorders.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Individuals could receive services in more than one service setting during the reporting period.

³ Mental health service setting category definitions can be found in Appendix C.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 2.7b. Individuals aged 17 years and younger, by mental health diagnosis, timing of admission, and mental health service setting: Column percent, 2020

	Mental health diagnosis ¹									
Timing of admission and mental health service setting	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Conduct disorder	Mood disorders		ODD	PDD	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Total individuals aged 0 to 17 years	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Timing of admission										
New individuals	47.9	53.4	48.0	37.7	45.8	35.3	52.4	40.2	32.3	43.7
Continuing individuals	52.1	46.6	52.0	62.3	54.2	64.7	47.6	59.8	67.7	56.3
Mental health service setting ^{2,3}										
State psychiatric hospital	†	†	†	†	†	2.4	†	†	†	†
Community-based program	99.2	99.4	99.5	99.4	98.5	97.8	98.4	99.1	99.6	98.7
Residential treatment center	†	†	†	†	1.4	3.3	1.3	1.3	†	1.4
Other psychiatric inpatient	1.8	1.4	2.1	1.4	1.7	5.9	4.3	1.7	†	3.0
Institutions under the justice system	†	†	†	†	1.1	†	†	†	†	†

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; ODD = oppositional defiant disorder; PDD = pervasive developmental disorders.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Percentages may not sum to 100 percent, because individuals could receive services in more than one service setting during the reporting period.

³ Mental health service setting category definitions can be found in Appendix C.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 2.7c. Individuals aged 17 years and younger, by mental health diagnosis, timing of admission, and mental health service setting: Row percent distribution, 2020

Timing of admission and mental health service setting	Total	Mental health diagnosis ¹								
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Conduct disorder	Mood disorders		ODD	PDD	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Total individuals aged 0 to 17 years	100.0	30.3	19.7	29.4	5.7	1.7	18.4	9.9	4.3	19.5
Timing of admission										
New individuals	100.0	33.8	19.7	23.2	5.4	1.2	20.1	8.3	2.9	17.8
Continuing individuals	100.0	27.1	19.7	35.2	5.9	2.1	16.8	11.4	5.6	21.1
Mental health service setting^{2,3}										
State psychiatric hospital	100.0	25.3	16.8	34.3	9.5	7.6	33.3	14.5	4.1	37.0
Community-based program	100.0	30.4	19.8	29.5	5.6	1.7	18.3	9.9	4.3	19.4
Residential treatment center	100.0	28.2	14.7	26.3	10.3	7.1	30.3	16.3	2.7	35.5
Other psychiatric inpatient	100.0	22.7	22.0	21.9	5.2	5.4	42.7	9.4	1.4	32.4
Institutions under the justice system	100.0	27.5	11.7	15.7	26.1	3.6	20.3	12.9	†	37.9

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; ODD = oppositional defiant disorder; PDD = pervasive developmental disorders.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Individuals could receive services in more than one service setting during the reporting period.

³ Mental health service setting category definitions can be found in Appendix C.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 2.8a. Individuals aged 17 years and younger with SED, by mental health diagnosis, timing of admission, and mental health service setting: Number, 2020

		Mental health diagnosis ¹								
Timing of admission and mental health service setting	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Conduct disorder	Mood disorders		ODD	PDD	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Total individuals with SED aged 0 to 17 years ²	1,155,930	332,184	264,993	339,561	66,916	23,982	256,237	123,208	42,263	233,570
Timing of admission										
New individuals	552,859	176,117	124,833	130,519	30,015	8,362	132,922	49,070	14,146	98,336
Continuing individuals	603,071	156,067	140,160	209,042	36,901	15,620	123,315	74,138	28,117	135,234
Mental health service setting ^{3,4}										
State psychiatric hospital	6,653	1,733	1,256	2,097	675	610	2,571	846	277	2,602
Community-based program	1,146,791	330,024	263,821	338,068	65,840	23,485	252,455	122,419	42,083	230,751
Residential treatment center	9,142	2,620	1,545	2,276	958	765	3,396	1,410	276	3,131
Other psychiatric inpatient	21,918	4,757	6,178	4,794	1,168	1,476	11,604	1,937	358	6,844
Institutions under the justice system	3,652	996	430	574	965	133	746	474	26	1,397

ADHD = attention deficit/hyperactivity disorder; ODD = oppositional defiant disorder; PDD = pervasive developmental disorders; SED = serious emotional disturbance.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SED status can be found in Appendix D.

³ Individuals could receive services in more than one service setting during the reporting period.

⁴ Mental health service setting category definitions can be found in Appendix C.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 2.8b. Individuals aged 17 years and younger with SED, by mental health diagnosis, timing of admission, and mental health service setting: Column percent, 2020

Timing of admission and mental health service setting	Mental health diagnosis ¹									
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Conduct disorder	Mood disorders		ODD	PDD	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Total individuals with SED aged 0 to 17 years²	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Timing of admission										
New individuals	47.8	53.0	47.1	38.4	44.9	34.9	51.9	39.8	33.5	42.1
Continuing individuals	52.2	47.0	52.9	61.6	55.1	65.1	48.1	60.2	66.5	57.9
Mental health service setting^{3,4}										
State psychiatric hospital	†	†	†	†	1.0	2.5	1.0	†	†	1.1
Community-based program	99.2	99.3	99.6	99.6	98.4	97.9	98.5	99.4	99.6	98.8
Residential treatment center	†	†	†	†	1.4	3.2	1.3	1.1	†	1.3
Other psychiatric inpatient	1.9	1.4	2.3	1.4	1.7	6.2	4.5	1.6	†	2.9
Institutions under the justice system	†	†	†	†	1.4	†	†	†	†	†

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; ODD = oppositional defiant disorder; PDD = pervasive developmental disorders; SED = serious emotional disturbance.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SED status can be found in Appendix D.

³ Percentages may not sum to 100 percent, because individuals could receive services in more than one service setting during the reporting period.

⁴ Mental health service setting category definitions can be found in Appendix C.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 2.8c. Individuals aged 17 years and younger with SED, by mental health diagnosis, timing of admission, and mental health service setting: Row percent distribution, 2020

	Mental health diagnosis ¹									
Timing of admission and mental health service setting	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Conduct disorder	Mood disorders		ODD	PDD	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Total individuals with SED aged 0 to 17 years ²	100.0	28.7	22.9	29.4	5.8	2.1	22.2	10.7	3.7	20.2
Timing of admission										
New individuals	100.0	31.9	22.6	23.6	5.4	1.5	24.0	8.9	2.6	17.8
Continuing individuals	100.0	25.9	23.2	34.7	6.1	2.6	20.4	12.3	4.7	22.4
Mental health service setting ^{3,4}										
State psychiatric hospital	100.0	26.0	18.9	31.5	10.1	9.2	38.6	12.7	4.2	39.1
Community-based program	100.0	28.8	23.0	29.5	5.7	2.0	22.0	10.7	3.7	20.1
Residential treatment center	100.0	28.7	16.9	24.9	10.5	8.4	37.1	15.4	3.0	34.2
Other psychiatric inpatient	100.0	21.7	28.2	21.9	5.3	6.7	52.9	8.8	1.6	31.2
Institutions under the justice system	100.0	27.3	11.8	15.7	26.4	3.6	20.4	13.0	†	38.3

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; ODD = oppositional defiant disorder; PDD = pervasive developmental disorders; SED = serious emotional disturbance.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SED status can be found in Appendix D.

³ Individuals could receive services in more than one service setting during the reporting period.

⁴ Mental health service setting category definitions can be found in Appendix C.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 2.9. Individuals aged 17 years and younger, by mental health service setting and SED status: Number and percent distribution, 2020

<i>Mental health service setting</i>	<i>SED status¹</i>								
	<i>Number of individuals</i>			<i>Column percent of individuals</i>			<i>Row percent distribution of individuals</i>		
	<i>Total</i>	<i>SED</i>	<i>Not SED</i>	<i>Total</i>	<i>SED</i>	<i>Not SED</i>	<i>Total</i>	<i>SED</i>	<i>Not SED</i>
Total individuals aged 0 to 17 years	1,844,026	1,333,084	510,942	100.0	100.0	100.0	100.0	72.3	27.7
Mental health service setting^{2,3}									
State psychiatric hospital	7,621	6,867	754	†	†	†	100.0	90.1	9.9
Community-based program	1,829,397	1,322,087	507,310	99.2	99.2	99.3	100.0	72.3	27.7
Residential treatment center	14,049	10,882	3,167	†	†	†	100.0	77.5	22.5
Other psychiatric inpatient	32,141	23,901	8,240	1.7	1.8	1.6	100.0	74.4	25.6
Institutions under the justice system	4,537	4,401	136	†	†	†	100.0	97.0	3.0

† Less than 1 percent of individuals.

SED = serious emotional disturbance.

¹ The federal definition for SED status can be found in Appendix D.

² Individuals could receive services in more than one mental health service setting during the reporting period.

³ Mental health service setting category definitions can be found in Appendix C.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 3.1a. Individuals aged 18 to 24 years, by gender, age, race, ethnicity, education, veteran status, and SMI status: Number, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Number of states reporting	45	47	49	49	48	50
Total individuals aged 18 to 24 years	521,586	573,628	626,984	653,224	676,862	724,047
Gender						
Male	260,665	278,684	306,378	312,838	315,325	335,403
Female	259,516	292,236	318,064	337,980	358,439	385,360
Total	520,181	570,920	624,442	650,818	673,764	720,763
Age¹						
18 to 20 years	225,984	250,346	273,807	293,478	310,263	325,599
21 to 24 years	295,602	323,282	353,177	359,746	366,599	398,448
Total	521,586	573,628	626,984	653,224	676,862	724,047
Race						
American Indian or Alaska Native	7,738	8,040	8,393	9,597	11,229	14,620
Asian ²	6,133	7,254	8,837	9,791	9,852	10,093
Black or African American	104,945	113,116	122,178	123,832	128,542	126,633
Native Hawaiian or Other Pacific Islander	1,278	1,468	1,305	1,593	1,652	1,808
White	306,198	339,619	365,694	380,801	388,124	417,017
Other	48,722	55,325	65,890	73,205	77,460	84,694
Total	475,014	524,822	572,297	598,819	616,859	654,865
Ethnicity						
Not of Hispanic or Latino origin	351,170	380,016	462,172	468,468	478,334	505,877
Hispanic or Latino origin	76,970	86,798	100,674	106,585	114,236	144,063
Total	428,140	466,814	562,846	575,053	592,570	649,940
Specific Hispanic or Latino origin						
Mexican	6,360	5,300	6,613	6,757	7,506	7,374
Puerto Rican	4,568	4,021	4,799	2,774	2,636	2,908
Cuban	620	609	647	685	756	897
Other specific	6,038	6,090	7,224	7,261	6,132	6,904
Total	17,586	16,020	19,283	17,477	17,030	18,083

Continued. See notes at end of table.

Table 3.1a. Individuals aged 18 to 24 years, by gender, age, race, ethnicity, education, veteran status, and SMI status: Number, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Highest school grade completed						
0 to 8	31,779	30,602	37,556	34,973	38,674	48,513
9 to 11	66,368	64,958	75,967	80,324	79,309	86,618
12 (or GED)	97,517	99,967	114,635	134,377	134,444	146,474
More than 12	26,264	27,467	34,241	40,187	41,051	46,041
Total	221,928	222,994	262,399	289,861	293,478	327,646
Veteran status						
Veteran	<i>n/c</i>	<i>n/c</i>	1,798	1,979	6,248	9,671
Not a veteran	<i>n/c</i>	<i>n/c</i>	164,762	238,905	260,248	279,194
Total	<i>n/c</i>	<i>n/c</i>	166,560	240,884	266,496	288,865
SMI status³						
SMI	310,841	344,604	379,997	418,175	454,966	470,276
Not SMI	163,004	177,510	194,318	183,453	165,334	198,709
Total	473,845	522,114	574,315	601,628	620,300	668,985

n/c Not collected.

SMI = serious mental illness; GED = general equivalency diploma.

¹ Age is based on the individual's age at the midpoint of the reporting period.

² For some states, "Asian" includes "Pacific Islanders." See Appendix C.

³ The federal definition for SMI status can be found in Appendix D.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 3.1b. Individuals aged 18 to 24 years, by gender, age, race, ethnicity, education, veteran status, and SMI status: Column percent distribution, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Number of states reporting	45	47	49	49	48	50
Total individuals aged 18 to 24 years	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	50.1	48.8	49.1	48.1	46.8	46.5
Female	49.9	51.2	50.9	51.9	53.2	53.5
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age¹						
18 to 20 years	43.3	43.6	43.7	44.9	45.8	45.0
21 to 24 years	56.7	56.4	56.3	55.1	54.2	55.0
Total	100.0	100.0	100.0	100.0	100.0	100.0
Race						
American Indian or Alaska Native	1.6	1.5	1.5	1.6	1.8	2.2
Asian ²	1.3	1.4	1.5	1.6	1.6	1.5
Black or African American	22.1	21.6	21.3	20.7	20.8	19.3
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	64.5	64.7	63.9	63.6	62.9	63.7
Other	10.3	10.5	11.5	12.2	12.6	12.9
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	82.0	81.4	82.1	81.5	80.7	77.8
Hispanic or Latino origin	18.0	18.6	17.9	18.5	19.3	22.2
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	36.2	33.1	34.3	38.7	44.1	40.8
Puerto Rican	26.0	25.1	24.9	15.9	15.5	16.1
Cuban	3.5	3.8	3.4	3.9	4.4	5.0
Other specific	34.3	38.0	37.5	41.5	36.0	38.2
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 3.1b. Individuals aged 18 to 24 years, by gender, age, race, ethnicity, education, veteran status, and SMI status: Column percent distribution, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Highest school grade completed						
0 to 8	14.3	13.7	14.3	12.1	13.2	14.8
9 to 11	29.9	29.1	29.0	27.7	27.0	26.4
12 (or GED)	43.9	44.8	43.7	46.4	45.8	44.7
More than 12	11.8	12.3	13.0	13.9	14.0	14.1
Total	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status						
Veteran	<i>n/c</i>	<i>n/c</i>	1.1	†	2.3	3.3
Not a veteran	<i>n/c</i>	<i>n/c</i>	98.9	99.2	97.7	96.7
Total	<i>n/c</i>	<i>n/c</i>	100.0	100.0	100.0	100.0
SMI status³						
SMI	65.6	66.0	66.2	69.5	73.3	70.3
Not SMI	34.4	34.0	33.8	30.5	26.7	29.7
Total	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals; *n/c* Not collected.

SMI = serious mental illness; GED = general equivalency diploma.

¹ Age is based on the individual's age at the midpoint of the reporting period.

² For some states, "Asian" includes "Pacific Islanders." See Appendix C.

³ The federal definition for SMI status can be found in Appendix D.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

**Table 3.2a. Individuals aged 18 to 24 years with SMI, by gender, age, race, ethnicity, education, and veteran status:
Number, 2015–2020**

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Number of states reporting	45	46	47	48	47	49
Total individuals with SMI aged 18 to 24 years¹	310,841	344,604	379,997	418,175	454,966	470,276
Gender						
Male	155,763	171,211	186,069	198,163	206,936	212,379
Female	154,340	172,271	192,878	219,049	246,544	256,744
Total	310,103	343,482	378,947	417,212	453,480	469,123
Age²						
18 to 20 years	126,211	145,956	161,181	183,931	200,657	205,035
21 to 24 years	184,630	198,648	218,816	234,244	254,309	265,241
Total	310,841	344,604	379,997	418,175	454,966	470,276
Race						
American Indian or Alaska Native	4,060	4,272	4,953	5,767	7,511	8,730
Asian ³	3,941	4,482	5,872	6,525	6,939	6,932
Black or African American	65,754	69,020	73,294	79,396	83,278	82,192
Native Hawaiian or Other Pacific Islander	862	995	910	1,112	1,208	1,234
White	176,765	203,084	218,145	238,466	260,103	269,821
Other	33,351	36,580	47,761	54,586	59,114	63,153
Total	284,733	318,433	350,935	385,852	418,153	432,062
Ethnicity						
Not of Hispanic or Latino origin	206,004	227,338	266,603	286,709	304,437	326,156
Hispanic or Latino origin	51,859	58,757	71,454	77,541	88,579	100,840
Total	257,863	286,095	338,057	364,250	393,016	426,996
Specific Hispanic or Latino origin						
Mexican	3,763	3,137	4,185	4,454	5,110	4,898
Puerto Rican	2,102	2,164	2,417	1,964	1,907	2,138
Cuban	425	524	559	622	623	761
Other specific	3,799	4,191	4,964	5,376	4,058	4,631
Total	10,089	10,016	12,125	12,416	11,698	12,428

Continued. See notes at end of table.

**Table 3.2a. Individuals aged 18 to 24 years with SMI, by gender, age, race, ethnicity, education, and veteran status:
Number, 2015–2020 (continued)**

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Highest school grade completed						
0 to 8	14,105	14,053	17,880	18,510	20,938	26,772
9 to 11	40,044	42,457	49,824	56,584	55,114	58,124
12 (or GED)	63,950	69,164	79,920	96,480	97,114	104,975
More than 12	17,281	19,159	23,883	28,533	29,835	32,282
Total	135,380	144,833	171,507	200,107	203,001	222,153
Veteran status						
Veteran	<i>n/c</i>	<i>n/c</i>	1,325	1,429	5,073	7,746
Not a veteran	<i>n/c</i>	<i>n/c</i>	83,526	149,371	166,533	168,916
Total	<i>n/c</i>	<i>n/c</i>	84,851	150,800	171,606	176,662

n/c Not collected.

SMI = serious mental illness; GED = general equivalency diploma.

¹ The federal definition for SMI status can be found in Appendix D.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

**Table 3.2b. Individuals aged 18 to 24 years with SMI, by gender, age, race, ethnicity, education, and veteran status:
Column percent distribution, 2015–2020**

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Number of states reporting	45	46	47	48	47	49
Total individuals with SMI aged 18 to 24 years¹	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	50.2	49.8	49.1	47.5	45.6	45.3
Female	49.8	50.2	50.9	52.5	54.4	54.7
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age²						
18 to 20 years	40.6	42.4	42.4	44.0	44.1	43.6
21 to 24 years	59.4	57.6	57.6	56.0	55.9	56.4
Total	100.0	100.0	100.0	100.0	100.0	100.0
Race						
American Indian or Alaska Native	1.4	1.3	1.4	1.5	1.8	2.0
Asian ³	1.4	1.4	1.7	1.7	1.7	1.6
Black or African American	23.1	21.7	20.9	20.6	19.9	19.0
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	62.1	63.8	62.2	61.8	62.2	62.4
Other	11.7	11.5	13.6	14.1	14.1	14.6
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	79.9	79.5	78.9	78.7	77.5	76.4
Hispanic or Latino origin	20.1	20.5	21.1	21.3	22.5	23.6
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	37.3	31.3	34.5	35.9	43.7	39.4
Puerto Rican	20.8	21.6	19.9	15.8	16.3	17.2
Cuban	4.2	5.2	4.6	5.0	5.3	6.1
Other specific	37.7	41.8	40.9	43.3	34.7	37.3
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

**Table 3.2b. Individuals aged 18 to 24 years with SMI, by gender, age, race, ethnicity, education, and veteran status:
Column percent distribution, 2015–2020 (continued)**

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Highest school grade completed						
0 to 8	10.4	9.7	10.4	9.3	10.3	12.1
9 to 11	29.6	29.3	29.1	28.3	27.1	26.2
12 (or GED)	47.2	47.8	46.6	48.2	47.8	47.3
More than 12	12.8	13.2	13.9	14.3	14.7	14.5
Total	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status						
Veteran	<i>n/c</i>	<i>n/c</i>	1.6	†	3.0	4.4
Not a veteran	<i>n/c</i>	<i>n/c</i>	98.4	99.1	97.0	95.6
Total	<i>n/c</i>	<i>n/c</i>	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals; *n/c* Not collected.

SMI = serious mental illness; GED = general equivalency diploma.

¹ The federal definition for SMI status can be found in Appendix D.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 3.3a. Individuals aged 18 to 24 years, by mental health diagnosis, gender, race, and ethnicity: Number, 2020

Gender, race, and ethnicity	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals aged 18 to 24 years	608,665	124,460	172,077	59,814	73,317	223,326	18,210	57,417	160,779
Gender									
Male	278,815	44,783	61,945	38,472	31,213	82,129	6,016	40,577	91,641
Female	327,743	79,136	109,638	21,170	41,754	140,509	12,112	16,653	68,662
Total	606,558	123,919	171,583	59,642	72,967	222,638	18,128	57,230	160,303
Race									
American Indian or Alaska Native	12,936	3,143	3,713	811	922	4,786	282	979	4,098
Asian ²	8,370	1,548	2,194	598	715	3,152	194	1,163	2,148
Black or African American	106,453	21,364	18,127	11,691	13,633	34,553	2,935	18,340	28,074
Native Hawaiian or Other Pacific Islander	1,495	332	439	98	149	600	33	175	376
White	352,582	72,984	114,632	36,449	46,463	136,033	11,959	23,270	88,939
Other	73,665	14,812	18,637	5,671	7,273	27,392	1,691	8,807	19,615
Total	555,501	114,183	157,742	55,318	69,155	206,516	17,094	52,734	143,250
Ethnicity									
Not of Hispanic or Latino origin	427,275	89,820	124,227	46,530	56,181	159,743	14,315	39,483	107,579
Hispanic or Latino origin	123,074	23,958	33,738	8,744	11,807	44,528	2,313	12,102	34,597
Total	550,349	113,778	157,965	55,274	67,988	204,271	16,628	51,585	142,176
Specific Hispanic or Latino origin									
Mexican	7,066	1,854	2,052	439	613	2,953	177	641	1,507
Puerto Rican	2,571	587	470	247	350	906	60	324	552
Cuban	863	122	158	59	97	242	14	96	222
Other specific	6,471	1,498	1,749	521	746	2,643	183	751	1,346
Total	16,971	4,061	4,429	1,266	1,806	6,744	434	1,812	3,627

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 3.3b. Individuals aged 18 to 24 years, by mental health diagnosis, gender, race, and ethnicity: Column percent distribution, 2020

Gender, race, and ethnicity	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals aged 18 to 24 years	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender									
Male	46.0	36.1	36.1	64.5	42.8	36.9	33.2	70.9	57.2
Female	54.0	63.9	63.9	35.5	57.2	63.1	66.8	29.1	42.8
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race									
American Indian or Alaska Native	2.3	2.8	2.4	1.5	1.3	2.3	1.6	1.9	2.9
Asian ²	1.5	1.4	1.4	1.1	1.0	1.5	1.1	2.2	1.5
Black or African American	19.2	18.7	11.5	21.1	19.7	16.7	17.2	34.8	19.6
Native Hawaiian or Other Pacific Islander	T	T	T	T	T	T	T	T	T
White	63.5	63.9	72.7	65.9	67.2	65.9	70.0	44.1	62.1
Other	13.3	13.0	11.8	10.3	10.5	13.3	9.9	16.7	13.7
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity									
Not of Hispanic or Latino origin	77.6	78.9	78.6	84.2	82.6	78.2	86.1	76.5	75.7
Hispanic or Latino origin	22.4	21.1	21.4	15.8	17.4	21.8	13.9	23.5	24.3
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	41.6	45.7	46.3	34.7	33.9	43.8	40.8	35.4	41.5
Puerto Rican	15.1	14.5	10.6	19.5	19.4	13.4	13.8	17.9	15.2
Cuban	5.1	3.0	3.6	4.7	5.4	3.6	3.2	5.3	6.1
Other specific	38.1	36.9	39.5	41.2	41.3	39.2	42.2	41.4	37.1
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 3.3c. Individuals aged 18 to 24 years, by mental health diagnosis, gender, race, and ethnicity: Row percent distribution, 2020

Gender, race, and ethnicity	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals aged 18 to 24 years	100.0	20.4	28.3	9.8	12.0	36.7	3.0	9.4	26.4
Gender									
Male	100.0	16.1	22.2	13.8	11.2	29.5	2.2	14.6	32.9
Female	100.0	24.1	33.5	6.5	12.7	42.9	3.7	5.1	20.9
Total	100.0	20.4	28.3	9.8	12.0	36.7	3.0	9.4	26.4
Race									
American Indian or Alaska Native	100.0	24.3	28.7	6.3	7.1	37.0	2.2	7.6	31.7
Asian ²	100.0	18.5	26.2	7.1	8.5	37.7	2.3	13.9	25.7
Black or African American	100.0	20.1	17.0	11.0	12.8	32.5	2.8	17.2	26.4
Native Hawaiian or Other Pacific Islander	100.0	22.2	29.4	6.6	10.0	40.1	2.2	11.7	25.2
White	100.0	20.7	32.5	10.3	13.2	38.6	3.4	6.6	25.2
Other	100.0	20.1	25.3	7.7	9.9	37.2	2.3	12.0	26.6
Total	100.0	20.6	28.4	10.0	12.4	37.2	3.1	9.5	25.8
Ethnicity									
Not of Hispanic or Latino origin	100.0	21.0	29.1	10.9	13.1	37.4	3.4	9.2	25.2
Hispanic or Latino origin	100.0	19.5	27.4	7.1	9.6	36.2	1.9	9.8	28.1
Total	100.0	20.7	28.7	10.0	12.4	37.1	3.0	9.4	25.8
Specific Hispanic or Latino origin									
Mexican	100.0	26.2	29.0	6.2	8.7	41.8	2.5	9.1	21.3
Puerto Rican	100.0	22.8	18.3	9.6	13.6	35.2	2.3	12.6	21.5
Cuban	100.0	14.1	18.3	6.8	11.2	28.0	1.6	11.1	25.7
Other specific	100.0	23.1	27.0	8.1	11.5	40.8	2.8	11.6	20.8
Total	100.0	23.9	26.1	7.5	10.6	39.7	2.6	10.7	21.4

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 3.4a. Individuals aged 18 to 24 years with SMI, by mental health diagnosis, gender, race, and ethnicity: Number, 2020

Gender, race, and ethnicity	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals with SMI aged 18 to 24 years²	414,898	76,426	123,667	34,255	61,291	177,560	14,075	50,920	91,152
Gender									
Male	186,163	25,792	44,007	21,571	25,985	65,262	4,535	36,039	50,918
Female	227,691	50,421	79,420	12,618	35,049	111,866	9,492	14,742	40,026
Total	413,854	76,213	123,427	34,189	61,034	177,128	14,027	50,781	90,944
Race									
American Indian or Alaska Native	8,026	1,964	2,540	460	758	3,701	204	833	1,945
Asian ³	6,139	974	1,666	315	618	2,615	146	1,078	1,339
Black or African American	74,179	12,281	13,280	6,866	11,467	27,964	2,247	16,105	16,260
Native Hawaiian or Other Pacific Islander	1,153	246	364	67	128	510	28	163	250
White	236,016	45,178	81,529	20,949	38,722	106,616	9,355	20,443	49,080
Other	58,443	10,761	15,220	3,420	6,307	23,736	1,416	8,298	14,042
Total	383,956	71,404	114,599	32,077	58,000	165,142	13,396	46,920	82,916
Ethnicity									
Not of Hispanic or Latino origin	288,498	55,330	88,515	26,616	46,223	123,482	11,283	34,521	61,966
Hispanic or Latino origin	89,653	15,356	25,736	4,981	10,656	39,074	1,777	11,379	20,153
Total	378,151	70,686	114,251	31,597	56,879	162,556	13,060	45,900	82,119
Specific Hispanic or Latino origin									
Mexican	4,844	1,154	1,363	234	533	2,480	136	573	809
Puerto Rican	2,015	411	376	150	328	790	55	310	323
Cuban	736	88	136	40	93	214	12	92	179
Other specific	4,539	951	1,202	282	626	2,130	145	673	766
Total	12,134	2,604	3,077	706	1,580	5,614	348	1,648	2,077

ADHD = attention deficit/hyperactivity disorder; SMI = serious mental illness.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 3.4b. Individuals aged 18 to 24 years with SMI, by mental health diagnosis, gender, race, and ethnicity: Column percent distribution, 2020

Gender, race, and ethnicity	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals with SMI aged 18 to 24 years²	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender									
Male	45.0	33.8	35.7	63.1	42.6	36.8	32.3	71.0	56.0
Female	55.0	66.2	64.3	36.9	57.4	63.2	67.7	29.0	44.0
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race									
American Indian or Alaska Native	2.1	2.8	2.2	1.4	1.3	2.2	1.5	1.8	2.3
Asian ³	1.6	1.4	1.5	†	1.1	1.6	1.1	2.3	1.6
Black or African American	19.3	17.2	11.6	21.4	19.8	16.9	16.8	34.3	19.6
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	†	†	†
White	61.5	63.3	71.1	65.3	66.8	64.6	69.8	43.6	59.2
Other	15.2	15.1	13.3	10.7	10.9	14.4	10.6	17.7	16.9
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity									
Not of Hispanic or Latino origin	76.3	78.3	77.5	84.2	81.3	76.0	86.4	75.2	75.5
Hispanic or Latino origin	23.7	21.7	22.5	15.8	18.7	24.0	13.6	24.8	24.5
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	39.9	44.3	44.3	33.1	33.7	44.2	39.1	34.8	39.0
Puerto Rican	16.6	15.8	12.2	21.2	20.8	14.1	15.8	18.8	15.6
Cuban	6.1	3.4	4.4	5.7	5.9	3.8	3.4	5.6	8.6
Other specific	37.4	36.5	39.1	39.9	39.6	37.9	41.7	40.8	36.9
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; SMI = serious mental illness.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 3.4c. Individuals aged 18 to 24 years with SMI, by mental health diagnosis, gender, race, and ethnicity: Row percent distribution, 2020

Gender, race, and ethnicity	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals with SMI aged 18 to 24 years ²	100.0	18.4	29.8	8.3	14.8	42.8	3.4	12.3	22.0
Gender									
Male	100.0	13.9	23.6	11.6	14.0	35.1	2.4	19.4	27.4
Female	100.0	22.1	34.9	5.5	15.4	49.1	4.2	6.5	17.6
Total	100.0	18.4	29.8	8.3	14.7	42.8	3.4	12.3	22.0
Race									
American Indian or Alaska Native	100.0	24.5	31.6	5.7	9.4	46.1	2.5	10.4	24.2
Asian ³	100.0	15.9	27.1	5.1	10.1	42.6	2.4	17.6	21.8
Black or African American	100.0	16.6	17.9	9.3	15.5	37.7	3.0	21.7	21.9
Native Hawaiian or Other Pacific Islander	100.0	21.3	31.6	5.8	11.1	44.2	2.4	14.1	21.7
White	100.0	19.1	34.5	8.9	16.4	45.2	4.0	8.7	20.8
Other	100.0	18.4	26.0	5.9	10.8	40.6	2.4	14.2	24.0
Total	100.0	18.6	29.8	8.4	15.1	43.0	3.5	12.2	21.6
Ethnicity									
Not of Hispanic or Latino origin	100.0	19.2	30.7	9.2	16.0	42.8	3.9	12.0	21.5
Hispanic or Latino origin	100.0	17.1	28.7	5.6	11.9	43.6	2.0	12.7	22.5
Total	100.0	18.7	30.2	8.4	15.0	43.0	3.5	12.1	21.7
Specific Hispanic or Latino origin									
Mexican	100.0	23.8	28.1	4.8	11.0	51.2	2.8	11.8	16.7
Puerto Rican	100.0	20.4	18.7	7.4	16.3	39.2	2.7	15.4	16.0
Cuban	100.0	12.0	18.5	5.4	12.6	29.1	1.6	12.5	24.3
Other specific	100.0	21.0	26.5	6.2	13.8	46.9	3.2	14.8	16.9
Total	100.0	21.5	25.4	5.8	13.0	46.3	2.9	13.6	17.1

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; SMI = serious mental illness.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 3.5a. Individuals aged 18 to 24 years, by mental health diagnosis, education, and marital status: Number, 2020

<i>Education and marital status</i>		<i>Mental health diagnosis</i> ¹							
					<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders</i> ¹
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
	<i>Total</i>	<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>ADHD</i>					
Total individuals aged 18 to 24 years	608,665	124,460	172,077	59,814	73,317	223,326	18,210	57,417	160,779
Highest school grade completed									
0 to 8	43,491	10,498	12,744	8,730	3,846	12,551	778	2,118	15,428
9 to 11	78,978	18,876	22,016	10,718	10,613	30,624	2,944	8,323	22,306
12 (or GED)	133,105	29,759	40,029	12,739	20,236	56,037	5,530	15,298	32,436
More than 12	42,008	9,149	15,370	2,673	6,501	19,955	1,929	3,706	7,130
Total	297,582	68,282	90,159	34,860	41,196	119,167	11,181	29,445	77,300
Marital status									
Never married	306,505	60,735	76,348	33,703	40,453	109,382	9,326	30,303	83,885
Currently married	12,429	3,365	3,883	556	1,964	5,602	596	595	2,218
Separated	3,266	959	940	142	603	1,457	174	217	536
Divorced	2,860	852	912	173	526	1,365	153	280	566
Widowed	394	136	86	20	77	139	17	31	95
Total	325,454	66,047	82,169	34,594	43,623	117,945	10,266	31,426	87,300

ADHD = attention deficit/hyperactivity disorder; GED = general equivalency diploma.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 3.5b. Individuals aged 18 to 24 years, by mental health diagnosis, education, and marital status: Column percent distribution, 2020

<i>Education and marital status</i>		<i>Mental health diagnosis</i> ¹							
		<i>Trauma- and stressor-related disorders</i> <i>Anxiety disorders</i> <i>ADHD</i>			<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders</i> ¹
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
	<i>Total</i>								
Total individuals aged 18 to 24 years	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Highest school grade completed									
0 to 8	14.6	15.4	14.1	25.0	9.3	10.5	7.0	7.2	20.0
9 to 11	26.5	27.6	24.4	30.7	25.8	25.7	26.3	28.3	28.9
12 (or GED)	44.7	43.6	44.4	36.5	49.1	47.0	49.5	52.0	42.0
More than 12	14.1	13.4	17.0	7.7	15.8	16.7	17.3	12.6	9.2
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Marital status									
Never married	94.2	92.0	92.9	97.4	92.7	92.7	90.8	96.4	96.1
Currently married	3.8	5.1	4.7	1.6	4.5	4.7	5.8	1.9	2.5
Separated	1.0	1.5	1.1	†	1.4	1.2	1.7	†	†
Divorced	†	1.3	1.1	†	1.2	1.2	1.5	†	†
Widowed	†	†	†	†	†	†	†	†	†
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; GED = general equivalency diploma.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total.

"Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 3.5c. Individuals aged 18 to 24 years, by mental health diagnosis, education, and marital status: Row percent distribution, 2020

<i>Education and marital status</i>		<i>Mental health diagnosis¹</i>							
					<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
	<i>Total</i>	<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>ADHD</i>					
Total individuals aged 18 to 24 years	100.0	20.4	28.3	9.8	12.0	36.7	3.0	9.4	26.4
Highest school grade completed									
0 to 8	100.0	24.1	29.3	20.1	8.8	28.9	1.8	4.9	35.5
9 to 11	100.0	23.9	27.9	13.6	13.4	38.8	3.7	10.5	28.2
12 (or GED)	100.0	22.4	30.1	9.6	15.2	42.1	4.2	11.5	24.4
More than 12	100.0	21.8	36.6	6.4	15.5	47.5	4.6	8.8	17.0
Total	100.0	22.9	30.3	11.7	13.8	40.0	3.8	9.9	26.0
Marital status									
Never married	100.0	19.8	24.9	11.0	13.2	35.7	3.0	9.9	27.4
Currently married	100.0	27.1	31.2	4.5	15.8	45.1	4.8	4.8	17.8
Separated	100.0	29.4	28.8	4.3	18.5	44.6	5.3	6.6	16.4
Divorced	100.0	29.8	31.9	6.0	18.4	47.7	5.3	9.8	19.8
Widowed	100.0	34.5	21.8	5.1	19.5	35.3	4.3	7.9	24.1
Total	100.0	20.3	25.2	10.6	13.4	36.2	3.2	9.7	26.8

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; GED = general equivalency diploma.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 3.6a. Individuals aged 18 to 24 years with SMI, by mental health diagnosis, education, and marital status: Number, 2020

Education and marital status	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals with SMI aged 18 to 24 years²	414,898	76,426	123,667	34,255	61,291	177,560	14,075	50,920	91,152
Highest school grade completed									
0 to 8	25,652	6,354	8,271	4,323	2,712	8,860	616	1,771	7,882
9 to 11	54,684	12,855	15,749	6,863	8,851	22,838	2,500	7,370	13,567
12 (or GED)	98,665	21,743	29,636	8,700	17,434	44,821	4,668	13,621	19,768
More than 12	30,586	6,429	10,807	1,762	5,535	15,499	1,532	3,293	4,522
Total	209,587	47,381	64,463	21,648	34,532	92,018	9,316	26,055	45,739
Marital status									
Never married	207,651	35,755	56,485	18,772	34,588	90,170	7,688	26,666	43,482
Currently married	8,952	2,286	2,917	383	1,682	4,496	493	530	1,244
Separated	2,517	684	766	108	544	1,265	146	192	306
Divorced	2,178	640	747	137	470	1,132	129	254	353
Widowed	285	103	62	13	67	113	11	29	56
Total	221,583	39,468	60,977	19,413	37,351	97,176	8,467	27,671	45,441

ADHD = attention deficit/hyperactivity disorder; GED = general equivalency diploma; SMI = serious mental illness.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 3.6b. Individuals aged 18 to 24 years with SMI, by mental health diagnosis, education, and marital status: Column percent distribution, 2020

Education and marital status	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals with SMI aged 18 to 24 years²	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Highest school grade completed									
0 to 8	12.2	13.4	12.8	20.0	7.9	9.6	6.6	6.8	17.2
9 to 11	26.1	27.1	24.4	31.7	25.6	24.8	26.8	28.3	29.7
12 (or GED)	47.1	45.9	46.0	40.2	50.5	48.7	50.1	52.3	43.2
More than 12	14.6	13.6	16.8	8.1	16.0	16.8	16.4	12.6	9.9
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Marital status									
Never married	93.7	90.6	92.6	96.7	92.6	92.8	90.8	96.4	95.7
Currently married	4.0	5.8	4.8	2.0	4.5	4.6	5.8	1.9	2.7
Separated	1.1	1.7	1.3	†	1.5	1.3	1.7	†	†
Divorced	†	1.6	1.2	†	1.3	1.2	1.5	†	†
Widowed	†	†	†	†	†	†	†	†	†
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; GED = general equivalency diploma; SMI = serious mental illness.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 3.6c. Individuals aged 18 to 24 years with SMI, by mental health diagnosis, education, marital status: Row percent distribution, 2020

Education and marital status	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals with SMI aged 18 to 24 years²	100.0	18.4	29.8	8.3	14.8	42.8	3.4	12.3	22.0
Highest school grade completed									
0 to 8	100.0	24.8	32.2	16.9	10.6	34.5	2.4	6.9	30.7
9 to 11	100.0	23.5	28.8	12.6	16.2	41.8	4.6	13.5	24.8
12 (or GED)	100.0	22.0	30.0	8.8	17.7	45.4	4.7	13.8	20.0
More than 12	100.0	21.0	35.3	5.8	18.1	50.7	5.0	10.8	14.8
Total	100.0	22.6	30.8	10.3	16.5	43.9	4.4	12.4	21.8
Marital status									
Never married	100.0	17.2	27.2	9.0	16.7	43.4	3.7	12.8	20.9
Currently married	100.0	25.5	32.6	4.3	18.8	50.2	5.5	5.9	13.9
Separated	100.0	27.2	30.4	4.3	21.6	50.3	5.8	7.6	12.2
Divorced	100.0	29.4	34.3	6.3	21.6	52.0	5.9	11.7	16.2
Widowed	100.0	36.1	21.8	4.6	23.5	39.6	3.9	10.2	19.6
Total	100.0	17.8	27.5	8.8	16.9	43.9	3.8	12.5	20.5

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; GED = general equivalency diploma; SMI = serious mental illness

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 3.7a. Individuals aged 18 to 24 years, by mental health diagnosis, living arrangement, SMI status, and veteran status: Number, 2020

Living arrangement, SMI status, and veteran status	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals aged 18 to 24 years	608,665	124,460	172,077	59,814	73,317	223,326	18,210	57,417	160,779
Living arrangement									
Homeless	12,492	2,802	2,278	593	1,938	4,278	575	2,454	3,986
Foster home/foster care	3,916	1,292	753	734	413	1,014	108	282	1,446
Residential care	7,816	1,814	1,539	939	1,273	2,089	369	1,592	3,191
Institutional setting	4,972	663	1,035	461	778	1,810	334	1,284	1,430
Jail/correctional facility	7,082	2,256	891	427	656	1,395	795	1,084	2,637
Private residence	331,720	71,376	99,404	36,547	43,206	129,864	10,810	32,015	88,333
Other	7,172	1,822	2,664	605	740	2,823	181	689	1,834
Total	375,170	82,025	108,564	40,306	49,004	143,273	13,172	39,400	102,857
SMI status²									
SMI	414,898	76,426	123,667	34,255	61,291	177,560	14,075	50,920	91,152
Not SMI	164,741	41,586	42,460	21,395	9,381	37,240	3,494	4,773	57,810
Total	579,639	118,012	166,127	55,650	70,672	214,800	17,569	55,693	148,962
Veteran status									
Veteran	8,471	1,498	1,094	354	1,035	3,468	100	920	923
Not a veteran	250,423	51,917	64,826	26,811	29,934	89,366	7,938	22,678	76,558
Total	258,894	53,415	65,920	27,165	30,969	92,834	8,038	23,598	77,481

ADHD = attention deficit/hyperactivity disorder; SMI = serious mental illness.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 3.7b. Individuals aged 18 to 24 years, by mental health diagnosis, living arrangement, SMI status, and veteran status: Column percent distribution, 2020

Living arrangement, SMI status, and veteran status	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals aged 18 to 24 years	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Living arrangement									
Homeless	3.3	3.4	2.1	1.5	4.0	3.0	4.4	6.2	3.9
Foster home/foster care	1.0	1.6	†	1.8	†	†	†	†	1.4
Residential care	2.1	2.2	1.4	2.3	2.6	1.5	2.8	4.0	3.1
Institutional setting	1.3	†	†	1.1	1.6	1.3	2.5	3.3	1.4
Jail/correctional facility	1.9	2.8	†	1.1	1.3	†	6.0	2.8	2.6
Private residence	88.4	87.0	91.6	90.7	88.2	90.6	82.1	81.3	85.9
Other	1.9	2.2	2.5	1.5	1.5	2.0	1.4	1.7	1.8
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
SMI status²									
SMI	71.6	64.8	74.4	61.6	86.7	82.7	80.1	91.4	61.2
Not SMI	28.4	35.2	25.6	38.4	13.3	17.3	19.9	8.6	38.8
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status									
Veteran	3.3	2.8	1.7	1.3	3.3	3.7	1.2	3.9	1.2
Not a veteran	96.7	97.2	98.3	98.7	96.7	96.3	98.8	96.1	98.8
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; SMI = serious mental illness.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 3.7c. Individuals aged 18 to 24 years, by mental health diagnosis, living arrangement, SMI status, and veteran status: Row percent distribution, 2020

<i>Living arrangement, SMI status, and veteran status</i>		<i>Mental health diagnosis¹</i>							
		<i>Trauma- and stressor-related disorders Anxiety disorders ADHD</i>			<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
	<i>Total</i>								
Total individuals aged 18 to 24 years	100.0	20.4	28.3	9.8	12.0	36.7	3.0	9.4	26.4
Living arrangement									
Homeless	100.0	22.4	18.2	4.7	15.5	34.2	4.6	19.6	31.9
Foster home/foster care	100.0	33.0	19.2	18.7	10.5	25.9	2.8	7.2	36.9
Residential care	100.0	23.2	19.7	12.0	16.3	26.7	4.7	20.4	40.8
Institutional setting	100.0	13.3	20.8	9.3	15.6	36.4	6.7	25.8	28.8
Jail/correctional facility	100.0	31.9	12.6	6.0	9.3	19.7	11.2	15.3	37.2
Private residence	100.0	21.5	30.0	11.0	13.0	39.1	3.3	9.7	26.6
Other	100.0	25.4	37.1	8.4	10.3	39.4	2.5	9.6	25.6
Total	100.0	21.9	28.9	10.7	13.1	38.2	3.5	10.5	27.4
SMI status²									
SMI	100.0	18.4	29.8	8.3	14.8	42.8	3.4	12.3	22.0
Not SMI	100.0	25.2	25.8	13.0	5.7	22.6	2.1	2.9	35.1
Total	100.0	20.4	28.7	9.6	12.2	37.1	3.0	9.6	25.7
Veteran status									
Veteran	100.0	17.7	12.9	4.2	12.2	40.9	1.2	10.9	10.9
Not a veteran	100.0	20.7	25.9	10.7	12.0	35.7	3.2	9.1	30.6
Total	100.0	20.6	25.5	10.5	12.0	35.9	3.1	9.1	29.9

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; SMI = serious mental illness.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 3.8a. Individuals aged 18 to 24 years with SMI, by mental health diagnosis, living arrangement and veteran status: Number, 2020

Living arrangement and veteran status	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals with SMI aged 18 to 24 years ²	414,898	76,426	123,667	34,255	61,291	177,560	14,075	50,920	91,152
Living arrangement									
Homeless	10,003	2,275	1,871	471	1,715	3,623	483	2,223	2,809
Foster home/foster care	2,746	839	606	478	318	836	97	251	965
Residential care	6,054	1,394	1,238	652	1,095	1,754	333	1,477	2,151
Institutional setting	4,145	532	920	383	689	1,602	220	1,112	1,003
Jail/correctional facility	5,018	1,464	711	309	547	1,111	740	938	1,784
Private residence	239,231	50,932	73,288	23,004	36,561	102,556	9,014	28,538	53,797
Other	5,928	1,276	2,387	428	662	2,588	167	646	1,413
Total	273,125	58,712	81,021	25,725	41,587	114,070	11,054	35,185	63,922
Veteran status									
Veteran	7,166	1,137	885	224	963	3,090	86	876	613
Not a veteran	162,149	33,152	43,115	15,487	24,060	67,227	6,030	19,416	40,124
Total	169,315	34,289	44,000	15,711	25,023	70,317	6,116	20,292	40,737

ADHD = attention deficit/hyperactivity disorder; SMI = serious mental illness.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 3.8b. Individuals aged 18 to 24 years with SMI, by mental health diagnosis, living arrangement and veteran status: Column percent distribution, 2020

		Mental health diagnosis ¹							
Living arrangement and veteran status		Trauma- and stressor-related disorders			Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
	Total	Anxiety disorders	ADHD	Bipolar disorders	Depressive disorders				
Total individuals with SMI aged 18 to 24 years ²	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Living arrangement									
Homeless	3.7	3.9	2.3	1.8	4.1	3.2	4.4	6.3	4.4
Foster home/foster care	1.0	1.4	†	1.9	†	†	†	†	1.5
Residential care	2.2	2.4	1.5	2.5	2.6	1.5	3.0	4.2	3.4
Institutional setting	1.5	†	1.1	1.5	1.7	1.4	2.0	3.2	1.6
Jail/correctional facility	1.8	2.5	†	1.2	1.3	†	6.7	2.7	2.8
Private residence	87.6	86.7	90.5	89.4	87.9	89.9	81.5	81.1	84.2
Other	2.2	2.2	2.9	1.7	1.6	2.3	1.5	1.8	2.2
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status									
Veteran	4.2	3.3	2.0	1.4	3.8	4.4	1.4	4.3	1.5
Not a veteran	95.8	96.7	98.0	98.6	96.2	95.6	98.6	95.7	98.5
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; SMI = serious mental illness.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 3.8c. Individuals aged 18 to 24 years with SMI, by mental health diagnosis, living arrangement and veteran status: Row percent distribution, 2020

Living arrangement and veteran status		Mental health diagnosis ¹									
		Trauma- and stressor-related disorders			Mood disorders		Personality disorders			Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders					
Total			Anxiety disorders	ADHD							
Total individuals with SMI aged 18 to 24 years ²		100.0	18.4	29.8	8.3	14.8	42.8	3.4	12.3	22.0	
Living arrangement											
Homeless		100.0	22.7	18.7	4.7	17.1	36.2	4.8	22.2	28.1	
Foster home/foster care		100.0	30.6	22.1	17.4	11.6	30.4	3.5	9.1	35.1	
Residential care		100.0	23.0	20.4	10.8	18.1	29.0	5.5	24.4	35.5	
Institutional setting		100.0	12.8	22.2	9.2	16.6	38.6	5.3	26.8	24.2	
Jail/correctional facility		100.0	29.2	14.2	6.2	10.9	22.1	14.7	18.7	35.6	
Private residence		100.0	21.3	30.6	9.6	15.3	42.9	3.8	11.9	22.5	
Other		100.0	21.5	40.3	7.2	11.2	43.7	2.8	10.9	23.8	
Total		100.0	21.5	29.7	9.4	15.2	41.8	4.0	12.9	23.4	
Veteran status											
Veteran		100.0	15.9	12.3	3.1	13.4	43.1	1.2	12.2	8.6	
Not a veteran		100.0	20.4	26.6	9.6	14.8	41.5	3.7	12.0	24.7	
Total		100.0	20.3	26.0	9.3	14.8	41.5	3.6	12.0	24.1	

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; SMI = serious mental illness.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 3.9a. Individuals aged 18 to 24 years, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Number, 2020

<i>Employment status and ‘detailed not in labor force’</i>		<i>Mental health diagnosis¹</i>							
		<i>Trauma- and stressor-related disorders</i>			<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
	<i>Total</i>	<i>disorders</i>	<i>disorders</i>	<i>ADHD</i>					
Total individuals aged 18 to 24 years	608,665	124,460	172,077	59,814	73,317	223,326	18,210	57,417	160,779
Employment status									
Employed	92,271	22,277	32,000	6,836	12,841	43,047	3,926	5,392	17,743
Full time	41,989	10,819	14,866	3,113	4,837	19,359	1,819	1,935	8,384
Part time	37,971	8,725	12,707	2,959	4,855	17,243	1,583	2,458	7,693
Unemployed	93,078	18,808	22,994	6,677	15,029	36,371	3,494	12,772	23,613
Not in labor force (see detail below)	137,816	29,194	41,673	19,121	16,640	51,630	4,282	14,369	40,387
Total	323,165	70,279	96,667	32,634	44,510	131,048	11,702	32,533	81,743
Detailed not in labor force									
Total individuals aged 18 to 24 years not in labor force	137,816	29,194	41,673	19,121	16,640	51,630	4,282	14,369	40,387
Disabled	18,023	2,969	4,635	2,799	3,275	4,770	986	4,087	7,689
Retired	134	34	28	11	11	58	3	25	28
Student	60,594	12,038	15,477	9,466	5,515	21,846	975	4,195	19,411
Homemaker	2,930	839	962	106	369	1,383	131	107	470
Sheltered/non-competitive employment	914	199	226	74	133	329	33	115	283
Other	55,221	13,115	20,345	6,665	7,337	23,244	2,154	5,840	12,506

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

NOTES: *Employed* includes “full-time/part-time not differentiated,” which is not presented separately; total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 3.9b. Individuals aged 18 to 24 years, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Column percent distribution, 2020

<i>Employment status and ‘detailed not in labor force’</i>	<i>Total</i>	<i>Mental health diagnosis¹</i>							
		<i>Trauma- and stressor-related disorders</i>			<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Total individuals aged 18 to 24 years	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Employment status									
Employed	28.6	31.7	33.1	20.9	28.8	32.8	33.5	16.6	21.7
Full time	13.0	15.4	15.4	9.5	10.9	14.8	15.5	5.9	10.3
Part time	11.7	12.4	13.1	9.1	10.9	13.2	13.5	7.6	9.4
Unemployed	28.8	26.8	23.8	20.5	33.8	27.8	29.9	39.3	28.9
Not in labor force (see detail below)	42.6	41.5	43.1	58.6	37.4	39.4	36.6	44.2	49.4
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Detailed not in labor force									
Total individuals aged 18 to 24 years not in labor force	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Disabled	13.1	10.2	11.1	14.6	19.7	9.2	23.0	28.4	19.0
Retired	†	†	†	†	†	†	†	†	†
Student	44.0	41.2	37.1	49.5	33.1	42.3	22.8	29.2	48.1
Homemaker	2.1	2.9	2.3	†	2.2	2.7	3.1	†	1.2
Sheltered/non-competitive employment	†	†	†	†	†	†	†	†	†
Other	40.1	44.9	48.8	34.9	44.1	45.0	50.3	40.6	31.0

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding. *Employed* includes “full-time/part-time not differentiated,” which is not presented separately.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 3.9c. Individuals aged 18 to 24 years, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Row percent distribution, 2020

Employment status and ‘detailed not in labor force’	Total	Mental health diagnosis [†]							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders [†]
					Bipolar disorders	Depressive disorders			
Total individuals aged 18 to 24 years	100.0	20.4	28.3	9.8	12.0	36.7	3.0	9.4	26.4
Employment status									
Employed	100.0	24.1	34.7	7.4	13.9	46.7	4.3	5.8	19.2
Full time	100.0	25.8	35.4	7.4	11.5	46.1	4.3	4.6	20.0
Part time	100.0	23.0	33.5	7.8	12.8	45.4	4.2	6.5	20.3
Unemployed	100.0	20.2	24.7	7.2	16.1	39.1	3.8	13.7	25.4
Not in labor force (see detail below)	100.0	21.2	30.2	13.9	12.1	37.5	3.1	10.4	29.3
Total	100.0	21.7	29.9	10.1	13.8	40.6	3.6	10.1	25.3
Detailed not in labor force									
Total individuals aged 18 to 24 years not in labor force	100.0	21.2	30.2	13.9	12.1	37.5	3.1	10.4	29.3
Disabled	100.0	16.5	25.7	15.5	18.2	26.5	5.5	22.7	42.7
Retired	100.0	25.4	20.9	8.2	8.2	43.3	2.2	18.7	20.9
Student	100.0	19.9	25.5	15.6	9.1	36.1	1.6	6.9	32.0
Homemaker	100.0	28.6	32.8	3.6	12.6	47.2	4.5	3.7	16.0
Sheltered/non-competitive employment	100.0	21.8	24.7	8.1	14.6	36.0	3.6	12.6	31.0
Other	100.0	23.8	36.8	12.1	13.3	42.1	3.9	10.6	22.6

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

[†] “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding. *Employed* includes “full-time/part-time not differentiated,” which is not presented separately.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 3.10a. Individuals aged 18 to 24 years with SMI, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Number, 2020

<i>Employment status and ‘detailed not in labor force’</i>		<i>Mental health diagnosis¹</i>							
		<i>Trauma- and stressor-related disorders Anxiety disorders ADHD</i>			<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
	<i>Total</i>								
Total individuals with SMI aged 18 to 24 years²	414,898	76,426	123,667	34,255	61,291	177,560	14,075	50,920	91,152
Employment status									
Employed	67,081	15,441	23,312	4,580	11,066	34,263	3,253	4,854	10,797
Full time	28,414	6,970	9,962	1,942	3,893	14,510	1,429	1,682	4,641
Part time	27,746	5,954	9,224	1,930	4,121	13,749	1,314	2,211	4,874
Unemployed	70,382	13,463	17,821	4,583	13,302	30,535	2,897	11,564	13,841
Not in labor force (see detail below)	95,574	19,921	29,466	11,409	13,890	39,373	3,541	12,755	23,958
Total	233,037	48,825	70,599	20,572	38,258	104,171	9,691	29,173	48,596
Detailed not in labor force									
Total individuals with SMI aged 18 to 24 years not in labor force	95,574	19,921	29,466	11,409	13,890	39,373	3,541	12,755	23,958
Disabled	13,726	2,500	3,802	2,017	2,921	4,124	909	3,794	4,684
Retired	116	28	25	9	11	53	3	23	21
Student	42,193	7,946	11,328	5,320	4,474	17,424	789	3,719	11,375
Homemaker	2,266	654	729	77	321	1,141	113	93	286
Sheltered/non-competitive employment	811	184	202	48	130	320	31	109	208
Other	36,462	8,609	13,380	3,938	6,033	16,311	1,696	5,017	7,384

ADHD = attention deficit/hyperactivity disorder; SMI = serious mental illness.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTES: *Employed* includes “full-time/part-time not differentiated,” which is not presented separately; total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 3.10b. Individuals aged 18 to 24 years with SMI, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Column percent distribution, 2020

<i>Employment status and ‘detailed not in labor force’</i>		<i>Mental health diagnosis¹</i>							
		<i>Trauma- and stressor-related disorders Anxiety disorders ADHD</i>			<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
	<i>Total</i>								
Total individuals with SMI aged 18 to 24 years²	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Employment status									
Employed	28.8	31.6	33.0	22.3	28.9	32.9	33.6	16.6	22.2
Full time	12.2	14.3	14.1	9.4	10.2	13.9	14.7	5.8	9.6
Part time	11.9	12.2	13.1	9.4	10.8	13.2	13.6	7.6	10.0
Unemployed	30.2	27.6	25.2	22.3	34.8	29.3	29.9	39.6	28.5
Not in labor force (see detail below)	41.0	40.8	41.7	55.5	36.3	37.8	36.5	43.7	49.3
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Detailed not in labor force									
Total individuals with SMI aged 18 to 24 years not in labor force	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Disabled	14.4	12.5	12.9	17.7	21.0	10.5	25.7	29.7	19.6
Retired	†	†	†	†	†	†	†	†	†
Student	44.1	39.9	38.4	46.6	32.2	44.3	22.3	29.2	47.5
Homemaker	2.4	3.3	2.5	†	2.3	2.9	3.2	†	1.2
Sheltered/non-competitive employment	†	†	†	†	†	†	†	†	†
Other	38.2	43.2	45.4	34.5	43.4	41.4	47.9	39.3	30.8

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; SMI = serious mental illness.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding. *Employed* includes “full-time/part-time not differentiated,” which is not presented separately.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 3.10c. Individuals aged 18 to 24 years with SMI, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Row percent distribution, 2020

<i>Employment status and ‘detailed not in labor force’</i>		<i>Mental health diagnosis[†]</i>							
		<i>Trauma- and stressor-related disorders Anxiety disorders ADHD</i>			<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
	<i>Total</i>								
Total individuals with SMI aged 18 to 24 years²	100.0	18.4	29.8	8.3	14.8	42.8	3.4	12.3	22.0
Employment status									
Employed	100.0	23.0	34.8	6.8	16.5	51.1	4.8	7.2	16.1
Full time	100.0	24.5	35.1	6.8	13.7	51.1	5.0	5.9	16.3
Part time	100.0	21.5	33.2	7.0	14.9	49.6	4.7	8.0	17.6
Unemployed	100.0	19.1	25.3	6.5	18.9	43.4	4.1	16.4	19.7
Not in labor force (see detail below)	100.0	20.8	30.8	11.9	14.5	41.2	3.7	13.3	25.1
Total	100.0	21.0	30.3	8.8	16.4	44.7	4.2	12.5	20.9
Detailed not in labor force									
Total individuals with SMI aged 18 to 24 years not in labor force	100.0	20.8	30.8	11.9	14.5	41.2	3.7	13.3	25.1
Disabled	100.0	18.2	27.7	14.7	21.3	30.0	6.6	27.6	34.1
Retired	100.0	24.1	21.6	7.8	9.5	45.7	2.6	19.8	18.1
Student	100.0	18.8	26.8	12.6	10.6	41.3	1.9	8.8	27.0
Homemaker	100.0	28.9	32.2	3.4	14.2	50.4	5.0	4.1	12.6
Sheltered/non-competitive employment	100.0	22.7	24.9	5.9	16.0	39.5	3.8	13.4	25.6
Other	100.0	23.6	36.7	10.8	16.5	44.7	4.7	13.8	20.3

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; SMI = serious mental illness.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding. *Employed* includes “full-time/part-time not differentiated,” which is not presented separately.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 3.11a. Individuals aged 18 to 24 years, by mental health diagnosis, timing of admission, and mental health service setting: Number, 2020

Timing of admission and mental health service setting	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals aged 18 to 24 years	608,665	124,460	172,077	59,814	73,317	223,326	18,210	57,417	160,779
Timing of admission									
New individuals	315,764	68,832	92,722	19,640	35,790	124,827	9,136	26,171	72,238
Continuing individuals	292,901	55,628	79,355	40,174	37,527	98,499	9,074	31,246	88,541
Mental health service setting^{2,3}									
State psychiatric hospital	12,271	1,626	1,333	520	2,286	3,006	1,092	5,160	3,593
Community-based program	590,350	121,415	170,677	59,134	71,041	217,797	17,553	53,378	154,590
Residential treatment center	6,342	1,190	885	715	1,011	1,697	320	1,558	2,436
Other psychiatric inpatient	32,958	4,938	5,388	1,257	5,460	13,395	1,385	7,262	7,196
Institutions under the justice system	7,489	2,640	804	273	419	1,112	837	1,001	3,212

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Individuals could receive services in more than one service setting during the reporting period.

³ Mental health service setting category definitions can be found in Appendix C.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 3.11b. Individuals aged 18 to 24 years, by mental health diagnosis, timing of admission, and mental health service setting: Column percent distribution, 2020

Mental health diagnosis ¹									
Timing of admission and mental health service setting	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals aged 18 to 24 years	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Timing of admission									
New individuals	51.9	55.3	53.9	32.8	48.8	55.9	50.2	45.6	44.9
Continuing individuals	48.1	44.7	46.1	67.2	51.2	44.1	49.8	54.4	55.1
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Column percent of individuals ²									
Mental health service setting ^{2,3}	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Bipolar disorders	Depressive disorders	Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
State psychiatric hospital	2.0	1.3	†	†	3.1	1.3	6.0	9.0	2.2
Community-based program	97.0	97.6	99.2	98.9	96.9	97.5	96.4	93.0	96.2
Residential treatment center	1.0	†	†	1.2	1.4	†	1.8	2.7	1.5
Other psychiatric inpatient	5.4	4.0	3.1	2.1	7.4	6.0	7.6	12.6	4.5
Institutions under the justice system	1.2	2.1	†	†	†	†	4.6	1.7	2.0

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Percentages may not sum to 100 percent, because individuals could receive services in more than one service setting during the reporting period.

³ Mental health service setting category definitions can be found in Appendix C.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 3.11c. Individuals aged 18 to 24 years, by mental health diagnosis, timing of admission, and mental health service setting: Row percent distribution, 2020

Timing of admission and mental health service setting	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals aged 18 to 24 years	100.0	20.4	28.3	9.8	12.0	36.7	3.0	9.4	26.4
Timing of admission									
New individuals	100.0	21.8	29.4	6.2	11.3	39.5	2.9	8.3	22.9
Continuing individuals	100.0	19.0	27.1	13.7	12.8	33.6	3.1	10.7	30.2
Mental health service setting ^{2,3}									
State psychiatric hospital	100.0	13.3	10.9	4.2	18.6	24.5	8.9	42.1	29.3
Community-based program	100.0	20.6	28.9	10.0	12.0	36.9	3.0	9.0	26.2
Residential treatment center	100.0	18.8	14.0	11.3	15.9	26.8	5.0	24.6	38.4
Other psychiatric inpatient	100.0	15.0	16.3	3.8	16.6	40.6	4.2	22.0	21.8
Institutions under the justice system	100.0	35.3	10.7	3.6	5.6	14.8	11.2	13.4	42.9

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total.

"Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Individuals could receive services in more than one service setting during the reporting period.

³ Mental health service setting category definitions can be found in Appendix C.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 3.12a. Individuals aged 18 to 24 years with SMI, by mental health diagnosis, timing of admission, and mental health service setting: Number, 2020

Timing of admission and mental health service setting	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals with SMI aged 18 to 24 years²	414,898	76,426	123,667	34,255	61,291	177,560	14,075	50,920	91,152
Timing of admission									
New individuals	214,766	41,792	66,307	11,435	30,024	98,847	6,685	22,584	40,231
Continuing individuals	200,132	34,634	57,360	22,820	31,267	78,713	7,390	28,336	50,921
Mental health service setting^{3,4}									
State psychiatric hospital	8,989	1,180	1,006	354	1,863	2,242	825	4,279	2,181
Community-based program	400,647	74,167	122,560	33,874	59,505	172,877	13,634	47,601	86,993
Residential treatment center	4,350	721	680	264	775	1,352	261	1,443	1,173
Other psychiatric inpatient	28,489	3,925	4,998	1,077	5,087	12,513	1,289	6,825	5,169
Institutions under the justice system	6,303	1,970	726	250	389	1,013	811	953	2,829

ADHD = attention deficit/hyperactivity disorder; SMI = serious mental illness.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

³ Individuals could receive services in more than one service setting during the reporting period.

⁴ Mental health service setting category definitions can be found in Appendix C.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 3.12b. Individuals aged 18 to 24 years with SMI, by mental health diagnosis, timing of admission, and mental health service setting: Column percent distribution, 2020

Timing of admission and mental health service setting	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals with SMI aged 18 to 24 years²	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Timing of admission									
New individuals	51.8	54.7	53.6	33.4	49.0	55.7	47.5	44.4	44.1
Continuing individuals	48.2	45.3	46.4	66.6	51.0	44.3	52.5	55.6	55.9
Mental health service setting^{3,4}									
State psychiatric hospital	2.2	1.5	†	1.0	3.0	1.3	5.9	8.4	2.4
Community-based program	96.6	97.0	99.1	98.9	97.1	97.4	96.9	93.5	95.4
Residential treatment center	1.0	†	†	†	1.3	†	1.9	2.8	1.3
Other psychiatric inpatient	6.9	5.1	4.0	3.1	8.3	7.0	9.2	13.4	5.7
Institutions under the justice system	1.5	2.6	†	†	†	†	5.8	1.9	3.1

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; SMI = serious mental illness.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

³ Percentages may not sum to 100 percent, because individuals could receive services in more than one service setting during the reporting period.

⁴ Mental health service setting category definitions can be found in Appendix C.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 3.12c. Individuals aged 18 to 24 years with SMI, by mental health diagnosis, timing of admission, and mental health service setting: Row percent distribution, 2020

Timing of admission and mental health service setting	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals with SMI aged 18 to 24 years²	100.0	18.4	29.8	8.3	14.8	42.8	3.4	12.3	22.0
Timing of admission									
New individuals	100.0	19.5	30.9	5.3	14.0	46.0	3.1	10.5	18.7
Continuing individuals	100.0	17.3	28.7	11.4	15.6	39.3	3.7	14.2	25.4
Mental health service setting^{3,4}									
State psychiatric hospital	100.0	13.1	11.2	3.9	20.7	24.9	9.2	47.6	24.3
Community-based program	100.0	18.5	30.6	8.5	14.9	43.1	3.4	11.9	21.7
Residential treatment center	100.0	16.6	15.6	6.1	17.8	31.1	6.0	33.2	27.0
Other psychiatric inpatient	100.0	13.8	17.5	3.8	17.9	43.9	4.5	24.0	18.1
Institutions under the justice system	100.0	31.3	11.5	4.0	6.2	16.1	12.9	15.1	44.9

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; SMI = serious mental illness.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

³ Individuals could receive services in more than one service setting during the reporting period.

⁴ Mental health service setting category definitions can be found in Appendix C.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 3.13. Individuals aged 18 to 24 years, by mental health service setting and SMI status: Number and percent distribution, 2020

<i>Mental health service setting</i>	<i>SMI status¹</i>								
	<i>Number of individuals</i>			<i>Column percent of individuals</i>			<i>Row percent distribution of individuals</i>		
	<i>Total</i>	<i>SMI</i>	<i>Not SMI</i>	<i>Total</i>	<i>SMI</i>	<i>Not SMI</i>	<i>Total</i>	<i>SMI</i>	<i>Not SMI</i>
Total individuals aged 18 to 24 years	668,985	470,276	198,709	100.0	100.0	100.0	100.0	70.3	29.7
Mental health service setting^{2,3}									
State psychiatric hospital	11,697	9,873	1,824	1.7	2.1	†	100.0	84.4	15.6
Community-based program	650,121	453,877	196,244	97.2	96.5	98.8	100.0	69.8	30.2
Residential treatment center	6,735	5,489	1,246	1.0	1.2	†	100.0	81.5	18.5
Other psychiatric inpatient	35,079	30,892	4,187	5.2	6.6	2.1	100.0	88.1	11.9
Institutions under the justice system	8,102	6,969	1,133	1.2	1.5	†	100.0	86.0	14.0

† Less than 1 percent of individuals.

SMI = serious mental illness.

¹ The federal definition for SMI status can be found in Appendix D.

² Individuals could receive services in more than one mental health service setting during the reporting period.

³ Mental health service setting category definitions can be found in Appendix C.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 4.1a. Individuals aged 25 to 54 years, by gender, age, race, ethnicity, education, veteran status, and SMI status: Number, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Number of states reporting	45	47	49	49	48	50
Total individuals aged 25 to 54 years	2,377,733	2,586,376	2,844,162	2,890,752	2,896,236	3,123,221
Gender						
Male	1,080,644	1,171,245	1,311,625	1,334,575	1,330,307	1,455,295
Female	1,293,752	1,408,469	1,523,861	1,551,119	1,560,085	1,661,394
Total	2,374,396	2,579,714	2,835,486	2,885,694	2,890,392	3,116,689
Age¹						
25 to 34 years	841,821	938,220	1,058,144	1,092,987	1,116,300	1,221,654
35 to 39 years	376,740	422,607	480,949	500,964	509,507	559,204
40 to 49 years	740,696	782,893	839,462	843,189	835,573	896,284
50 to 54 years	418,476	442,656	465,607	453,612	434,856	446,079
Total	2,377,733	2,586,376	2,844,162	2,890,752	2,896,236	3,123,221
Race						
American Indian or Alaska Native	32,157	33,510	37,644	42,189	47,532	63,295
Asian ²	31,706	36,415	40,995	45,277	41,129	41,416
Black or African American	446,713	485,914	530,175	546,511	557,975	570,602
Native Hawaiian or Other Pacific Islander	6,565	7,399	5,126	6,368	6,546	7,157
White	1,520,013	1,659,872	1,789,361	1,824,514	1,800,040	1,946,322
Other	140,857	160,979	193,512	217,188	228,239	251,296
Total	2,178,011	2,384,089	2,596,813	2,682,047	2,681,461	2,880,088
Ethnicity						
Not of Hispanic or Latino origin	1,680,884	1,809,217	2,199,958	2,233,243	2,223,030	2,364,429
Hispanic or Latino origin	249,253	279,388	330,395	341,987	350,612	486,305
Total	1,930,137	2,088,605	2,530,353	2,575,230	2,573,642	2,850,734
Specific Hispanic or Latino origin						
Mexican	18,516	15,379	17,769	17,082	17,647	17,364
Puerto Rican	23,361	19,927	23,955	13,628	13,921	14,128
Cuban	4,290	3,895	3,784	3,759	3,822	4,257
Other specific	21,045	21,136	24,324	24,842	20,335	22,282
Total	67,212	60,337	69,832	59,311	55,725	58,031

Continued. See notes at end of table.

Table 4.1a. Individuals aged 25 to 54 years, by gender, age, race, ethnicity, education, veteran status, and SMI status: Number, 2015–2020
(continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Highest school grade completed						
0 to 8	123,090	124,987	131,314	126,206	133,509	125,173
9 to 11	206,859	204,520	236,622	240,303	236,089	253,598
12 (or GED)	516,915	531,605	605,040	671,080	661,261	710,614
More than 12	237,997	247,657	317,116	351,127	350,040	387,543
Total	1,084,861	1,108,769	1,290,092	1,388,716	1,380,899	1,476,928
Veteran status						
Veteran	<i>n/c</i>	<i>n/c</i>	22,872	25,438	45,035	59,166
Not a veteran	<i>n/c</i>	<i>n/c</i>	780,726	1,126,091	1,172,637	1,207,174
Total	<i>n/c</i>	<i>n/c</i>	803,598	1,151,529	1,217,672	1,266,340
SMI status³						
SMI	1,626,431	1,735,632	1,901,453	2,019,609	2,111,153	2,164,629
Not SMI	637,034	695,917	775,584	738,580	644,685	815,832
Total	2,263,465	2,431,549	2,677,037	2,758,189	2,755,838	2,980,461

n/c Not collected.

GED = general equivalency diploma; SMI = serious mental illness.

¹ Age is based on the individual's age at the midpoint of the reporting period.

² For some states, "Asian" includes "Pacific Islanders." See Appendix C.

³ The federal definition for SMI status can be found in Appendix D.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 4.1b. Individuals aged 25 to 54 years, by gender, age, race, ethnicity, education, veteran status, and SMI status: Column percent distribution, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Number of states reporting	45	47	49	49	48	50
Total individuals aged 25 to 54 years	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	45.5	45.4	46.3	46.2	46.0	46.7
Female	54.5	54.6	53.7	53.8	54.0	53.3
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age¹						
25 to 34 years	35.4	36.3	37.2	37.8	38.5	39.1
35 to 39 years	15.8	16.3	16.9	17.3	17.6	17.9
40 to 49 years	31.2	30.3	29.5	29.2	28.9	28.7
50 to 54 years	17.6	17.1	16.4	15.7	15.0	14.3
Total	100.0	100.0	100.0	100.0	100.0	100.0
Race						
American Indian or Alaska Native	1.5	1.4	1.4	1.6	1.8	2.2
Asian ²	1.5	1.5	1.6	1.7	1.5	1.4
Black or African American	20.5	20.4	20.4	20.4	20.8	19.8
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	69.8	69.6	68.9	68.0	67.1	67.6
Other	6.5	6.8	7.5	8.1	8.5	8.7
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	87.1	86.6	86.9	86.7	86.4	82.9
Hispanic or Latino origin	12.9	13.4	13.1	13.3	13.6	17.1
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	27.5	25.5	25.4	28.8	31.7	29.9
Puerto Rican	34.8	33.0	34.3	23.0	25.0	24.3
Cuban	6.4	6.5	5.4	6.3	6.9	7.3
Other specific	31.3	35.0	34.8	41.9	36.5	38.4
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 4.1b. Individuals aged 25 to 54 years, by gender, age, race, ethnicity, education, veteran status, and SMI status: Column percent distribution, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Highest school grade completed						
0 to 8	11.3	11.3	10.2	9.1	9.7	8.5
9 to 11	19.1	18.4	18.3	17.3	17.1	17.2
12 (or GED)	47.6	47.9	46.9	48.3	47.9	48.1
More than 12	21.9	22.3	24.6	25.3	25.3	26.2
Total	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status						
Veteran	<i>n/c</i>	<i>n/c</i>	2.8	2.2	3.7	4.7
Not a veteran	<i>n/c</i>	<i>n/c</i>	97.2	97.8	96.3	95.3
Total	<i>n/c</i>	<i>n/c</i>	100.0	100.0	100.0	100.0
SMI status³						
SMI	71.9	71.4	71.0	73.2	76.6	72.6
Not SMI	28.1	28.6	29.0	26.8	23.4	27.4
Total	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals; *n/c* Not collected.

GED = general equivalency diploma; SMI = serious mental illness.

¹ Age is based on the individual's age at the midpoint of the reporting period.

² For some states, "Asian" includes "Pacific Islanders." See Appendix C.

³ The federal definition for SMI status can be found in Appendix D.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 4.2a. Individuals aged 25 to 54 years with SMI, by gender, age, race, ethnicity, education, and veteran status: Number, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Number of states reporting	45	46	47	48	47	49
Total individuals with SMI aged 25 to 54 years¹	1,626,431	1,735,632	1,901,453	2,019,609	2,111,153	2,164,629
Gender						
Male	738,630	788,436	875,885	927,416	955,390	990,654
Female	886,088	945,165	1,023,468	1,090,070	1,152,695	1,171,165
Total	1,624,718	1,733,601	1,899,353	2,017,486	2,108,085	2,161,819
Age²						
25 to 34 years	547,207	597,705	672,646	731,259	791,205	821,922
35 to 39 years	254,741	281,245	318,077	347,171	368,849	383,176
40 to 49 years	520,657	541,236	579,192	605,568	620,787	634,205
50 to 54 years	303,826	315,446	331,538	335,611	330,312	325,326
Total	1,626,431	1,735,632	1,901,453	2,019,609	2,111,153	2,164,629
Race						
American Indian or Alaska Native	19,305	19,904	24,127	26,769	32,570	37,939
Asian ³	23,878	25,884	30,272	33,884	32,507	31,318
Black or African American	327,328	341,690	370,984	398,793	412,202	414,328
Native Hawaiian or Other Pacific Islander	5,317	5,723	3,914	4,838	5,266	5,410
White	1,016,729	1,106,461	1,190,531	1,250,088	1,304,679	1,337,345
Other	107,912	118,255	152,299	174,968	186,689	197,996
Total	1,500,469	1,617,917	1,772,127	1,889,340	1,973,913	2,024,336
Ethnicity						
Not of Hispanic or Latino origin	1,164,655	1,244,086	1,436,901	1,517,954	1,569,642	1,640,713
Hispanic or Latino origin	183,735	207,173	248,058	264,227	285,410	345,699
Total	1,348,390	1,451,259	1,684,959	1,782,181	1,855,052	1,986,412
Specific Hispanic or Latino origin						
Mexican	13,141	10,695	12,682	12,606	13,152	12,923
Puerto Rican	15,941	14,023	15,886	11,093	11,293	11,790
Cuban	4,006	3,642	3,473	3,540	3,575	3,998
Other specific	16,507	16,180	18,784	19,864	15,328	16,645
Total	49,595	44,540	50,825	47,103	43,348	45,356

Continued. See notes at end of table.

Table 4.2a. Individuals aged 25 to 54 years with SMI, by gender, age, race, ethnicity, education, and veteran status: Number, 2015–2020
(continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Highest school grade completed						
0 to 8	70,718	73,983	83,394	80,988	86,650	77,796
9 to 11	148,845	153,750	175,923	181,143	178,204	186,767
12 (or GED)	359,499	387,635	443,814	505,102	497,970	527,833
More than 12	175,195	190,436	232,982	260,482	262,500	279,535
Total	754,257	805,804	936,113	1,027,715	1,025,324	1,071,931
Veteran status						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	19,338	37,251	49,754
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	768,295	821,757	809,124
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	787,633	859,008	858,878

n/c Not collected.

GED = general equivalency diploma; SMI = serious mental illness.

¹ The federal definition for SMI status can be found in Appendix D.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 4.2b. Individuals aged 25 to 54 years with SMI, by gender, age, race, ethnicity, education, and veteran status: Column percent distribution, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Number of states reporting	45	46	47	48	47	49
Total individuals with SMI aged 25 to 54 years¹	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	45.5	45.5	46.1	46.0	45.3	45.8
Female	54.5	54.5	53.9	54.0	54.7	54.2
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age²						
25 to 34 years	33.6	34.4	35.4	36.2	37.5	38.0
35 to 39 years	15.7	16.2	16.7	17.2	17.5	17.7
40 to 49 years	32.0	31.2	30.5	30.0	29.4	29.3
50 to 54 years	18.7	18.2	17.4	16.6	15.6	15.0
Total	100.0	100.0	100.0	100.0	100.0	100.0
Race						
American Indian or Alaska Native	1.3	1.2	1.4	1.4	1.7	1.9
Asian ³	1.6	1.6	1.7	1.8	1.6	1.5
Black or African American	21.8	21.1	20.9	21.1	20.9	20.5
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	67.8	68.4	67.2	66.2	66.1	66.1
Other	7.2	7.3	8.6	9.3	9.5	9.8
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	86.4	85.7	85.3	85.2	84.6	82.6
Hispanic or Latino origin	13.6	14.3	14.7	14.8	15.4	17.4
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	26.5	24.0	25.0	26.8	30.3	28.5
Puerto Rican	32.1	31.5	31.3	23.6	26.1	26.0
Cuban	8.1	8.2	6.8	7.5	8.2	8.8
Other specific	33.3	36.3	37.0	42.2	35.4	36.7
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 4.2b. Individuals aged 25 to 54 years with SMI, by gender, age, race, ethnicity, education, and veteran status: Column percent distribution, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Highest school grade completed						
0 to 8	9.4	9.2	8.9	7.9	8.5	7.3
9 to 11	19.7	19.1	18.8	17.6	17.4	17.4
12 (or GED)	47.7	48.1	47.4	49.1	48.6	49.2
More than 12	23.2	23.6	24.9	25.3	25.6	26.1
Total	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	2.5	4.3	5.8
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	97.5	95.7	94.2
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	100.0	100.0	100.0

† Less than 1 percent of individuals; *n/c* Not collected.

GED = general equivalency diploma; SMI = serious mental illness.

¹ The federal definition for SMI status can be found in Appendix D.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 4.3a. Individuals aged 25 to 54 years, by mental health diagnosis, gender, race, and ethnicity: Number, 2020

Gender, race, and ethnicity	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals aged 25 to 54 years	2,667,813	532,647	697,741	101,545	463,761	914,629	105,153	460,519	619,606
Gender									
Male	1,221,454	193,373	255,235	51,267	183,600	354,183	43,223	303,982	338,210
Female	1,441,995	338,316	441,418	50,124	279,383	559,030	61,822	155,994	280,339
Total	2,663,449	531,689	696,653	101,391	462,983	913,213	105,045	459,976	618,549
Race									
American Indian or Alaska Native	57,135	13,427	14,823	1,553	6,098	18,512	1,552	6,450	19,662
Asian ²	36,310	6,093	7,314	892	3,830	11,631	767	11,582	7,673
Black or African American	487,726	89,277	78,648	9,777	76,536	156,148	18,815	147,254	103,525
Islander	6,123	1,334	1,473	198	804	2,007	152	1,447	1,405
White	1,663,532	342,025	497,084	75,316	320,878	590,653	71,637	217,492	377,105
Other	224,725	43,622	50,407	6,243	32,618	77,745	7,341	47,069	49,183
Total	2,475,551	495,778	649,749	93,979	440,764	856,696	100,264	431,294	558,553
Ethnicity									
Not of Hispanic or Latino origin	2,017,914	413,793	540,655	82,507	374,657	708,363	86,619	349,991	438,254
Hispanic or Latino origin	427,781	79,916	106,488	12,484	57,927	138,262	11,532	68,402	119,963
Total	2,445,695	493,709	647,143	94,991	432,584	846,625	98,151	418,393	558,217
Specific Hispanic or Latino origin									
Mexican	16,602	4,749	4,108	463	2,231	6,106	480	2,452	3,148
Puerto Rican	12,170	2,416	2,210	242	2,263	4,322	372	2,733	1,693
Cuban	3,984	460	573	69	584	1,232	36	794	839
Other specific	20,562	5,142	5,065	652	3,337	7,598	745	3,558	3,228
Total	53,318	12,767	11,956	1,426	8,415	19,258	1,633	9,537	8,908

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 4.3b. Individuals aged 25 to 54 years, by mental health diagnosis, gender, race, and ethnicity: Column percent distribution, 2020

Gender, race, and ethnicity	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals aged 25 to 54 years	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender									
Male	45.9	36.4	36.6	50.6	39.7	38.8	41.1	66.1	54.7
Female	54.1	63.6	63.4	49.4	60.3	61.2	58.9	33.9	45.3
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race									
American Indian or Alaska Native	2.3	2.7	2.3	1.7	1.4	2.2	1.5	1.5	3.5
Asian ²	1.5	1.2	1.1	†	†	1.4	†	2.7	1.4
Black or African American	19.7	18.0	12.1	10.4	17.4	18.2	18.8	34.1	18.5
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	†	†	†
White	67.2	69.0	76.5	80.1	72.8	68.9	71.4	50.4	67.5
Other	9.1	8.8	7.8	6.6	7.4	9.1	7.3	10.9	8.8
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity									
Not of Hispanic or Latino origin	82.5	83.8	83.5	86.9	86.6	83.7	88.3	83.7	78.5
Hispanic or Latino origin	17.5	16.2	16.5	13.1	13.4	16.3	11.7	16.3	21.5
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	31.1	37.2	34.4	32.5	26.5	31.7	29.4	25.7	35.3
Puerto Rican	22.8	18.9	18.5	17.0	26.9	22.4	22.8	28.7	19.0
Cuban	7.5	3.6	4.8	4.8	6.9	6.4	2.2	8.3	9.4
Other specific	38.6	40.3	42.4	45.7	39.7	39.5	45.6	37.3	36.2
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

Other mental disorders¹

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 4.3c. Individuals aged 25 to 54 years, by mental health diagnosis, gender, race, and ethnicity: Row percent distribution, 2020

Gender, race, and ethnicity		Mental health diagnosis ¹								
		Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Total individuals aged 25 to 54 years		100.0	20.0	26.2	3.8	17.4	34.3	3.9	17.3	23.2
Gender										
Male		100.0	15.8	20.9	4.2	15.0	29.0	3.5	24.9	27.7
Female		100.0	23.5	30.6	3.5	19.4	38.8	4.3	10.8	19.4
Total		100.0	20.0	26.2	3.8	17.4	34.3	3.9	17.3	23.2
Race										
American Indian or Alaska Native		100.0	23.5	25.9	2.7	10.7	32.4	2.7	11.3	34.4
Asian ²		100.0	16.8	20.1	2.5	10.5	32.0	2.1	31.9	21.1
Black or African American		100.0	18.3	16.1	2.0	15.7	32.0	3.9	30.2	21.2
Native Hawaiian or Other Pacific Islander		100.0	21.8	24.1	3.2	13.1	32.8	2.5	23.6	22.9
White		100.0	20.6	29.9	4.5	19.3	35.5	4.3	13.1	22.7
Other		100.0	19.4	22.4	2.8	14.5	34.6	3.3	20.9	21.9
Total		100.0	20.0	26.2	3.8	17.8	34.6	4.1	17.4	22.6
Ethnicity										
Not of Hispanic or Latino origin		100.0	20.5	26.8	4.1	18.6	35.1	4.3	17.3	21.7
Hispanic or Latino origin		100.0	18.7	24.9	2.9	13.5	32.3	2.7	16.0	28.0
Total		100.0	20.2	26.5	3.9	17.7	34.6	4.0	17.1	22.8
Specific Hispanic or Latino origin										
Mexican		100.0	28.6	24.7	2.8	13.4	36.8	2.9	14.8	19.0
Puerto Rican		100.0	19.9	18.2	2.0	18.6	35.5	3.1	22.5	13.9
Cuban		100.0	11.5	14.4	1.7	14.7	30.9	†	19.9	21.1
Other specific		100.0	25.0	24.6	3.2	16.2	37.0	3.6	17.3	15.7
Total		100.0	23.9	22.4	2.7	15.8	36.1	3.1	17.9	16.7

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 4.4a. Individuals aged 25 to 54 years with SMI, gender, race, and ethnicity: Number, 2020

Gender, race, and ethnicity	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals with SMI aged 25 to 54 years²	1,947,307	351,778	518,721	61,183	396,179	743,169	85,573	415,559	354,901
Gender									
Male	885,736	121,735	187,050	29,737	155,980	286,832	34,761	274,548	191,902
Female	1,059,100	229,631	331,091	31,367	239,616	455,376	50,751	140,570	162,486
Total	1,944,836	351,366	518,141	61,104	395,596	742,208	85,512	415,118	354,388
Race									
American Indian or Alaska Native	35,284	8,582	10,232	955	5,017	14,350	1,199	5,622	8,359
Asian ³	28,633	4,112	5,600	502	3,339	9,676	619	10,821	4,956
Black or African American	378,748	58,938	61,502	6,057	67,058	131,293	15,677	133,048	64,518
Native Hawaiian or Other Pacific Islander	4,926	1,004	1,203	143	719	1,781	133	1,356	967
White	1,195,991	227,616	368,434	45,681	273,081	475,481	58,461	194,944	211,738
Other	186,045	32,918	41,591	4,161	28,910	68,418	6,365	44,509	35,324
Total	1,829,627	333,170	488,562	57,499	378,124	700,999	82,454	390,300	325,862
Ethnicity									
Not of Hispanic or Latino origin	1,477,661	276,070	403,575	50,794	316,360	565,747	71,278	313,848	259,878
Hispanic or Latino origin	313,136	51,587	79,057	6,600	53,105	121,646	9,441	64,576	61,840
Total	1,790,797	327,657	482,632	57,394	369,465	687,393	80,719	378,424	321,718
Specific Hispanic or Latino origin									
Mexican	12,691	3,371	2,946	306	2,049	5,506	390	2,278	1,754
Puerto Rican	10,674	1,881	1,906	191	2,150	4,022	312	2,659	1,129
Cuban	3,787	401	517	55	569	1,191	31	781	779
Other specific	16,106	3,605	3,767	447	3,021	6,677	628	3,270	1,980
Total	43,258	9,258	9,136	999	7,789	17,396	1,361	8,988	5,642

ADHD = attention deficit/hyperactivity disorder; SMI = serious mental illness.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 4.4b. Individuals aged 25 to 54 years with SMI, by mental health diagnosis, gender, race, and ethnicity: Column percent distribution, 2020

Gender, race, and ethnicity	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals with SMI aged 25 to 54 years²	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender									
Male	45.5	34.6	36.1	48.7	39.4	38.6	40.7	66.1	54.2
Female	54.5	65.4	63.9	51.3	60.6	61.4	59.3	33.9	45.8
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race									
American Indian or Alaska Native	1.9	2.6	2.1	1.7	1.3	2.0	1.5	1.4	2.6
Asian ³	1.6	1.2	1.1	†	†	1.4	†	2.8	1.5
Black or African American	20.7	17.7	12.6	10.5	17.7	18.7	19.0	34.1	19.8
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	†	†	†
White	65.4	68.3	75.4	79.4	72.2	67.8	70.9	49.9	65.0
Other	10.2	9.9	8.5	7.2	7.6	9.8	7.7	11.4	10.8
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity									
Not of Hispanic or Latino origin	82.5	84.3	83.6	88.5	85.6	82.3	88.3	82.9	80.8
Hispanic or Latino origin	17.5	15.7	16.4	11.5	14.4	17.7	11.7	17.1	19.2
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	29.3	36.4	32.2	30.6	26.3	31.7	28.7	25.3	31.1
Puerto Rican	24.7	20.3	20.9	19.1	27.6	23.1	22.9	29.6	20.0
Cuban	8.8	4.3	5.7	5.5	7.3	6.8	2.3	8.7	13.8
Other specific	37.2	38.9	41.2	44.7	38.8	38.4	46.1	36.4	35.1
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; SMI = serious mental illness.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 4.4c. Individuals aged 25 to 54 years with SMI, by mental health diagnosis, gender, race, and ethnicity: Row percent distribution, 2020

Gender, race, and ethnicity	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals with SMI aged 25 to 54 years²	100.0	18.1	26.6	3.1	20.3	38.2	4.4	21.3	18.2
Gender									
Male	100.0	13.7	21.1	3.4	17.6	32.4	3.9	31.0	21.7
Female	100.0	21.7	31.3	3.0	22.6	43.0	4.8	13.3	15.3
Total	100.0	18.1	26.6	3.1	20.3	38.2	4.4	21.3	18.2
Race									
American Indian or Alaska Native	100.0	24.3	29.0	2.7	14.2	40.7	3.4	15.9	23.7
Asian ³	100.0	14.4	19.6	1.8	11.7	33.8	2.2	37.8	17.3
Black or African American	100.0	15.6	16.2	1.6	17.7	34.7	4.1	35.1	17.0
Native Hawaiian or Other Pacific Islander	100.0	20.4	24.4	2.9	14.6	36.2	2.7	27.5	19.6
White	100.0	19.0	30.8	3.8	22.8	39.8	4.9	16.3	17.7
Other	100.0	17.7	22.4	2.2	15.5	36.8	3.4	23.9	19.0
Total	100.0	18.2	26.7	3.1	20.7	38.3	4.5	21.3	17.8
Ethnicity									
Not of Hispanic or Latino origin	100.0	18.7	27.3	3.4	21.4	38.3	4.8	21.2	17.6
Hispanic or Latino origin	100.0	16.5	25.2	2.1	17.0	38.8	3.0	20.6	19.7
Total	100.0	18.3	27.0	3.2	20.6	38.4	4.5	21.1	18.0
Specific Hispanic or Latino origin									
Mexican	100.0	26.6	23.2	2.4	16.1	43.4	3.1	17.9	13.8
Puerto Rican	100.0	17.6	17.9	1.8	20.1	37.7	2.9	24.9	10.6
Cuban	100.0	10.6	13.7	1.5	15.0	31.4	†	20.6	20.6
Other specific	100.0	22.4	23.4	2.8	18.8	41.5	3.9	20.3	12.3
Total	100.0	21.4	21.1	2.3	18.0	40.2	3.1	20.8	13.0

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; SMI = serious mental illness.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 4.5a. Individuals aged 25 to 54 years, by mental health diagnosis, education, and marital status: Number, 2020

Mental health diagnosis ¹									
Education and marital status	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals aged 25 to 54 years	2,667,813	532,647	697,741	101,545	463,761	914,629	105,153	460,519	619,606
Highest school grade completed									
0 to 8	113,179	29,616	30,407	8,901	14,338	38,135	4,404	18,304	34,409
9 to 11	229,309	50,139	51,449	7,356	43,093	77,682	12,781	55,219	60,305
12 (or GED)	648,164	142,946	177,316	24,946	128,633	241,387	35,144	122,830	149,872
More than 12	359,951	86,868	113,598	17,134	77,695	141,952	19,489	50,345	67,799
Total	1,350,603	309,569	372,770	58,337	263,759	499,156	71,818	246,698	312,385
Marital status									
Never married	953,573	164,765	195,505	37,330	165,136	291,045	36,992	210,044	251,513
Currently married	229,768	54,503	66,777	7,425	44,900	91,757	8,935	21,246	44,411
Separated	98,217	24,203	24,947	2,431	22,216	39,339	4,398	11,641	17,303
Divorced	183,527	43,242	50,706	5,609	44,869	72,680	10,379	24,892	32,263
Widowed	30,659	8,859	7,474	781	6,238	12,212	1,142	3,436	5,793
Total	1,495,744	295,572	345,409	53,576	283,359	507,033	61,846	271,259	351,283

ADHD = attention deficit/hyperactivity disorder; GED = general equivalency diploma.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 4.5b. Individuals aged 25 to 54 years, by mental health diagnosis, education, and marital status: Column percent distribution, 2020

<i>Education and marital status</i>		<i>Mental health diagnosis¹</i>							
		<i>Trauma- and stressor-related disorders</i>			<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
	<i>Total</i>			<i>ADHD</i>					
Total individuals aged 25 to 54 years	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Highest school grade completed									
0 to 8	8.4	9.6	8.2	15.3	5.4	7.6	6.1	7.4	11.0
9 to 11	17.0	16.2	13.8	12.6	16.3	15.6	17.8	22.4	19.3
12 (or GED)	48.0	46.2	47.6	42.8	48.8	48.4	48.9	49.8	48.0
More than 12	26.7	28.1	30.5	29.4	29.5	28.4	27.1	20.4	21.7
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Marital status									
Never married	63.8	55.7	56.6	69.7	58.3	57.4	59.8	77.4	71.6
Currently married	15.4	18.4	19.3	13.9	15.8	18.1	14.4	7.8	12.6
Separated	6.6	8.2	7.2	4.5	7.8	7.8	7.1	4.3	4.9
Divorced	12.3	14.6	14.7	10.5	15.8	14.3	16.8	9.2	9.2
Widowed	2.0	3.0	2.2	1.5	2.2	2.4	1.8	1.3	1.6
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

ADHD = attention deficit/hyperactivity disorder; GED = general equivalency diploma.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 4.5c. Individuals aged 25 to 54 years, by mental health diagnosis, education, and marital status: Row percent distribution, 2020

<i>Education and marital status</i>	<i>Total</i>	<i>Mental health diagnosis¹</i>							
		<i>Trauma- and stressor- related disorders</i>	<i>Anxiety disorders</i>	<i>ADHD</i>	<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Total individuals aged 25 to 54 years	100.0	20.0	26.2	3.8	17.4	34.3	3.9	17.3	23.2
Highest school grade completed									
0 to 8	100.0	26.2	26.9	7.9	12.7	33.7	3.9	16.2	30.4
9 to 11	100.0	21.9	22.4	3.2	18.8	33.9	5.6	24.1	26.3
12 (or GED)	100.0	22.1	27.4	3.8	19.8	37.2	5.4	19.0	23.1
More than 12	100.0	24.1	31.6	4.8	21.6	39.4	5.4	14.0	18.8
Total	100.0	22.9	27.6	4.3	19.5	37.0	5.3	18.3	23.1
Marital status									
Never married	100.0	17.3	20.5	3.9	17.3	30.5	3.9	22.0	26.4
Currently married	100.0	23.7	29.1	3.2	19.5	39.9	3.9	9.2	19.3
Separated	100.0	24.6	25.4	2.5	22.6	40.1	4.5	11.9	17.6
Divorced	100.0	23.6	27.6	3.1	24.4	39.6	5.7	13.6	17.6
Widowed	100.0	28.9	24.4	2.5	20.3	39.8	3.7	11.2	18.9
Total	100.0	19.8	23.1	3.6	18.9	33.9	4.1	18.1	23.5

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; GED = general equivalency diploma.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 4.6a. Individuals aged 25 to 54 years with SMI, by mental health diagnosis, education, and marital status: Number, 2020

Education and marital status	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals with SMI aged 25 to 54 years²	1,947,307	351,778	518,721	61,183	396,179	743,169	85,573	415,559	354,901
Highest school grade completed									
0 to 8	73,194	18,036	20,042	4,052	11,595	26,961	3,594	16,157	17,379
9 to 11	174,735	36,842	40,025	4,903	36,756	61,856	11,050	50,409	36,437
12 (or GED)	493,036	106,612	134,180	17,007	110,074	192,854	30,074	110,971	89,019
More than 12	266,363	62,144	83,156	11,399	66,083	110,489	16,396	45,200	40,822
Total	1,007,328	223,634	277,403	37,361	224,508	392,160	61,114	222,737	183,657
Marital status									
Never married	700,063	102,890	152,545	21,870	143,085	243,792	30,837	188,909	132,342
Currently married	165,673	35,237	50,234	4,694	38,382	74,261	7,461	18,735	23,727
Separated	77,424	17,074	20,771	1,723	19,918	34,102	3,883	10,660	10,294
Divorced	146,512	32,875	42,025	4,174	39,599	61,599	9,096	22,283	20,172
Widowed	23,835	6,490	5,646	513	5,707	10,776	936	3,181	3,474
Total	1,113,507	194,566	271,221	32,974	246,691	424,530	52,213	243,768	190,009

ADHD = attention deficit/hyperactivity disorder; GED = general equivalency diploma; SMI = serious mental illness.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 4.6b. Individuals aged 25 to 54 years with SMI, by mental health diagnosis, education, and marital status: Column percent distribution, 2020

<i>Education and marital status</i>		<i>Mental health diagnosis¹</i>							
		<i>Trauma- and stressor-related disorders</i> <i>Anxiety disorders</i> <i>ADHD</i>			<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
	<i>Total</i>								
Total individuals with SMI aged 25 to 54 years²	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Highest school grade completed									
0 to 8	7.3	8.1	7.2	10.8	5.2	6.9	5.9	7.3	9.5
9 to 11	17.3	16.5	14.4	13.1	16.4	15.8	18.1	22.6	19.8
12 (or GED)	48.9	47.7	48.4	45.5	49.0	49.2	49.2	49.8	48.5
More than 12	26.4	27.8	30.0	30.5	29.4	28.2	26.8	20.3	22.2
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Marital status									
Never married	62.9	52.9	56.2	66.3	58.0	57.4	59.1	77.5	69.7
Currently married	14.9	18.1	18.5	14.2	15.6	17.5	14.3	7.7	12.5
Separated	7.0	8.8	7.7	5.2	8.1	8.0	7.4	4.4	5.4
Divorced	13.2	16.9	15.5	12.7	16.1	14.5	17.4	9.1	10.6
Widowed	2.1	3.3	2.1	1.6	2.3	2.5	1.8	1.3	1.8
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

ADHD = attention deficit/hyperactivity disorder; GED = general equivalency diploma; SMI = serious mental illness.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 4.6c. Individuals aged 25 to 54 years with SMI, by mental health diagnosis, education, and marital status: Row percent distribution, 2020

Education and marital status		Mental health diagnosis ¹								
		Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Total individuals with SMI aged 25 to 54 years ²		100.0	18.1	26.6	3.1	20.3	38.2	4.4	21.3	18.2
Highest school grade completed										
0 to 8	100.0	24.6	27.4	5.5	15.8	36.8	4.9	22.1	23.7	
9 to 11	100.0	21.1	22.9	2.8	21.0	35.4	6.3	28.8	20.9	
12 (or GED)	100.0	21.6	27.2	3.4	22.3	39.1	6.1	22.5	18.1	
More than 12	100.0	23.3	31.2	4.3	24.8	41.5	6.2	17.0	15.3	
Total	100.0	22.2	27.5	3.7	22.3	38.9	6.1	22.1	18.2	
Marital status										
Never married	100.0	14.7	21.8	3.1	20.4	34.8	4.4	27.0	18.9	
Currently married	100.0	21.3	30.3	2.8	23.2	44.8	4.5	11.3	14.3	
Separated	100.0	22.1	26.8	2.2	25.7	44.0	5.0	13.8	13.3	
Divorced	100.0	22.4	28.7	2.8	27.0	42.0	6.2	15.2	13.8	
Widowed	100.0	27.2	23.7	2.2	23.9	45.2	3.9	13.3	14.6	
Total	100.0	17.5	24.4	3.0	22.2	38.1	4.7	21.9	17.1	

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; GED = general equivalency diploma; SMI = serious mental illness.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 4.7a. Individuals aged 25 to 54 years, by mental health diagnosis, living arrangement, SMI status, and veteran status: Number, 2020

Living arrangement, SMI status, and veteran status	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals aged 25 to 54 years	2,667,813	532,647	697,741	101,545	463,761	914,629	105,153	460,519	619,606
Living arrangement									
Homeless	112,266	22,675	19,351	2,146	20,737	36,552	5,012	27,949	33,397
Foster home/foster care	7,064	1,441	1,713	664	878	1,785	471	1,960	3,070
Residential care	57,826	7,867	9,372	1,716	9,903	12,620	3,324	24,800	18,949
Institutional setting	27,487	2,877	5,015	705	4,600	8,041	1,800	11,353	7,168
Jail/correctional facility	38,116	10,860	5,108	705	4,495	7,246	5,631	8,524	12,471
Private residence	1,426,011	312,423	399,827	59,766	267,205	518,175	64,430	242,232	336,543
Other	29,766	7,962	10,229	1,315	4,369	10,410	1,149	5,861	6,518
Total	1,698,536	366,105	450,615	67,017	312,187	594,829	81,817	322,679	418,116
SMI status²									
SMI	1,947,307	351,778	518,721	61,183	396,179	743,169	85,573	415,559	354,901
Not SMI	650,237	169,134	166,419	38,283	57,784	151,162	17,071	34,459	236,363
Total	2,597,544	520,912	685,140	99,466	453,963	894,331	102,644	450,018	591,264
Veteran status									
Veteran	50,594	9,941	7,414	1,076	8,215	18,606	1,108	8,420	7,020
Not a veteran	1,091,578	221,794	255,071	40,897	187,070	364,055	46,006	184,804	324,854
Total	1,142,172	231,735	262,485	41,973	195,285	382,661	47,114	193,224	331,874

ADHD = attention deficit/hyperactivity disorder; SMI = serious mental illness.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 4.7b. Individuals aged 25 to 54 years, by mental health diagnosis, living arrangement, SMI status, and veteran status: Column percent distribution, 2020

Living arrangement, SMI status, and veteran status	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals aged 25 to 54 years	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Living arrangement									
Homeless	6.6	6.2	4.3	3.2	6.6	6.1	6.1	8.7	8.0
Foster home/foster care	†	†	†	†	†	†	†	†	†
Residential care	3.4	2.1	2.1	2.6	3.2	2.1	4.1	7.7	4.5
Institutional setting	1.6	†	1.1	1.1	1.5	1.4	2.2	3.5	1.7
Jail/correctional facility	2.2	3.0	1.1	1.1	1.4	1.2	6.9	2.6	3.0
Private residence	84.0	85.3	88.7	89.2	85.6	87.1	78.7	75.1	80.5
Other	1.8	2.2	2.3	2.0	1.4	1.8	1.4	1.8	1.6
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
SMI status²									
SMI	75.0	67.5	75.7	61.5	87.3	83.1	83.4	92.3	60.0
Not SMI	25.0	32.5	24.3	38.5	12.7	16.9	16.6	7.7	40.0
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status									
Veteran	4.4	4.3	2.8	2.6	4.2	4.9	2.4	4.4	2.1
Not a veteran	95.6	95.7	97.2	97.4	95.8	95.1	97.6	95.6	97.9
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; SMI = serious mental illness.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 4.7c. Individuals aged 25 to 54 years, by mental health diagnosis, living arrangement, SMI status, and veteran status: Row percent distribution, 2020

Living arrangement, SMI status, and veteran status	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals aged 25 to 54 years	100.0	20.0	26.2	3.8	17.4	34.3	3.9	17.3	23.2
Living arrangement									
Homeless	100.0	20.2	17.2	1.9	18.5	32.6	4.5	24.9	29.7
Foster home/foster care	100.0	20.4	24.2	9.4	12.4	25.3	6.7	27.7	43.5
Residential care	100.0	13.6	16.2	3.0	17.1	21.8	5.7	42.9	32.8
Institutional setting	100.0	10.5	18.2	2.6	16.7	29.3	6.5	41.3	26.1
Jail/correctional facility	100.0	28.5	13.4	1.8	11.8	19.0	14.8	22.4	32.7
Private residence	100.0	21.9	28.0	4.2	18.7	36.3	4.5	17.0	23.6
Other	100.0	26.7	34.4	4.4	14.7	35.0	3.9	19.7	21.9
Total	100.0	21.6	26.5	3.9	18.4	35.0	4.8	19.0	24.6
SMI status²									
SMI	100.0	18.1	26.6	3.1	20.3	38.2	4.4	21.3	18.2
Not SMI	100.0	26.0	25.6	5.9	8.9	23.2	2.6	5.3	36.4
Total	100.0	20.1	26.4	3.8	17.5	34.4	4.0	17.3	22.8
Veteran status									
Veteran	100.0	19.6	14.7	2.1	16.2	36.8	2.2	16.6	13.9
Not a veteran	100.0	20.3	23.4	3.7	17.1	33.4	4.2	16.9	29.8
Total	100.0	20.3	23.0	3.7	17.1	33.5	4.1	16.9	29.1

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; SMI = serious mental illness.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 4.8a. Individuals aged 25 to 54 years with SMI, by mental health diagnosis, living arrangement, and veteran status: Number, 2020

Living arrangement and veteran status	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals with SMI aged 25 to 54 years²	1,947,307	351,778	518,721	61,183	396,179	743,169	85,573	415,559	354,901
Living arrangement									
Homeless	92,198	18,914	16,050	1,697	18,301	31,519	4,371	25,680	23,479
Foster home/foster care	4,544	909	1,088	388	647	1,289	399	1,646	1,638
Residential care	47,730	6,445	7,694	1,239	8,772	10,798	3,016	23,296	12,235
Institutional setting	23,180	2,360	4,400	596	4,103	7,008	1,275	10,156	4,928
Jail/correctional facility	27,963	7,100	4,069	554	3,780	5,907	5,341	7,549	8,483
Private residence	1,078,720	231,391	304,779	38,629	228,908	413,747	54,573	218,803	203,605
Other	25,114	5,820	9,030	887	4,046	9,457	1,040	5,485	5,137
Total	1,299,449	272,939	347,110	43,990	268,557	479,725	70,015	292,615	259,505
Veteran status									
Veteran	43,837	7,842	5,955	762	7,641	16,950	920	7,994	4,568
Not a veteran	770,224	152,483	182,953	25,744	156,380	284,537	36,868	162,495	175,598
Total	814,061	160,325	188,908	26,506	164,021	301,487	37,788	170,489	180,166

ADHD = attention deficit/hyperactivity disorder; SMI = serious mental illness.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 4.8b. Individuals aged 25 to 54 years with SMI, by mental health diagnosis, living arrangement, and veteran status: Column percent distribution, 2020

<i>Living arrangement and veteran status</i>	<i>Total</i>	<i>Mental health diagnosis¹</i>							
		<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>ADHD</i>	<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Total individuals with SMI aged 25 to 54 years²	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Living arrangement									
Homeless	7.1	6.9	4.6	3.9	6.8	6.6	6.2	8.8	9.0
Foster home/foster care	†	†	†	†	†	†	†	†	†
Residential care	3.7	2.4	2.2	2.8	3.3	2.3	4.3	8.0	4.7
Institutional setting	1.8	†	1.3	1.4	1.5	1.5	1.8	3.5	1.9
Jail/correctional facility	2.2	2.6	1.2	1.3	1.4	1.2	7.6	2.6	3.3
Private residence	83.0	84.8	87.8	87.8	85.2	86.2	77.9	74.8	78.5
Other	1.9	2.1	2.6	2.0	1.5	2.0	1.5	1.9	2.0
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status									
Veteran	5.4	4.9	3.2	2.9	4.7	5.6	2.4	4.7	2.5
Not a veteran	94.6	95.1	96.8	97.1	95.3	94.4	97.6	95.3	97.5
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; SMI = serious mental illness.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 4.8c. Individuals aged 25 to 54 years with SMI, by mental health diagnosis, living arrangement, and veteran status: Row percent distribution, 2020

Living arrangement and veteran status		Mental health diagnosis ¹								
		Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Total individuals with SMI aged 25 to 54 years ²		100.0	18.1	26.6	3.1	20.3	38.2	4.4	21.3	18.2
Living arrangement										
Homeless	100.0	20.5	17.4	1.8	19.8	34.2	4.7	27.9	25.5	
Foster home/foster care	100.0	20.0	23.9	8.5	14.2	28.4	8.8	36.2	36.0	
Residential care	100.0	13.5	16.1	2.6	18.4	22.6	6.3	48.8	25.6	
Institutional setting	100.0	10.2	19.0	2.6	17.7	30.2	5.5	43.8	21.3	
Jail/correctional facility	100.0	25.4	14.6	2.0	13.5	21.1	19.1	27.0	30.3	
Private residence	100.0	21.5	28.3	3.6	21.2	38.4	5.1	20.3	18.9	
Other	100.0	23.2	36.0	3.5	16.1	37.7	4.1	21.8	20.5	
Total	100.0	21.0	26.7	3.4	20.7	36.9	5.4	22.5	20.0	
Veteran status										
Veteran	100.0	17.9	13.6	1.7	17.4	38.7	2.1	18.2	10.4	
Not a veteran	100.0	19.8	23.8	3.3	20.3	36.9	4.8	21.1	22.8	
Total	100.0	19.7	23.2	3.3	20.1	37.0	4.6	20.9	22.1	

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; SMI = serious mental illness.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 4.9a. Individuals aged 25 to 54 years, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Number, 2020

Employment status and ‘detailed not in labor force’	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals aged 25 to 54 years	2,667,813	532,647	697,741	101,545	463,761	914,629	105,153	460,519	619,606
Employment status									
Employed	396,992	98,422	131,064	20,859	70,526	163,109	17,018	34,809	75,968
Full time	219,748	58,439	76,786	13,089	31,824	91,819	9,085	13,865	43,011
Part time	122,893	27,659	36,224	6,194	22,909	47,568	5,614	14,293	25,954
Unemployed	481,264	97,500	112,532	14,893	96,209	177,175	18,352	83,637	112,666
Not in labor force (see detail below)	515,690	110,545	151,500	20,761	106,841	181,623	33,029	130,083	105,322
Total	1,393,946	306,467	395,096	56,513	273,576	521,907	68,399	248,529	293,956
Detailed not in labor force									
Total individuals aged 25 to 54 years not in labor force	515,690	110,545	151,500	20,761	106,841	181,623	33,029	130,083	105,322
Disabled	231,293	44,532	62,608	8,005	51,619	73,452	18,777	78,203	50,457
Retired	3,018	778	609	50	509	1,049	84	771	580
Student	20,473	3,611	4,768	1,323	3,083	6,463	606	3,565	5,053
Homemaker	26,920	7,327	9,410	881	4,711	11,962	1,108	1,767	4,189
Sheltered/non-competitive employment	5,966	880	1,219	255	943	1,586	226	1,204	2,810
Other	228,020	53,417	72,886	10,247	45,976	87,111	12,228	44,573	42,233

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

NOTES: *Employed* includes “full-time/part-time not differentiated,” which is not presented separately; total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 4.9b. Individuals aged 25 to 54 years, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Column percent distribution, 2020

<i>Employment status and ‘detailed not in labor force’</i>	<i>Total</i>	<i>Mental health diagnosis¹</i>							
		<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>ADHD</i>	<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Total individuals aged 25 to 54 years	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Employment status									
Employed	28.5	32.1	33.2	36.9	25.8	31.3	24.9	14.0	25.8
Full time	15.8	19.1	19.4	23.2	11.6	17.6	13.3	5.6	14.6
Part time	8.8	9.0	9.2	11.0	8.4	9.1	8.2	5.8	8.8
Unemployed	34.5	31.8	28.5	26.4	35.2	33.9	26.8	33.7	38.3
Not in labor force (see detail below)	37.0	36.1	38.3	36.7	39.1	34.8	48.3	52.3	35.8
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Detailed not in labor force									
Total individuals aged 25 to 54 years not in labor force	44.2	48.3	48.1	49.4	43.0	48.0	37.0	34.3	40.1
Disabled	44.9	40.3	41.3	38.6	48.3	40.4	56.9	60.1	47.9
Retired	†	†	†	†	†	†	†	†	†
Student	4.0	3.3	3.1	6.4	2.9	3.6	1.8	2.7	4.8
Homemaker	5.2	6.6	6.2	4.2	4.4	6.6	3.4	1.4	4.0
Sheltered/non-competitive employment	1.2	†	†	1.2	†	†	†	†	2.7
Other	44.2	48.3	48.1	49.4	43.0	48.0	37.0	34.3	40.1

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding. *Employed* includes “full-time/part-time not differentiated,” which is not presented separately.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 4.9c. Individuals aged 25 to 54 years, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Row percent distribution, 2020

<i>Employment status and ‘detailed not in labor force’</i>		<i>Mental health diagnosis¹</i>							
		<i>Trauma- and stressor-related disorders</i>			<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
	<i>Total</i>		<i>Anxiety disorders</i>	<i>ADHD</i>					
Total individuals aged 25 to 54 years	100.0	20.0	26.2	3.8	17.4	34.3	3.9	17.3	23.2
Employment status									
Employed	100.0	24.8	33.0	5.3	17.8	41.1	4.3	8.8	19.1
Full time	100.0	26.6	34.9	6.0	14.5	41.8	4.1	6.3	19.6
Part time	100.0	22.5	29.5	5.0	18.6	38.7	4.6	11.6	21.1
Unemployed	100.0	20.3	23.4	3.1	20.0	36.8	3.8	17.4	23.4
Not in labor force (see detail below)	100.0	21.4	29.4	4.0	20.7	35.2	6.4	25.2	20.4
Total	100.0	22.0	28.3	4.1	19.6	37.4	4.9	17.8	21.1
Detailed not in labor force									
Total individuals aged 25 to 54 years not in labor force	100.0	21.4	29.4	4.0	20.7	35.2	6.4	25.2	20.4
Disabled	100.0	19.3	27.1	3.5	22.3	31.8	8.1	33.8	21.8
Retired	100.0	25.8	20.2	1.7	16.9	34.8	2.8	25.5	19.2
Student	100.0	17.6	23.3	6.5	15.1	31.6	3.0	17.4	24.7
Homemaker	100.0	27.2	35.0	3.3	17.5	44.4	4.1	6.6	15.6
Sheltered/non-competitive employment	100.0	14.8	20.4	4.3	15.8	26.6	3.8	20.2	47.1
Other	100.0	23.4	32.0	4.5	20.2	38.2	5.4	19.5	18.5

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding. *Employed* includes “full-time/part-time not differentiated,” which is not presented separately.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 4.10a. Individuals aged 25 to 54 years with SMI, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Number, 2020

Employment status and 'detailed not in labor force'		Mental health diagnosis ¹							
		Trauma- and stressor-related disorders			Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
	Total								
Total individuals with SMI aged 25 to 54 years ²	1,947,307	351,778	518,721	61,183	396,179	743,169	85,573	415,559	354,901
Employment status									
Employed	286,776	67,428	94,905	13,075	60,790	129,762	14,213	31,188	42,493
Full time	146,729	36,930	50,984	7,600	25,682	68,372	7,268	11,819	22,262
Part time	93,127	19,500	27,376	4,106	19,897	39,550	4,706	12,978	15,691
Unemployed	379,021	73,552	89,423	10,741	85,762	151,830	15,511	76,056	67,122
Not in labor force (see detail below)	406,348	86,664	116,340	13,864	93,193	146,230	29,016	119,024	67,147
Total	1,072,145	227,644	300,668	37,680	239,745	427,822	58,740	226,268	176,762
Detailed not in labor force									
Total individuals with SMI aged 25 to 54 years not in labor force	406,348	86,664	116,340	13,864	93,193	146,230	29,016	119,024	67,147
Disabled	199,280	38,992	53,823	6,380	47,312	64,385	17,352	72,981	35,705
Retired	2,605	648	507	38	479	922	78	737	421
Student	16,845	2,804	3,829	803	2,838	5,858	536	3,380	3,341
Homemaker	21,577	5,681	7,349	646	4,157	10,068	957	1,623	2,751
Sheltered/non-competitive employment	4,318	782	957	140	850	1,419	191	1,054	1,354
Other	161,723	37,757	49,875	5,857	37,557	63,578	9,902	39,249	23,575

ADHD = attention deficit/hyperactivity disorder; SMI = serious mental illness.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTES: *Employed* includes “full-time/part-time not differentiated,” which is not presented separately; total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 4.10b. Individuals aged 25 to 54 years with SMI, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Column percent distribution, 2020

<i>Employment status and ‘detailed not in labor force’</i>		<i>Mental health diagnosis¹</i>							
		<i>Trauma- and stressor-related disorders Anxiety disorders ADHD</i>			<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
	<i>Total</i>								
Total individuals with SMI aged 25 to 54 years²	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Employment status									
Employed	26.7	29.6	31.6	34.7	25.4	30.3	24.2	13.8	24.0
Full time	13.7	16.2	17.0	20.2	10.7	16.0	12.4	5.2	12.6
Part time	8.7	8.6	9.1	10.9	8.3	9.2	8.0	5.7	8.9
Unemployed	35.4	32.3	29.7	28.5	35.8	35.5	26.4	33.6	38.0
Not in labor force (see detail below)	37.9	38.1	38.7	36.8	38.9	34.2	49.4	52.6	38.0
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Detailed not in labor force									
Total individuals with SMI aged 25 to 54 years not in labor force	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Disabled	49.0	45.0	46.3	46.0	50.8	44.0	59.8	61.3	53.2
Retired	†	†	†	†	†	†	†	†	†
Student	4.1	3.2	3.3	5.8	3.0	4.0	1.8	2.8	5.0
Homemaker	5.3	6.6	6.3	4.7	4.5	6.9	3.3	1.4	4.1
Sheltered/non-competitive employment	1.1	†	†	1.0	†	†	†	†	2.0
Other	39.8	43.6	42.9	42.2	40.3	43.5	34.1	33.0	35.1

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; SMI = serious mental illness.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding. *Employed* includes “full-time/part-time not differentiated,” which is not presented separately.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 4.10c. Individuals aged 25 to 54 years with SMI, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Row percent distribution, 2020

Employment status and ‘detailed not in labor force’		Mental health diagnosis ¹							
		Trauma- and stressor-related disorders			Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total									
Total individuals with SMI aged 25 to 54 years ²	100.0	18.1	26.6	3.1	20.3	38.2	4.4	21.3	18.2
Employment status									
Employed	100.0	23.5	33.1	4.6	21.2	45.2	5.0	10.9	14.8
Full time	100.0	25.2	34.7	5.2	17.5	46.6	5.0	8.1	15.2
Part time	100.0	20.9	29.4	4.4	21.4	42.5	5.1	13.9	16.8
Unemployed	100.0	19.4	23.6	2.8	22.6	40.1	4.1	20.1	17.7
Not in labor force (see detail below)	100.0	21.3	28.6	3.4	22.9	36.0	7.1	29.3	16.5
Total	100.0	21.2	28.0	3.5	22.4	39.9	5.5	21.1	16.5
Detailed not in labor force									
Total individuals with SMI aged 25 to 54 years not in labor force	100.0	21.3	28.6	3.4	22.9	36.0	7.1	29.3	16.5
Disabled	100.0	19.6	27.0	3.2	23.7	32.3	8.7	36.6	17.9
Retired	100.0	24.9	19.5	1.5	18.4	35.4	3.0	28.3	16.2
Student	100.0	16.6	22.7	4.8	16.8	34.8	3.2	20.1	19.8
Homemaker	100.0	26.3	34.1	3.0	19.3	46.7	4.4	7.5	12.7
Sheltered/non-competitive employment	100.0	18.1	22.2	3.2	19.7	32.9	4.4	24.4	31.4
Other	100.0	23.3	30.8	3.6	23.2	39.3	6.1	24.3	14.6

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; SMI = serious mental illness.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding. *Employed* includes “full-time/part-time not differentiated,” which is not presented separately.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 4.11a. Individuals aged 25 to 54 years, by mental health diagnosis, timing of admission, and mental health service setting: Number, 2020

Timing of admission and mental health service setting	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals aged 25 to 54 years	2,667,813	532,647	697,741	101,545	463,761	914,629	105,153	460,519	619,606
Timing of admission									
New individuals	1,182,230	260,601	328,756	37,011	180,024	432,658	38,334	149,345	254,758
Continuing individuals	1,485,583	272,046	368,985	64,534	283,737	481,971	66,819	311,174	364,848
Mental health service setting^{2,3}									
State psychiatric hospital	64,905	5,993	5,695	851	11,965	10,893	6,325	36,789	17,274
Community-based program	2,578,850	522,393	692,050	100,540	451,116	892,217	100,897	430,015	589,933
Residential treatment center	28,074	3,299	3,760	532	5,094	6,811	1,377	11,557	8,167
Other psychiatric inpatient	138,453	17,594	20,240	2,523	27,648	50,376	5,647	39,853	28,120
Institutions under the justice system	38,594	12,171	4,093	510	2,553	5,559	5,937	7,495	15,847

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Individuals could receive services in more than one service setting during the reporting period.

³ Mental health service setting category definitions can be found in Appendix C.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 4.11b. Individuals aged 25 to 54 years, by mental health diagnosis, timing of admission, and mental health service setting: Column percent distribution, 2020

Timing of admission and mental health service setting	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals aged 25 to 54 years	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Timing of admission									
New individuals	44.3	48.9	47.1	36.4	38.8	47.3	36.5	32.4	41.1
Continuing individuals	55.7	51.1	52.9	63.6	61.2	52.7	63.5	67.6	58.9
Mental health service setting^{2,3}									
State psychiatric hospital	2.4	1.1	†	†	2.6	1.2	6.0	8.0	2.8
Community-based program	96.7	98.1	99.2	99.0	97.3	97.5	96.0	93.4	95.2
Residential treatment center	1.1	†	†	†	1.1	†	1.3	2.5	1.3
Other psychiatric inpatient	5.2	3.3	2.9	2.5	6.0	5.5	5.4	8.7	4.5
Institutions under the justice system	1.4	2.3	†	†	†	†	5.6	1.6	2.6

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Percentages may not sum to 100 percent, because individuals could receive services in more than one service setting during the reporting period.

³ Mental health service setting category definitions can be found in Appendix C.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 4.11c. Individuals aged 25 to 54 years, by mental health diagnosis, timing of admission, and mental health service setting: Row percent distribution, 2020

Timing of admission and mental health service setting	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals aged 25 to 54 years	100.0	20.0	26.2	3.8	17.4	34.3	3.9	17.3	23.2
Timing of admission									
New individuals	100.0	22.0	27.8	3.1	15.2	36.6	3.2	12.6	21.5
Continuing individuals	100.0	18.3	24.8	4.3	19.1	32.4	4.5	20.9	24.6
Mental health service setting^{2,3}									
State psychiatric hospital	100.0	9.2	8.8	1.3	18.4	16.8	9.7	56.7	26.6
Community-based program	100.0	20.3	26.8	3.9	17.5	34.6	3.9	16.7	22.9
Residential treatment center	100.0	11.8	13.4	1.9	18.1	24.3	4.9	41.2	29.1
Other psychiatric inpatient	100.0	12.7	14.6	1.8	20.0	36.4	4.1	28.8	20.3
Institutions under the justice system	100.0	31.5	10.6	1.3	6.6	14.4	15.4	19.4	41.1

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Individuals could receive services in more than one service setting during the reporting period.

³ Mental health service setting category definitions can be found in Appendix C.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 4.12a. Individuals aged 25 to 54 years with SMI, by mental health diagnosis, timing of admission, and mental health service setting: Number, 2020

Timing of admission and mental health service setting	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals with SMI aged 25 to 54 years ²	1,947,307	351,778	518,721	61,183	396,179	743,169	85,573	415,559	354,901
Timing of admission									
New individuals	845,104	168,902	243,172	22,902	153,945	354,721	29,534	130,126	142,997
Continuing individuals	1,102,203	182,876	275,549	38,281	242,234	388,448	56,039	285,433	211,904
Mental health service setting ^{3,4}									
State psychiatric hospital	51,182	4,627	4,306	618	9,797	8,353	5,142	31,843	12,199
Community-based program	1,873,409	343,397	514,214	60,457	385,829	723,684	82,346	389,018	332,435
Residential treatment center	23,890	2,715	3,217	418	4,498	5,914	1,218	10,771	6,052
Other psychiatric inpatient	121,026	14,638	18,712	2,191	25,382	46,648	5,189	37,426	20,052
Institutions under the justice system	32,341	8,810	3,693	449	2,361	5,119	5,826	7,274	13,402

ADHD = attention deficit/hyperactivity disorder; SMI = serious mental illness.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

³ Individuals could receive services in more than one service setting during the reporting period.

⁴ Mental health service setting category definitions can be found in Appendix C.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 4.12b. Individuals aged 25 to 54 years with SMI, by mental health diagnosis, timing of admission, and mental health service setting: Column percent, 2020

Timing of admission and mental health service setting	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals with SMI aged 25 to 54 years²	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Timing of admission									
New individuals	43.4	48.0	46.9	37.4	38.9	47.7	34.5	31.3	40.3
Continuing individuals	56.6	52.0	53.1	62.6	61.1	52.3	65.5	68.7	59.7
Mental health service setting^{3,4}									
State psychiatric hospital	2.6	1.3	†	1.0	2.5	1.1	6.0	7.7	3.4
Community-based program	96.2	97.6	99.1	98.8	97.4	97.4	96.2	93.6	93.7
Residential treatment center	1.2	†	†	†	1.1	†	1.4	2.6	1.7
Other psychiatric inpatient	6.2	4.2	3.6	3.6	6.4	6.3	6.1	9.0	5.7
Institutions under the justice system	1.7	2.5	†	†	†	†	6.8	1.8	3.8

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; SMI = serious mental illness.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

³ Percentages may not sum to 100 percent, because individuals could receive services in more than one service setting during the reporting period.

⁴ Mental health service setting category definitions can be found in Appendix C.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 4.12c. Individuals aged 25 to 54 years with SMI, by mental health diagnosis, timing of admission, and mental health service setting: Row percent distribution, 2020

Timing of admission and mental health service setting	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals with SMI aged 25 to 54 years ²	100.0	18.1	26.6	3.1	20.3	38.2	4.4	21.3	18.2
Timing of admission									
New individuals	100.0	20.0	28.8	2.7	18.2	42.0	3.5	15.4	16.9
Continuing individuals	100.0	16.6	25.0	3.5	22.0	35.2	5.1	25.9	19.2
Mental health service setting ^{3,4}									
State psychiatric hospital	100.0	9.0	8.4	1.2	19.1	16.3	10.0	62.2	23.8
Community-based program	100.0	18.3	27.4	3.2	20.6	38.6	4.4	20.8	17.7
Residential treatment center	100.0	11.4	13.5	1.7	18.8	24.8	5.1	45.1	25.3
Other psychiatric inpatient	100.0	12.1	15.5	1.8	21.0	38.5	4.3	30.9	16.6
Institutions under the justice system	100.0	27.2	11.4	1.4	7.3	15.8	18.0	22.5	41.4

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; SMI = serious mental illness.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

³ Individuals could receive services in more than one service setting during the reporting period.

⁴ Mental health service setting category definitions can be found in Appendix C.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 4.13. Individuals aged 25 to 54 years, by mental health service setting and SMI status: Number and percent distribution, 2020

<i>Mental health service setting</i>	<i>SMI status</i> ¹								
	<i>Number of individuals</i>			<i>Column percent of individuals</i>			<i>Row percent distribution of individuals</i>		
	<i>Total</i>	<i>SMI</i>	<i>Not SMI</i>	<i>Total</i>	<i>SMI</i>	<i>Not SMI</i>	<i>Total</i>	<i>SMI</i>	<i>Not SMI</i>
Total individuals aged 25 to 54 years	2,980,461	2,164,629	815,832	100.0	100.0	100.0	100.0	72.6	27.4
Mental health service setting^{2,3}									
State psychiatric hospital	63,827	56,372	7,455	2.1	2.6	†	100.0	88.3	11.7
Community-based program	2,886,446	2,079,516	806,930	96.8	96.1	98.9	100.0	72.0	28.0
Residential treatment center	34,458	29,858	4,600	1.2	1.4	†	100.0	86.7	13.3
Other psychiatric inpatient	149,624	132,634	16,990	5.0	6.1	2.1	100.0	88.6	11.4
Institutions under the justice system	40,939	34,831	6,108	1.4	1.6	†	100.0	85.1	14.9

† Less than 1 percent of individuals.

SMI = serious emotional disturbance.

¹ The federal definition for SMI status can be found in Appendix D.

² Individuals could receive services in more than one mental health service setting during the reporting period.

³ Mental health service setting category definitions can be found in Appendix C.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 5.1a. Individuals aged 55 years and older, by gender, age, race, ethnicity, education, veteran status, and SMI status: Number, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Number of states reporting	45	47	49	49	48	50
Total individuals aged 55 years and older	781,018	871,999	1,019,548	1,046,659	1,078,578	1,178,713
Gender						
Male	320,355	359,807	428,257	440,285	455,336	499,199
Female	459,352	509,527	586,930	604,806	621,280	676,894
Total	779,707	869,334	1,015,187	1,045,091	1,076,616	1,176,093
Age¹						
55 to 59 years	342,046	377,066	418,358	429,411	429,053	451,653
60 to 64 years	215,213	242,842	278,831	297,124	307,640	336,524
65 years and older	223,759	252,091	322,359	320,124	341,885	390,536
Total	781,018	871,999	1,019,548	1,046,659	1,078,578	1,178,713
Race						
American Indian or Alaska Native	7,145	7,933	9,160	10,146	11,594	15,543
Asian ²	14,166	16,369	18,254	20,308	19,577	19,923
Black or African American	136,570	155,274	184,016	195,721	206,917	220,420
Native Hawaiian or Other Pacific Islander	1,935	2,214	1,317	1,575	1,868	1,930
White	514,386	575,530	655,447	680,170	689,714	750,151
Other	37,330	43,541	55,499	59,968	63,987	69,464
Total	711,532	800,861	923,693	967,888	993,657	1,077,431
Ethnicity						
Not of Hispanic or Latino origin	523,793	588,211	803,827	836,447	857,721	916,771
Hispanic or Latino origin	64,105	72,745	98,715	99,334	102,831	152,404
Total	587,898	660,956	902,542	935,781	960,552	1,069,175
Specific Hispanic or Latino origin						
Mexican	2,854	2,568	2,976	3,013	3,136	3,022
Puerto Rican	11,020	9,775	12,832	5,015	5,400	5,541
Cuban	3,312	3,161	3,076	3,734	3,469	4,194
Other specific	5,738	6,278	6,944	7,370	5,657	6,190
Total	22,924	21,782	25,828	19,132	17,662	18,947

Continued. See notes at end of table.

Table 5.1a. Individuals aged 55 years and older, by gender, age, race, ethnicity, education, veteran status, and SMI status: Number, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Highest school grade completed						
0 to 8	63,840	65,476	63,672	62,348	61,750	62,425
9 to 11	56,867	60,012	74,128	78,492	80,153	89,497
12 (or GED)	180,488	194,742	226,992	252,629	256,583	285,256
More than 12	77,395	85,120	126,035	139,354	142,371	161,153
Total	378,590	405,350	490,827	532,823	540,857	598,331
Veteran status						
Veteran	<i>n/c</i>	<i>n/c</i>	16,313	16,893	27,392	32,010
Not a veteran	<i>n/c</i>	<i>n/c</i>	297,222	445,461	441,660	471,476
Total	<i>n/c</i>	<i>n/c</i>	313,535	462,354	469,052	503,486
SMI status³						
SMI	551,786	602,110	696,724	748,945	781,925	824,495
Not SMI	193,842	223,215	264,483	254,464	246,630	300,380
Total	745,628	825,325	961,207	1,003,409	1,028,555	1,124,875

n/c Not collected.

GED = general equivalency diploma; SMI = serious mental illness.

¹ Age is based on the individual's age at the midpoint of the reporting period.

² For some states, "Asian" includes "Pacific Islanders." See Appendix C.

³ The federal definition for SMI status can be found in Appendix D.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 5.1b. Individuals aged 55 years and older, by gender, age, race, ethnicity, education, veteran status, and SMI status: Column percent distribution, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Number of states reporting	45	47	49	49	48	50
Total individuals aged 55 years and older	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	41.1	41.4	42.2	42.1	42.3	42.4
Female	58.9	58.6	57.8	57.9	57.7	57.6
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age¹						
55 to 59 years	43.8	43.2	41.0	41.0	39.8	38.3
60 to 64 years	27.6	27.8	27.3	28.4	28.5	28.6
65 years and older	28.6	28.9	31.6	30.6	31.7	33.1
Total	100.0	100.0	100.0	100.0	100.0	100.0
Race						
American Indian or Alaska Native	1.0	†	†	1.0	1.2	1.4
Asian ²	2.0	2.0	2.0	2.1	2.0	1.8
Black or African American	19.2	19.4	19.9	20.2	20.8	20.5
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	72.3	71.9	71.0	70.3	69.4	69.6
Other	5.2	5.4	6.0	6.2	6.4	6.4
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	89.1	89.0	89.1	89.4	89.3	85.7
Hispanic or Latino origin	10.9	11.0	10.9	10.6	10.7	14.3
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	12.4	11.8	11.5	15.7	17.8	15.9
Puerto Rican	48.1	44.9	49.7	26.2	30.6	29.2
Cuban	14.4	14.5	11.9	19.5	19.6	22.1
Other specific	25.0	28.8	26.9	38.5	32.0	32.7
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 5.1b. Individuals aged 55 years and older, by gender, age, race, ethnicity, education, veteran status, and SMI status: Column percent distribution, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Highest school grade completed						
0 to 8	16.9	16.2	13.0	11.7	11.4	10.4
9 to 11	15.0	14.8	15.1	14.7	14.8	15.0
12 (or GED)	47.7	48.0	46.2	47.4	47.4	47.7
More than 12	20.4	21.0	25.7	26.2	26.3	26.9
Total	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status						
Veteran	<i>n/c</i>	<i>n/c</i>	5.2	3.7	5.8	6.4
Not a veteran	<i>n/c</i>	<i>n/c</i>	94.8	96.3	94.2	93.6
Total	<i>n/c</i>	<i>n/c</i>	100.0	100.0	100.0	100.0
SMI status³						
SMI	74.0	73.0	72.5	74.6	76.0	73.3
Not SMI	26.0	27.0	27.5	25.4	24.0	26.7
Total	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals; *n/c* Not collected.

GED = general equivalency diploma; SMI = serious mental illness.

¹ Age is based on the individual's age at the midpoint of the reporting period.

² For some states, "Asian" includes "Pacific Islanders." See Appendix C.

³ The federal definition for SMI status can be found in Appendix D.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 5.2a. Individuals aged 55 years and older with SMI, by gender, age, race, ethnicity, education, and veteran status: Number, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Number of states reporting	45	46	47	48	46	49
Total individuals with SMI aged 55 years and older¹	551,786	602,110	696,724	748,945	781,925	824,495
Gender						
Male	225,104	246,881	290,889	313,493	327,051	344,982
Female	326,036	354,522	405,142	434,854	454,074	478,834
Total	551,140	601,403	696,031	748,347	781,125	823,816
Age²						
55 to 59 years	250,165	270,694	299,467	320,124	326,348	330,839
60 to 64 years	155,747	172,454	196,797	218,031	230,008	243,086
65 years and older	145,874	158,962	200,460	210,790	225,569	250,570
Total	551,786	602,110	696,724	748,945	781,925	824,495
Race						
American Indian or Alaska Native	4,499	5,017	6,129	6,840	7,962	9,419
Asian ³	11,155	11,650	13,312	14,979	14,842	14,566
Black or African American	104,275	113,681	132,883	146,994	154,898	163,741
Native Hawaiian or Other Pacific Islander	1,702	1,874	1,115	1,334	1,621	1,556
White	353,186	392,267	448,796	479,324	495,054	518,339
Other	29,581	32,747	43,090	48,020	52,264	55,395
Total	504,398	557,236	645,325	697,491	726,641	763,016
Ethnicity						
Not of Hispanic or Latino origin	387,324	426,268	541,998	584,307	606,111	638,038
Hispanic or Latino origin	50,253	57,012	74,399	78,009	81,346	110,651
Total	437,577	483,280	616,397	662,316	687,457	748,689
Specific Hispanic or Latino origin						
Mexican	2,287	1,994	2,375	2,420	2,550	2,439
Puerto Rican	8,377	7,541	9,390	4,176	4,439	4,784
Cuban	3,200	3,052	2,954	3,639	3,324	4,050
Other specific	4,907	5,287	5,764	6,325	4,539	5,009
Total	18,771	17,874	20,483	16,560	14,852	16,282

Continued. See notes at end of table.

Table 5.2a. Individuals aged 55 years and older with SMI, by gender, age, race, ethnicity, education, and veteran status: Number, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Highest school grade completed						
0 to 8	39,968	41,557	44,471	44,204	44,204	44,181
9 to 11	42,874	46,384	56,832	60,547	61,761	67,551
12 (or GED)	121,956	136,267	161,434	184,119	185,399	202,774
More than 12	60,861	69,164	90,180	100,157	101,643	110,367
Total	265,659	293,372	352,917	389,027	393,007	424,873
Veteran status						
Veteran	<i>n/c</i>	<i>n/c</i>	11,077	13,102	19,996	23,598
Not a veteran	<i>n/c</i>	<i>n/c</i>	189,251	299,219	311,623	322,458
Total	<i>n/c</i>	<i>n/c</i>	200,328	312,321	331,619	346,056

n/c Not collected.

GED = general equivalency diploma; SMI = serious mental illness.

¹ The federal definition for SMI status can be found in Appendix D.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 5.2b. Individuals aged 55 years and older with SMI, by gender, age, race, ethnicity, education, and veteran status: Column percent distribution, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Number of states reporting	45	46	47	48	46	49
Total individuals with SMI aged 55 years and older¹	100	100	100	100	100	100
Gender						
Male	40.8	41.1	41.8	41.9	41.9	41.9
Female	59.2	58.9	58.2	58.1	58.1	58.1
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age²						
55 to 59 years	45.3	45.0	43.0	42.7	41.7	40.1
60 to 64 years	28.2	28.6	28.2	29.1	29.4	29.5
65 years and older	26.4	26.4	28.8	28.1	28.8	30.4
Total	100.0	100.0	100.0	100.0	100.0	100.0
Race						
American Indian or Alaska Native	†	†	†	†	1.1	1.2
Asian ³	2.2	2.1	2.1	2.1	2.0	1.9
Black or African American	20.7	20.4	20.6	21.1	21.3	21.5
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	70.0	70.4	69.5	68.7	68.1	67.9
Other	5.9	5.9	6.7	6.9	7.2	7.3
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	88.5	88.2	87.9	88.2	88.2	85.2
Hispanic or Latino origin	11.5	11.8	12.1	11.8	11.8	14.8
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	12.2	11.2	11.6	14.6	17.2	15.0
Puerto Rican	44.6	42.2	45.8	25.2	29.9	29.4
Cuban	17.0	17.1	14.4	22.0	22.4	24.9
Other specific	26.1	29.6	28.1	38.2	30.6	30.8
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 5.2b. Individuals aged 55 years and older with SMI, by gender, age, race, ethnicity, education, and veteran status: Column percent distribution, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Highest school grade completed						
0 to 8	15.0	14.2	12.6	11.4	11.2	10.4
9 to 11	16.1	15.8	16.1	15.6	15.7	15.9
12 (or GED)	45.9	46.4	45.7	47.3	47.2	47.7
More than 12	22.9	23.6	25.6	25.7	25.9	26.0
Total	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status						
Veteran	<i>n/c</i>	<i>n/c</i>	5.5	4.2	6.0	6.8
Not a veteran	<i>n/c</i>	<i>n/c</i>	94.5	95.8	94.0	93.2
Total	<i>n/c</i>	<i>n/c</i>	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals; *n/c* Not collected.

GED = general equivalency diploma; SMI = serious mental illness.

¹ The federal definition for SMI status can be found in Appendix D.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 5.3a. Individuals aged 55 years and older, by mental health diagnosis, gender, race, and ethnicity: Number, 2020

Gender, race, and ethnicity	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	Delirium/dementia	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals aged 55 years and older	1,010,080	140,825	212,400	22,719	148,935	394,254	35,333	236,477	224,006
Gender									
Male	420,602	50,023	69,463	9,929	53,012	140,744	14,037	125,434	110,376
Female	587,812	90,544	142,644	12,768	95,777	252,897	21,265	110,815	113,181
Total	1,008,414	140,567	212,107	22,697	148,789	393,641	35,302	236,249	223,557
Race									
American Indian or Alaska Native	13,953	2,617	3,194	610	1,457	5,209	395	2,269	3,879
Asian ²	16,647	2,771	2,682	438	1,191	7,663	221	4,681	2,573
Black or African American	189,901	22,685	23,746	3,153	21,406	65,533	5,759	72,803	40,298
Native Hawaiian or Other Pacific Islander	1,629	226	296	54	190	650	40	494	266
White	645,748	93,462	155,388	14,735	108,556	257,540	25,782	127,309	143,370
Other	62,272	8,212	11,682	1,075	7,785	27,211	1,500	14,791	10,507
Total	930,150	129,973	196,988	20,065	140,585	363,806	33,697	222,347	200,893
Ethnicity									
Not of Hispanic or Latino origin	787,527	112,484	167,483	15,970	122,563	307,071	30,290	190,916	172,886
Hispanic or Latino origin	134,443	17,905	29,218	4,053	15,210	56,392	2,977	24,185	31,570
Total	921,970	130,389	196,701	20,023	137,773	363,463	33,267	215,101	204,456
Specific Hispanic or Latino origin									
Mexican	2,898	584	610	28	394	1,218	63	608	398
Puerto Rican	4,705	498	825	52	652	1,947	83	1,459	506
Cuban	3,848	200	483	27	369	1,721	29	609	769
Other specific	5,724	962	1,214	90	838	2,691	153	1,042	687
Total	17,175	2,244	3,132	197	2,253	7,577	328	3,718	2,360

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 5.3b. Individuals aged 55 years and older, by mental health diagnosis, gender, race, and ethnicity: Column percent distribution, 2020

Gender, race, and ethnicity	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	Delirium/dementia	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals aged 55 years and older	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender									
Male	41.7	35.6	32.7	43.7	35.6	35.8	39.8	53.1	49.4
Female	58.3	64.4	67.3	56.3	64.4	64.2	60.2	46.9	50.6
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race									
American Indian or Alaska Native	1.5	2.0	1.6	3.0	1.0	1.4	1.2	1.0	1.9
Asian ²	1.8	2.1	1.4	2.2	†	2.1	†	2.1	1.3
Black or African American	20.4	17.5	12.1	15.7	15.2	18.0	17.1	32.7	20.1
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	†	†	†
White	69.4	71.9	78.9	73.4	77.2	70.8	76.5	57.3	71.4
Other	6.7	6.3	5.9	5.4	5.5	7.5	4.5	6.7	5.2
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity									
Not of Hispanic or Latino origin	85.4	86.3	85.1	79.8	89.0	84.5	91.1	88.8	84.6
Hispanic or Latino origin	14.6	13.7	14.9	20.2	11.0	15.5	8.9	11.2	15.4
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	16.9	26.0	19.5	14.2	17.5	16.1	19.2	16.4	16.9
Puerto Rican	27.4	22.2	26.3	26.4	28.9	25.7	25.3	39.2	21.4
Cuban	22.4	8.9	15.4	13.7	16.4	22.7	8.8	16.4	32.6
Other specific	33.3	42.9	38.8	45.7	37.2	35.5	46.6	28.0	29.1
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 5.3c. Individuals aged 55 years and older, by mental health diagnosis, gender, race, and ethnicity: Row percent distribution, 2020

Gender, race, and ethnicity	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	Delirium/dementia	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals aged 55 years and older	100.0	13.9	21.0	2.2	14.7	39.0	3.5	23.4	22.2
Gender									
Male	100.0	11.9	16.5	2.4	12.6	33.5	3.3	29.8	26.2
Female	100.0	15.4	24.3	2.2	16.3	43.0	3.6	18.9	19.3
Total	100.0	13.9	21.0	2.3	14.8	39.0	3.5	23.4	22.2
Race									
American Indian or Alaska Native	100.0	18.8	22.9	4.4	10.4	37.3	2.8	16.3	27.8
Asian ²	100.0	16.6	16.1	2.6	7.2	46.0	1.3	28.1	15.5
Black or African American	100.0	11.9	12.5	1.7	11.3	34.5	3.0	38.3	21.2
Native Hawaiian or Other Pacific Islander	100.0	13.9	18.2	3.3	11.7	39.9	2.5	30.3	16.3
White	100.0	14.5	24.1	2.3	16.8	39.9	4.0	19.7	22.2
Other	100.0	13.2	18.8	1.7	12.5	43.7	2.4	23.8	16.9
Total	100.0	14.0	21.2	2.2	15.1	39.1	3.6	23.9	21.6
Ethnicity									
Not of Hispanic or Latino origin	100.0	14.3	21.3	2.0	15.6	39.0	3.8	24.2	22.0
Hispanic or Latino origin	100.0	13.3	21.7	3.0	11.3	41.9	2.2	18.0	23.5
Total	100.0	14.1	21.3	2.2	14.9	39.4	3.6	23.3	22.2
Specific Hispanic or Latino origin									
Mexican	100.0	20.2	21.0	†	13.6	42.0	2.2	21.0	13.7
Puerto Rican	100.0	10.6	17.5	1.1	13.9	41.4	1.8	31.0	10.8
Cuban	100.0	5.2	12.6	†	9.6	44.7	†	15.8	20.0
Other specific	100.0	16.8	21.2	1.6	14.6	47.0	2.7	18.2	12.0
Total	100.0	13.1	18.2	1.1	13.1	44.1	1.9	21.6	13.7

† Less than 1 percent of individuals.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 5.4a. Individuals aged 55 years and older with SMI, by mental health diagnosis, gender, race, and ethnicity: Number, 2020

Gender, race, and ethnicity	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	Delirium/ dementia	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals with SMI aged 55 years and older²	744,318	86,667	156,372	13,273	125,391	313,555	27,018	208,978	126,612
Gender									
Male	308,648	29,267	50,486	5,947	44,214	112,400	10,715	111,295	61,735
Female	435,074	57,335	105,800	7,316	81,095	200,893	16,289	97,530	64,739
Total	743,722	86,602	156,286	13,263	125,309	313,293	27,004	208,825	126,474
Race									
American Indian or Alaska Native	8,818	1,595	2,142	140	1,147	3,966	283	1,898	1,790
Asian ³	13,574	2,007	2,099	297	1,049	6,605	178	4,346	1,522
Black or African American	150,577	14,903	18,557	2,204	18,566	54,657	4,690	65,199	25,014
Native Hawaiian or Other Pacific Islander	1,391	174	245	43	164	594	37	473	191
White	466,437	57,022	113,944	8,884	90,872	201,106	19,575	111,276	79,771
Other	52,649	6,115	9,828	779	6,861	24,282	1,303	13,810	7,137
Total	693,446	81,816	146,815	12,347	118,659	291,210	26,066	197,002	115,425
Ethnicity									
Not of Hispanic or Latino origin	578,203	70,526	123,708	10,578	101,766	238,483	23,204	167,487	99,191
Hispanic or Latino origin	101,083	10,112	21,472	1,587	13,976	49,162	2,323	22,383	16,599
Total	679,286	80,638	145,180	12,165	115,742	287,645	25,527	189,870	115,790
Specific Hispanic or Latino origin									
Mexican	2,379	403	459	22	371	1,104	55	571	265
Puerto Rican	4,247	418	724	45	624	1,801	73	1,426	342
Cuban	3,740	185	456	22	356	1,681	28	594	748
Other specific	4,829	699	979	61	767	2,460	135	962	467
Total	15,195	1,705	2,618	150	2,118	7,046	291	3,553	1,822

SMI = serious mental illness.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 5.4b. Individuals aged 55 years and older with SMI, by mental health diagnosis, gender, race, and ethnicity: Column percent distribution, 2020

Gender, race, and ethnicity	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	Delirium/ dementia	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals with SMI aged 55 years and older²	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender									
Male	41.5	33.8	32.3	44.8	35.3	35.9	39.7	53.3	48.8
Female	58.5	66.2	67.7	55.2	64.7	64.1	60.3	46.7	51.2
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race									
American Indian or Alaska Native	1.3	1.9	1.5	1.1	†	1.4	1.1	†	1.6
Asian ³	2.0	2.5	1.4	2.4	†	2.3	†	2.2	1.3
Black or African American	21.7	18.2	12.6	17.9	15.6	18.8	18.0	33.1	21.7
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	†	†	†
White	67.3	69.7	77.6	72.0	76.6	69.1	75.1	56.5	69.1
Other	7.6	7.5	6.7	6.3	5.8	8.3	5.0	7.0	6.2
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity									
Not of Hispanic or Latino origin	85.1	87.5	85.2	87.0	87.9	82.9	90.9	88.2	85.7
Hispanic or Latino origin	14.9	12.5	14.8	13.0	12.1	17.1	9.1	11.8	14.3
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	15.7	23.6	17.5	14.7	17.5	15.7	18.9	16.1	14.5
Puerto Rican	27.9	24.5	27.7	30.0	29.5	25.6	25.1	40.1	18.8
Cuban	24.6	10.9	17.4	14.7	16.8	23.9	9.6	16.7	41.1
Other specific	31.8	41.0	37.4	40.7	36.2	34.9	46.4	27.1	25.6
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

SMI = serious mental illness.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 5.4c. Individuals aged 55 years and older with SMI, by mental health diagnosis, gender, race, and ethnicity: Row percent distribution, 2020

		Mental health diagnosis ¹								
Gender, race, and ethnicity		Total	Trauma- and stressor-related disorders	Anxiety disorders	Delirium/ dementia	Mood disorders		Personality disorders	Schizophrenia, other psychotics disorders	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Total individuals with SMI aged 55 years and older ²		100.0	11.6	21.0	1.8	16.8	42.1	3.6	28.1	17.0
Gender										
Male		100.0	9.5	16.4	1.9	14.3	36.4	3.5	36.1	20.0
Female		100.0	13.2	24.3	1.7	18.6	46.2	3.7	22.4	14.9
Total		100.0	11.6	21.0	1.8	16.8	42.1	3.6	28.1	17.0
Race										
American Indian or Alaska Native		100.0	18.1	24.3	1.6	13.0	45.0	3.2	21.5	20.3
Asian ³		100.0	14.8	15.5	2.2	7.7	48.7	1.3	32.0	11.2
Black or African American		100.0	9.9	12.3	1.5	12.3	36.3	3.1	43.3	16.6
Native Hawaiian or Other Pacific Islander		100.0	12.5	17.6	3.1	11.8	42.7	2.7	34.0	13.7
White		100.0	12.2	24.4	1.9	19.5	43.1	4.2	23.9	17.1
Other		100.0	11.6	18.7	1.5	13.0	46.1	2.5	26.2	13.6
Total		100.0	11.8	21.2	1.8	17.1	42.0	3.8	28.4	16.6
Ethnicity										
Not of Hispanic or Latino origin		100.0	12.2	21.4	1.8	17.6	41.2	4.0	29.0	17.2
Hispanic or Latino origin		100.0	10.0	21.2	1.6	13.8	48.6	2.3	22.1	16.4
Total		100.0	11.9	21.4	1.8	17.0	42.3	3.8	28.0	17.0
Specific Hispanic or Latino origin										
Mexican		100.0	16.9	19.3	†	15.6	46.4	2.3	24.0	11.1
Puerto Rican		100.0	9.8	17.0	1.1	14.7	42.4	1.7	33.6	8.1
Cuban		100.0	4.9	12.2	†	9.5	44.9	†	15.9	20.0
Other specific		100.0	14.5	20.3	1.3	15.9	50.9	2.8	19.9	9.7
Total		100.0	11.2	17.2	†	13.9	46.4	1.9	23.4	12.0

† Less than 1 percent of individuals.

SMI = serious mental illness.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 5.5a. Individuals aged 55 years and older, by mental health diagnosis, education, and marital status: Number, 2020

<i>Education and marital status</i>	<i>Total</i>	<i>Mental health diagnosis¹</i>							
		<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>Delirium/ dementia</i>	<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Total individuals aged 55 years and older	1,010,080	140,825	212,400	22,719	148,935	394,254	35,333	236,477	224,006
Highest school grade completed									
0 to 8	56,275	9,044	12,975	1,326	5,812	24,011	1,934	14,723	14,942
9 to 11	80,577	10,853	15,687	1,242	11,475	31,208	3,687	26,058	20,474
12 (or GED)	254,616	38,753	55,798	3,901	40,449	102,851	11,732	63,044	62,340
More than 12	146,322	26,077	34,476	1,688	29,098	60,132	7,716	28,164	31,604
Total	537,790	84,727	118,936	8,157	86,834	218,202	25,069	131,989	129,360
Marital status									
Never married	243,304	23,966	36,400	5,115	31,739	73,933	8,402	87,291	64,459
Currently married	119,017	19,573	25,852	2,967	17,642	49,742	3,583	15,459	31,169
Separated	42,129	6,910	7,635	491	7,170	17,708	1,631	8,240	9,646
Divorced	146,359	21,785	32,099	2,469	29,423	61,027	6,934	29,780	29,591
Widowed	53,594	8,507	11,601	3,185	7,963	24,290	1,562	8,802	10,263
Total	604,403	80,741	113,587	14,227	93,937	226,700	22,112	149,572	145,128

GED = general equivalency diploma.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 5.5b. Individuals aged 55 years and older, by mental health diagnosis, education, and marital status: Column percent distribution, 2020

Education and marital status	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	Delirium/ dementia	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals aged 55 years and older	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Highest school grade completed									
0 to 8	10.5	10.7	10.9	16.3	6.7	11.0	7.7	11.2	11.6
9 to 11	15.0	12.8	13.2	15.2	13.2	14.3	14.7	19.7	15.8
12 (or GED)	47.3	45.7	46.9	47.8	46.6	47.1	46.8	47.8	48.2
More than 12	27.2	30.8	29.0	20.7	33.5	27.6	30.8	21.3	24.4
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Marital status									
Never married	40.3	29.7	32.0	36.0	33.8	32.6	38.0	58.4	44.4
Currently married	19.7	24.2	22.8	20.9	18.8	21.9	16.2	10.3	21.5
Separated	7.0	8.6	6.7	3.5	7.6	7.8	7.4	5.5	6.6
Divorced	24.2	27.0	28.3	17.4	31.3	26.9	31.4	19.9	20.4
Widowed	8.9	10.5	10.2	22.4	8.5	10.7	7.1	5.9	7.1
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

GED = general equivalency diploma.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 5.5c. Individuals aged 55 years and older, by mental health diagnosis, education, and marital status: Row percent distribution, 2020

<i>Education and marital status</i>		<i>Mental health diagnosis¹</i>							
		<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>Delirium/dementia</i>	<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
	<i>Total</i>								
Total individuals aged 55 years and older	100.0	13.9	21.0	2.2	14.7	39.0	3.5	23.4	22.2
Highest school grade completed									
0 to 8	100.0	16.1	23.1	2.4	10.3	42.7	3.4	26.2	26.6
9 to 11	100.0	13.5	19.5	1.5	14.2	38.7	4.6	32.3	25.4
12 (or GED)	100.0	15.2	21.9	1.5	15.9	40.4	4.6	24.8	24.5
More than 12	100.0	17.8	23.6	1.2	19.9	41.1	5.3	19.2	21.6
Total	100.0	15.8	22.1	1.5	16.1	40.6	4.7	24.5	24.1
Marital status									
Never married	100.0	9.9	15.0	2.1	13.0	30.4	3.5	35.9	26.5
Currently married	100.0	16.4	21.7	2.5	14.8	41.8	3.0	13.0	26.2
Separated	100.0	16.4	18.1	1.2	17.0	42.0	3.9	19.6	22.9
Divorced	100.0	14.9	21.9	1.7	20.1	41.7	4.7	20.3	20.2
Widowed	100.0	15.9	21.6	5.9	14.9	45.3	2.9	16.4	19.1
Total	100.0	13.4	18.8	2.4	15.5	37.5	3.7	24.7	24.0

† Less than 1 percent of individuals.

GED = general equivalency diploma.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 5.6a. Individuals aged 55 years and older with SMI, by mental health diagnosis, education, and marital status: Number, 2020

Education and marital status	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	Delirium/ dementia	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals with SMI aged 55 years and older²	744,318	86,667	156,372	13,273	125,391	313,555	27,018	208,978	126,612
Highest school grade completed									
0 to 8	40,994	6,338	9,056	980	4,813	17,901	1,495	12,663	8,767
9 to 11	62,763	7,861	12,149	1,033	9,739	24,450	2,974	23,289	13,079
12 (or GED)	186,859	25,409	41,075	3,020	33,887	77,930	9,163	55,537	35,052
More than 12	104,053	16,195	24,980	1,327	24,176	44,567	5,897	24,639	17,117
Total	394,669	55,803	87,260	6,360	72,615	164,848	19,529	116,128	74,015
Marital status									
Never married	184,622	14,238	27,533	2,916	27,052	60,851	6,389	77,384	35,922
Currently married	74,688	8,919	17,588	1,796	13,987	35,772	2,277	12,971	12,962
Separated	31,302	3,935	6,153	342	6,185	14,303	1,174	7,371	5,106
Divorced	113,407	14,905	26,035	1,771	25,220	49,652	5,588	26,081	17,557
Widowed	39,540	5,328	8,764	1,950	6,875	19,571	1,251	7,659	5,865
Total	443,559	47,325	86,073	8,775	79,319	180,149	16,679	131,466	77,412

GED = general equivalency diploma; SMI = serious mental illness.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 5.6b. Individuals aged 55 years and older with SMI, by mental health diagnosis, education, and marital status: Column percent distribution, 2020

		Mental health diagnosis ¹								
Education and marital status		Total	Trauma- and stressor-related disorders	Anxiety disorders	Delirium/ dementia	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Total individuals with SMI aged 55 years and older ²		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Highest school grade completed										
0 to 8	10.4	11.4	10.4	15.4	6.6	10.9	7.7	10.9	11.8	
9 to 11	15.9	14.1	13.9	16.2	13.4	14.8	15.2	20.1	17.7	
12 (or GED)	47.3	45.5	47.1	47.5	46.7	47.3	46.9	47.8	47.4	
More than 12	26.4	29.0	28.6	20.9	33.3	27.0	30.2	21.2	23.1	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	
Marital status										
Never married	41.6	30.1	32.0	33.2	34.1	33.8	38.3	58.9	46.4	
Currently married	16.8	18.8	20.4	20.5	17.6	19.9	13.7	9.9	16.7	
Separated	7.1	8.3	7.1	3.9	7.8	7.9	7.0	5.6	6.6	
Divorced	25.6	31.5	30.2	20.2	31.8	27.6	33.5	19.8	22.7	
Widowed	8.9	11.3	10.2	22.2	8.7	10.9	7.5	5.8	7.6	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	

GED = general equivalency diploma; SMI = serious mental illness.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 5.6c. Individuals aged 55 years and older with SMI, by mental health diagnosis, education, and marital status: Row percent distribution, 2020

Education and marital status		Mental health diagnosis ¹								
		Total	Trauma- and stressor-related disorders	Anxiety disorders	Delirium/ dementia	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Total individuals with SMI aged 55 years and older ²		100.0	11.6	21.0	1.8	16.8	42.1	3.6	28.1	17.0
Highest school grade completed										
0 to 8	100.0	15.5	22.1	2.4	11.7	43.7	3.6	30.9	21.4	
9 to 11	100.0	12.5	19.4	1.6	15.5	39.0	4.7	37.1	20.8	
12 (or GED)	100.0	13.6	22.0	1.6	18.1	41.7	4.9	29.7	18.8	
More than 12	100.0	15.6	24.0	1.3	23.2	42.8	5.7	23.7	16.5	
Total	100.0	14.1	22.1	1.6	18.4	41.8	4.9	29.4	18.8	
Marital status										
Never married	100.0	7.7	14.9	1.6	14.7	33.0	3.5	41.9	19.5	
Currently married	100.0	11.9	23.5	2.4	18.7	47.9	3.0	17.4	17.4	
Separated	100.0	12.6	19.7	1.1	19.8	45.7	3.8	23.5	16.3	
Divorced	100.0	13.1	23.0	1.6	22.2	43.8	4.9	23.0	15.5	
Widowed	100.0	13.5	22.2	4.9	17.4	49.5	3.2	19.4	14.8	
Total	100.0	10.7	19.4	2.0	17.9	40.6	3.8	29.6	17.5	

† Less than 1 percent of individuals.

GED = general equivalency diploma; SMI = serious mental illness.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 5.7a. Individuals aged 55 years and older, by mental health diagnosis, living arrangement, SMI status, and veteran status: Number, 2020

Living arrangement, SMI status, and veteran status	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	Delirium/dementia	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals aged 55 years and older	1,010,080	140,825	212,400	22,719	148,935	394,254	35,333	236,477	224,006
Living arrangement									
Homeless	30,476	4,872	4,168	178	4,745	11,961	1,144	8,123	8,558
Foster home/foster care	3,719	294	926	124	442	934	210	1,875	1,109
Residential care	44,236	2,437	4,816	1,334	5,325	8,151	2,038	25,468	14,052
Institutional setting	28,852	2,697	5,082	2,393	3,313	10,475	952	10,185	5,466
Jail/correctional facility	5,074	1,103	483	91	578	1,042	786	1,601	1,551
Private residence	545,541	85,713	125,616	6,165	88,469	226,939	22,495	117,611	129,783
Other	11,446	2,364	3,275	355	1,509	4,736	385	2,982	1,952
Total	669,344	99,480	144,366	10,640	104,381	264,238	28,010	167,845	162,471
SMI status²									
SMI	744,318	86,667	156,372	13,273	125,391	313,555	27,018	208,978	126,612
Not SMI	237,805	50,772	51,622	8,720	20,034	71,546	7,432	21,809	87,260
Total	982,123	137,439	207,994	21,993	145,425	385,101	34,450	230,787	213,872
Veteran status									
Veteran	26,381	4,357	3,550	583	3,533	9,850	814	6,009	6,178
Not a veteran	425,374	59,805	83,354	6,513	63,023	160,753	16,662	98,525	123,453
Total	451,755	64,162	86,904	7,096	66,556	170,603	17,476	104,534	129,631

SMI = serious mental illness.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 5.7b. Individuals aged 55 years and older, by mental health diagnosis, living arrangement, SMI status, and veteran status: Column percent distribution, 2020

Living arrangement, SMI status, and veteran status	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	Delirium/dementia	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals aged 55 years and older	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Living arrangement									
Homeless	4.6	4.9	2.9	1.7	4.5	4.5	4.1	4.8	5.3
Foster home/foster care	†	†	†	1.2	†	†	†	1.1	†
Residential care	6.6	2.4	3.3	12.5	5.1	3.1	7.3	15.2	8.6
Institutional setting	4.3	2.7	3.5	22.5	3.2	4.0	3.4	6.1	3.4
Jail/correctional facility	†	1.1	†	†	†	†	2.8	†	†
Private residence	81.5	86.2	87.0	57.9	84.8	85.9	80.3	70.1	79.9
Other	1.7	2.4	2.3	3.3	1.4	1.8	1.4	1.8	1.2
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
SMI status²									
SMI	75.8	63.1	75.2	60.4	86.2	81.4	78.4	90.6	59.2
Not SMI	24.2	36.9	24.8	39.6	13.8	18.6	21.6	9.4	40.8
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status									
Veteran	5.8	6.8	4.1	8.2	5.3	5.8	4.7	5.7	4.8
Not a veteran	94.2	93.2	95.9	91.8	94.7	94.2	95.3	94.3	95.2
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

SMI = serious mental illness.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 5.7c. Individuals aged 55 years and older, by mental health diagnosis, living arrangement, SMI status, and veteran status: Row percent distribution, 2020

Living arrangement, SMI status, and veteran status	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	Delirium/dementia	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals aged 55 years and older	100.0	13.9	21.0	2.2	14.7	39.0	3.5	23.4	22.2
Living arrangement									
Homeless	100.0	16.0	13.7	†	15.6	39.2	3.8	26.7	28.1
Foster home/foster care	100.0	7.9	24.9	3.3	11.9	25.1	5.6	50.4	29.8
Residential care	100.0	5.5	10.9	3.0	12.0	18.4	4.6	57.6	31.8
Institutional setting	100.0	9.3	17.6	8.3	11.5	36.3	3.3	35.3	18.9
Jail/correctional facility	100.0	21.7	9.5	1.8	11.4	20.5	15.5	31.6	30.6
Private residence	100.0	15.7	23.0	1.1	16.2	41.6	4.1	21.6	23.8
Other	100.0	20.7	28.6	3.1	13.2	41.4	3.4	26.1	17.1
Total	100.0	14.9	21.6	1.6	15.6	39.5	4.2	25.1	24.3
SMI status²									
SMI	100.0	11.6	21.0	1.8	16.8	42.1	3.6	28.1	17.0
Not SMI	100.0	21.4	21.7	3.7	8.4	30.1	3.1	9.2	36.7
Total	100.0	14.0	21.2	2.2	14.8	39.2	3.5	23.5	21.8
Veteran status									
Veteran	100.0	16.5	13.5	2.2	13.4	37.3	3.1	22.8	23.4
Not a veteran	100.0	14.1	19.6	1.5	14.8	37.8	3.9	23.2	29.0
Total	100.0	14.2	19.2	1.6	14.7	37.8	3.9	23.1	28.7

† Less than 1 percent of individuals.

SMI = serious mental illness.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 5.8a. Individuals aged 55 years and older with SMI, by mental health diagnosis, living arrangement, and veteran status: Number, 2020

Mental health diagnosis ¹									
Living arrangement and veteran status	Total	Trauma- and stressor-related disorders	Anxiety disorders	Delirium/ dementia	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals with SMI aged 55 years and older ²	744,318	86,667	156,372	13,273	125,391	313,555	27,018	208,978	126,612
Living arrangement									
Homeless	25,416	3,988	3,464	137	4,176	10,204	983	7,363	6,362
Foster home/foster care	2,504	211	562	93	327	644	166	1,525	555
Residential care	36,803	1,958	3,812	1,030	4,623	6,894	1,826	23,415	9,203
Institutional setting	22,619	1,727	4,054	1,668	2,827	8,286	768	8,968	3,845
Jail/correctional facility	3,892	687	404	61	503	898	723	1,426	1,128
Private residence	408,393	58,611	95,076	4,751	74,763	175,502	17,963	104,052	75,171
Other	9,464	1,663	2,780	200	1,365	4,158	338	2,579	1,470
Total	509,091	68,845	110,152	7,940	88,584	206,586	22,767	149,328	97,734
Veteran status									
Veteran	20,109	2,814	2,594	456	2,970	8,050	570	5,334	3,528
Not a veteran	302,953	37,117	59,885	4,934	51,986	121,231	12,625	85,324	67,664
Total	323,062	39,931	62,479	5,390	54,956	129,281	13,195	90,658	71,192

SMI = serious mental illness.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 5.8b. Individuals aged 55 years and older with SMI, by mental health diagnosis, living arrangement, and veteran status: Column percent distribution, 2020

Living arrangement and veteran status		Mental health diagnosis ¹								
		Total	Trauma- and stressor-related disorders	Anxiety disorders	Delirium/ dementia	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Total individuals with SMI aged 55 years and older ²		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Living arrangement										
Homeless	5.0	5.8	3.1	1.7	4.7	4.9	4.3	4.9	6.5	
Foster home/foster care	†	†	†	1.2	†	†	†	1.0	†	
Residential care	7.2	2.8	3.5	13.0	5.2	3.3	8.0	15.7	9.4	
Institutional setting	4.4	2.5	3.7	21.0	3.2	4.0	3.4	6.0	3.9	
Jail/correctional facility	†	†	†	†	†	†	3.2	†	1.2	
Private residence	80.2	85.1	86.3	59.8	84.4	85.0	78.9	69.7	76.9	
Other	1.9	2.4	2.5	2.5	1.5	2.0	1.5	1.7	1.5	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status										
Veteran	6.2	7.0	4.2	8.5	5.4	6.2	4.3	5.9	5.0	
Not a veteran	93.8	93.0	95.8	91.5	94.6	93.8	95.7	94.1	95.0	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

SMI = serious mental illness.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 5.8c. Individuals aged 55 years and older with SMI, by mental health diagnosis, living arrangement, and veteran status: Row percent distribution, 2020

		Mental health diagnosis ¹								
Living arrangement and veteran status		Total	Trauma- and stressor-related disorders	Anxiety disorders	Delirium/ dementia	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Total individuals with SMI aged 55 years and older ²		100.0	11.6	21.0	1.8	16.8	42.1	3.6	28.1	17.0
Living arrangement										
Homeless	100.0	100.0	15.7	13.6	†	16.4	40.1	3.9	29.0	25.0
Foster home/foster care	100.0	100.0	8.4	22.4	3.7	13.1	25.7	6.6	60.9	22.2
Residential care	100.0	100.0	5.3	10.4	2.8	12.6	18.7	5.0	63.6	25.0
Institutional setting	100.0	100.0	7.6	17.9	7.4	12.5	36.6	3.4	39.6	17.0
Jail/correctional facility	100.0	100.0	17.7	10.4	1.6	12.9	23.1	18.6	36.6	29.0
Private residence	100.0	100.0	14.4	23.3	1.2	18.3	43.0	4.4	25.5	18.4
Other	100.0	100.0	17.6	29.4	2.1	14.4	43.9	3.6	27.3	15.5
	Total	100.0	13.5	21.6	1.6	17.4	40.6	4.5	29.3	19.2
Veteran status										
Veteran	100.0	100.0	14.0	12.9	2.3	14.8	40.0	2.8	26.5	17.5
Not a veteran	100.0	100.0	12.3	19.8	1.6	17.2	40.0	4.2	28.2	22.3
	Total	100.0	12.4	19.3	1.7	17.0	40.0	4.1	28.1	22.0

† Less than 1 percent of individuals.

SMI = serious mental illness.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 5.9a. Individuals aged 55 years and older, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Number, 2020

<i>Employment status and ‘detailed not in labor force’</i>		<i>Mental health diagnosis¹</i>							
		<i>Trauma- and stressor-related disorders Anxiety disorders Delirium/ dementia</i>			<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
	<i>Total</i>								
Total individuals aged 55 years and older	1,010,080	140,825	212,400	22,719	148,935	394,254	35,333	236,477	224,006
Employment status									
Employed	61,144	11,308	17,748	234	10,258	28,240	2,443	8,444	12,164
Full time	28,911	6,083	9,300	72	4,002	14,098	1,130	2,984	5,996
Part time	23,259	3,790	5,936	124	4,125	9,952	986	3,971	4,856
Unemployed	103,309	15,041	19,833	1,118	17,687	42,465	3,370	23,834	22,290
Not in labor force (see detail below)	303,227	45,734	76,761	5,309	52,999	126,427	15,318	84,567	51,089
Total	467,680	72,083	114,342	6,661	80,944	197,132	21,131	116,845	85,543
Detailed not in labor force									
Total individuals aged 55 years and older not in labor force	303,227	45,734	76,761	5,309	52,999	126,427	15,318	84,567	51,089
Disabled	170,360	25,318	44,893	2,628	31,857	68,400	10,860	55,477	29,698
Retired	40,548	6,096	9,810	1,542	6,153	19,122	1,092	6,589	5,532
Student	992	167	193	7	182	359	37	254	213
Homemaker	6,218	989	1,771	64	969	3,187	199	887	862
Sheltered/non-competitive employment	2,363	187	375	37	282	559	65	732	1,226
Other	82,746	12,977	19,719	1,031	13,556	34,800	3,065	20,628	13,558

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

NOTES: *Employed* includes “full-time/part-time not differentiated,” which is not presented separately; total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 5.9b. Individuals aged 55 years and older, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Column percent distribution, 2020

Employment status and ‘detailed not in labor force’	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	Delirium/ dementia	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals aged 55 years and older	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Employment status									
Employed	13.1	15.7	15.5	3.5	12.7	14.3	11.6	7.2	14.2
Full time	6.2	8.4	8.1	1.1	4.9	7.2	5.3	2.6	7.0
Part time	5.0	5.3	5.2	1.9	5.1	5.0	4.7	3.4	5.7
Unemployed	22.1	20.9	17.3	16.8	21.9	21.5	15.9	20.4	26.1
Not in labor force (see detail below)	64.8	63.4	67.1	79.7	65.5	64.1	72.5	72.4	59.7
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Detailed not in labor force									
Total individuals aged 55 years and older not in labor force	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Disabled	56.2	55.4	58.5	49.5	60.1	54.1	70.9	65.6	58.1
Retired	13.4	13.3	12.8	29.0	11.6	15.1	7.1	7.8	10.8
Student	†	†	†	†	†	†	†	†	†
Homemaker	2.1	2.2	2.3	1.2	1.8	2.5	1.3	1.0	1.7
Sheltered/non-competitive employment	†	†	†	†	†	†	†	†	2.4
Other	27.3	28.4	25.7	19.4	25.6	27.5	20.0	24.4	26.5

† Less than 1 percent of individuals.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding. *Employed* includes “full-time/part-time not differentiated,” which is not presented separately.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 5.9c. Individuals aged 55 years and older, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Row percent distribution, 2020

Employment status and 'detailed not in labor force'		Mental health diagnosis ¹								
		Total	Trauma- and stressor-related disorders	Anxiety disorders	Delirium/ dementia	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Total individuals aged 55 years and older		100.0	13.9	21.0	2.2	14.7	39.0	3.5	23.4	22.2
Employment status										
Employed	100.0	18.5	29.0	†	16.8	46.2	4.0	13.8	19.9	
Full time	100.0	21.0	32.2	†	13.8	48.8	3.9	10.3	20.7	
Part time	100.0	16.3	25.5	†	17.7	42.8	4.2	17.1	20.9	
Unemployed	100.0	14.6	19.2	1.1	17.1	41.1	3.3	23.1	21.6	
Not in labor force (see detail below)	100.0	15.1	25.3	1.8	17.5	41.7	5.1	27.9	16.8	
Total	100.0	15.4	24.4	1.4	17.3	42.2	4.5	25.0	18.3	
Detailed not in labor force										
Total individuals aged 55 years and older not in labor force	100.0	15.1	25.3	1.8	17.5	41.7	5.1	27.9	16.8	
Disabled	100.0	14.9	26.4	1.5	18.7	40.2	6.4	32.6	17.4	
Retired	100.0	15.0	24.2	3.8	15.2	47.2	2.7	16.2	13.6	
Student	100.0	16.8	19.5	†	18.3	36.2	3.7	25.6	21.5	
Homemaker	100.0	15.9	28.5	1.0	15.6	51.3	3.2	14.3	13.9	
Sheltered/non-competitive employment	100.0	7.9	15.9	1.6	11.9	23.7	2.8	31.0	51.9	
Other	100.0	15.7	23.8	1.2	16.4	42.1	3.7	24.9	16.4	

† Less than 1 percent of individuals.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding. *Employed* includes “full-time/part-time not differentiated,” which is not presented separately.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 5.10a. Individuals aged 55 years and older with SMI, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Number, 2020

<i>Employment status and ‘detailed not in labor force’</i>		<i>Mental health diagnosis¹</i>							
		<i>Trauma- and stressor-related disorders Anxiety disorders Delirium/dementia</i>			<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
	<i>Total</i>								
Total individuals with SMI aged 55 years and older²	744,318	86,667	156,372	13,273	125,391	313,555	27,018	208,978	126,612
Employment status									
Employed	45,065	7,436	12,800	145	8,804	22,170	1,991	7,353	7,255
Full time	19,128	3,607	6,016	41	3,148	10,081	849	2,367	3,256
Part time	18,090	2,606	4,502	75	3,607	8,213	828	3,544	3,080
Unemployed	84,287	11,177	15,921	849	15,853	36,844	2,777	21,329	14,690
Not in labor force (see detail below)	246,529	35,209	60,182	4,058	46,931	103,723	13,575	77,190	34,440
Total	375,881	53,822	88,903	5,052	71,588	162,737	18,343	105,872	56,385
Detailed not in labor force									
Total individuals with SMI aged 55 years and older not in labor force	246,529	35,209	60,182	4,058	46,931	103,723	13,575	77,190	34,440
Disabled	147,449	21,499	37,889	2,135	29,194	59,441	9,846	51,209	21,968
Retired	33,896	4,401	7,976	1,191	5,692	16,841	943	6,133	3,857
Student	853	130	153	6	176	326	31	232	157
Homemaker	5,267	769	1,417	45	880	2,741	171	820	644
Sheltered/non-competitive employment	1,566	163	243	22	239	473	52	644	507
Other	57,498	8,247	12,504	659	10,750	23,901	2,532	18,152	7,307

SMI = serious mental illness.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTES: *Employed* includes “full-time/part-time not differentiated,” which is not presented separately; total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 5.10b. Individuals aged 55 years and older with SMI, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Column percent distribution, 2020

<i>Employment status and ‘detailed not in labor force’</i>		<i>Mental health diagnosis¹</i>							
		<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>Delirium/ dementia</i>	<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
	<i>Total</i>								
Total individuals with SMI aged 55 years and older²	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Employment status									
Employed	12.0	13.8	14.4	2.9	12.3	13.6	10.9	6.9	12.9
Full time	5.1	6.7	6.8	†	4.4	6.2	4.6	2.2	5.8
Part time	4.8	4.8	5.1	1.5	5.0	5.0	4.5	3.3	5.5
Unemployed	22.4	20.8	17.9	16.8	22.1	22.6	15.1	20.1	26.1
Not in labor force (see detail below)	65.6	65.4	67.7	80.3	65.6	63.7	74.0	72.9	61.1
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Detailed not in labor force									
Total individuals with SMI aged 55 years and older not in labor force	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Disabled	59.8	61.1	63.0	52.6	62.2	57.3	72.5	66.3	63.8
Retired	13.7	12.5	13.3	29.3	12.1	16.2	6.9	7.9	11.2
Student	†	†	†	†	†	†	†	†	†
Homemaker	2.1	2.2	2.4	1.1	1.9	2.6	1.3	1.1	1.9
Sheltered/non-competitive employment	†	†	†	†	†	†	†	†	1.5
Other	23.3	23.4	20.8	16.2	22.9	23.0	18.7	23.5	21.2

† Less than 1 percent of individuals.

SMI = serious mental illness.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding. *Employed* includes “full-time/part-time not differentiated,” which is not presented separately.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 5.10c. Individuals aged 55 years and older with SMI, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Row percent distribution, 2020

Employment status and 'detailed not in labor force'		Mental health diagnosis ¹								
		Total	Trauma- and stressor-related disorders	Anxiety disorders	Delirium/ dementia	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Total individuals with SMI aged 55 years and older ²		100.0	11.6	21.0	1.8	16.8	42.1	3.6	28.1	17.0
Employment status										
Employed	100.0	16.5	28.4	†	19.5	49.2	4.4	16.3	16.1	
Full time	100.0	18.9	31.5	†	16.5	52.7	4.4	12.4	17.0	
Part time	100.0	14.4	24.9	†	19.9	45.4	4.6	19.6	17.0	
Unemployed	100.0	13.3	18.9	1.0	18.8	43.7	3.3	25.3	17.4	
Not in labor force (see detail below)	100.0	14.3	24.4	1.6	19.0	42.1	5.5	31.3	14.0	
	Total	100.0	14.3	23.7	1.3	19.0	43.3	4.9	28.2	15.0
Detailed not in labor force										
Total individuals with SMI aged 55 years and older not in labor force		100.0	14.3	24.4	1.6	19.0	42.1	5.5	31.3	14.0
Disabled	100.0	14.6	25.7	1.4	19.8	40.3	6.7	34.7	14.9	
Retired	100.0	13.0	23.5	3.5	16.8	49.7	2.8	18.1	11.4	
Student	100.0	15.2	17.9	†	20.6	38.2	3.6	27.2	18.4	
Homemaker	100.0	14.6	26.9	†	16.7	52.0	3.2	15.6	12.2	
Sheltered/non-competitive employment	100.0	10.4	15.5	1.4	15.3	30.2	3.3	41.1	32.4	
Other	100.0	14.3	21.7	1.1	18.7	41.6	4.4	31.6	12.7	

† Less than 1 percent of individuals.

SMI = serious mental illness.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding. *Employed* includes “full-time/part-time not differentiated,” which is not presented separately.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 5.11a. Individuals aged 55 years and older, by mental health diagnosis, timing of admission, and mental health service setting: Number, 2020

<i>Timing of admission and mental health service setting</i>	<i>Total</i>	<i>Mental health diagnosis¹</i>							
		<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>Delirium/ dementia</i>	<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Total individuals aged 55 years and older	1,010,080	140,825	212,400	22,719	148,935	394,254	35,333	236,477	224,006
Timing of admission									
New individuals	313,381	52,337	73,697	10,230	37,668	131,014	7,519	50,506	58,417
Continuing individuals	696,699	88,488	138,703	12,489	111,267	263,240	27,814	185,971	165,589
Mental health service setting^{2,3}									
State psychiatric hospital	21,009	1,018	1,317	1,831	2,940	3,067	1,690	12,439	5,069
Community-based program	979,949	139,188	210,629	20,834	145,467	387,496	33,889	222,775	214,647
Residential treatment center	14,413	782	1,692	293	1,884	3,591	462	7,423	3,454
Other psychiatric inpatient	38,778	3,441	5,248	1,101	5,983	15,604	1,273	13,168	8,892
Institutions under the justice system	5,191	1,222	411	57	411	922	805	1,384	2,129

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Individuals could receive services in more than one service setting during the reporting period.

³ Mental health service setting category definitions can be found in Appendix C.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 5.11b. Individuals aged 55 years and older, by mental health diagnosis, timing of admission, and mental health service setting: Column percent, 2020

<i>Timing of admission and mental health service setting</i>	<i>Mental health diagnosis¹</i>								
	<i>Total</i>	<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>Delirium/ dementia</i>	<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Total individuals aged 55 years and older	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Timing of admission									
New individuals	31.0	37.2	34.7	45.0	25.3	33.2	21.3	21.4	26.1
Continuing individuals	69.0	62.8	65.3	55.0	74.7	66.8	78.7	78.6	73.9
Mental health service setting^{2,3}									
State psychiatric hospital	2.1	†	†	8.1	2.0	†	4.8	5.3	2.3
Community-based program	97.0	98.8	99.2	91.7	97.7	98.3	95.9	94.2	95.8
Residential treatment center	1.4	†	†	1.3	1.3	†	1.3	3.1	1.5
Other psychiatric inpatient	3.8	2.4	2.5	4.8	4.0	4.0	3.6	5.6	4.0
Institutions under the justice system	†	†	†	†	†	†	2.3	†	†

† Less than 1 percent of individuals.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Percentages may not sum to 100 percent, because individuals could receive services in more than one service setting during the reporting period.

³ Mental health service setting category definitions can be found in Appendix C.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 5.11c. Individuals aged 55 years and older, by mental health diagnosis, timing of admission, and mental health service setting: Row percent distribution, 2020

Timing of admission and mental health service setting	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	Delirium/ dementia	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals aged 55 years and older	100.0	13.9	21.0	2.2	14.7	39.0	3.5	23.4	22.2
Timing of admission									
New individuals	100.0	16.7	23.5	3.3	12.0	41.8	2.4	16.1	18.6
Continuing individuals	100.0	12.7	19.9	1.8	16.0	37.8	4.0	26.7	23.8
Mental health service setting^{2,3}									
State psychiatric hospital	100.0	4.8	6.3	8.7	14.0	14.6	8.0	59.2	24.1
Community-based program	100.0	14.2	21.5	2.1	14.8	39.5	3.5	22.7	21.9
Residential treatment center	100.0	5.4	11.7	2.0	13.1	24.9	3.2	51.5	24.0
Other psychiatric inpatient	100.0	8.9	13.5	2.8	15.4	40.2	3.3	34.0	22.9
Institutions under the justice system	100.0	23.5	7.9	1.1	7.9	17.8	15.5	26.7	41.0

† Less than 1 percent of individuals.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

³ Mental health service setting category definitions can be found in Appendix C.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 5.12a. Individuals aged 55 years and older with SMI, by mental health diagnosis, timing of admission, and mental health service setting: Number, 2020

Timing of admission and mental health service setting	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	Delirium/ dementia	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals with SMI aged 55 years and older²	744,318	86,667	156,372	13,273	125,391	313,555	27,018	208,978	126,612
Timing of admission									
New individuals	223,368	31,565	53,736	4,664	31,622	107,058	5,494	42,644	32,240
Continuing individuals	520,950	55,102	102,636	8,609	93,769	206,497	21,524	166,334	94,372
Mental health service setting^{3,4}									
State psychiatric hospital	16,888	768	938	1,495	2,347	2,210	1,400	10,920	3,852
Community-based program	719,162	85,403	154,934	11,739	122,605	307,850	25,884	196,802	119,369
Residential treatment center	12,284	622	1,527	238	1,549	3,114	366	6,581	2,509
Other psychiatric inpatient	34,677	2,969	4,951	830	5,498	14,717	1,158	12,357	7,002
Institutions under the justice system	4,391	847	383	45	386	874	780	1,317	1,770

SMI = serious mental illness.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

³ Individuals could receive services in more than one service setting during the reporting period.

⁴ Mental health service setting category definitions can be found in Appendix C.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 5.12b. Individuals aged 55 years and older with SMI, by mental health diagnosis, timing of admission, and mental health service setting: Column percent, 2020

Timing of admission and mental health service setting	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	Delirium/dementia	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals with SMI aged 55 years and older²	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Timing of admission									
New individuals	30.0	36.4	34.4	35.1	25.2	34.1	20.3	20.4	25.5
Continuing individuals	70.0	63.6	65.6	64.9	74.8	65.9	79.7	79.6	74.5
Mental health service setting^{3,4}									
State psychiatric hospital	2.3	†	†	11.3	1.9	†	5.2	5.2	3.0
Community-based program	96.6	98.5	99.1	88.4	97.8	98.2	95.8	94.2	94.3
Residential treatment center	1.7	†	†	1.8	1.2	†	1.4	3.1	2.0
Other psychiatric inpatient	4.7	3.4	3.2	6.3	4.4	4.7	4.3	5.9	5.5
Institutions under the justice system	†	†	†	†	†	†	2.9	†	1.4

† Less than 1 percent of individuals.

SMI = serious mental illness.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

³ Percentages may not sum to 100 percent, because individuals could receive services in more than one service setting during the reporting period.

⁴ Mental health service setting category definitions can be found in Appendix C.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 5.12c. Individuals aged 55 years and older with SMI, by mental health diagnosis, timing of admission, and mental health service setting: Row percent distribution, 2020

Mental health diagnosis ¹									
Timing of admission and mental health service setting	Total	Trauma- and stressor-related disorders	Anxiety disorders	Delirium/ dementia	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals with SMI aged 55 years and older ²	100.0	11.6	21.0	1.8	16.8	42.1	3.6	28.1	17.0
Timing of admission									
New individuals	100.0	14.1	24.1	2.1	14.2	47.9	2.5	19.1	14.4
Continuing individuals	100.0	10.6	19.7	1.7	18.0	39.6	4.1	31.9	18.1
Mental health service setting ^{3,4}									
State psychiatric hospital	100.0	4.5	5.6	8.9	13.9	13.1	8.3	64.7	22.8
Community-based program	100.0	11.9	21.5	1.6	17.0	42.8	3.6	27.4	16.6
Residential treatment center	100.0	5.1	12.4	1.9	12.6	25.4	3.0	53.6	20.4
Other psychiatric inpatient	100.0	8.6	14.3	2.4	15.9	42.4	3.3	35.6	20.2
Institutions under the justice system	100.0	19.3	8.7	1.0	8.8	19.9	17.8	30.0	40.3

† Less than 1 percent of individuals.

SMI = serious mental illness.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

³ Individuals could receive services in more than one service setting during the reporting period.

⁴ Mental health service setting category definitions can be found in Appendix C.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 5.13. Individuals aged 55 years and older, by mental health service setting and SMI status: Number and percent distribution, 2020

<i>Mental health service setting</i>	<i>SMI status</i>								
	<i>Number of individuals</i>			<i>Column percent of individuals</i>			<i>Row percent distribution of individuals</i>		
	<i>Total</i>	<i>SMI</i>	<i>Not SMI</i>	<i>Total</i>	<i>SMI</i>	<i>Not SMI</i>	<i>Total</i>	<i>SMI</i>	<i>Not SMI</i>
Total individuals aged 55 years and older	1,124,875	824,495	300,380	100.0	100.0	100.0	100.0	73.3	26.7
Mental health service setting^{2,3}									
State psychiatric hospital	20,750	18,709	2,041	1.8	2.3	†	100.0	90.2	9.8
Community-based program	1,091,109	793,833	297,276	97.0	96.3	99.0	100.0	72.8	27.2
Residential treatment center	18,242	16,064	2,178	1.6	1.9	†	100.0	88.1	11.9
Other psychiatric inpatient	42,420	38,518	3,902	3.8	4.7	1.3	100.0	90.8	9.2
Institutions under the justice system	5,474	4,688	786	†	†	†	100.0	85.6	14.4

† Less than 1 percent of individuals.

SMI = serious emotional disturbance.

¹ The federal definition for SMI status can be found in Appendix D.

² Individuals could receive services in more than one mental health service setting during the reporting period.

³ Mental health service setting category definitions can be found in Appendix C.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 6.1a. Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by gender, age, race, ethnicity, education, veteran status, and SMI/SED status: Number, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Number of states reporting	44	44	46	46	46	49
Total individuals with co-occurring MH and SU disorders aged 12 years and older¹	747,400	682,142	903,371	1,048,708	1,263,731	1,663,657
Gender						
Male	417,766	378,504	483,709	558,791	655,115	825,644
Female	329,193	303,163	417,199	487,983	606,022	835,488
Total	746,959	681,667	900,908	1,046,774	1,261,137	1,661,132
Age²						
12 to 14 years	7,915	7,468	19,190	18,617	36,682	72,138
15 to 17 years	35,092	33,046	45,987	48,803	64,160	93,684
18 to 20 years	30,768	28,262	36,644	42,613	56,194	77,477
21 to 24 years	57,227	50,825	64,267	73,144	89,569	117,078
25 to 29 years	90,205	84,149	111,447	132,700	157,854	197,270
30 to 34 years	95,197	87,726	114,073	134,794	159,794	201,555
35 to 39 years	80,659	77,483	104,408	125,281	147,478	185,646
40 to 44 years	75,035	65,677	82,694	96,492	114,562	149,637
45 to 49 years	81,331	71,122	88,735	99,947	111,954	137,622
50 to 54 years	85,803	75,755	93,412	103,115	113,457	138,263
55 to 59 years	60,064	56,406	76,167	90,658	104,980	134,590
60 to 64 years	30,343	29,050	41,409	51,217	63,255	88,164
65 years and older	17,761	15,173	24,938	31,327	43,792	70,533
Total	747,400	682,142	903,371	1,048,708	1,263,731	1,663,657
Race						
American Indian or Alaska Native	13,661	13,178	15,488	19,307	26,283	32,555
Asian ³	6,245	7,143	7,800	9,610	9,764	13,796
Black or African American	135,044	122,226	170,612	201,631	248,557	356,196
Native Hawaiian or Other Pacific Islander	2,137	1,671	1,753	2,229	2,905	3,208
White	477,078	438,662	580,245	666,494	800,442	1,043,122
Other	60,888	57,802	76,690	95,120	120,381	144,754
Total	695,053	640,682	852,588	994,391	1,208,332	1,593,631
Ethnicity						
Not of Hispanic or Latino origin	555,462	528,360	737,108	847,676	1,024,187	1,367,223
Hispanic or Latino origin	91,305	90,317	100,154	111,733	145,445	203,385
Total	646,767	618,677	837,262	959,409	1,169,632	1,570,608
Specific Hispanic or Latino origin						
Mexican	5,885	2,702	4,431	5,056	8,343	10,105
Puerto Rican	6,889	5,007	6,314	6,686	14,157	16,102
Cuban	849	633	877	950	6,641	8,132
Other specific	7,414	5,485	6,768	7,285	12,080	15,700
Total	21,037	13,827	18,390	19,977	41,221	50,039

Continued. See notes at end of table.

Table 6.1a. Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by gender, age, race, ethnicity, education, veteran status, and SMI/SED status: Number, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Highest school grade completed						
0 to 8	40,059	31,453	46,067	57,271	91,737	117,479
9 to 11	98,280	79,964	109,153	139,011	176,447	218,794
12 (or GED)	189,080	154,506	213,419	288,832	375,613	468,406
More than 12	78,984	67,341	95,379	125,562	169,728	217,805
Total	406,403	333,264	464,018	610,676	813,525	1,022,484
Veteran status⁴						
Veteran	<i>n/c</i>	<i>n/c</i>	9,652	13,735	40,948	60,699
Not a veteran	<i>n/c</i>	<i>n/c</i>	213,578	381,265	557,595	700,215
Total	<i>n/c</i>	<i>n/c</i>	223,230	395,000	598,543	760,914
SMI/SED status⁵						
SMI	540,056	515,516	643,140	757,824	946,913	1,180,301
SED	34,843	34,031	51,944	55,495	70,905	97,266
Not SMI/SED	155,537	122,120	187,862	209,276	205,176	315,606
Total	730,436	671,667	882,946	1,022,595	1,222,994	1,593,173

n/c Not collected.

GED = general equivalency diploma; SED = serious emotional disturbance; SMI = serious mental illness.

¹ Total individuals with co-occurring mental health and substance use disorders who had at least one valid mental health diagnosis other than an alcohol- or substance-related mental health disorder.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

⁵ The federal definition for SMI and SED status can be found in Appendix D.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 6.1b. Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by gender, age, race, ethnicity, education, veteran status, and SMI/SED status: Column percent distribution, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Number of states reporting	44	44	46	46	46	49
Total individuals with co-occurring MH and SU disorders aged 12 years and older¹	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	55.9	55.5	53.7	53.4	51.9	49.7
Female	44.1	44.5	46.3	46.6	48.1	50.3
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age²						
12 to 14 years	1.1	1.1	2.1	1.8	2.9	4.3
15 to 17 years	4.7	4.8	5.1	4.7	5.1	5.6
18 to 20 years	4.1	4.1	4.1	4.1	4.4	4.7
21 to 24 years	7.7	7.5	7.1	7.0	7.1	7.0
25 to 29 years	12.1	12.3	12.3	12.7	12.5	11.9
30 to 34 years	12.7	12.9	12.6	12.9	12.6	12.1
35 to 39 years	10.8	11.4	11.6	11.9	11.7	11.2
40 to 44 years	10.0	9.6	9.2	9.2	9.1	9.0
45 to 49 years	10.9	10.4	9.8	9.5	8.9	8.3
50 to 54 years	11.5	11.1	10.3	9.8	9.0	8.3
55 to 59 years	8.0	8.3	8.4	8.6	8.3	8.1
60 to 64 years	4.1	4.3	4.6	4.9	5.0	5.3
65 years and older	2.4	2.2	2.8	3.0	3.5	4.2
Total	100.0	100.0	100.0	100.0	100.0	100.0
Race						
American Indian or Alaska Native	2.0	2.1	1.8	1.9	2.2	2.0
Asian ³	†	1.1	†	†	†	†
Black or African American	19.4	19.1	20.0	20.3	20.6	22.4
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	68.6	68.5	68.1	67.0	66.2	65.5
Other	8.8	9.0	9.0	9.6	10.0	9.1
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	85.9	85.4	88.0	88.4	87.6	87.1
Hispanic or Latino origin	14.1	14.6	12.0	11.6	12.4	12.9
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	28.0	19.5	24.1	25.3	20.2	20.2
Puerto Rican	32.7	36.2	34.3	33.5	34.3	32.2
Cuban	4.0	4.6	4.8	4.8	16.1	16.3
Other specific	35.2	39.7	36.8	36.5	29.3	31.4
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 6.1b. Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by gender, age, race, ethnicity, education, veteran status, and SMI/SED status: Column percent distribution, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Highest school grade completed						
0 to 8	9.9	9.4	9.9	9.4	11.3	11.5
9 to 11	24.2	24.0	23.5	22.8	21.7	21.4
12 (or GED)	46.5	46.4	46.0	47.3	46.2	45.8
More than 12	19.4	20.2	20.6	20.6	20.9	21.3
Total	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁴						
Veteran	<i>n/c</i>	<i>n/c</i>	4.3	3.5	6.8	8.0
Not a veteran	<i>n/c</i>	<i>n/c</i>	95.7	96.5	93.2	92.0
Total	<i>n/c</i>	<i>n/c</i>	100.0	100.0	100.0	100.0
SMI/SED status⁵						
SMI	73.9	76.8	72.8	74.1	77.4	74.1
SED	4.8	5.1	5.9	5.4	5.8	6.1
Not SMI/SED	21.3	18.2	21.3	20.5	16.8	19.8
Total	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals; *n/c* Not collected.

GED = general equivalency diploma; SED = serious emotional disturbance; SMI = serious mental illness.

¹ Total individuals with co-occurring mental health and substance use disorders who had at least one valid mental health diagnosis other than an alcohol- or substance-related mental health disorder.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

⁵ The federal definition for SMI and SED status can be found in Appendix D.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 6.2a. Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, gender, and age: Number, 2020

Gender and age		Mental health diagnosis ¹								
		Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Total individuals with co-occurring MH and SU disorders aged 12 years and older ²		1,663,657	341,389	394,514	90,755	292,252	604,444	72,546	295,895	433,149
Gender										
Male	825,644	141,808	163,142	56,285	125,696	263,039	33,436	198,872	247,488	
Female	835,488	198,923	230,726	34,328	166,094	340,541	39,025	96,740	185,010	
Total	1,661,132	340,731	393,868	90,613	291,790	603,580	72,461	295,612	432,498	
Age ³										
12 to 14 years	72,138	20,831	10,678	20,224	1,618	17,312	133	524	25,223	
15 to 17 years	93,684	24,281	18,620	17,389	4,522	32,542	532	1,724	34,927	
18 to 20 years	77,477	17,174	19,017	8,183	9,667	30,190	2,494	6,639	22,092	
21 to 24 years	117,078	24,649	29,729	6,908	19,961	43,665	5,522	17,946	29,734	
25 to 29 years	197,270	41,956	51,878	9,270	36,900	69,671	9,310	34,650	49,506	
30 to 34 years	201,555	44,218	53,745	8,433	40,045	69,470	10,117	36,659	50,126	
35 to 39 years	185,646	40,634	48,906	6,874	38,019	64,071	9,760	36,005	45,073	
40 to 44 years	149,637	31,885	38,143	4,409	32,349	53,131	7,911	30,120	35,955	
45 to 49 years	137,622	28,021	33,309	3,259	30,239	51,659	7,167	28,254	32,570	
50 to 54 years	138,263	25,800	31,322	2,376	28,620	55,006	6,867	29,740	33,176	
55 to 59 years	134,590	22,049	28,991	1,863	25,608	54,977	6,377	31,780	32,410	
60 to 64 years	88,164	12,152	17,756	1,041	14,946	35,865	3,818	22,911	21,602	
65 years and older	70,533	7,739	12,420	526	9,758	26,885	2,538	18,943	20,755	

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category. Mental health diagnostic category definitions can be found in Appendix E.

² Total individuals with co-occurring mental health and substance use disorders who had at least one valid mental health diagnosis other than an alcohol- or substance-related mental health disorder.

³ Age is based on the individual's age at the midpoint of the reporting period.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 6.2b. Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, gender, and age: Column percent distribution, 2020

Gender and age	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals with co-occurring MH and SU disorders aged 12 years and older²	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender									
Male	49.7	41.6	41.4	62.1	43.1	43.6	46.1	67.3	57.2
Female	50.3	58.4	58.6	37.9	56.9	56.4	53.9	32.7	42.8
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age³									
12 to 14 years	4.3	6.1	2.7	22.3	†	2.9	†	†	5.8
15 to 17 years	5.6	7.1	4.7	19.2	1.5	5.4	†	†	8.1
18 to 20 years	4.7	5.0	4.8	9.0	3.3	5.0	3.4	2.2	5.1
21 to 24 years	7.0	7.2	7.5	7.6	6.8	7.2	7.6	6.1	6.9
25 to 29 years	11.9	12.3	13.1	10.2	12.6	11.5	12.8	11.7	11.4
30 to 34 years	12.1	13.0	13.6	9.3	13.7	11.5	13.9	12.4	11.6
35 to 39 years	11.2	11.9	12.4	7.6	13.0	10.6	13.5	12.2	10.4
40 to 44 years	9.0	9.3	9.7	4.9	11.1	8.8	10.9	10.2	8.3
45 to 49 years	8.3	8.2	8.4	3.6	10.3	8.5	9.9	9.5	7.5
50 to 54 years	8.3	7.6	7.9	2.6	9.8	9.1	9.5	10.1	7.7
55 to 59 years	8.1	6.5	7.3	2.1	8.8	9.1	8.8	10.7	7.5
60 to 64 years	5.3	3.6	4.5	1.1	5.1	5.9	5.3	7.7	5.0
65 years and older	4.2	2.3	3.1	†	3.3	4.4	3.5	6.4	4.8
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category. Mental health diagnostic category definitions can be found in Appendix E.

² Total individuals with co-occurring mental health and substance use disorders who had at least one valid mental health diagnosis other than an alcohol- or substance-related mental health disorder.

³ Age is based on the individual's age at the midpoint of the reporting period.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 6.2c. Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, gender, and age: Row percent distribution, 2020

Gender and age		Mental health diagnosis ¹								
		Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Total individuals with co-occurring MH and SU disorders aged 12 years and older ²		100.0	20.5	23.7	5.5	17.6	36.3	4.4	17.8	26.0
Gender										
Male		100.0	17.2	19.8	6.8	15.2	31.9	4.0	24.1	30.0
Female		100.0	23.8	27.6	4.1	19.9	40.8	4.7	11.6	22.1
	Total	100.0	20.5	23.7	5.5	17.6	36.3	4.4	17.8	26.0
Age ³										
12 to 14 years		100.0	28.9	14.8	28.0	2.2	24.0	†	†	35.0
15 to 17 years		100.0	25.9	19.9	18.6	4.8	34.7	†	1.8	37.3
18 to 20 years		100.0	22.2	24.5	10.6	12.5	39.0	3.2	8.6	28.5
21 to 25 years		100.0	21.1	25.4	5.9	17.0	37.3	4.7	15.3	25.4
26 to 29 years		100.0	21.3	26.3	4.7	18.7	35.3	4.7	17.6	25.1
30 to 34 years		100.0	21.9	26.7	4.2	19.9	34.5	5.0	18.2	24.9
35 to 39 years		100.0	21.9	26.3	3.7	20.5	34.5	5.3	19.4	24.3
40 to 44 years		100.0	21.3	25.5	2.9	21.6	35.5	5.3	20.1	24.0
45 to 49 years		100.0	20.4	24.2	2.4	22.0	37.5	5.2	20.5	23.7
50 to 54 years		100.0	18.7	22.7	1.7	20.7	39.8	5.0	21.5	24.0
55 to 59 years		100.0	16.4	21.5	1.4	19.0	40.8	4.7	23.6	24.1
60 to 64 years		100.0	13.8	20.1	1.2	17.0	40.7	4.3	26.0	24.5
65 years and older		100.0	11.0	17.6	†	13.8	38.1	3.6	26.9	29.4

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category. Mental health diagnostic category definitions can be found in Appendix E.

² Total individuals with co-occurring mental health and substance use disorders who had at least one valid mental health diagnosis other than an alcohol- or substance-related mental health disorder.

³ Age is based on the individual's age at the midpoint of the reporting period.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 6.3a. Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, race, and ethnicity: Number, 2020

Race and ethnicity	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders			Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals with co-occurring MH and SU disorders aged 12 years and older²	1,663,657	341,389	394,514	90,755	292,252	604,444	72,546	295,895	433,149
Race									
American Indian or Alaska Native	32,555	9,253	9,739	1,211	3,734	13,823	1,162	4,386	9,252
Asian ³	13,796	2,595	2,472	563	1,614	4,357	365	3,956	4,093
Black or African American	356,196	64,960	51,157	18,683	52,134	114,249	14,217	106,345	92,002
Native Hawaiian or Other Pacific Islander	3,208	683	615	121	460	1,093	96	739	867
White	1,043,122	219,110	281,786	57,970	203,766	395,209	48,776	141,261	254,128
Other	144,754	30,750	30,273	8,685	20,691	51,566	5,385	26,473	43,977
Total	1,593,631	327,351	376,042	87,233	282,399	580,297	70,001	283,160	404,319
Ethnicity									
Not of Hispanic or Latino origin	1,367,223	285,261	331,856	77,296	247,951	500,157	61,101	238,311	328,830
Hispanic or Latino origin	203,385	39,231	44,311	9,249	30,364	75,931	7,020	34,669	64,962
Total	1,570,608	324,492	376,167	86,545	278,315	576,088	68,121	272,980	393,792
Specific Hispanic or Latino origin									
Mexican	10,105	2,709	2,262	735	1,166	3,798	298	1,269	2,772
Puerto Rican	16,102	2,745	2,246	735	2,443	5,621	327	3,711	2,292
Cuban	8,132	614	969	199	886	2,927	44	1,241	1,893
Other specific	15,700	3,190	3,146	841	2,444	6,103	477	2,557	2,900
Total	50,039	9,258	8,623	2,510	6,939	18,449	1,146	8,778	9,857

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category. Mental health diagnostic category definitions can be found in Appendix E.

² Total individuals with co-occurring mental health and substance use disorders who had at least one valid mental health diagnosis other than an alcohol- or substance-related mental health disorder.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 6.3b. Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, race, and ethnicity: Column percent distribution, 2020

Race and ethnicity	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals with co-occurring MH and SU disorders aged 12 years and older²	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race									
American Indian or Alaska Native	2.0	2.8	2.6	1.4	1.3	2.4	1.7	1.5	2.3
Asian ³	†	†	†	†	†	†	†	1.4	1.0
Black or African American	22.4	19.8	13.6	21.4	18.5	19.7	20.3	37.6	22.8
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	†	†	†
White	65.5	66.9	74.9	66.5	72.2	68.1	69.7	49.9	62.9
Other	9.1	9.4	8.1	10.0	7.3	8.9	7.7	9.3	10.9
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity									
Not of Hispanic or Latino origin	87.1	87.9	88.2	89.3	89.1	86.8	89.7	87.3	83.5
Hispanic or Latino origin	12.9	12.1	11.8	10.7	10.9	13.2	10.3	12.7	16.5
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	20.2	29.3	26.2	29.3	16.8	20.6	26.0	14.5	28.1
Puerto Rican	32.2	29.7	26.0	29.3	35.2	30.5	28.5	42.3	23.3
Cuban	16.3	6.6	11.2	7.9	12.8	15.9	3.8	14.1	19.2
Other specific	31.4	34.5	36.5	33.5	35.2	33.1	41.6	29.1	29.4
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category. Mental health diagnostic category definitions can be found in Appendix E.

² Total individuals with co-occurring mental health and substance use disorders who had at least one valid mental health diagnosis other than an alcohol- or substance-related mental health disorder.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 6.3c. Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, race, and ethnicity: Row percent distribution, 2020

Race and ethnicity	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other
					Bipolar disorders	Depressive disorders			
Total individuals with co-occurring MH and SU disorders aged 12 years and older²	100.0	20.5	23.7	5.5	17.6	36.3	4.4	17.8	26.0
Race									
American Indian or Alaska Native	100.0	28.4	29.9	3.7	11.5	42.5	3.6	13.5	28.4
Asian ³	100.0	18.8	17.9	4.1	11.7	31.6	2.6	28.7	29.7
Black or African American	100.0	18.2	14.4	5.2	14.6	32.1	4.0	29.9	25.8
Native Hawaiian or Other Pacific Islander	100.0	21.3	19.2	3.8	14.3	34.1	3.0	23.0	27.0
White	100.0	21.0	27.0	5.6	19.5	37.9	4.7	13.5	24.4
Other	100.0	21.2	20.9	6.0	14.3	35.6	3.7	18.3	30.4
Total	100.0	20.5	23.6	5.5	17.7	36.4	4.4	17.8	25.4
Ethnicity									
Not of Hispanic or Latino origin	100.0	20.9	24.3	5.7	18.1	36.6	4.5	17.4	24.1
Hispanic or Latino origin	100.0	19.3	21.8	4.5	14.9	37.3	3.5	17.0	31.9
Total	100.0	20.7	24.0	5.5	17.7	36.7	4.3	17.4	25.1
Specific Hispanic or Latino origin									
Mexican	100.0	26.8	22.4	7.3	11.5	37.6	2.9	12.6	27.4
Puerto Rican	100.0	17.0	13.9	4.6	15.2	34.9	2.0	23.0	14.2
Cuban	100.0	7.6	11.9	2.4	10.9	36.0	†	15.3	23.3
Other specific	100.0	20.3	20.0	5.4	15.6	38.9	3.0	16.3	18.5
Total	100.0	18.5	17.2	5.0	13.9	36.9	2.3	17.5	19.7

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category. Mental health diagnostic category definitions can be found in Appendix E.

² Total individuals with co-occurring mental health and substance use disorders who had at least one valid mental health diagnosis other than an alcohol- or substance-related mental health disorder.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

**Table 6.4a. Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis and substance use diagnosis:
Number, 2020**

<i>Substance use diagnosis¹</i>	<i>Mental health diagnosis²</i>								
	<i>Total</i>	<i>Trauma- and stressor- related disorders</i>	<i>Anxiety disorders</i>	<i>ADHD</i>	<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders²</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Total individuals with co-occurring MH and SU disorders aged 12 years and older³	1,663,657	341,389	394,514	90,755	292,252	604,444	72,546	295,895	433,149
Alcoholic psychoses	10,000	2,033	2,912	320	1,880	4,431	680	1,861	3,289
Alcohol intoxication	8,040	1,584	2,499	286	1,115	3,660	288	1,057	3,010
Drug psychoses	30,528	6,539	8,505	1,582	6,158	10,670	2,473	7,615	10,472
Dependence									
Alcohol	161,392	35,494	45,334	5,983	32,121	69,456	11,376	26,656	51,905
Cocaine	27,284	6,422	5,611	604	6,956	10,093	2,097	6,618	7,561
Marijuana	85,659	21,142	22,520	5,319	19,257	29,367	6,875	18,899	29,389
Opioid	93,655	23,847	32,870	5,178	17,762	39,019	4,598	7,249	24,570
Other	169,131	38,496	51,648	7,616	35,456	59,338	11,272	39,747	52,713
Non-dependent substance abuse									
Alcohol	56,713	11,401	16,041	1,937	12,528	22,282	4,080	13,204	16,855
Cocaine	12,048	2,228	2,401	276	2,995	4,291	965	3,507	3,624
Marijuana	80,664	17,501	22,687	5,654	17,574	29,316	5,162	17,539	27,134
Opioid	10,495	2,335	3,488	443	2,205	4,535	586	1,297	3,501
Other	50,501	9,926	12,726	1,984	10,555	16,888	3,375	13,972	19,870
None reported	867,547	162,441	165,272	53,573	125,690	301,098	18,719	136,674	179,256

ADHD = attention deficit/hyperactivity disorder.

¹ Substance use diagnostic category definitions can be found in Appendix E.

² "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category. Mental health diagnostic category definitions can be found in Appendix E.

³ Total individuals with co-occurring mental health and substance use disorders who had at least one valid mental health diagnosis other than an alcohol- or substance-related mental health disorder.

NOTE: Total individuals with co-occurring mental health and substance use disorders who had at least one valid mental health diagnosis other than an alcohol- or substance-related disorder.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

**Table 6.4b. Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis and substance use diagnosis:
Column percent distribution, 2020**

<i>Substance use diagnosis¹</i>	<i>Mental health diagnosis²</i>								
	<i>Total</i>	<i>Trauma- and stressor-related disorders</i> <i>Anxiety disorders</i> <i>ADHD</i>			<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders²</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Total individuals with co-occurring MH and SU disorders aged 12 years and older³	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Alcoholic psychoses	†	†	†	†	†	†	†	†	†
Alcohol intoxication	†	†	†	†	†	†	†	†	†
Drug psychoses	1.8	1.9	2.2	1.7	2.1	1.8	3.4	2.6	2.4
Substance dependence									
Alcohol	9.7	10.4	11.5	6.6	11.0	11.5	15.7	9.0	12.0
Cocaine	1.6	1.9	1.4	†	2.4	1.7	2.9	2.2	1.7
Marijuana	5.1	6.2	5.7	5.9	6.6	4.9	9.5	6.4	6.8
Opioid	5.6	7.0	8.3	5.7	6.1	6.5	6.3	2.4	5.7
Other	10.2	11.3	13.1	8.4	12.1	9.8	15.5	13.4	12.2
Non-dependent substance abuse									
Alcohol	3.4	3.3	4.1	2.1	4.3	3.7	5.6	4.5	3.9
Cocaine	†	†	†	†	1.0	†	1.3	1.2	†
Marijuana	4.8	5.1	5.8	6.2	6.0	4.9	7.1	5.9	6.3
Opioid	†	†	†	†	†	†	†	†	†
Other	3.0	2.9	3.2	2.2	3.6	2.8	4.7	4.7	4.6
None reported	52.1	47.6	41.9	59.0	43.0	49.8	25.8	46.2	41.4

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ Substance use diagnostic category definitions can be found in Appendix E.

² “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category. Mental health diagnostic category definitions can be found in Appendix E.

³ Total individuals with co-occurring mental health and substance use disorders who had at least one valid mental health diagnosis other than an alcohol- or substance-related mental health disorder.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

**Table 6.4c. Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis and substance use diagnosis:
Row percent distribution, 2020**

Substance use diagnosis ¹	Total	Mental health diagnosis ²							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ²
					Bipolar disorders	Depressive disorders			
Total individuals with co-occurring MH and SU disorders aged 12 years and older ³	100.0	20.5	23.7	5.5	17.6	36.3	4.4	17.8	26.0
Alcoholic psychoses	100.0	20.3	29.1	3.2	18.8	44.3	6.8	18.6	32.9
Alcohol intoxication	100.0	19.7	31.1	3.6	13.9	45.5	3.6	13.1	37.4
Drug psychoses	100.0	21.4	27.9	5.2	20.2	35.0	8.1	24.9	34.3
Substance dependence									
Alcohol	100.0	22.0	28.1	3.7	19.9	43.0	7.0	16.5	32.2
Cocaine	100.0	23.5	20.6	2.2	25.5	37.0	7.7	24.3	27.7
Marijuana	100.0	24.7	26.3	6.2	22.5	34.3	8.0	22.1	34.3
Opioid	100.0	25.5	35.1	5.5	19.0	41.7	4.9	7.7	26.2
Other	100.0	22.8	30.5	4.5	21.0	35.1	6.7	23.5	31.2
Non-dependent substance abuse									
Alcohol	100.0	20.1	28.3	3.4	22.1	39.3	7.2	23.3	29.7
Cocaine	100.0	18.5	19.9	2.3	24.9	35.6	8.0	29.1	30.1
Marijuana	100.0	21.7	28.1	7.0	21.8	36.3	6.4	21.7	33.6
Opioid	100.0	22.2	33.2	4.2	21.0	43.2	5.6	12.4	33.4
Other	100.0	19.7	25.2	3.9	20.9	33.4	6.7	27.7	39.3
None reported	100.0	18.7	19.1	6.2	14.5	34.7	2.2	15.8	20.7

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ Substance use diagnostic category definitions can be found in Appendix E.

² “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category. Mental health diagnostic category definitions can be found in Appendix E.

³ Total individuals with co-occurring mental health and substance use disorders who had at least one valid mental health diagnosis other than an alcohol- or substance-related mental health disorder.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

**Table 6.5a. Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, education, and marital status:
Number, 2020**

		Mental health diagnosis ¹								
Education and marital status		Total	Trauma- and stressor-related disorders			Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Total individuals with co-occurring MH and SU disorders aged 12 years and older ²		1,663,657	341,389	394,514	90,755	292,252	604,444	72,546	295,895	433,149
Highest school grade completed										
0 to 8	117,479	27,659	22,827	19,731	12,295	36,961	3,545	17,122	36,994	
9 to 11	218,794	47,247	47,170	15,479	35,886	79,347	11,300	48,197	60,926	
12 (or GED)	468,406	97,404	114,929	17,708	94,721	180,378	26,969	97,453	108,139	
More than 12	217,805	47,497	60,793	8,525	51,413	87,779	13,320	36,138	44,192	
Total	1,022,484	219,807	245,719	61,443	194,315	384,465	55,134	198,910	250,251	
Marital status										
Never married	819,360	157,534	157,774	59,966	122,809	264,415	28,701	159,778	210,929	
Currently married	141,705	30,845	35,992	3,639	27,763	57,921	5,682	15,639	29,639	
Separated	66,391	14,864	15,268	1,383	14,612	27,095	3,055	9,296	13,108	
Divorced	160,210	33,124	38,532	3,778	37,993	65,246	9,043	25,632	30,942	
Widowed	28,651	5,656	6,289	347	5,092	12,813	1,025	4,454	6,803	
Total	1,216,317	242,023	253,855	69,113	208,269	427,490	47,506	214,799	291,421	

ADHD = attention deficit/hyperactivity disorder; GED = general equivalency diploma.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category. Mental health diagnostic category definitions can be found in Appendix E.

² Total individuals with co-occurring mental health and substance use disorders who had at least one valid mental health diagnosis other than an alcohol- or substance-related mental health disorder.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

**Table 6.5b. Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, education, and marital status:
Column percent distribution, 2020**

Education and marital status		Mental health diagnosis ¹							
		Trauma- and stressor-related disorders			Mood disorders		Schizophrenia, other psychotic disorders		
					Bipolar disorders	Depressive disorders			
Total			Anxiety disorders	ADHD			Personality disorders		Other mental disorders ¹
Total individuals with co-occurring MH and SU disorders aged 12 years and older ²		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Highest school grade completed									
0 to 8	11.5	12.6	9.3	32.1	6.3	9.6	6.4	8.6	14.8
9 to 11	21.4	21.5	19.2	25.2	18.5	20.6	20.5	24.2	24.3
12 (or GED)	45.8	44.3	46.8	28.8	48.7	46.9	48.9	49.0	43.2
More than 12	21.3	21.6	24.7	13.9	26.5	22.8	24.2	18.2	17.7
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Marital status									
Never married	67.4	65.1	62.2	86.8	59.0	61.9	60.4	74.4	72.4
Currently married	11.7	12.7	14.2	5.3	13.3	13.5	12.0	7.3	10.2
Separated	5.5	6.1	6.0	2.0	7.0	6.3	6.4	4.3	4.5
Divorced	13.2	13.7	15.2	5.5	18.2	15.3	19.0	11.9	10.6
Widowed	2.4	2.3	2.5	†	2.4	3.0	2.2	2.1	2.3
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; GED = general equivalency diploma.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category. Mental health diagnostic category definitions can be found in Appendix E.

² Total individuals with co-occurring mental health and substance use disorders who had at least one valid mental health diagnosis other than an alcohol- or substance-related mental health disorder.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

**Table 6.5c. Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, education, and marital status:
Row percent distribution, 2020**

<i>Education and marital status</i>		<i>Mental health diagnosis¹</i>							
					<i>Mood disorders</i>				
	<i>Total</i>	<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>ADHD</i>	<i>Bipolar disorders</i>	<i>Depressive disorders</i>	<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
Total individuals with co-occurring MH and SU disorders aged 12 years and older²	100.0	20.5	23.7	5.5	17.6	36.3	4.4	17.8	26.0
Highest school grade completed									
0 to 8	100.0	23.5	19.4	16.8	10.5	31.5	3.0	14.6	31.5
9 to 11	100.0	21.6	21.6	7.1	16.4	36.3	5.2	22.0	27.8
12 (or GED)	100.0	20.8	24.5	3.8	20.2	38.5	5.8	20.8	23.1
More than 12	100.0	21.8	27.9	3.9	23.6	40.3	6.1	16.6	20.3
Total	100.0	21.5	24.0	6.0	19.0	37.6	5.4	19.5	24.5
Marital status									
Never married	100.0	19.2	19.3	7.3	15.0	32.3	3.5	19.5	25.7
Currently married	100.0	21.8	25.4	2.6	19.6	40.9	4.0	11.0	20.9
Separated	100.0	22.4	23.0	2.1	22.0	40.8	4.6	14.0	19.7
Divorced	100.0	20.7	24.1	2.4	23.7	40.7	5.6	16.0	19.3
Widowed	100.0	19.7	22.0	1.2	17.8	44.7	3.6	15.5	23.7
Total	100.0	19.9	20.9	5.7	17.1	35.1	3.9	17.7	24.0

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; GED = general equivalency diploma.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category. Mental health diagnostic category definitions can be found in Appendix E.

² Total individuals with co-occurring mental health and substance use disorders who had at least one valid mental health diagnosis other than an alcohol- or substance-related mental health disorder.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 6.6a. Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, living arrangement, SMI/SED status, and veteran status: Number, 2020

Living arrangement, SMI/SED status, and veteran status	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals with co-occurring MH and SU disorders aged 12 years and older²	1,663,657	341,389	394,514	90,755	292,252	604,444	72,546	295,895	433,149
Living arrangement									
Homeless	81,994	17,360	14,769	1,785	16,196	31,296	4,015	20,392	24,743
Foster home/foster care	5,686	1,748	1,258	645	639	1,625	353	1,624	1,913
Residential care	46,352	7,098	7,701	1,585	7,850	11,520	3,015	20,715	15,014
Institutional setting	23,842	2,826	4,969	1,048	4,195	7,963	1,658	8,960	7,220
Jail/correctional facility	30,749	9,466	4,368	1,105	3,891	6,505	5,143	6,864	10,344
Private residence	959,881	202,301	238,685	59,940	181,761	364,807	45,352	168,770	255,826
Other	19,905	5,399	6,765	957	3,270	7,335	897	4,155	5,055
Total	1,168,409	246,198	278,515	67,065	217,802	431,051	60,433	231,480	320,115
SMI/SED status³									
SMI	1,180,301	208,540	290,102	35,892	249,593	463,694	61,709	265,623	257,703
SED	97,266	20,965	23,201	20,582	5,289	40,546	520	1,811	37,129
Not SMI/SED	315,606	97,520	67,858	29,051	29,274	78,595	7,640	18,612	110,960
Total	1,593,173	327,025	381,161	85,525	284,156	582,835	69,869	286,046	405,792
Veteran status⁴									
Veteran	60,699	9,823	7,183	1,030	9,565	23,971	1,077	10,977	7,614
Not a veteran	700,215	130,731	149,866	23,912	131,735	257,361	31,302	145,780	200,864
Total	760,914	140,554	157,049	24,942	141,300	281,332	32,379	156,757	208,478

ADHD = attention deficit/hyperactivity disorder; SED = serious emotional disturbance; SMI = serious mental illness.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category. Mental health diagnostic category definitions can be found in Appendix E.

² Total individuals with co-occurring mental health and substance use disorders who had at least one valid mental health diagnosis other than an alcohol- or substance-related mental health disorder.

³ The federal definitions for SMI and SED statuses can be found in Appendix D.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 6.6b. Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, living arrangement, SMI/SED status, and veteran status: Column percent distribution, 2020

Living arrangement, SMI/SED status, and veteran status		Mental health diagnosis ¹							
		Trauma- and stressor-related disorders			Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
	Total		Anxiety disorders	ADHD					
Total individuals with co-occurring MH and SU disorders aged 12 years and older²	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Living arrangement									
Homeless	7.0	7.1	5.3	2.7	7.4	7.3	6.6	8.8	7.7
Foster home/foster care	†	†	†	†	†	†	†	†	†
Residential care	4.0	2.9	2.8	2.4	3.6	2.7	5.0	8.9	4.7
Institutional setting	2.0	1.1	1.8	1.6	1.9	1.8	2.7	3.9	2.3
Jail/correctional facility	2.6	3.8	1.6	1.6	1.8	1.5	8.5	3.0	3.2
Private residence	82.2	82.2	85.7	89.4	83.5	84.6	75.0	72.9	79.9
Other	1.7	2.2	2.4	1.4	1.5	1.7	1.5	1.8	1.6
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
SMI/SED status³									
SMI	74.1	63.8	76.1	42.0	87.8	79.6	88.3	92.9	63.5
SED	6.1	6.4	6.1	24.1	1.9	7.0	†	†	9.1
Not SMI/SED	19.8	29.8	17.8	34.0	10.3	13.5	10.9	6.5	27.3
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁴									
Veteran	8.0	7.0	4.6	4.1	6.8	8.5	3.3	7.0	3.7
Not a veteran	92.0	93.0	95.4	95.9	93.2	91.5	96.7	93.0	96.3
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; SED = serious emotional disturbance; SMI = serious mental illness.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category. Mental health diagnostic category definitions can be found in Appendix E.

² Total individuals with co-occurring mental health and substance use disorders who had at least one valid mental health diagnosis other than an alcohol- or substance-related mental health disorder.

³ The federal definitions for SMI and SED statuses can be found in Appendix D.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 6.6c. Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, living arrangement, SMI/SED status, and veteran status: Row percent distribution, 2020

Living arrangement, SMI/SED status, and veteran status		Mental health diagnosis ¹									
		Trauma- and stressor-related disorders			Mood disorders		Personality disorders				
					Schizophrenia, other psychotic disorders						
Total					Bipolar disorders		Depressive disorders		Other mental disorders ¹		
Total individuals with co-occurring MH and SU disorders aged 12 years and older ²		100.0	20.5	23.7	5.5	17.6	36.3	4.4	17.8	26.0	
Living arrangement											
Homeless	100.0	21.2	18.0	2.2	19.8	38.2	4.9	24.9	30.2		
Foster home/foster care	100.0	30.7	22.1	11.3	11.2	28.6	6.2	28.6	33.6		
Residential care	100.0	15.3	16.6	3.4	16.9	24.9	6.5	44.7	32.4		
Institutional setting	100.0	11.9	20.8	4.4	17.6	33.4	7.0	37.6	30.3		
Jail/correctional facility	100.0	30.8	14.2	3.6	12.7	21.2	16.7	22.3	33.6		
Private residence	100.0	21.1	24.9	6.2	18.9	38.0	4.7	17.6	26.7		
Other	100.0	27.1	34.0	4.8	16.4	36.9	4.5	20.9	25.4		
Total	100.0	21.1	23.8	5.7	18.6	36.9	5.2	19.8	27.4		
SMI/SED status ³											
SMI	100.0	17.7	24.6	3.0	21.1	39.3	5.2	22.5	21.8		
SED	100.0	21.6	23.9	21.2	5.4	41.7	†	1.9	38.2		
Not SMI/SED	100.0	30.9	21.5	9.2	9.3	24.9	2.4	5.9	35.2		
Total	100.0	20.5	23.9	5.4	17.8	36.6	4.4	18.0	25.5		
Veteran status ⁴											
Veteran	100.0	16.2	11.8	1.7	15.8	39.5	1.8	18.1	12.5		
Not a veteran	100.0	18.7	21.4	3.4	18.8	36.8	4.5	20.8	28.7		
Total	100.0	18.5	20.6	3.3	18.6	37.0	4.3	20.6	27.4		

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; SED = serious emotional disturbance; SMI = serious mental illness.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category. Mental health diagnostic category definitions can be found in Appendix E.

² Total individuals with co-occurring mental health and substance use disorders who had at least one valid mental health diagnosis other than an alcohol- or substance-related mental health disorder.

³ The federal definitions for SMI and SED statuses can be found in Appendix D.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 6.7a. Individuals aged 18 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Number, 2020

Employment status and 'detailed not in labor force'		Mental health diagnosis ¹								
		Trauma- and stressor-related disorders			Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹	
					Bipolar disorders	Depressive disorders				
Total			Anxiety disorders	ADHD						
Total individuals with co-occurring MH and SU disorders aged 12 years and older ²		1,497,835	296,277	365,216	53,142	286,112	554,590	71,881	293,647	372,999
Employment status										
Employed		218,800	53,721	66,691	10,590	42,684	95,090	10,876	21,200	43,557
Full time		118,811	30,966	37,804	6,273	19,583	52,961	5,653	9,188	23,159
Part time		78,860	17,350	21,310	3,727	15,442	32,772	3,883	9,373	16,893
Unemployed		346,096	69,590	78,875	11,736	70,873	136,735	14,228	62,681	81,445
Not in labor force (see detail below)		333,196	67,304	87,780	13,756	71,804	122,491	25,336	90,800	75,359
Total		898,092	190,615	233,346	36,082	185,361	354,316	50,440	174,681	200,361
Detailed not in labor force										
Disabled		169,126	31,074	42,477	5,401	39,273	57,091	15,642	58,718	39,263
Retired		15,346	2,580	2,984	151	2,393	6,850	351	2,484	2,905
Student		22,083	4,443	4,363	2,579	2,624	7,416	506	2,068	7,739
Homemaker		14,017	3,369	3,846	340	2,605	6,077	557	1,122	2,797
Sheltered/non-competitive employment		2,313	276	453	89	360	638	108	533	1,335
Other		110,311	25,562	33,657	5,196	24,549	44,419	8,172	25,875	21,320
Total		333,196	67,304	87,780	13,756	71,804	122,491	25,336	90,800	75,359

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category. Mental health diagnostic category definitions can be found in Appendix E.

² Total individuals with co-occurring mental health and substance use disorders who had at least one valid mental health diagnosis other than an alcohol- or substance-related mental health disorder.

NOTES: *Employed* includes “full-time/part-time not differentiated,” which is not presented separately; total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 6.7b. Individuals aged 18 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Column percent distribution, 2020

<i>Employment status and ‘detailed not in labor force’</i>		<i>Mental health diagnosis¹</i>							
		<i>Trauma- and stressor-related disorders</i>			<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
	<i>Total</i>		<i>Anxiety disorders</i>	<i>ADHD</i>					
Total individuals with co-occurring MH and SU disorders aged 12 years and older²	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Employment status									
Employed	24.4	28.2	28.6	29.3	23.0	26.8	21.6	12.1	21.7
Full time	13.2	16.2	16.2	17.4	10.6	14.9	11.2	5.3	11.6
Part time	8.8	9.1	9.1	10.3	8.3	9.2	7.7	5.4	8.4
Unemployed	38.5	36.5	33.8	32.5	38.2	38.6	28.2	35.9	40.6
Not in labor force (see detail below)	37.1	35.3	37.6	38.1	38.7	34.6	50.2	52.0	37.6
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Detailed not in labor force									
Disabled	50.8	46.2	48.4	39.3	54.7	46.6	61.7	64.7	52.1
Retired	4.6	3.8	3.4	1.1	3.3	5.6	1.4	2.7	3.9
Student	6.6	6.6	5.0	18.7	3.7	6.1	2.0	2.3	10.3
Homemaker	4.2	5.0	4.4	2.5	3.6	5.0	2.2	1.2	3.7
Sheltered/non-competitive employment	†	†	†	†	†	†	†	†	1.8
Other	33.1	38.0	38.3	37.8	34.2	36.3	32.3	28.5	28.3
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category. Mental health diagnostic category definitions can be found in Appendix E.

² Total individuals with co-occurring mental health and substance use disorders who had at least one valid mental health diagnosis other than an alcohol- or substance-related mental health disorder.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding. *Employed* includes “full-time/part-time not differentiated,” which is not presented separately.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 6.7c. Individuals aged 18 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Row percent distribution, 2020

<i>Employment status and ‘detailed not in labor force’</i>		<i>Mental health diagnosis¹</i>							
					<i>Mood disorders</i>				
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
	<i>Total</i>	<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>ADHD</i>			<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
Total individuals with co-occurring MH and SU disorders aged 12 years and older²	100.0	19.8	24.4	3.5	19.1	37.0	4.8	19.6	24.9
Employment status									
Employed	100.0	24.6	30.5	4.8	19.5	43.5	5.0	9.7	19.9
Full time	100.0	26.1	31.8	5.3	16.5	44.6	4.8	7.7	19.5
Part time	100.0	22.0	27.0	4.7	19.6	41.6	4.9	11.9	21.4
Unemployed	100.0	20.1	22.8	3.4	20.5	39.5	4.1	18.1	23.5
Not in labor force (see detail below)	100.0	20.2	26.3	4.1	21.6	36.8	7.6	27.3	22.6
Total	100.0	21.2	26.0	4.0	20.6	39.5	5.6	19.5	22.3
Detailed not in labor force									
Disabled	100.0	18.4	25.1	3.2	23.2	33.8	9.2	34.7	23.2
Retired	100.0	16.8	19.4	†	15.6	44.6	2.3	16.2	18.9
Student	100.0	20.1	19.8	11.7	11.9	33.6	2.3	9.4	35.0
Homemaker	100.0	24.0	27.4	2.4	18.6	43.4	4.0	8.0	20.0
Sheltered/non-competitive employment	100.0	11.9	19.6	3.8	15.6	27.6	4.7	23.0	57.7
Other	100.0	23.2	30.5	4.7	22.3	40.3	7.4	23.5	19.3
Total	100.0	20.2	26.3	4.1	21.6	36.8	7.6	27.3	22.6

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category. Mental health diagnostic category definitions can be found in Appendix E.

² Total individuals with co-occurring mental health and substance use disorders who had at least one valid mental health diagnosis other than an alcohol- or substance-related mental health disorder.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding. *Employed* includes “full-time/part-time not differentiated,” which is not presented separately.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 6.8a. Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, timing of admission, and mental health service setting: Number, 2020

Timing of admission and mental health service setting	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals with co-occurring MH and SU disorders aged 12 years and older²	1,663,657	341,389	394,514	90,755	292,252	604,444	72,546	295,895	433,149
Timing of admission									
New individuals	853,712	190,403	207,825	42,727	128,733	324,683	28,179	104,195	205,464
Continuing individuals	809,945	150,986	186,689	48,028	163,519	279,761	44,367	191,700	227,685
Mental health service setting^{3,4}									
State psychiatric hospital	54,030	5,655	5,181	1,518	10,253	10,480	5,777	29,611	16,644
Community-based program	1,586,320	332,284	389,195	89,205	282,081	582,323	68,866	270,617	406,434
Residential treatment center	24,244	3,639	3,823	1,167	4,369	6,760	1,139	9,688	7,114
Other psychiatric inpatient	143,038	20,492	20,856	4,346	25,784	56,225	5,433	35,050	30,176
Institutions under the justice system	27,526	9,596	3,601	791	2,111	5,117	5,505	4,827	12,276

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category. Mental health diagnostic category definitions can be found in Appendix E.

² Total individuals with co-occurring mental health and substance use disorders who had at least one valid mental health diagnosis other than an alcohol- or substance-related mental health disorder.

³ Individuals could receive services in more than one service setting during the reporting period.

⁴ Mental health service setting category definitions can be found in Appendix C.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 6.8b. Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, timing of admission, and mental health service setting: Column percent, 2020

Timing of admission and mental health service setting	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals with co-occurring MH and SU disorders aged 12 years and older ²	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Timing of admission									
New individuals	51.3	55.8	52.7	47.1	44.0	53.7	38.8	35.2	47.4
Continuing individuals	48.7	44.2	47.3	52.9	56.0	46.3	61.2	64.8	52.6
Mental health service setting ^{3,4}									
State psychiatric hospital	3.2	1.7	1.3	1.7	3.5	1.7	8.0	10.0	3.8
Community-based program	95.4	97.3	98.7	98.3	96.5	96.3	94.9	91.5	93.8
Residential treatment center	1.5	1.1	†	1.3	1.5	1.1	1.6	3.3	1.6
Other psychiatric inpatient	8.6	6.0	5.3	4.8	8.8	9.3	7.5	11.8	7.0
Institutions under the justice system	1.7	2.8	†	†	†	†	7.6	1.6	2.8

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category. Mental health diagnostic category definitions can be found in Appendix E.

² Total individuals with co-occurring mental health and substance use disorders who had at least one valid mental health diagnosis other than an alcohol- or substance-related mental health disorder.

³ Percentages may not sum to 100 percent, because individuals could receive services in more than one service setting during the reporting period.

⁴ Mental health service setting category definitions can be found in Appendix C.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 6.8c. Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, timing of admission, and mental health service setting: Row percent distribution, 2020

Timing of admission and mental health service setting		Mental health diagnosis ¹									
		Trauma- and stressor-related disorders			Mood disorders		Personality disorders			Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders					
Total			Anxiety disorders	ADHD							
Total individuals with co-occurring MH and SU disorders aged 12 years and older ²		100.0	20.5	23.7	5.5	17.6	36.3	4.4	17.8	26.0	
Timing of admission											
New individuals	100.0	22.3	24.3	5.0	15.1	38.0	3.3	12.2	24.1		
Continuing individuals	100.0	18.6	23.0	5.9	20.2	34.5	5.5	23.7	28.1		
Mental health service setting ^{3,4}											
State psychiatric hospital	100.0	10.5	9.6	2.8	19.0	19.4	10.7	54.8	30.8		
Community-based program	100.0	20.9	24.5	5.6	17.8	36.7	4.3	17.1	25.6		
Residential treatment center	100.0	15.0	15.8	4.8	18.0	27.9	4.7	40.0	29.3		
Other psychiatric inpatient	100.0	14.3	14.6	3.0	18.0	39.3	3.8	24.5	21.1		
Institutions under the justice system	100.0	34.9	13.1	2.9	7.7	18.6	20.0	17.5	44.6		

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category. Mental health diagnostic category definitions can be found in Appendix E.

² Total individuals with co-occurring mental health and substance use disorders who had at least one valid mental health diagnosis other than an alcohol- or substance-related mental health disorder.

³ Individuals could receive services in more than one service setting during the reporting period.

⁴ Mental health service setting category definitions can be found in Appendix C.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 7.1a. Individuals served in both 2017 and 2020, by living arrangement and selected characteristics in 2017: Number, 2017

Selected characteristics		Living arrangement							
		Total	Homeless	Foster home/care	Residential care	Institutional setting	Jail/ correctional facility	Private residence	Other
Number of states reporting	45		42	35	41	39	41	44	36
Total individuals ¹	1,296,090		40,547	18,217	71,547	21,682	15,777	1,108,799	19,521
Gender									
Male	624,529		23,568	9,813	42,687	12,584	12,407	514,492	8,978
Female	670,972		16,958	8,396	28,817	9,063	3,363	593,839	10,536
Total	1,295,501		40,526	18,209	71,504	21,647	15,770	1,108,331	19,514
Age ²									
0 to 11 years	131,573		387	5,512	961	152	39	123,613	909
12 to 14 years	64,683		149	2,020	1,188	201	391	59,993	741
15 to 17 years	47,341		197	1,446	1,382	223	760	42,657	676
18 to 20 years	38,571		789	678	1,220	352	607	34,179	746
21 to 25 years	64,621		2,142	828	2,593	965	1,380	55,634	1,079
26 to 29 years	100,733		3,972	1,102	4,571	1,629	2,383	85,304	1,772
30 to 34 years	105,088		4,559	982	5,146	1,761	2,290	88,453	1,897
35 to 39 years	105,848		4,487	842	5,526	1,769	2,094	89,202	1,928
40 to 44 years	95,908		4,280	715	5,101	1,479	1,496	81,228	1,609
45 to 49 years	113,124		5,343	726	6,687	1,719	1,442	95,324	1,883
50 to 54 years	128,804		5,813	893	8,806	2,170	1,253	107,833	2,036
55 to 59 years	121,261		4,497	885	10,018	2,494	858	100,607	1,902
60 to 64 years	81,736		2,345	724	7,882	2,085	432	67,018	1,250
65 years and older	96,727		1,586	862	10,458	4,676	349	77,705	1,091
Total	1,296,018		40,546	18,215	71,539	21,675	15,774	1,108,750	19,519
SED status ³									
SED	184,702		594	8,036	3,090	522	1,095	169,236	2,129
Not SED	51,335		97	811	298	29	68	49,872	160
Total	236,037		691	8,847	3,388	551	1,163	219,108	2,289
SMI status ³									
SMI	752,493		33,150	5,059	54,635	16,319	11,003	617,804	14,523
Not SMI	261,382		4,801	3,936	11,068	2,947	2,580	233,643	2,407
Total	1,013,875		37,951	8,995	65,703	19,266	13,583	851,447	16,930

Continued. See notes at end of table.

Table 7.1a. Individuals served in both 2017 and 2020, by living arrangement and selected characteristics in 2017: Number, 2017 (continued)

Selected characteristics	Total	Living arrangement						
		Homeless	Foster home/care	Residential care	Institutional setting	Jail/correctional facility	Private residence	Other
Substance use diagnosis								
Total individuals aged 12 years and older with co-occurring MH and SU disorders	260,770	16,336	1,797	20,224	6,091	7,684	202,093	6,545
Alcoholic psychoses	3,019	163	15	249	115	52	2,360	65
Alcohol intoxication	1,182	165	10	79	36	35	786	71
Drug psychoses	6,165	613	28	377	346	182	4,513	106
Dependence								
Alcohol	37,650	2,381	157	2,840	809	1,642	29,202	619
Cocaine	8,208	610	17	565	107	310	6,509	90
Marijuana	17,512	908	92	954	245	1,086	13,976	251
Opioid	11,366	752	23	547	171	675	8,868	330
Other	41,586	2,853	329	3,645	1,087	1,164	31,608	900
Non-dependent abuse								
Alcohol	16,930	784	76	1,433	414	496	13,446	281
Cocaine	4,650	325	19	375	84	106	3,682	59
Marijuana	19,241	888	137	1,069	367	542	15,960	278
Opioid	2,459	131	15	131	47	61	2,011	63
Other	18,010	1,111	89	1,497	851	422	13,733	307
None reported	72,792	4,652	790	6,463	1,412	911	55,439	3,125

Continued. See notes at end of table.

Table 7.1a. Individuals served in both 2017 and 2020, by living arrangement and selected characteristics in 2017: Number, 2017 (continued)

		Living arrangement						
Selected characteristics	Total	Homeless	Foster home/care	Residential care	Institutional setting	Jail/ correctional facility	Private residence	Other
Mental health diagnosis								
Trauma- and stressor-related disorders	201,505	6,143	5,432	5,244	1,375	3,093	176,001	4,217
Anxiety disorders	240,510	5,341	2,975	7,275	1,911	1,564	216,791	4,653
Attention deficit/hyperactivity disorder	135,914	859	3,002	2,399	376	519	128,126	633
Mood disorders	520,146	18,702	3,931	18,920	5,871	4,426	460,255	8,041
Bipolar disorders	188,719	7,737	1,217	8,956	2,601	1,887	163,437	2,884
Depressive disorders	342,349	11,568	2,811	10,428	3,437	2,673	306,115	5,317
Personality disorders	61,706	2,439	596	4,002	1,499	2,587	49,718	865
Schizophrenia and other psychotic disorders	261,679	11,370	2,881	34,841	10,082	4,161	193,078	5,266
Alcohol or substance-related disorders	72,131	5,870	199	3,759	1,528	1,387	58,692	696
Other mental disorders ⁴	322,706	5,770	6,474	21,707	4,307	3,939	277,765	2,744
No/deferred diagnosis	106,616	4,011	1,236	5,388	3,436	1,639	90,158	748

MH = mental health; SED = serious emotional disturbance; SMI = serious mental illness; SU = substance use.

¹ Of 2,291,260 individuals who were served in both the 2017 and 2020 reporting periods, the total number of individuals presented in this table consists only of those with known living arrangement in both reporting years.

² Age is based on the individual's age at the midpoint of the reporting period in 2017.

³ The federal definitions for SED and SMI status can be found in Appendix D.

⁴ "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol or substance-related disorders. See Appendix E.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2017 and 2020 reporting periods. Data as of 03.03.2022.

Table 7.1b. Individuals served in both 2017 and 2020, by living arrangement and selected characteristics in 2017: Number, 2017

Selected characteristics	Living arrangement							
	Total	Homeless	Foster home/care	Residential care	Institutional setting	Jail/correctional facility	Private residence	Other
Number of states reporting	45	42	35	41	39	41	44	36
Total individuals¹	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender								
Male	48.2	58.2	53.9	59.7	58.1	78.7	46.4	46.0
Female	51.8	41.8	46.1	40.3	41.9	21.3	53.6	54.0
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age²								
0 to 11 years	10.2	†	30.3	1.3	†	†	11.1	4.7
12 to 14 years	5.0	†	11.1	1.7	†	2.5	5.4	3.8
15 to 17 years	3.7	†	7.9	1.9	1.0	4.8	3.8	3.5
18 to 20 years	3.0	1.9	3.7	1.7	1.6	3.8	3.1	3.8
21 to 25 years	5.0	5.3	4.5	3.6	4.5	8.7	5.0	5.5
26 to 29 years	7.8	9.8	6.0	6.4	7.5	15.1	7.7	9.1
30 to 34 years	8.1	11.2	5.4	7.2	8.1	14.5	8.0	9.7
35 to 39 years	8.2	11.1	4.6	7.7	8.2	13.3	8.0	9.9
40 to 44 years	7.4	10.6	3.9	7.1	6.8	9.5	7.3	8.2
45 to 49 years	8.7	13.2	4.0	9.3	7.9	9.1	8.6	9.6
50 to 54 years	9.9	14.3	4.9	12.3	10.0	7.9	9.7	10.4
55 to 59 years	9.4	11.1	4.9	14.0	11.5	5.4	9.1	9.7
60 to 64 years	6.3	5.8	4.0	11.0	9.6	2.7	6.0	6.4
65 years and older	7.5	3.9	4.7	14.6	21.6	2.2	7.0	5.6
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
SED status³								
SED	78.3	86.0	90.8	91.2	94.7	94.2	77.2	93.0
Not SED	21.7	14.0	9.2	8.8	5.3	5.8	22.8	7.0
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
SMI status³								
SMI	74.2	87.3	56.2	83.2	84.7	81.0	72.6	85.8
Not SMI	25.8	12.7	43.8	16.8	15.3	19.0	27.4	14.2
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 7.1b. Individuals served in both 2017 and 2020, by living arrangement and selected characteristics in 2017: Column percent distribution, 2017 (continued)

<i>Selected characteristics</i>	<i>Total</i>	<i>Living arrangement</i>						
		<i>Homeless</i>	<i>Foster home/care</i>	<i>Residential care</i>	<i>Institutional setting</i>	<i>Jail/correctional facility</i>	<i>Private residence</i>	<i>Other</i>
Substance use diagnosis								
Total individuals aged 12 years and older with co-occurring MH and SU disorders	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Alcoholic psychoses	1.2	†	†	1.2	1.9	†	1.2	†
Alcohol intoxication	†	1.0	†	†	†	†	†	1.1
Drug psychoses	2.4	3.8	1.6	1.9	5.7	2.4	2.2	1.6
Dependence								
Alcohol	14.4	14.6	8.7	14.0	13.3	21.4	14.4	9.5
Cocaine	3.1	3.7	†	2.8	1.8	4.0	3.2	1.4
Marijuana	6.7	5.6	5.1	4.7	4.0	14.1	6.9	3.8
Opioid	4.4	4.6	1.3	2.7	2.8	8.8	4.4	5.0
Other	15.9	17.5	18.3	18.0	17.8	15.1	15.6	13.8
Non-dependent abuse								
Alcohol	6.5	4.8	4.2	7.1	6.8	6.5	6.7	4.3
Cocaine	1.8	2.0	1.1	1.9	1.4	1.4	1.8	†
Marijuana	7.4	5.4	7.6	5.3	6.0	7.1	7.9	4.2
Opioid	†	†	†	†	†	†	†	†
Other	6.9	6.8	5.0	7.4	14.0	5.5	6.8	4.7
None reported	27.9	28.5	44.0	32.0	23.2	11.9	27.4	47.7

Continued. See notes at end of table.

Table 7.1b. Individuals served in both 2017 and 2020, by living arrangement and selected characteristics in 2017: Column percent distribution, 2017 (continued)

		Living arrangement						
Selected characteristics	Total	Homeless	Foster home/care	Residential care	Institutional setting	Jail/ correctional facility	Private residence	Other
Mental health diagnosis								
Trauma- and stressor-related disorders	15.5	15.2	29.8	7.3	6.3	19.6	15.9	21.6
Anxiety disorders	18.6	13.2	16.3	10.2	8.8	9.9	19.6	23.8
Attention deficit/hyperactivity disorder	10.5	2.1	16.5	3.4	1.7	3.3	11.6	3.2
Mood disorders	40.1	46.1	21.6	26.4	27.1	28.1	41.5	41.2
Bipolar disorders	14.6	19.1	6.7	12.5	12.0	12.0	14.7	14.8
Depressive disorders	26.4	28.5	15.4	14.6	15.9	16.9	27.6	27.2
Personality disorders	4.8	6.0	3.3	5.6	6.9	16.4	4.5	4.4
Schizophrenia and other psychotic disorders	20.2	28.0	15.8	48.7	46.5	26.4	17.4	27.0
Alcohol or substance-related disorders	5.6	14.5	1.1	5.3	7.0	8.8	5.3	3.6
Other mental disorders ⁴	24.9	14.2	35.5	30.3	19.9	25.0	25.1	14.1
No/deferred diagnosis	8.2	9.9	6.8	7.5	15.8	10.4	8.1	3.8

† Less than 1 percent of individuals.

MH = mental health; SED = serious emotional disturbance; SMI = serious mental illness; SU = substance use.

¹ Of 2,291,260 individuals who were served in both the 2017 and 2020 reporting periods, the total number of individuals presented in this table consists only of those with known living arrangement in both reporting years.

² Age is based on the individual's age at the midpoint of the reporting period in 2017.

³ The federal definitions for SED and SMI status can be found in Appendix D.

⁴ "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol or substance-related disorders. See Appendix E.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2017 and 2020 reporting periods. Data as of 03.03.2022.

Table 7.1c. Individuals served in both 2017 and 2020, by living arrangement and selected characteristics in 2017: Column percent distribution, 2017

Selected characteristics	Total	Living arrangement						
		Homeless	Foster home/care	Residential care	Institutional setting	Jail/correctional facility	Private residence	Other
Number of states reporting	45	42	35	41	39	41	44	36
Total individuals¹	100.0	3.1	1.4	5.5	1.7	1.2	85.5	1.5
Gender								
Male	100.0	3.8	1.6	6.8	2.0	2.0	82.4	1.4
Female	100.0	2.5	1.3	4.3	1.4	†	88.5	1.6
Total	100.0	3.1	1.4	5.5	1.7	1.2	85.6	1.5
Age²								
0 to 11 years	100.0	†	4.2	†	†	†	94.0	†
12 to 14 years	100.0	†	3.1	1.8	†	†	92.7	1.1
15 to 17 years	100.0	†	3.1	2.9	†	1.6	90.1	1.4
18 to 20 years	100.0	2.0	1.8	3.2	†	1.6	88.6	1.9
21 to 25 years	100.0	3.3	1.3	4.0	1.5	2.1	86.1	1.7
26 to 29 years	100.0	3.9	1.1	4.5	1.6	2.4	84.7	1.8
30 to 34 years	100.0	4.3	†	4.9	1.7	2.2	84.2	1.8
35 to 39 years	100.0	4.2	†	5.2	1.7	2.0	84.3	1.8
40 to 44 years	100.0	4.5	†	5.3	1.5	1.6	84.7	1.7
45 to 49 years	100.0	4.7	†	5.9	1.5	1.3	84.3	1.7
50 to 54 years	100.0	4.5	†	6.8	1.7	†	83.7	1.6
55 to 59 years	100.0	3.7	†	8.3	2.1	†	83.0	1.6
60 to 64 years	100.0	2.9	†	9.6	2.6	†	82.0	1.5
65 years and older	100.0	1.6	†	10.8	4.8	†	80.3	1.1
Total	100.0	3.1	1.4	5.5	1.7	1.2	85.6	1.5
SED status³								
SED	100.0	†	4.4	1.7	†	†	91.6	1.2
Not SED	100.0	†	1.6	†	†	†	97.2	†
Total	100.0	†	3.7	1.4	†	†	92.8	†
SMI status³								
SMI	100.0	4.4	†	7.3	2.2	1.5	82.1	1.9
Not SMI	100.0	1.8	1.5	4.2	1.1	†	89.4	†
Total	100.0	3.7	†	6.5	1.9	1.3	84.0	1.7

Continued. See notes at end of table.

Table 7.1c. Individuals served in both 2017 and 2020, by living arrangement and selected characteristics in 2017: Row percent distribution, 2017 (continued)

		Living arrangement							
Selected characteristics	Total	Homeless	Foster home/care	Residential care	Institutional setting	Jail/ correctional facility	Private residence	Other	
Substance use diagnosis									
Total individuals aged 12 years and older with co-occurring MH and SU disorders		100.0	6.3	†	7.8	2.3	2.9	77.5	2.5
Alcoholic psychoses		100.0	5.4	†	8.2	3.8	1.7	78.2	2.2
Alcohol intoxication		100.0	14.0	†	6.7	3.0	3.0	66.5	6.0
Drug psychoses		100.0	9.9	†	6.1	5.6	3.0	73.2	1.7
Dependence									
Alcohol		100.0	6.3	†	7.5	2.1	4.4	77.6	1.6
Cocaine		100.0	7.4	†	6.9	1.3	3.8	79.3	1.1
Marijuana		100.0	5.2	†	5.4	1.4	6.2	79.8	1.4
Opioid		100.0	6.6	†	4.8	1.5	5.9	78.0	2.9
Other		100.0	6.9	†	8.8	2.6	2.8	76.0	2.2
Non-dependent abuse									
Alcohol		100.0	4.6	†	8.5	2.4	2.9	79.4	1.7
Cocaine		100.0	7.0	†	8.1	1.8	2.3	79.2	1.3
Marijuana		100.0	4.6	†	5.6	1.9	2.8	82.9	1.4
Opioid		100.0	5.3	†	5.3	1.9	2.5	81.8	2.6
Other		100.0	6.2	†	8.3	4.7	2.3	76.3	1.7
None reported		100.0	6.4	1.1	8.9	1.9	1.3	76.2	4.3

Continued. See notes at end of table.

Table 7.1c. Individuals served in both 2017 and 2020, by living arrangement and selected characteristics in 2017: Row percent distribution, 2017 (continued)

		Living arrangement						
Selected characteristics	Total	Homeless	Foster home/care	Residential care	Institutional setting	Jail/ correctional facility	Private residence	Other
Mental health diagnosis								
Trauma- and stressor-related disorders	100.0	3.0	2.7	2.6	†	1.5	87.3	2.1
Anxiety disorders	100.0	2.2	1.2	3.0	†	†	90.1	1.9
Attention deficit/hyperactivity disorder	100.0	†	2.2	1.8	†	†	94.3	†
Mood disorders	100.0	3.6	†	3.6	1.1	†	88.5	1.5
Bipolar disorders	100.0	4.1	†	4.7	1.4	†	86.6	1.5
Depressive disorders	100.0	3.4	†	3.0	1.0	†	89.4	1.6
Personality disorders	100.0	4.0	†	6.5	2.4	4.2	80.6	1.4
Schizophrenia and other psychotic disorders	100.0	4.3	1.1	13.3	3.9	1.6	73.8	2.0
Alcohol or substance-related disorders	100.0	8.1	†	5.2	2.1	1.9	81.4	†
Other mental disorders ⁴	100.0	1.8	2.0	6.7	1.3	1.2	86.1	†
No/deferred diagnosis	100.0	3.8	1.2	5.1	3.2	1.5	84.6	†

† Less than 1 percent of individuals.

MH = mental health; SED = serious emotional disturbance; SMI = serious mental illness; SU = substance use.

¹ Of 2,291,260 individuals who were served in both the 2017 and 2020 reporting periods, the total number of individuals presented in this table consists only of those with known living arrangement in both reporting years.

² Age is based on the individual's age at the midpoint of the reporting period in 2017.

³ The federal definitions for SED and SMI status can be found in Appendix D.

⁴ "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol or substance-related disorders. See Appendix E.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2017 and 2020 reporting periods. Data as of 03.03.2022.

Table 7.2a. Individuals served in both 2017 and 2020, by living arrangement and selected characteristics in 2020: Number, 2020

Selected characteristics	Living arrangement							
	Total	Homeless	Foster home/care	Residential care	Institutional setting	Jail/ correctional facility	Private residence	Other
Number of states reporting	45	42	34	40	41	40	44	39
Total individuals¹	1,296,090	43,985	17,667	73,371	22,555	15,329	1,105,041	18,142
Gender								
Male	624,529	25,801	9,656	43,873	12,766	12,286	511,216	8,931
Female	670,972	18,159	8,003	29,454	9,757	3,035	593,358	9,206
Total	1,295,501	43,960	17,659	73,327	22,523	15,321	1,104,574	18,137
Age²								
0 to 11 years	70,655	198	2,951	472	72	22	66,388	552
12 to 14 years	61,734	149	1,991	861	122	135	57,863	613
15 to 17 years	64,738	210	2,168	1,500	217	780	59,004	859
18 to 20 years	46,820	716	1,054	1,158	242	763	41,989	898
21 to 25 years	52,906	1,629	784	1,867	601	854	46,360	811
26 to 29 years	89,617	3,747	1,151	4,067	1,383	1,890	76,057	1,322
30 to 34 years	103,309	4,860	1,061	5,036	1,497	2,283	86,979	1,593
35 to 39 years	107,139	5,309	996	5,453	1,647	2,199	89,776	1,759
40 to 44 years	99,907	4,799	797	5,314	1,475	1,617	84,352	1,553
45 to 49 years	102,682	4,997	743	5,740	1,540	1,301	86,843	1,518
50 to 54 years	117,518	5,697	781	7,573	1,873	1,261	98,675	1,658
55 to 59 years	131,636	5,476	980	9,936	2,526	1,059	109,794	1,865
60 to 64 years	107,556	3,511	876	9,480	2,645	630	88,924	1,490
65 years and older	139,801	2,685	1,332	14,906	6,709	530	111,989	1,650
Total	1,296,018	43,983	17,665	73,363	22,549	15,324	1,104,993	18,141
SED status³								
SED	156,539	428	6,399	2,509	370	866	144,126	1,841
Not SED	33,208	93	561	145	22	44	32,237	106
Total	189,747	521	6,960	2,654	392	910	176,363	1,947
SMI status³								
SMI	798,918	36,265	6,656	57,277	17,682	11,243	655,495	14,300
Not SMI	261,467	5,314	3,617	11,379	2,943	2,122	234,608	1,484
Total	1,060,385	41,579	10,273	68,656	20,625	13,365	890,103	15,784

Continued. See notes at end of table.

Table 7.2a. Individuals served in both 2017 and 2020, by living arrangement and selected characteristics in 2020: Number, 2020 (continued)

Selected characteristics	Total	Living arrangement						
		Homeless	Foster home/care	Residential care	Institutional setting	Jail/ correctional facility	Private residence	Other
Substance use diagnosis								
Total individuals aged 12 years and older with co-occurring MH and SU disorders	339,870	21,725	2,786	23,801	6,733	8,469	269,734	6,622
Alcoholic psychoses	2,677	139	22	258	111	44	2,050	53
Alcohol intoxication	1,347	187	14	71	68	23	928	56
Drug psychoses	7,606	872	71	525	338	220	5,408	172
Dependence								
Alcohol	45,374	3,005	292	3,144	831	1,863	35,386	853
Cocaine	9,473	825	30	609	102	323	7,488	96
Marijuana	22,513	1,224	180	1,243	264	1,147	18,081	374
Opioid	15,131	1,097	60	690	183	745	11,961	395
Other	54,228	4,085	821	4,579	1,285	1,145	40,287	2,026
Non-dependent abuse								
Alcohol	17,839	906	120	1,462	418	433	14,223	277
Cocaine	4,842	393	18	374	84	84	3,826	63
Marijuana	22,957	1,036	231	1,157	400	530	19,232	371
Opioid	2,756	145	12	136	65	72	2,272	54
Other	20,343	1,469	202	1,620	815	375	15,434	428
None reported	112,784	6,342	713	7,933	1,769	1,465	93,158	1,404

Continued. See notes at end of table.

Table 7.2a. Individuals served in both 2017 and 2020, by living arrangement and selected characteristics in 2020: Number, 2020 (continued)

Selected characteristics	Total	Living arrangement						
		Homeless	Foster home/care	Residential care	Institutional setting	Jail/ correctional facility	Private residence	Other
Mental health diagnosis								
Trauma- and stressor-related disorders	217,182	7,256	4,793	5,920	1,554	3,216	190,828	3,615
Anxiety disorders	282,023	6,692	3,751	8,439	2,434	1,614	254,145	4,948
Attention deficit/hyperactivity disorder	145,115	1,132	3,314	2,803	395	636	135,448	1,387
Mood disorders	543,850	19,681	4,696	19,560	6,369	4,121	481,524	7,899
Bipolar disorders	198,412	8,318	1,415	9,181	2,666	1,773	172,287	2,772
Depressive disorders	358,383	12,030	3,433	10,893	3,940	2,505	320,226	5,356
Personality disorders	65,046	2,683	693	4,023	1,379	2,759	52,649	860
Schizophrenia and other psychotic disorders	276,080	13,323	3,386	36,657	10,448	4,021	202,736	5,509
Alcohol or substance-related disorders	79,488	7,275	213	4,285	1,647	1,386	63,599	1,083
Other mental disorders ⁴	331,484	6,350	6,739	22,624	4,732	4,149	283,538	3,352
No/deferred diagnosis	98,104	4,130	1,199	5,293	3,546	1,530	81,844	562

MH = mental health; SED = serious emotional disturbance; SMI = serious mental illness; SU = substance use.

¹ Of 2,291,260 individuals who were served in both the 2017 and 2020 reporting periods, the total number of individuals presented in this table consists only of those with known living arrangement in both reporting years.

² Age is based on the individual's age at the midpoint of the reporting period in 2020.

³ The federal definitions for SED and SMI status can be found in Appendix D.

⁴ "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol or substance-related disorders. See Appendix E.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2017 and 2020 reporting periods. Data as of 03.03.2022.

Table 7.2b. Individuals served in both 2017 and 2020, by living arrangement and selected characteristics in 2020: Column percent distribution, 2020

Selected characteristics	Total	Living arrangement						
		Homeless	Foster home/care	Residential care	Institutional setting	Jail/correctional facility	Private residence	Other
Number of states reporting	45	42	34	40	41	40	44	39
Total individuals¹	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender								
Male	48.2	58.7	54.7	59.8	56.7	80.2	46.3	49.2
Female	51.8	41.3	45.3	40.2	43.3	19.8	53.7	50.8
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age²								
0 to 11 years	5.5	†	16.7	†	†	†	6.0	3.0
12 to 14 years	4.8	†	11.3	1.2	†	†	5.2	3.4
15 to 17 years	5.0	†	12.3	2.0	†	5.1	5.3	4.7
18 to 20 years	3.6	1.6	6.0	1.6	1.1	5.0	3.8	5.0
21 to 25 years	4.1	3.7	4.4	2.5	2.7	5.6	4.2	4.5
26 to 29 years	6.9	8.5	6.5	5.5	6.1	12.3	6.9	7.3
30 to 34 years	8.0	11.0	6.0	6.9	6.6	14.9	7.9	8.8
35 to 39 years	8.3	12.1	5.6	7.4	7.3	14.4	8.1	9.7
40 to 44 years	7.7	10.9	4.5	7.2	6.5	10.6	7.6	8.6
45 to 49 years	7.9	11.4	4.2	7.8	6.8	8.5	7.9	8.4
50 to 54 years	9.1	13.0	4.4	10.3	8.3	8.2	8.9	9.1
55 to 59 years	10.2	12.5	5.5	13.5	11.2	6.9	9.9	10.3
60 to 64 years	8.3	8.0	5.0	12.9	11.7	4.1	8.0	8.2
65 years and older	10.8	6.1	7.5	20.3	29.8	3.5	10.1	9.1
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
SED status³								
SED	82.5	82.1	91.9	94.5	94.4	95.2	81.7	94.6
Not SED	17.5	17.9	8.1	5.5	5.6	4.8	18.3	5.4
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
SMI status³								
SMI	75.3	87.2	64.8	83.4	85.7	84.1	73.6	90.6
Not SMI	24.7	12.8	35.2	16.6	14.3	15.9	26.4	9.4
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 7.2b. Individuals served in both 2017 and 2020, by living arrangement and selected characteristics in 2020: Column percent distribution, 2020 (continued)

Selected characteristics	Total	Living arrangement						
		Homeless	Foster home/care	Residential care	Institutional setting	Jail/ correctional facility	Private residence	Other
Substance use diagnosis								
Total individuals aged 12 years and older with co-occurring MH and SU disorders	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Alcoholic psychoses	†	†	†	1.1	1.6	†	†	†
Alcohol intoxication	†	†	†	†	1.0	†	†	†
Drug psychoses	2.2	4.0	2.5	2.2	5.0	2.6	2.0	2.6
Dependence								
Alcohol	13.4	13.8	10.5	13.2	12.3	22.0	13.1	12.9
Cocaine	2.8	3.8	1.1	2.6	1.5	3.8	2.8	1.4
Marijuana	6.6	5.6	6.5	5.2	3.9	13.5	6.7	5.6
Opioid	4.5	5.0	2.2	2.9	2.7	8.8	4.4	6.0
Other	16.0	18.8	29.5	19.2	19.1	13.5	14.9	30.6
Non-dependent abuse								
Alcohol	5.2	4.2	4.3	6.1	6.2	5.1	5.3	4.2
Cocaine	1.4	1.8	†	1.6	1.2	†	1.4	†
Marijuana	6.8	4.8	8.3	4.9	5.9	6.3	7.1	5.6
Opioid	†	†	†	†	†	†	†	†
Other	6.0	6.8	7.3	6.8	12.1	4.4	5.7	6.5
None reported	33.2	29.2	25.6	33.3	26.3	17.3	34.5	21.2

Continued. See notes at end of table.

Table 7.2b. Individuals served in both 2017 and 2020, by living arrangement and selected characteristics in 2020: Column percent distribution, 2020 (continued)

Selected characteristics	Total	Living arrangement						
		Homeless	Foster home/care	Residential care	Institutional setting	Jail/correctional facility	Private residence	Other
Mental health diagnosis								
Trauma- and stressor-related disorders	16.8	16.5	27.1	8.1	6.9	21.0	17.3	19.9
Anxiety disorders	21.8	15.2	21.2	11.5	10.8	10.5	23.0	27.3
Attention deficit/hyperactivity disorder	11.2	2.6	18.8	3.8	1.8	4.1	12.3	7.6
Mood disorders	42.0	44.7	26.6	26.7	28.2	26.9	43.6	43.5
Bipolar disorders	15.3	18.9	8.0	12.5	11.8	11.6	15.6	15.3
Depressive disorders	27.7	27.4	19.4	14.8	17.5	16.3	29.0	29.5
Personality disorders	5.0	6.1	3.9	5.5	6.1	18.0	4.8	4.7
Schizophrenia and other psychotic disorders	21.3	30.3	19.2	50.0	46.3	26.2	18.3	30.4
Alcohol or substance-related disorders	6.1	16.5	1.2	5.8	7.3	9.0	5.8	6.0
Other mental disorders ⁴	25.6	14.4	38.1	30.8	21.0	27.1	25.7	18.5
No/deferred diagnosis	7.6	9.4	6.8	7.2	15.7	10.0	7.4	3.1

† Less than 1 percent of individuals.

MH = mental health; SED = serious emotional disturbance; SMI = serious mental illness; SU = substance use.

¹ Of 2,291,260 individuals who were served in both the 2017 and 2020 reporting periods, the total number of individuals presented in this table consists only of those with known living arrangement in both reporting years.

² Age is based on the individual's age at the midpoint of the reporting period in 2020.

³ The federal definitions for SED and SMI status can be found in Appendix D.

⁴ "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol or substance-related disorders. See Appendix E.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2017 and 2020 reporting periods. Data as of 03.03.2022.

Table 7.2c. Individuals served in both 2017 and 2020, by living arrangement and selected characteristics in 2020: Row percent distribution, 2020

Selected characteristics		Living arrangement							
		Total	Homeless	Foster home/care	Residential care	Institutional setting	Jail/ correctional facility	Private residence	Other
Number of states reporting	45		42	34	40	41	40	44	39
Total individuals¹	100.0		3.4	1.4	5.7	1.7	1.2	85.3	1.4
Gender									
Male	100.0		4.1	1.5	7.0	2.0	2.0	81.9	1.4
Female	100.0		2.7	1.2	4.4	1.5	†	88.4	1.4
Total	100.0		3.4	1.4	5.7	1.7	1.2	85.3	1.4
Age²									
0 to 11 years	100.0		†	4.2	†	†	†	94.0	†
12 to 15 years	100.0		†	3.2	1.4	†	†	93.7	†
15 to 17 years	100.0		†	3.3	2.3	†	1.2	91.1	1.3
18 to 20 years	100.0		1.5	2.3	2.5	†	1.6	89.7	1.9
21 to 25 years	100.0		3.1	1.5	3.5	1.1	1.6	87.6	1.5
26 to 29 years	100.0		4.2	1.3	4.5	1.5	2.1	84.9	1.5
30 to 34 years	100.0		4.7	1.0	4.9	1.4	2.2	84.2	1.5
35 to 39 years	100.0		5.0	†	5.1	1.5	2.1	83.8	1.6
40 to 44 years	100.0		4.8	†	5.3	1.5	1.6	84.4	1.6
45 to 49 years	100.0		4.9	†	5.6	1.5	1.3	84.6	1.5
50 to 54 years	100.0		4.8	†	6.4	1.6	1.1	84.0	1.4
55 to 59 years	100.0		4.2	†	7.5	1.9	†	83.4	1.4
60 to 64 years	100.0		3.3	†	8.8	2.5	†	82.7	1.4
65 years and older	100.0		1.9	†	10.7	4.8	†	80.1	1.2
Total	100.0		3.4	1.4	5.7	1.7	1.2	85.3	1.4
SED status³									
SED	100.0		†	4.1	1.6	†	†	92.1	1.2
Not SED	100.0		†	1.7	†	†	†	97.1	†
Total	100.0		†	3.7	1.4	†	†	92.9	1.0
SMI status³									
SMI	100.0		4.5	†	7.2	2.2	1.4	82.0	1.8
Not SMI	100.0		2.0	1.4	4.4	1.1	†	89.7	†
Total	100.0		3.9	†	6.5	1.9	1.3	83.9	1.5

Continued. See notes at end of table.

Table 7.2c. Individuals served in both 2017 and 2020, by living arrangement and selected characteristics in 2020: Row percent distribution, 2020 (continued)

		Living arrangement							
Selected characteristics	Total	Homeless	Foster home/care	Residential care	Institutional setting	Jail/ correctional facility	Private residence	Other	
Substance use diagnosis									
Total individuals aged 12 and older with co-occurring MH and SU disorders		100.0	6.4	†	7.0	2.0	2.5	79.4	1.9
Alcoholic psychoses		100.0	5.2	†	9.6	4.1	1.6	76.6	2.0
Alcohol intoxication		100.0	13.9	1.0	5.3	5.0	1.7	68.9	4.2
Drug psychoses		100.0	11.5	†	6.9	4.4	2.9	71.1	2.3
Dependence									
Alcohol		100.0	6.6	†	6.9	1.8	4.1	78.0	1.9
Cocaine		100.0	8.7	†	6.4	1.1	3.4	79.0	1.0
Marijuana		100.0	5.4	†	5.5	1.2	5.1	80.3	1.7
Opioid		100.0	7.3	†	4.6	1.2	4.9	79.0	2.6
Other		100.0	7.5	1.5	8.4	2.4	2.1	74.3	3.7
Non-dependent abuse									
Alcohol		100.0	5.1	†	8.2	2.3	2.4	79.7	1.6
Cocaine		100.0	8.1	†	7.7	1.7	1.7	79.0	1.3
Marijuana		100.0	4.5	1.0	5.0	1.7	2.3	83.8	1.6
Opioid		100.0	5.3	†	4.9	2.4	2.6	82.4	2.0
Other		100.0	7.2	†	8.0	4.0	1.8	75.9	2.1
None reported		100.0	5.6	†	7.0	1.6	1.3	82.6	1.2

Continued. See notes at end of table.

Table 7.2c. Individuals served in both 2017 and 2020, by living arrangement and selected characteristics in 2020: Row percent distribution, 2020 (continued)

Selected characteristics	Total	Living arrangement						
		Homeless	Foster home/care	Residential care	Institutional setting	Jail/correctional facility	Private residence	Other
Mental health diagnosis								
Trauma- and stressor-related disorders	100.0	3.3	2.2	2.7	†	1.5	87.9	1.7
Anxiety disorders	100.0	2.4	1.3	3.0	†	†	90.1	1.8
Attention deficit/hyperactivity disorder	100.0	†	2.3	1.9	†	†	93.3	†
Mood disorders	100.0	3.6	†	3.6	1.2	†	88.5	1.5
Bipolar disorders	100.0	4.2	†	4.6	1.3	†	86.8	1.4
Depressive disorders	100.0	3.4	†	3.0	1.1	†	89.4	1.5
Personality disorders	100.0	4.1	1.1	6.2	2.1	4.2	80.9	1.3
Schizophrenia and other psychotic disorders	100.0	4.8	1.2	13.3	3.8	1.5	73.4	2.0
Alcohol or substance-related disorders	100.0	9.2	†	5.4	2.1	1.7	80.0	1.4
Other mental disorders ⁴	100.0	1.9	2.0	6.8	1.4	1.3	85.5	1.0
No/deferred diagnosis	100.0	4.2	1.2	5.4	3.6	1.6	83.4	†

† Less than 1 percent of individuals.

MH = mental health; SED = serious emotional disturbance; SMI = serious mental illness; SU = substance use.

¹ Of 2,291,260 individuals who were served in both the 2017 and 2020 reporting periods, the total number of individuals presented in this table consists only of those with known living arrangement in both reporting years.

² Age is based on the individual's age at the midpoint of the reporting period in 2020.

³ The federal definitions for SED and SMI status can be found in Appendix D.

⁴ "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol or substance-related disorders. See Appendix E.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2017 and 2020 reporting periods. Data as of 03.03.2022.

Table 7.3a. Individuals served in both 2017 and 2020, by employment status and selected characteristics in 2017: Number, 2017

<i>Selected characteristics</i>	<i>Employment status</i>			
	<i>Total</i>	<i>Employed</i>	<i>Unemployed</i>	<i>Not in labor force</i>
Number of states reporting	43	42	43	43
Total individuals aged 18 years and older¹	665,210	124,918	152,346	387,946
Gender				
Male	297,767	51,423	73,131	173,213
Female	367,270	73,457	79,139	214,674
Total	665,037	124,880	152,270	387,887
Age²				
18 to 20 years	26,047	4,454	5,587	16,006
21 to 25 years	42,134	10,338	12,258	19,538
26 to 29 years	67,626	17,846	19,492	30,288
30 to 34 years	70,586	18,010	19,639	32,937
35 to 39 years	70,634	16,335	18,451	35,848
40 to 44 years	64,134	13,252	15,723	35,159
45 to 49 years	74,187	13,306	17,087	43,794
50 to 54 years	83,482	12,718	17,057	53,707
55 to 59 years	76,588	10,445	13,536	52,607
60 to 64 years	48,266	5,522	7,275	35,469
65 years and older	41,526	2,692	6,241	32,593
Total	665,210	124,918	152,346	387,946
SMI status³				
SMI	534,305	93,568	122,434	318,303
Not SMI	112,240	28,807	23,686	59,747
Total	646,545	122,375	146,120	378,050

Continued. See notes at end of table.

Table 7.3a. Individuals served in both 2017 and 2020, by employment status and selected characteristics in 2017: Number, 2017 (continued)

<i>Selected characteristics</i>	<i>Employment status</i>			
	<i>Total</i>	<i>Employed</i>	<i>Unemployed</i>	<i>Not in labor force</i>
Substance use diagnosis				
Total individuals aged 18 years and older with co-occurring MH and SU disorders	183,453	32,797	43,677	106,979
Alcoholic psychoses	2,530	524	529	1,477
Alcohol intoxication	884	163	280	441
Drug psychoses	4,853	815	1,504	2,534
Dependence				
Alcohol	25,874	5,103	6,011	14,760
Cocaine	5,182	696	1,339	3,147
Marijuana	11,842	2,365	3,001	6,476
Opioid	9,120	1,907	2,882	4,331
Other	27,523	5,384	6,339	15,800
Non-dependent abuse				
Alcohol	13,772	2,645	2,680	8,447
Cocaine	2,992	423	662	1,907
Marijuana	13,070	2,481	3,075	7,514
Opioid	1,715	306	501	908
Other	8,862	1,638	2,044	5,180
None reported	55,234	8,347	12,830	34,057

Continued. See notes at end of table.

Table 7.3a. Individuals served in both 2017 and 2020, by employment status and selected characteristics in 2017: Number, 2017 (continued)

<i>Selected characteristics</i>	<i>Employment status</i>			
	<i>Total</i>	<i>Employed</i>	<i>Unemployed</i>	<i>Not in labor force</i>
Mental health diagnosis				
Trauma- and stressor-related disorders	92,555	20,796	21,257	50,502
Anxiety disorders	155,069	38,099	29,738	87,232
Attention deficit/hyperactivity disorder	28,191	8,487	5,175	14,529
Mood disorders	348,057	72,484	81,610	193,963
Bipolar disorders	137,392	26,459	32,996	77,937
Depressive disorders	220,309	48,028	51,393	120,888
Personality disorders	46,749	9,052	8,839	28,858
Schizophrenia and other psychotic disorders	180,074	20,565	38,059	121,450
Alcohol or substance use disorders	26,280	4,394	10,992	10,894
Other mental disorders ⁴	107,220	18,418	23,459	65,343
No/deferred diagnosis	35,335	6,341	8,612	20,382

MH = mental health; SMI = serious mental illness; SU = substance use.

¹ Of 2,291,260 individuals who were served in both the 2017 and 2020 reporting periods, the total number of individuals presented in this table consists only of those with known employment status in both reporting years. The total number of individuals aged 18 years and older in Table 7.3a is lower than that shown in Table 7.4a because of the progression in age of individuals who were served in 2017, but who did not become 18 years or older until 2020.

² Age is based on the individual's age at the midpoint of the reporting period in 2017.

³ The federal definition for SMI status can be found in Appendix D.

⁴ "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol or substance-related disorders. See Appendix E.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2017 and 2020 reporting periods. Data as of 03.03.2022.

**Table 7.3b. Individuals served in both 2017 and 2020, by employment status and selected characteristics
in 2017: Column percent distribution, 2020**

<i>Selected characteristics</i>	<i>Employment status</i>			
	<i>Total</i>	<i>Employed</i>	<i>Unemployed</i>	<i>Not in labor force</i>
Number of states reporting	43	42	43	43
Total individuals aged 18 years and older¹	100.0	100.0	100.0	100.0
Gender				
Male	44.8	41.2	48.0	44.7
Female	55.2	58.8	52.0	55.3
Total	100.0	100.0	100.0	100.0
Age²				
18 to 20 years	3.9	3.6	3.7	4.1
21 to 25 years	6.3	8.3	8.0	5.0
26 to 29 years	10.2	14.3	12.8	7.8
30 to 34 years	10.6	14.4	12.9	8.5
35 to 39 years	10.6	13.1	12.1	9.2
40 to 44 years	9.6	10.6	10.3	9.1
45 to 49 years	11.2	10.7	11.2	11.3
50 to 54 years	12.5	10.2	11.2	13.8
55 to 59 years	11.5	8.4	8.9	13.6
60 to 64 years	7.3	4.4	4.8	9.1
65 years and older	6.2	2.2	4.1	8.4
Total	100.0	100.0	100.0	100.0
SMI status³				
SMI	82.6	76.5	83.8	84.2
Not SMI	17.4	23.5	16.2	15.8
Total	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 7.3b. Individuals served in both 2017 and 2020, by employment status and selected characteristics in 2017: Column percent distribution, 2020 (continued)

<i>Selected characteristics</i>	<i>Employment status</i>			
	<i>Total</i>	<i>Employed</i>	<i>Unemployed</i>	<i>Not in labor force</i>
Substance use diagnosis				
Total individuals aged 18 years and older with co-occurring MH and SU disorders	100.0	100.0	100.0	100.0
Alcoholic psychoses	1.4	1.6	1.2	1.4
Alcohol intoxication	†	†	†	†
Drug psychoses	2.6	2.5	3.4	2.4
Dependence				
Alcohol	14.1	15.6	13.8	13.8
Cocaine	2.8	2.1	3.1	2.9
Marijuana	6.5	7.2	6.9	6.1
Opioid	5.0	5.8	6.6	4.0
Other	15.0	16.4	14.5	14.8
Non-dependent abuse				
Alcohol	7.5	8.1	6.1	7.9
Cocaine	1.6	1.3	1.5	1.8
Marijuana	7.1	7.6	7.0	7.0
Opioid	†	†	1.1	†
Other	4.8	5.0	4.7	4.8
None reported	30.1	25.5	29.4	31.8

Continued. See notes at end of table.

Table 7.3b. Individuals served in both 2017 and 2020, by employment status and selected characteristics in 2017: Column percent distribution, 2020 (continued)

<i>Selected characteristics</i>	<i>Employment status</i>			
	<i>Total</i>	<i>Employed</i>	<i>Unemployed</i>	<i>Not in labor force</i>
Mental health diagnosis				
Trauma- and stressor-related disorders	13.9	16.6	14.0	13.0
Anxiety disorders	23.3	30.5	19.5	22.5
Attention deficit/hyperactivity disorder	4.2	6.8	3.4	3.7
Mood disorders	52.3	58.0	53.6	50.0
Bipolar disorders	20.7	21.2	21.7	20.1
Depressive disorders	33.1	38.4	33.7	31.2
Personality disorders	7.0	7.2	5.8	7.4
Schizophrenia and other psychotic disorders	27.1	16.5	25.0	31.3
Alcohol or substance-related disorders	4.0	3.5	7.2	2.8
Other mental disorders ⁴	16.1	14.7	15.4	16.8
No/deferred diagnosis	5.3	5.1	5.7	5.3

† Less than 1 percent of individuals.

MH = mental health; SMI = serious mental illness; SU = substance use.

¹ Of 2,291,260 individuals who were served in both the 2017 and 2020 reporting periods, the total number of individuals presented in this table consists only of those with known employment status in both reporting years. The total number of individuals aged 18 years and older in Table 7.3a is lower than that shown in Table 7.4a because of the progression in age of individuals who were served in 2017, but who did not become 18 years or older until 2020.

² Age is based on the individual's age at the midpoint of the reporting period in 2017.

³ The federal definition for SMI status can be found in Appendix D.

⁴ "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol or substance-related disorders. See Appendix E.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2017 and 2020 reporting periods. Data as of 03.03.2022.

**Table 7.3c. Individuals served in both 2017 and 2020, by employment status and selected characteristics
in 2017: Row percent distribution, 2020**

<i>Selected characteristics</i>	<i>Employment status</i>			
	<i>Total</i>	<i>Employed</i>	<i>Unemployed</i>	<i>Not in labor force</i>
Number of states reporting	43	42	43	43
Total individuals aged 18 years and older¹	100.0	18.8	22.9	58.3
Gender				
Male	100.0	17.3	24.6	58.2
Female	100.0	20.0	21.5	58.5
Total	100.0	18.8	22.9	58.3
Age²				
18 to 20 years	100.0	17.1	21.4	61.5
21 to 25 years	100.0	24.5	29.1	46.4
26 to 29 years	100.0	26.4	28.8	44.8
30 to 34 years	100.0	25.5	27.8	46.7
35 to 39 years	100.0	23.1	26.1	50.8
40 to 44 years	100.0	20.7	24.5	54.8
45 to 49 years	100.0	17.9	23.0	59.0
50 to 54 years	100.0	15.2	20.4	64.3
55 to 59 years	100.0	13.6	17.7	68.7
60 to 64 years	100.0	11.4	15.1	73.5
65 years and older	100.0	6.5	15.0	78.5
Total	100.0	18.8	22.9	58.3
SMI status³				
SMI	100.0	17.5	22.9	59.6
Not SMI	100.0	25.7	21.1	53.2
Total	100.0	18.9	22.6	58.5
Continued. See notes at end of table.				

Table 7.3c. Individuals served in both 2017 and 2020, by employment status and selected characteristics in 2017: Row percent distribution, 2020 (continued)

<i>Selected characteristics</i>	<i>Employment status</i>			
	<i>Total</i>	<i>Employed</i>	<i>Unemployed</i>	<i>Not in labor force</i>
Substance use diagnosis				
Total individuals aged 18 years and older with co-occurring MH and SU disorders	100.0	17.9	23.8	58.3
Alcoholic psychoses	100.0	20.7	20.9	58.4
Alcohol intoxication	100.0	18.4	31.7	49.9
Drug psychoses	100.0	16.8	31.0	52.2
Dependence				
Alcohol	100.0	19.7	23.2	57.0
Cocaine	100.0	13.4	25.8	60.7
Marijuana	100.0	20.0	25.3	54.7
Opioid	100.0	20.9	31.6	47.5
Other	100.0	19.6	23.0	57.4
Non-dependent abuse				
Alcohol	100.0	19.2	19.5	61.3
Cocaine	100.0	14.1	22.1	63.7
Marijuana	100.0	19.0	23.5	57.5
Opioid	100.0	17.8	29.2	52.9
Other	100.0	18.5	23.1	58.5
None reported	100.0	15.1	23.2	61.7

Continued. See notes at end of table.

Table 7.3c. Individuals served in both 2017 and 2020, by employment status and selected characteristics in 2017: Row percent distribution, 2020 (continued)

<i>Selected characteristics</i>	<i>Employment status</i>			
	<i>Total</i>	<i>Employed</i>	<i>Unemployed</i>	<i>Not in labor force</i>
Mental health diagnosis				
Trauma- and stressor-related disorders	100.0	22.5	23.0	54.6
Anxiety disorders	100.0	24.6	19.2	56.3
Attention deficit/hyperactivity disorder	100.0	30.1	18.4	51.5
Mood disorders	100.0	20.8	23.4	55.7
Bipolar disorders	100.0	19.3	24.0	56.7
Depressive disorders	100.0	21.8	23.3	54.9
Personality disorders	100.0	19.4	18.9	61.7
Schizophrenia and other psychotic disorders	100.0	11.4	21.1	67.4
Alcohol or substance-related disorders	100.0	16.7	41.8	41.5
Other mental disorders ⁴	100.0	17.2	21.9	60.9
No/deferred diagnosis	100.0	17.9	24.4	57.7

MH = mental health; SMI = serious mental illness; SU = substance use.

¹ Of 2,291,260 individuals who were served in both the 2017 and 2020 reporting periods, the total number of individuals presented in this table consists only of those with known employment status in both reporting years. The total number of individuals aged 18 years and older in Table 7.3a is lower than that shown in Table 7.4a because of the progression in age of individuals who were served in 2017, but who did not become 18 years or older until 2020.

² Age is based on the individual's age at the midpoint of the reporting period in 2017.

³ The federal definition for SMI status can be found in Appendix D.

⁴ "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol or substance-related disorders. See Appendix E.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2017 and 2020 reporting periods. Data as of 03.03.2022.

Table 7.4a. Individuals served in both 2017 and 2020, by employment status and selected characteristics in 2020: Number, 2020

<i>Selected characteristics</i>	<i>Employment status</i>			
	<i>Total</i>	<i>Employed</i>	<i>Unemployed</i>	<i>Not in labor force</i>
Number of states reporting	43	42	43	42
Total individuals aged 18 years and older¹	683,855	131,123	152,781	399,951
Gender				
Male	306,481	53,458	73,668	179,355
Female	377,199	77,626	79,037	220,536
Total	683,680	131,084	152,705	399,891
Age²				
18 to 20 years	18,829	3,316	3,036	12,477
21 to 25 years	35,303	8,704	9,003	17,596
26 to 29 years	59,238	16,249	16,847	26,142
30 to 34 years	69,395	18,972	19,457	30,966
35 to 39 years	71,859	18,291	19,108	34,460
40 to 44 years	66,816	15,399	16,465	34,952
45 to 49 years	68,060	13,346	15,851	38,863
50 to 54 years	76,500	12,540	16,388	47,572
55 to 59 years	84,904	11,890	15,774	57,240
60 to 64 years	66,459	7,833	10,818	47,808
65 years and older	66,492	4,583	10,034	51,875
Total	683,855	131,123	152,781	399,951
SMI status³				
SMI	560,225	103,141	124,481	332,603
Not SMI	106,264	25,729	23,268	57,267
Total	666,489	128,870	147,749	389,870

Continued. See notes at end of table.

Table 7.4a. Individuals served in both 2017 and 2020, by employment status and selected characteristics in 2020: Number, 2020 (continued)

<i>Selected characteristics</i>	<i>Employment status</i>			
	<i>Total</i>	<i>Employed</i>	<i>Unemployed</i>	<i>Not in labor force</i>
Substance use diagnosis				
Total individuals aged 18 years and older with co-occurring MH and SU disorders	236,837	43,692	55,930	137,215
Alcoholic psychoses	2,236	412	405	1,419
Alcohol intoxication	919	173	288	458
Drug psychoses	5,623	980	1,522	3,121
Dependence				
Alcohol	31,970	6,418	7,215	18,337
Cocaine	6,378	863	1,447	4,068
Marijuana	14,601	3,107	3,478	8,016
Opioid	12,142	2,576	3,678	5,888
Other	35,835	7,440	8,024	20,371
Non-dependent abuse				
Alcohol	14,696	2,963	2,696	9,037
Cocaine	3,104	389	673	2,042
Marijuana	15,246	3,091	3,323	8,832
Opioid	1,952	377	470	1,105
Other	10,524	2,194	2,260	6,070
None reported	81,611	12,709	20,451	48,451

Continued. See notes at end of table.

Table 7.4a. Individuals served in both 2017 and 2020, by employment status and selected characteristics in 2020: Number, 2020 (continued)

<i>Selected characteristics</i>	<i>Employment status</i>			
	<i>Total</i>	<i>Employed</i>	<i>Unemployed</i>	<i>Not in labor force</i>
Mental health diagnosis				
Trauma- and stressor-related disorders	111,518	26,016	24,027	61,475
Anxiety disorders	185,822	46,418	33,229	106,175
Attention deficit/hyperactivity disorder	34,382	10,223	6,526	17,633
Mood disorders	361,310	77,919	79,401	203,990
Bipolar disorders	144,421	29,233	33,670	81,518
Depressive disorders	228,277	51,211	48,631	128,435
Personality disorders	49,179	10,130	9,182	29,867
psychotic disorders	190,447	21,439	40,421	128,587
Alcohol or substance-related disorders	29,834	5,067	12,396	12,371
Other mental disorders ⁴	118,165	20,094	26,590	71,481
No/deferred diagnosis	32,664	6,720	8,290	17,654

MH = mental health; SMI = serious mental illness; SU = substance use.

¹ Of 2,291,260 individuals who were served in both the 2017 and 2020 reporting periods, the total number of individuals presented in this table consists only of those with known employment status in both reporting years. The total number of individuals aged 18 years and older in Table 7.3a is lower than that shown in Table 7.4a because of the progression in age of individuals who were served in 2017, but who did not become 18 years or older until 2020.

² Age is based on the individual's age at the midpoint of the reporting period in 2020.

³ The federal definition for SMI status can be found in Appendix D.

⁴ "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol or substance-related disorders. See Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2017 and 2020 reporting periods. Data as of 03.03.2022.

Table 7.4b. Individuals served in both 2017 and 2020, by employment status and selected characteristics in 2020: Column percent distribution, 2020

Selected characteristics	Employment status			
	Total	Employed	Unemployed	Not in labor force
Number of states reporting	43	42	43	42
Total individuals aged 18 years and older¹	100.0	100.0	100.0	100.0
Gender				
Male	44.8	40.8	48.2	44.9
Female	55.2	59.2	51.8	55.1
Total	100.0	100.0	100.0	100.0
Age²				
18 to 20 years	2.8	2.5	2.0	3.1
21 to 25 years	5.2	6.6	5.9	4.4
26 to 29 years	8.7	12.4	11.0	6.5
30 to 34 years	10.1	14.5	12.7	7.7
35 to 39 years	10.5	13.9	12.5	8.6
40 to 44 years	9.8	11.7	10.8	8.7
45 to 49 years	10.0	10.2	10.4	9.7
50 to 54 years	11.2	9.6	10.7	11.9
55 to 59 years	12.4	9.1	10.3	14.3
60 to 64 years	9.7	6.0	7.1	12.0
65 years and older	9.7	3.5	6.6	13.0
Total	100.0	100.0	100.0	100.0
SMI status³				
SMI	84.1	80.0	84.3	85.3
Not SMI	15.9	20.0	15.7	14.7
Total	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 7.4b. Individuals served in both 2017 and 2020, by employment status and selected characteristics in 2020: Column percent distribution, 2020 (continued)

Selected characteristics	Employment status			
	Total	Employed	Unemployed	Not in labor force
Substance use diagnosis				
Total individuals aged 18 years and older with co-occurring MH and SU disorders	100.0	100.0	100.0	100.0
Alcoholic psychoses	†	†	†	1.0
Alcohol intoxication	†	†	†	†
Drug psychoses	2.4	2.2	2.7	2.3
Dependence				
Alcohol	13.5	14.7	12.9	13.4
Cocaine	2.7	2.0	2.6	3.0
Marijuana	6.2	7.1	6.2	5.8
Opioid	5.1	5.9	6.6	4.3
Other	15.1	17.0	14.3	14.8
Non-dependent abuse				
Alcohol	6.2	6.8	4.8	6.6
Cocaine	1.3	†	1.2	1.5
Marijuana	6.4	7.1	5.9	6.4
Opioid	†	†	†	†
Other	4.4	5.0	4.0	4.4
None reported	34.5	29.1	36.6	35.3

Continued. See notes at end of table.

Table 7.4b. Individuals served in both 2017 and 2020, by employment status and selected characteristics in 2020: Column percent distribution, 2020 (continued)

Selected characteristics	Employment status			
	Total	Employed	Unemployed	Not in labor force
Mental health diagnosis				
Trauma- and stressor-related disorders	16.3	19.8	15.7	15.4
Anxiety disorders	27.2	35.4	21.7	26.5
Attention deficit/hyperactivity disorder	5.0	7.8	4.3	4.4
Mood disorders	52.8	59.4	52.0	51.0
Bipolar disorders	21.1	22.3	22.0	20.4
Depressive disorders	33.4	39.1	31.8	32.1
Personality disorders	7.2	7.7	6.0	7.5
Schizophrenia and other psychotic disorders	27.8	16.4	26.5	32.2
Alcohol or substance-related disorders	4.4	3.9	8.1	3.1
Other mental disorders ⁴	17.3	15.3	17.4	17.9
No/deferred diagnosis	4.8	5.1	5.4	4.4

† Less than 1 percent of individuals.

MH = mental health; SMI = serious mental illness; SU = substance use.

¹ Of 2,291,260 individuals who were served in both the 2017 and 2020 reporting periods, the total number of individuals presented in this table consists only of those with known employment status in both reporting years. The total number of individuals aged 18 years and older in Table 7.3a is lower than that shown in Table 7.4a because of the progression in age of individuals who were served in 2017, but who did not become 18 years or older until 2020.

² Age is based on the individual's age at the midpoint of the reporting period in 2020.

³ The federal definition for SMI status can be found in Appendix D.

⁴ "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol or substance-related disorders. See Appendix E.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2017 and 2020 reporting periods. Data as of 03.03.2022.

Table 7.4c. Individuals served in both 2017 and 2020, by employment status and selected characteristics in 2020: Row percent distribution, 2020

Selected characteristics	Employment status			
	Total	Employed	Unemployed	Not in labor force
Number of states reporting	43	42	43	42
Total individuals aged 18 years and older¹	100.0	19.2	22.3	58.5
Gender				
Male	100.0	17.4	24.0	58.5
Female	100.0	20.6	21.0	58.5
Total	100.0	19.2	22.3	58.5
Age²				
18 to 20 years	100.0	17.6	16.1	66.3
21 to 25 years	100.0	24.7	25.5	49.8
26 to 29 years	100.0	27.4	28.4	44.1
30 to 34 years	100.0	27.3	28.0	44.6
35 to 39 years	100.0	25.5	26.6	48.0
40 to 44 years	100.0	23.0	24.6	52.3
45 to 49 years	100.0	19.6	23.3	57.1
50 to 54 years	100.0	16.4	21.4	62.2
55 to 59 years	100.0	14.0	18.6	67.4
60 to 64 years	100.0	11.8	16.3	71.9
65 years and older	100.0	6.9	15.1	78.0
Total	100.0	19.2	22.3	58.5
SMI status³				
SMI	100.0	18.4	22.2	59.4
Not SMI	100.0	24.2	21.9	53.9
Total	100.0	19.3	22.2	58.5

Continued. See notes at end of table.

Table 7.4c. Individuals served in both 2017 and 2020, by employment status and selected characteristics in 2020: Row percent distribution, 2020 (continued)

<i>Selected characteristics</i>	<i>Employment status</i>			
	<i>Total</i>	<i>Employed</i>	<i>Unemployed</i>	<i>Not in labor force</i>
Substance use diagnosis				
Total individuals aged 18 years and older with co-occurring MH and SU disorders	100.0	18.4	23.6	57.9
Alcoholic psychoses	100.0	18.4	18.1	63.5
Alcohol intoxication	100.0	18.8	31.3	49.8
Drug psychoses	100.0	17.4	27.1	55.5
Dependence				
Alcohol	100.0	20.1	22.6	57.4
Cocaine	100.0	13.5	22.7	63.8
Marijuana	100.0	21.3	23.8	54.9
Opioid	100.0	21.2	30.3	48.5
Other	100.0	20.8	22.4	56.8
Non-dependent abuse				
Alcohol	100.0	20.2	18.3	61.5
Cocaine	100.0	12.5	21.7	65.8
Marijuana	100.0	20.3	21.8	57.9
Opioid	100.0	19.3	24.1	56.6
Other	100.0	20.8	21.5	57.7
None reported	100.0	15.6	25.1	59.4

Continued. See notes at end of table.

Table 7.4c. Individuals served in both 2017 and 2020, by employment status and selected characteristics in 2020: Row percent distribution, 2020 (continued)

Selected characteristics	Employment status			
	Total	Employed	Unemployed	Not in labor force
Mental health diagnosis				
Trauma- and stressor-related disorders	100.0	23.3	21.5	55.1
Anxiety disorders	100.0	25.0	17.9	57.1
Attention deficit/hyperactivity disorder	100.0	29.7	19.0	51.3
Mood disorders	100.0	21.6	22.0	56.5
Bipolar disorders	100.0	20.2	23.3	56.4
Depressive disorders	100.0	22.4	21.3	56.3
Personality disorders	100.0	20.6	18.7	60.7
Schizophrenia and other psychotic disorders	100.0	11.3	21.2	67.5
Alcohol or substance-related disorders	100.0	17.0	41.5	41.5
Other mental disorders ⁴	100.0	17.0	22.5	60.5
No/deferred diagnosis	100.0	20.6	25.4	54.0

MH = mental health; SMI = serious mental illness; SU = substance use.

¹ Of 2,291,260 individuals who were served in both the 2017 and 2020 reporting periods, the total number of individuals presented in this table consists only of those with known employment status in both reporting years. The total number of individuals aged 18 years and older in Table 7.3a is lower than that shown in Table 7.4a because of the progression in age of individuals who were served in 2017, but who did not become 18 years or older until 2020.

² Age is based on the individual's age at the midpoint of the reporting period in 2020.

³ The federal definition for SMI status can be found in Appendix D.

⁴ "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol or substance-related disorders. See Appendix E.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2017 and 2020 reporting periods. Data as of 03.03.2022.

Table 7.5a. Individuals aged 18 years and older served in both 2017 and 2020, by mental health diagnosis in 2017 and living arrangement in 2017 and 2020: Number

			Living arrangement in 2020							
Living arrangement in 2017										
Diagnostic group in 2017 ¹			Total	Homeless	Foster home/care	Residential care	Institutional setting	Jail/ correctional facility	Private residence	Other
Homeless	Total	35,858	19,628	68	1,544	469	496	12,868	785	
	Bipolar disorders	7,717	3,859	9	294	89	104	3,192	170	
	Depressive disorders	11,452	6,081	16	376	73	114	4,569	223	
	Schizophrenia and other psychotic disorders	11,354	6,131	42	750	270	212	3,682	267	
	Alcohol or substance-related disorders	5,861	4,018	2	197	70	73	1,400	101	
Foster home/care	Total	8,635	83	6,623	471	89	11	1,209	149	
	Bipolar disorders	996	22	662	59	13	5	211	24	
	Depressive disorders	1,901	27	1,364	94	21	2	368	25	
	Schizophrenia and other psychotic disorders	2,823	32	2,026	230	45	1	416	73	
	Alcohol or substance-related disorders	163	4	143	1	--	1	14	--	
Residential care	Total	62,828	1,448	644	47,322	1,185	313	11,158	758	
	Bipolar disorders	8,701	302	89	5,753	135	44	2,259	119	
	Depressive disorders	9,755	340	88	6,446	123	45	2,558	155	
	Schizophrenia and other psychotic disorders	34,738	673	340	26,899	898	193	5,287	448	
	Alcohol or substance-related disorders	3,683	206	2	2,774	59	30	573	39	
Institutional settings	Total	17,698	486	136	1,432	11,409	244	3,634	357	
	Bipolar disorders	2,527	73	14	192	1,462	41	693	52	
	Depressive disorders	3,280	91	22	148	2,036	26	904	53	
	Schizophrenia and other psychotic disorders	10,053	269	99	1,050	6,491	156	1,748	240	
	Alcohol or substance-related disorders	1,520	100	1	89	845	23	443	19	

Continued. See notes at end of table.

Table 7.5a. Individuals aged 18 years and older served in both 2017 and 2020, by mental health diagnosis in 2017 and living arrangement in 2017 and 2020: Number (continued)

Living arrangement in 2017		Living arrangement in 2020							
		Total	Homeless	Foster home/care	Residential care	Institutional setting	Jail/ correctional facility	Private residence	Other
Diagnostic group in 2017¹		Total	Homeless	Foster home/care	Residential care	Institutional setting	Jail/ correctional facility	Private residence	Other
Jail/correctional facility	Total	13,076	735	8	602	378	8,028	3,131	194
	Bipolar disorders	1,842	144	2	89	52	825	693	37
	Depressive disorders	2,497	142	1	83	40	1,422	764	45
	Schizophrenia and other psychotic disorders	4,143	313	4	371	238	2,085	1,063	69
	Alcohol or substance-related disorders	1,146	98	--	37	25	638	325	23
Private residence	Total	809,103	15,100	1,186	11,656	4,627	3,070	767,764	5,700
	Bipolar disorders	156,344	3,627	214	2,249	814	680	147,643	1,117
	Depressive disorders	275,585	4,948	351	2,610	1,290	804	263,726	1,856
	Schizophrenia and other psychotic disorders	190,283	4,475	516	5,717	2,108	1,113	174,486	1,868
	Alcohol or substance-related disorders	57,586	1,800	11	710	383	321	53,957	404
Other	Total	16,500	1,069	232	842	342	148	7,240	6,627
	Bipolar disorders	2,819	201	23	121	57	24	1,432	961
	Depressive disorders	4,873	292	45	161	79	49	2,233	2,014
	Schizophrenia and other psychotic disorders	5,236	372	134	471	174	47	1,870	2,168
	Alcohol or substance-related disorders	681	126	--	36	8	13	285	213

-- Quantity is zero.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2017 and 2020 reporting periods. Data as of 03.03.2022.

Table 7.5b. Individuals aged 18 years and older served in both 2017 and 2020, by mental health diagnosis in 2017 and living arrangement in 2017 and 2020: Row percent distribution, 2020

				Living arrangement in 2020						
Living arrangement in 2017		Diagnostic group in 2017 ¹	Total	Homeless	Foster home/care	Residential care	Institutional setting	Jail/ correctional facility	Private residence	Other
Homeless	Total	100.0	54.7	†	4.3	1.3	1.4	35.9	2.2	
	Bipolar disorders	100.0	50.0	†	3.8	1.2	1.3	41.4	2.2	
	Depressive disorders	100.0	53.1	†	3.3	†	†	39.9	1.9	
	Schizophrenia and other psychotic disorders	100.0	54.0	†	6.6	2.4	1.9	32.4	2.4	
	Alcohol or substance-related disorders	100.0	68.6	†	3.4	1.2	1.2	23.9	1.7	
Foster home/care	Total	100.0	†	76.7	5.5	1.0	†	14.0	1.7	
	Bipolar disorders	100.0	2.2	66.5	5.9	1.3	†	21.2	2.4	
	Depressive disorders	100.0	1.4	71.8	4.9	1.1	†	19.4	1.3	
	Schizophrenia and other psychotic disorders	100.0	1.1	71.8	8.1	1.6	†	14.7	2.6	
	Alcohol or substance-related disorders	100.0	2.5	87.7	†	--	†	8.6	--	
Residential care	Total	100.0	2.3	1.0	75.3	1.9	†	17.8	1.2	
	Bipolar disorders	100.0	3.5	1.0	66.1	1.6	†	26.0	1.4	
	Depressive disorders	100.0	3.5	†	66.1	1.3	†	26.2	1.6	
	Schizophrenia and other psychotic disorders	100.0	1.9	†	77.4	2.6	†	15.2	1.3	
	Alcohol or substance-related disorders	100.0	5.6	†	75.3	1.6	†	15.6	1.1	
Institutional settings	Total	100.0	2.7	†	8.1	64.5	1.4	20.5	2.0	
	Bipolar disorders	100.0	2.9	†	7.6	57.9	1.6	27.4	2.1	
	Depressive disorders	100.0	2.8	†	4.5	62.1	†	27.6	1.6	
	Schizophrenia and other psychotic disorders	100.0	2.7	†	10.4	64.6	1.6	17.4	2.4	
	Alcohol or substance-related disorders	100.0	6.6	†	5.9	55.6	1.5	29.1	1.3	

Continued. See notes at end of table.

Table 7.5b. Individuals aged 18 years and older served in both 2017 and 2020, by mental health diagnosis in 2017 and living arrangement in 2017 and 2020: Row percent distribution, 2020 (continued)

<i>Living arrangement in 2017</i>		<i>Diagnostic group in 2017¹</i>	<i>Total</i>	<i>Living arrangement in 2020</i>						
				<i>Homeless</i>	<i>Foster home/care</i>	<i>Residential care</i>	<i>Institutional setting</i>	<i>Jail/correctional facility</i>	<i>Private residence</i>	<i>Other</i>
Jail/correctional facility	Total		100.0	5.6	†	4.6	2.9	61.4	23.9	1.5
	Bipolar disorders		100.0	7.8	†	4.8	2.8	44.8	37.6	2.0
	Depressive disorders		100.0	5.7	†	3.3	1.6	56.9	30.6	1.8
	Schizophrenia and other psychotic disorders		100.0	7.6	†	9.0	5.7	50.3	25.7	1.7
	Alcohol or substance-related disorders		100.0	8.6	- -	3.2	2.2	55.7	28.4	2.0
Private residence	Total		100.0	1.9	†	1.4	†	†	94.9	†
	Bipolar disorders		100.0	2.3	†	1.4	†	†	94.4	†
	Depressive disorders		100.0	1.8	†	†	†	†	95.7	†
	Schizophrenia and other psychotic disorders		100.0	2.4	†	3.0	1.1	†	91.7	†
	Alcohol or substance-related disorders		100.0	3.1	†	1.2	†	†	93.7	†
Other	Total		100.0	6.5	1.4	5.1	2.1	†	43.9	40.2
	Bipolar disorders		100.0	7.1	†	4.3	2.0	†	50.8	34.1
	Depressive disorders		100.0	6.0	†	3.3	1.6	1.0	45.8	41.3
	Schizophrenia and other psychotic disorders		100.0	7.1	2.6	9.0	3.3	†	35.7	41.4
	Alcohol or substance-related disorders		100.0	18.5	- -	5.3	1.2	1.9	41.9	31.3

- - Quantity is zero; † Less than 1 percent individuals.

¹ "Diagnostic group" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2017 and 2020 reporting periods. Data as of 03.03.2022.

Table 7.6a. Individuals aged 18 years and older served in both 2017 and 2020, by mental health diagnosis in 2017 and employment status in 2017 and 2020: Number

			Employment status in 2020		
Employment status in 2017	Diagnostic group in 2017 ¹	Total	Employed	Unemployed	Not in labor force
Employed	Total	118,577	81,644	13,482	23,451
	Bipolar disorders	26,459	17,398	3,413	5,648
	Depressive disorders	48,028	33,211	5,399	9,418
	Schizophrenia and other psychotic	20,565	13,157	2,101	5,307
	Alcohol or substance-related	4,394	3,017	950	427
Unemployed	Total	143,734	15,577	99,445	28,712
	Bipolar disorders	32,996	4,044	22,088	6,864
	Depressive disorders	51,393	6,579	34,504	10,310
	Schizophrenia and other psychotic	38,059	2,299	27,155	8,605
	Alcohol or substance-related	10,992	963	8,793	1,236
Not in labor force	Total	367,564	24,202	28,461	314,901
	Bipolar disorders	77,937	5,992	6,503	65,442
	Depressive disorders	120,888	9,489	8,732	102,667
	Schizophrenia and other psychotic	121,450	4,854	8,292	108,304
	Alcohol or substance-related	10,894	476	1,285	9,133

¹ "Diagnostic group" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2017 and 2020 reporting periods. Data as of 03.03.2022.

Table 7.6b. Individuals aged 18 years and older served in both 2017 and 2020, by mental health diagnosis in 2017 and employment status in 2017 and 2020: Row percent

<i>Employment status in 2017</i>		<i>Employment status in 2020</i>			
		<i>Total</i>	<i>Employed</i>	<i>Unemployed</i>	<i>Not in labor force</i>
<i>Diagnostic group in 2017¹</i>					
Employed	Total	100.0	68.9	11.4	19.8
	Bipolar disorders	100.0	65.8	12.9	21.3
	Depressive disorders	100.0	69.1	11.2	19.6
	Schizophrenia and other psychotic disorders	100.0	64.0	10.2	25.8
	Alcohol or substance-related disorders	100.0	68.7	21.6	9.7
Unemployed	Total	100.0	10.8	69.2	20.0
	Bipolar disorders	100.0	12.3	66.9	20.8
	Depressive disorders	100.0	12.8	67.1	20.1
	Schizophrenia and other psychotic disorders	100.0	6.0	71.3	22.6
	Alcohol or substance-related disorders	100.0	8.8	80.0	11.2
Not in labor force	Total	100.0	6.6	7.7	85.7
	Bipolar disorders	100.0	7.7	8.3	84.0
	Depressive disorders	100.0	7.8	7.2	84.9
	Schizophrenia and other psychotic disorders	100.0	4.0	6.8	89.2
	Alcohol or substance-related disorders	100.0	4.4	11.8	83.8

¹ "Diagnostic group" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2017 and 2020 reporting periods. Data as of 03.03.2022.

Table 7.7. Individuals aged 18 years and older served in consecutive reporting periods, by mental health diagnosis and living arrangement: Number, 2015–2020

Living arrangement	Diagnostic group ¹	Reporting Periods				
		2015	2015–2017	2015–2018	2015–2019	2015–2020
Homeless	Total	98,376	9,237	5,367	2,937	1,913
	Bipolar disorders	25,802	1,321	700	373	234
	Depressive disorders	34,629	2,013	1,108	497	303
	Schizophrenia and other psychotic	24,692	2,673	1,593	959	619
	Alcohol or substance-related	15,334	616	248	121	72
Foster home/care	Total	14,625	4,622	3,628	2,921	2,527
	Bipolar disorders	3,012	444	329	253	210
	Depressive disorders	4,335	807	619	447	358
	Schizophrenia and other psychotic	3,643	1,439	1,088	820	690
	Alcohol or substance-related	530	12	7	--	--
Residential care	Total	95,878	29,332	22,493	17,145	12,478
	Bipolar disorders	18,921	2,824	1,957	1,343	925
	Depressive disorders	20,345	2,972	2,067	1,355	951
	Schizophrenia and other psychotic	47,806	18,639	14,647	11,549	8,398
	Alcohol or substance-related	7,101	544	329	204	142
Institutional settings	Total	43,874	7,931	5,510	3,330	2,388
	Bipolar disorders	7,531	678	420	187	142
	Depressive disorders	10,488	992	648	345	224
	Schizophrenia and other psychotic	21,506	4,856	3,442	2,209	1,600
	Alcohol or substance-related	2,699	106	39	12	10

Continued. See notes at end of table.

Table 7.7. Individuals aged 18 years and older served in consecutive reporting periods, by mental health diagnosis and living arrangement: Number, 2015–2020 (continued)

<i>Living arrangement</i>	<i>Diagnostic group¹</i>	<i>Reporting Periods</i>				
		<i>2015</i>	<i>2015–2017</i>	<i>2015–2018</i>	<i>2015–2019</i>	<i>2015–2020</i>
Jail/correctional facility	Total	48,144	7,169	4,496	3,114	2,277
	Bipolar disorders	8,819	422	228	145	105
	Depressive disorders	10,635	1,160	707	452	326
	Schizophrenia and other psychotic disorders	9,831	1,521	999	754	576
	Alcohol or substance-related disorders	7,227	196	48	22	9
Private residence	Total	1,721,653	464,796	339,433	245,209	185,552
	Bipolar disorders	425,682	82,537	57,892	40,358	29,401
	Depressive disorders	696,635	134,034	92,041	61,842	44,784
	Schizophrenia and other psychotic disorders	318,490	111,212	86,856	66,542	52,145
	Alcohol or substance-related disorders	86,654	4,962	2,542	1,608	1,185
Other	Total	91,348	10,275	3,366	1,974	1,333
	Bipolar disorders	18,407	1,302	327	199	128
	Depressive disorders	33,378	2,005	502	312	218
	Schizophrenia and other psychotic disorders	16,947	3,522	1,670	917	627
	Alcohol or substance-related disorders	5,008	163	68	30	16

- - Quantity is zero.

¹ "Diagnostic group" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.
 SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 7.8a. Individuals aged 18 years and older served in both 2015 and 2017, by mental health diagnosis in 2015 and living arrangement in 2015 and 2017: Number

<i>Living arrangement in 2015</i>		<i>Diagnostic group in 2015¹</i>	<i>Total</i>	<i>Living arrangement in 2017</i>						
				<i>Homeless</i>	<i>Foster home/care</i>	<i>Residential care</i>	<i>Institutional setting</i>	<i>Jail/ correctional facility</i>	<i>Private residence</i>	<i>Other</i>
Homeless	Total		32,888	14,906	52	1,754	609	576	13,032	1,959
	Bipolar disorders		8,856	3,972	13	426	118	150	3,706	471
	Depressive disorders		10,968	4,722	14	427	113	119	4,789	784
	Schizophrenia and other psychotic		10,604	5,109	26	871	359	246	3,470	523
	Alcohol or substance-related		4,862	2,871	1	246	69	112	1,370	193
Foster home/care	Total		8,258	78	5,504	313	105	23	1,913	322
	Bipolar disorders		1,674	16	1,171	66	22	3	349	47
	Depressive disorders		2,356	26	1,518	74	26	4	637	71
	Schizophrenia and other psychotic		2,772	25	1,974	157	57	9	427	123
	Alcohol or substance-related		245	2	66	5	- -	2	166	4
Residential care	Total		54,657	1,705	404	37,878	1,535	330	10,727	2,078
	Bipolar disorders		9,739	417	56	5,903	224	77	2,625	437
	Depressive disorders		9,246	427	82	5,492	205	50	2,567	423
	Schizophrenia and other psychotic		33,062	693	219	24,908	1,150	184	4,978	930
	Alcohol or substance-related		3,251	371	3	1,950	78	55	716	78
Institutional settings	Total		18,771	534	96	1,943	10,757	307	4,512	622
	Bipolar disorders		3,123	120	13	290	1,527	52	1,021	100
	Depressive disorders		3,515	99	25	202	2,003	27	1,052	107
	Schizophrenia and other psychotic		11,542	303	60	1,416	6,902	200	2,272	389
	Alcohol or substance-related		1,107	68	3	110	506	31	377	12

Continued. See notes at end of table.

Table 7.8a. Individuals aged 18 years and older served in both 2015 and 2017, by mental health diagnosis in 2015 and living arrangement in 2015 and 2017: Number (continued)

Living arrangement in 2015			Living arrangement in 2017						
			Homeless	Foster home/care	Residential care	Institutional setting	Jail/ correctional facility	Private residence	Other
Diagnostic group in 2015¹		Total							
Jail/correctional facility	Total	14,831	741	13	739	382	9,458	3,226	272
	Bipolar disorders	2,650	178	3	144	67	1,311	890	57
	Depressive disorders	3,112	133	6	114	20	2,029	765	45
	Schizophrenia and other psychotic disorders	4,548	329	4	418	273	2,330	1,082	112
	Alcohol or substance-related disorders	1,797	161	1	95	22	1,070	413	35
Private residence	Total	659,484	13,834	890	12,326	5,445	3,430	600,943	22,616
	Bipolar disorders	168,995	4,133	190	2,850	1,272	1,025	153,866	5,659
	Depressive disorders	249,398	4,504	283	2,902	1,449	889	230,668	8,703
	Schizophrenia and other psychotic disorders	175,146	3,925	342	6,035	2,647	1,142	157,253	3,802
	Alcohol or substance-related disorders	26,927	1,708	22	727	224	379	23,282	585
Other	Total	34,712	1,683	244	1,638	637	254	13,120	17,136
	Bipolar disorders	7,476	424	35	302	122	64	3,193	3,336
	Depressive disorders	11,496	597	36	355	138	67	4,742	5,561
	Schizophrenia and other psychotic disorders	9,659	459	145	812	381	105	2,708	5,049
	Alcohol or substance-related disorders	1,688	212	- -	75	18	35	605	743

- - Quantity is zero.

¹ "Diagnostic group" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015 and 2017 reporting periods. Data as of 03.03.2022.

Table 7.8b. Individuals aged 18 years and older served in both 2015 and 2017, by mental health diagnosis in 2015 and living arrangement in 2015 and 2017: Row percent distribution

<i>Living arrangement in 2015</i>		<i>Diagnostic group in 2015¹</i>	<i>Total</i>	<i>Living arrangement in 2017</i>						
				<i>Homeless</i>	<i>Foster home/care</i>	<i>Residential care</i>	<i>Institutional setting</i>	<i>Jail/correctional facility</i>	<i>Private residence</i>	<i>Other</i>
Homeless	Total		100.0	45.3	†	5.3	1.9	1.8	39.6	6.0
	Bipolar disorders		100.0	44.9	†	4.8	1.3	1.7	41.8	5.3
	Depressive disorders		100.0	43.1	†	3.9	1.0	1.1	43.7	7.1
	Schizophrenia and other psychotic disorders		100.0	48.2	†	8.2	3.4	2.3	32.7	4.9
	Alcohol or substance-related disorders		100.0	59.0	†	5.1	1.4	2.3	28.2	4.0
Foster home/care	Total		100.0	†	66.7	3.8	1.3	†	23.2	3.9
	Bipolar disorders		100.0	†	70.0	3.9	1.3	†	20.8	2.8
	Depressive disorders		100.0	1.1	64.4	3.1	1.1	†	27.0	3.0
	Schizophrenia and other psychotic disorders		100.0	†	71.2	5.7	2.1	†	15.4	4.4
	Alcohol or substance-related disorders		100.0	†	26.9	2.0	- -	†	67.8	1.6
Residential care	Total		100.0	3.1	†	69.3	2.8	†	19.6	3.8
	Bipolar disorders		100.0	4.3	†	60.6	2.3	†	27.0	4.5
Institutional settings	Depressive disorders		100.0	4.6	†	59.4	2.2	†	27.8	4.6
	Schizophrenia and other psychotic disorders		100.0	2.1	†	75.3	3.5	†	15.1	2.8
	Alcohol or substance-related disorders		100.0	11.4	†	60.0	2.4	1.7	22.0	2.4
	Total		100.0	2.8	†	10.4	57.3	1.6	24.0	3.3
	Bipolar disorders		100.0	3.8	†	9.3	48.9	1.7	32.7	3.2
	Depressive disorders		100.0	2.8	†	5.7	57.0	†	29.9	3.0
	Schizophrenia and other psychotic disorders		100.0	2.6	†	12.3	59.8	1.7	19.7	3.4
	Alcohol or substance-related disorders		100.0	6.1	†	9.9	45.7	2.8	34.1	1.1

Continued. See notes at end of table.

Table 7.8b. Individuals aged 18 years and older served in both 2015 and 2017, by mental health diagnosis in 2015 and living arrangement in 2015 and 2017: Row percent distribution (continued)

<i>Living arrangement in 2015</i>		<i>Diagnostic group in 2015¹</i>	<i>Total</i>	<i>Living arrangement in 2017</i>						
				<i>Homeless</i>	<i>Foster home/care</i>	<i>Residential care</i>	<i>Institutional setting</i>	<i>Jail/correctional facility</i>	<i>Private residence</i>	<i>Other</i>
Jail/correctional facility	Total		100.0	5.0	†	5.0	2.6	63.8	21.8	1.8
	Bipolar disorders		100.0	6.7	†	5.4	2.5	49.5	33.6	2.2
	Depressive disorders		100.0	4.3	†	3.7	†	65.2	24.6	1.4
	Schizophrenia and other psychotic disorders		100.0	7.2	†	9.2	6.0	51.2	23.8	2.5
	Alcohol or substance-related disorders		100.0	9.0	†	5.3	1.2	59.5	23.0	1.9
Private residence	Total		100.0	2.1	†	1.9	†	†	91.1	3.4
	Bipolar disorders		100.0	2.4	†	1.7	†	†	91.0	3.3
	Depressive disorders		100.0	1.8	†	1.2	†	†	92.5	3.5
	Schizophrenia and other psychotic disorders		100.0	2.2	†	3.4	1.5	†	89.8	2.2
	Alcohol or substance-related disorders		100.0	6.3	†	2.7	†	1.4	86.5	2.2
Other	Total		100.0	4.8	†	4.7	1.8	†	37.8	49.4
	Bipolar disorders		100.0	5.7	†	4.0	1.6	†	42.7	44.6
	Depressive disorders		100.0	5.2	†	3.1	1.2	†	41.2	48.4
	Schizophrenia and other psychotic disorders		100.0	4.8	1.5	8.4	3.9	1.1	28.0	52.3
	Alcohol or substance-related disorders		100.0	12.6	--	4.4	1.1	2.1	35.8	44.0

-- Quantity is zero; † Less than 1 percent individuals.

¹ "Diagnostic group" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015 and 2017 reporting periods. Data as of 03.03.2022.

Table 7.9a. Individuals aged 18 years and older served in both 2015 and 2018, by mental health diagnosis in 2015 and living arrangement in 2015 and 2018: Number

<i>Living arrangement in 2015</i>		<i>Diagnostic group in 2015¹</i>	<i>Total</i>	<i>Living arrangement in 2018</i>						
				<i>Homeless</i>	<i>Foster home/care</i>	<i>Residential care</i>	<i>Institutional setting</i>	<i>Jail/ correctional facility</i>	<i>Private residence</i>	<i>Other</i>
Homeless	Total		28,643	12,201	48	1,660	569	544	12,625	996
	Bipolar disorders		7,740	3,221	12	371	98	132	3,649	257
	Depressive disorders		9,480	3,875	12	386	102	129	4,611	365
	Schizophrenia and other psychotic disorders		9,396	4,259	27	828	362	239	3,345	336
	Alcohol or substance-related disorders		4,379	2,428	- -	272	73	99	1,368	139
Foster home/care	Total		7,539	119	4,639	402	123	17	2,074	165
	Bipolar disorders		1,488	25	977	71	25	5	357	28
	Depressive disorders		2,195	27	1,314	98	27	1	689	39
	Schizophrenia and other psychotic disorders		2,573	51	1,634	207	69	6	529	77
	Alcohol or substance-related disorders		236	4	62	5	- -	1	162	2
Residential care	Total		47,620	1,772	505	30,966	1,452	305	11,443	1,177
	Bipolar disorders		8,423	426	87	4,692	194	59	2,729	236
	Depressive disorders		8,137	434	95	4,481	176	51	2,697	203
	Schizophrenia and other psychotic disorders		28,791	753	272	20,356	1,103	184	5,448	675
	Alcohol or substance-related disorders		2,979	394	3	1,669	85	51	708	69
Institutional settings	Total		16,006	499	107	1,837	8,394	291	4,469	409
	Bipolar disorders		2,674	113	22	268	1,144	50	1,022	55
	Depressive disorders		2,879	95	26	191	1,459	30	999	79
	Schizophrenia and other psychotic		10,059	290	69	1,353	5,557	194	2,331	265
	Alcohol or substance-related		1,021	56	2	113	458	24	358	10

Continued. See notes at end of table.

**Table 7.9a. Individuals aged 18 years and older served in both 2015 and 2018, by mental health diagnosis in 2015 and living arrangement in 2015 and 2018:
Number (continued)**

			Living arrangement in 2018						
Living arrangement in 2015	Diagnostic group in 2015 ¹	Total	Homeless	Foster home/care	Residential care	Institutional setting	Jail/ correctional facility	Private residence	Other
Jail/correctional facility	Total	12,653	768	12	696	367	7,544	3,064	202
	Bipolar disorders	2,200	188	2	141	61	942	832	34
	Depressive disorders	2,661	121	7	110	35	1,618	733	37
	Schizophrenia and other psychotic disorders	3,961	345	2	388	252	1,830	1,061	83
	Alcohol or substance-related disorders	1,397	177	--	96	25	658	410	31
Private residence	Total	569,054	14,502	1,045	13,088	5,617	3,280	524,375	7,147
	Bipolar disorders	145,038	4,417	220	3,067	1,246	926	133,405	1,757
	Depressive disorders	210,718	4,637	320	3,009	1,489	864	198,052	2,347
	Schizophrenia and other psychotic disorders	159,839	4,250	431	6,348	2,778	1,110	142,673	2,249
	Alcohol or substance-related disorders	23,823	1,809	11	736	253	331	20,225	458
Other	Total	26,629	1,789	258	1,765	589	261	15,564	6,403
	Bipolar disorders	5,812	451	32	352	99	65	3,668	1,145
	Depressive disorders	8,874	619	34	389	115	74	5,807	1,836
	Schizophrenia and other psychotic disorders	7,473	485	154	808	358	95	2,906	2,667
	Alcohol or substance-related disorders	1,439	213	--	74	14	43	575	520

-- Quantity is zero.

¹ "Diagnostic group" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015 and 2018 reporting periods. Data as of 03.03.2022.

Table 7.9b. Individuals aged 18 years and older served in both 2015 and 2018, by mental health diagnosis in 2015 and living arrangement in 2015 and 2018:
Row percent distribution

<i>Living arrangement in 2015</i>		<i>Diagnostic group in 2015¹</i>	<i>Total</i>	<i>Living arrangement in 2018</i>						
				<i>Homeless</i>	<i>Foster home/care</i>	<i>Residential care</i>	<i>Institutional setting</i>	<i>Jail/correctional facility</i>	<i>Private residence</i>	<i>Other</i>
Homeless	Total		100.0	42.6	†	5.8	2.0	1.9	44.1	3.5
	Bipolar disorders		100.0	41.6	†	4.8	1.3	1.7	47.1	3.3
	Depressive disorders		100.0	40.9	†	4.1	1.1	1.4	48.6	3.9
	Schizophrenia and other psychotic disorders		100.0	45.3	†	8.8	3.9	2.5	35.6	3.6
	Alcohol or substance-related disorders		100.0	55.4	--	6.2	1.7	2.3	31.2	3.2
Foster home/care	Total		100.0	1.6	61.5	5.3	1.6	†	27.5	2.2
	Bipolar disorders		100.0	1.7	65.7	4.8	1.7	†	24.0	1.9
	Depressive disorders		100.0	1.2	59.9	4.5	1.2	†	31.4	1.8
	Schizophrenia and other psychotic disorders		100.0	2.0	63.5	8.0	2.7	†	20.6	3.0
	Alcohol or substance-related disorders		100.0	1.7	26.3	2.1	--	†	68.6	†
Residential care	Total		100.0	3.7	1.1	65.0	3.0	†	24.0	2.5
	Bipolar disorders		100.0	5.1	1.0	55.7	2.3	†	32.4	2.8
	Depressive disorders		100.0	5.3	1.2	55.1	2.2	†	33.1	2.5
	Schizophrenia and other psychotic disorders		100.0	2.6	†	70.7	3.8	†	18.9	2.3
	Alcohol or substance-related disorders		100.0	13.2	†	56.0	2.9	1.7	23.8	2.3
Institutional settings	Total		100.0	3.1	†	11.5	52.4	1.8	27.9	2.6
	Bipolar disorders		100.0	4.2	†	10.0	42.8	1.9	38.2	2.1
	Depressive disorders		100.0	3.3	†	6.6	50.7	1.0	34.7	2.7
	Schizophrenia and other psychotic disorders		100.0	2.9	†	13.5	55.2	1.9	23.2	2.6
	Alcohol or substance-related disorders		100.0	5.5	†	11.1	44.9	2.4	35.1	†

Continued. See notes at end of table.

**Table 7.9b. Individuals aged 18 years and older served in both 2015 and 2018, by mental health diagnosis in 2015 and living arrangement in 2015 and 2018:
Row percent distribution (continued)**

			Living arrangement in 2018						
Living arrangement in 2015	Diagnostic group in 2015 ¹	Total	Homeless	Foster home/care	Residential care	Institutional setting	Jail/ correctional facility	Private residence	Other
Jail/correctional facility	Total	100.0	6.1	†	5.5	2.9	59.6	24.2	1.6
	Bipolar disorders	100.0	8.5	†	6.4	2.8	42.8	37.8	1.5
	Depressive disorders	100.0	4.5	†	4.1	1.3	60.8	27.5	1.4
	Schizophrenia and other psychotic disorders	100.0	8.7	†	9.8	6.4	46.2	26.8	2.1
	Alcohol or substance-related disorders	100.0	12.7	--	6.9	1.8	47.1	29.3	2.2
Private residence	Total	100.0	2.5	†	2.3	†	†	92.1	1.3
	Bipolar disorders	100.0	3.0	†	2.1	†	†	92.0	1.2
	Depressive disorders	100.0	2.2	†	1.4	†	†	94.0	1.1
	Schizophrenia and other psychotic disorders	100.0	2.7	†	4.0	1.7	†	89.3	1.4
	Alcohol or substance-related disorders	100.0	7.6	†	3.1	1.1	1.4	84.9	1.9
Other	Total	100.0	6.7	†	6.6	2.2	†	58.4	24.0
	Bipolar disorders	100.0	7.8	†	6.1	1.7	1.1	63.1	19.7
	Depressive disorders	100.0	7.0	†	4.4	1.3	†	65.4	20.7
	Schizophrenia and other psychotic disorders	100.0	6.5	2.1	10.8	4.8	1.3	38.9	35.7
	Alcohol or substance-related disorders	100.0	14.8	--	5.1	†	3.0	40.0	36.1

-- Quantity is zero; † Less than 1 percent individuals.

¹ "Diagnostic group" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015 and 2018 reporting periods. Data as of 03.03.2022.

Table 7.10a. Individuals aged 18 years and older served in both 2015 and 2019, by mental health diagnosis in 2015 and living arrangement in 2015 and 2019: Number

<i>Living arrangement in 2015</i>		<i>Diagnostic group in 2015¹</i>	<i>Total</i>	<i>Living arrangement in 2019</i>						
				<i>Homeless</i>	<i>Foster home/care</i>	<i>Residential care</i>	<i>Institutional setting</i>	<i>Jail/correctional facility</i>	<i>Private residence</i>	<i>Other</i>
Homeless	Total		23,016	8,796	61	1,545	490	403	10,771	950
	Bipolar disorders		6,347	2,340	16	393	112	87	3,153	246
	Depressive disorders		7,322	2,644	10	359	76	87	3,828	318
	Schizophrenia and other psychotic disorders		7,816	3,151	39	768	295	197	3,042	324
	Alcohol or substance-related disorders		3,772	1,906	3	257	77	87	1,289	153
Foster home/care	Total		6,919	100	4,010	388	122	25	2,076	198
	Bipolar disorders		1,399	25	885	65	23	7	357	37
	Depressive disorders		1,961	28	1,099	92	33	4	664	41
	Schizophrenia and other psychotic disorders		2,359	39	1,396	216	63	7	537	101
	Alcohol or substance-related disorders		213	6	44	4	2	-	156	1
Residential care	Total		41,175	1,709	524	25,219	1,341	299	11,042	1,041
	Bipolar disorders		7,217	410	100	3,669	193	53	2,592	200
	Depressive disorders		6,841	385	102	3,424	149	33	2,538	210
	Schizophrenia and other psychotic disorders		25,348	741	276	17,073	1,027	192	5,434	605
	Alcohol or substance-related disorders		2,736	362	3	1,408	86	55	763	59
Institutional settings	Total		12,946	586	130	1,653	5,845	311	4,043	378
	Bipolar disorders		2,135	130	28	228	761	39	891	58
	Depressive disorders		2,248	99	27	177	963	28	885	69
	Schizophrenia and other psychotic disorders		8,264	335	87	1,233	4,001	220	2,143	245
	Alcohol or substance-related disorders		938	91	1	121	327	36	348	14

Continued. See notes at end of table.

Table 7.10a. Individuals aged 18 years and older served in both 2015 and 2019, by mental health diagnosis in 2015 and living arrangement in 2015 and 2019: Number (continued)

			Living arrangement in 2019							
Living arrangement in 2015										
Diagnostic group in 2015 ¹			Total	Homeless	Foster home/care	Residential care	Institutional setting	Jail/ correctional facility	Private residence	Other
Jail/correctional facility	Total	10,935	713	12	635	301	6,103	3,000	171	
	Bipolar disorders	1,941	180	2	140	46	745	782	46	
	Depressive disorders	2,219	127	5	99	26	1,224	701	37	
	Schizophrenia and other psychotic disorders	3,402	264	4	357	214	1,521	984	58	
	Alcohol or substance-related disorders	1,322	191	- -	92	22	524	461	32	
Private residence	Total	466,842	13,268	1,176	12,671	5,079	2,715	423,812	8,121	
	Bipolar disorders	118,636	3,981	225	2,936	1,127	765	107,636	1,966	
	Depressive disorders	169,750	4,099	367	2,793	1,389	690	157,588	2,824	
	Schizophrenia and other psychotic disorders	136,408	4,006	511	6,346	2,489	962	119,531	2,563	
	Alcohol or substance-related disorders	21,977	1,815	17	828	281	355	18,148	533	
Other	Total	23,719	1,528	263	1,700	481	189	14,403	5,155	
	Bipolar disorders	5,227	407	36	318	97	47	3,355	967	
	Depressive disorders	8,101	518	43	365	119	46	5,273	1,737	
	Schizophrenia and other psychotic disorders	6,162	375	154	796	246	73	2,726	1,792	
	Alcohol or substance-related disorders	1,448	224	1	98	23	35	658	409	

- - Quantity is zero.

¹ "Diagnostic group" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015 and 2019 reporting periods. Data as of 03.03.2022.

Table 7.10b. Individuals aged 18 years and older served in both 2015 and 2019, by mental health diagnosis in 2015 and living arrangement in 2015 and 2019: Row percent distribution

<i>Living arrangement in 2015</i>		<i>Diagnostic group in 2015¹</i>	<i>Total</i>	<i>Living arrangement in 2019</i>						
				<i>Homeless</i>	<i>Foster home/care</i>	<i>Residential care</i>	<i>Institutional setting</i>	<i>Jail/correctional facility</i>	<i>Private residence</i>	<i>Other</i>
Homeless	Total		100.0	38.2	†	6.7	2.1	1.8	46.8	4.1
	Bipolar disorders		100.0	36.9	†	6.2	1.8	1.4	49.7	3.9
	Depressive disorders		100.0	36.1	†	4.9	1.0	1.2	52.3	4.3
	Schizophrenia and other psychotic disorders		100.0	40.3	†	9.8	3.8	2.5	38.9	4.1
	Alcohol or substance-related disorders		100.0	50.5	†	6.8	2.0	2.3	34.2	4.1
Foster home/care	Total		100.0	1.4	58.0	5.6	1.8	†	30.0	2.9
	Bipolar disorders		100.0	1.8	63.3	4.6	1.6	†	25.5	2.6
	Depressive disorders		100.0	1.4	56.0	4.7	1.7	†	33.9	2.1
	Schizophrenia and other psychotic disorders		100.0	1.7	59.2	9.2	2.7	†	22.8	4.3
	Alcohol or substance-related disorders		100.0	2.8	20.7	1.9	†	--	73.2	†
Residential care	Total		100.0	4.2	1.3	61.2	3.3	†	26.8	2.5
	Bipolar disorders		100.0	5.7	1.4	50.8	2.7	†	35.9	2.8
Institutional settings	Depressive disorders		100.0	5.6	1.5	50.1	2.2	†	37.1	3.1
	Schizophrenia and other psychotic disorders		100.0	2.9	1.1	67.4	4.1	†	21.4	2.4
	Alcohol or substance-related disorders		100.0	13.2	†	51.5	3.1	2.0	27.9	2.2
	Total		100.0	4.5	1.0	12.8	45.1	2.4	31.2	2.9
	Bipolar disorders		100.0	6.1	1.3	10.7	35.6	1.8	41.7	2.7
Institutional settings	Depressive disorders		100.0	4.4	1.2	7.9	42.8	1.2	39.4	3.1
	Schizophrenia and other psychotic disorders		100.0	4.1	1.1	14.9	48.4	2.7	25.9	3.0
	Alcohol or substance-related disorders		100.0	9.7	†	12.9	34.9	3.8	37.1	1.5

Continued. See notes at end of table.

Table 7.10b. Individuals aged 18 years and older served in both 2015 and 2019, by mental health diagnosis in 2015 and living arrangement in 2015 and 2019: Row percent distribution (continued)

			Living arrangement in 2019							
Living arrangement in 2015										
Diagnostic group in 2015¹			Total	Homeless	Foster home/care	Residential care	Institutional setting	Jail/ correctional facility	Private residence	Other
Jail/correctional facility	Total	100.0	6.5	†	5.8	2.8	55.8	27.4	1.6	
	Bipolar disorders	100.0	9.3	†	7.2	2.4	38.4	40.3	2.4	
	Depressive disorders	100.0	5.7	†	4.5	1.2	55.2	31.6	1.7	
	Schizophrenia and other psychotic disorders	100.0	7.8	†	10.5	6.3	44.7	28.9	1.7	
	Alcohol or substance-related disorders	100.0	14.4	--	7.0	1.7	39.6	34.9	2.4	
Private residence	Total	100.0	2.8	†	2.7	1.1	†	90.8	1.7	
	Bipolar disorders	100.0	3.4	†	2.5	†	†	90.7	1.7	
	Depressive disorders	100.0	2.4	†	1.6	†	†	92.8	1.7	
	Schizophrenia and other psychotic disorders	100.0	2.9	†	4.7	1.8	†	87.6	1.9	
	Alcohol or substance-related disorders	100.0	8.3	†	3.8	1.3	1.6	82.6	2.4	
Other	Total	100.0	6.4	1.1	7.2	2.0	†	60.7	21.7	
	Bipolar disorders	100.0	7.8	†	6.1	1.9	†	64.2	18.5	
	Depressive disorders	100.0	6.4	†	4.5	1.5	†	65.1	21.4	
	Schizophrenia and other psychotic disorders	100.0	6.1	2.5	12.9	4.0	1.2	44.2	29.1	
	Alcohol or substance-related disorders	100.0	15.5	†	6.8	1.6	2.4	45.4	28.2	

-- Quantity is zero; † Less than 1 percent individuals.

¹ "Diagnostic group" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015 and 2019 reporting periods. Data as of 03.03.2022.

Table 7.11a. Individuals aged 18 years and older served in both 2015 and 2020, by mental health diagnosis in 2015 and living arrangement in 2015 and 2020: Number

<i>Living arrangement in 2015</i>		<i>Diagnostic group in 2015¹</i>	<i>Total</i>	<i>Living arrangement in 2020</i>						
				<i>Homeless</i>	<i>Foster home/care</i>	<i>Residential care</i>	<i>Institutional setting</i>	<i>Jail/correctional facility</i>	<i>Private residence</i>	<i>Other</i>
Homeless	Total		18,664	7,227	66	1,208	431	345	8,744	643
	Bipolar disorders		5,079	1,912	18	291	95	83	2,501	179
	Depressive disorders		5,932	2,201	20	249	72	67	3,097	226
	Schizophrenia and other psychotic disorders		6,486	2,598	38	644	264	167	2,564	211
	Alcohol or substance-related disorders		2,675	1,408	4	170	72	71	877	73
Foster home/care	Total		6,708	90	3,787	405	108	28	2,120	170
	Bipolar disorders		1,370	19	840	79	24	7	365	36
	Depressive disorders		1,939	25	1,062	108	21	4	684	35
	Schizophrenia and other psychotic disorders		2,297	33	1,321	211	62	8	571	91
	Alcohol or substance-related disorders		211	6	38	6	-	1	159	1
Residential care	Total		33,984	1,351	543	20,378	1,222	229	9,517	744
	Bipolar disorders		5,907	286	101	2,965	165	43	2,220	127
	Depressive disorders		5,841	312	107	2,915	129	33	2,192	153
	Schizophrenia and other psychotic disorders		20,929	598	300	13,669	952	146	4,802	462
	Alcohol or substance-related disorders		1,910	239	4	1,005	67	31	522	42
Institutional settings	Total		12,058	467	139	1,526	5,276	257	4,067	326
	Bipolar disorders		2,095	100	28	213	753	25	926	50
	Depressive disorders		2,208	80	27	160	918	29	929	65
	Schizophrenia and other psychotic disorders		7,524	274	98	1,146	3,524	180	2,098	204
	Alcohol or substance-related disorders		790	64	1	74	295	23	324	9

Continued. See notes at end of table.

Table 7.11a. Individuals aged 18 years and older served in both 2015 and 2020, by mental health diagnosis in 2015 and living arrangement in 2015 and 2020: Number (continued)

			Living arrangement in 2020						
Living arrangement in 2015									
	Diagnostic group in 2015¹	Total	Homeless	Foster home/care	Residential care	Institutional setting	Jail/ correctional facility	Private residence	Other
Jail/correctional facility	Total	9,168	593	14	562	288	4,957	2,607	147
	Bipolar disorders	1,692	157	3	123	43	615	714	37
	Depressive disorders	1,840	102	7	95	29	978	592	37
	Schizophrenia and other psychotic disorders	2,923	241	4	324	196	1,222	889	47
	Alcohol or substance-related disorders	927	116	- -	48	16	376	354	17
Private residence	Total	400,436	11,431	1,264	10,907	4,984	2,326	363,798	5,726
	Bipolar disorders	99,703	3,387	250	2,465	1,088	674	90,531	1,308
	Depressive disorders	146,988	3,582	396	2,475	1,395	568	136,572	2,000
	Schizophrenia and other psychotic disorders	120,939	3,557	560	5,629	2,421	880	105,976	1,916
	Alcohol or substance-related disorders	17,359	1,389	16	629	237	275	14,465	348
Other	Total	12,566	898	240	838	420	124	6,559	3,487
	Bipolar disorders	2,670	220	26	126	88	29	1,558	623
	Depressive disorders	4,095	287	42	175	87	26	2,288	1,190
	Schizophrenia and other psychotic disorders	4,314	298	158	486	222	54	1,829	1,267
	Alcohol or substance-related disorders	864	141	1	42	26	31	369	254

- - Quantity is zero.

¹ "Diagnostic group" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015 and 2020 reporting periods. Data as of 03.03.2022.

Table 7.11b. Individuals aged 18 years and older served in both 2015 and 2020, by mental health diagnosis in 2015 and living arrangement in 2015 and 2020: Row percent distribution

<i>Living arrangement in 2015</i>		<i>Diagnostic group in 2015¹</i>	<i>Total</i>	<i>Living arrangement in 2020</i>						
				<i>Homeless</i>	<i>Foster home/care</i>	<i>Residential care</i>	<i>Institutional setting</i>	<i>Jail/correctional facility</i>	<i>Private residence</i>	<i>Other</i>
Homeless	Total		100.0	38.7	†	6.5	2.3	1.8	46.8	3.4
	Bipolar disorders		100.0	37.6	†	5.7	1.9	1.6	49.2	3.5
	Depressive disorders		100.0	37.1	†	4.2	1.2	1.1	52.2	3.8
	Schizophrenia and other psychotic disorders		100.0	40.1	†	9.9	4.1	2.6	39.5	3.3
	Alcohol or substance-related disorders		100.0	52.6	†	6.4	2.7	2.7	32.8	2.7
Foster home/care	Total		100.0	1.3	56.5	6.0	1.6	†	31.6	2.5
	Bipolar disorders		100.0	1.4	61.3	5.8	1.8	†	26.6	2.6
	Depressive disorders		100.0	1.3	54.8	5.6	1.1	†	35.3	1.8
	Schizophrenia and other psychotic disorders		100.0	1.4	57.5	9.2	2.7	†	24.9	4.0
	Alcohol or substance-related disorders		100.0	2.8	18.0	2.8	- -	†	75.4	†
Residential care	Total		100.0	4.0	1.6	60.0	3.6	†	28.0	2.2
	Bipolar disorders		100.0	4.8	1.7	50.2	2.8	†	37.6	2.1
	Depressive disorders		100.0	5.3	1.8	49.9	2.2	†	37.5	2.6
	Schizophrenia and other psychotic disorders		100.0	2.9	1.4	65.3	4.5	†	22.9	2.2
	Alcohol or substance-related disorders		100.0	12.5	†	52.6	3.5	1.6	27.3	2.2
Institutional settings	Total		100.0	3.9	1.2	12.7	43.8	2.1	33.7	2.7
	Bipolar disorders		100.0	4.8	1.3	10.2	35.9	1.2	44.2	2.4
	Depressive disorders		100.0	3.6	1.2	7.2	41.6	1.3	42.1	2.9
	Schizophrenia and other psychotic disorders		100.0	3.6	1.3	15.2	46.8	2.4	27.9	2.7
	Alcohol or substance-related disorders		100.0	8.1	†	9.4	37.3	2.9	41.0	1.1

Continued. See notes at end of table.

Table 7.11b. Individuals aged 18 years and older served in both 2015 and 2020, by mental health diagnosis in 2015 and living arrangement in 2015 and 2020: Row percent distribution (continued)

<i>Living arrangement in 2015</i>		<i>Diagnostic group in 2015¹</i>	<i>Total</i>	<i>Living arrangement in 2020</i>						
				<i>Homeless</i>	<i>Foster home/care</i>	<i>Residential care</i>	<i>Institutional setting</i>	<i>Jail/correctional facility</i>	<i>Private residence</i>	<i>Other</i>
Jail/correctional facility	Total		100.0	6.5	†	6.1	3.1	54.1	28.4	1.6
	Bipolar disorders		100.0	9.3	†	7.3	2.5	36.3	42.2	2.2
	Depressive disorders		100.0	5.5	†	5.2	1.6	53.2	32.2	2.0
	Schizophrenia and other psychotic disorders		100.0	8.2	†	11.1	6.7	41.8	30.4	1.6
	Alcohol or substance-related disorders		100.0	12.5	--	5.2	1.7	40.6	38.2	1.8
	Total		100.0	2.9	†	2.7	1.2	†	90.9	1.4
Private residence	Bipolar disorders		100.0	3.4	†	2.5	1.1	†	90.8	1.3
	Depressive disorders		100.0	2.4	†	1.7	†	†	92.9	1.4
	Schizophrenia and other psychotic disorders		100.0	2.9	†	4.7	2.0	†	87.6	1.6
	Alcohol or substance-related disorders		100.0	8.0	†	3.6	1.4	1.6	83.3	2.0
	Total		100.0	7.1	1.9	6.7	3.3	†	52.2	27.7
	Bipolar disorders		100.0	8.2	†	4.7	3.3	1.1	58.4	23.3
Other	Depressive disorders		100.0	7.0	1.0	4.3	2.1	†	55.9	29.1
	Schizophrenia and other psychotic disorders		100.0	6.9	3.7	11.3	5.1	1.3	42.4	29.4
	Alcohol or substance-related disorders		100.0	16.3	†	4.9	3.0	3.6	42.7	29.4
	Total		100.0	7.1	1.9	6.7	3.3	†	52.2	27.7
	Bipolar disorders		100.0	8.2	†	4.7	3.3	1.1	58.4	23.3
	Depressive disorders		100.0	7.0	1.0	4.3	2.1	†	55.9	29.1

-- Quantity is zero; † Less than 1 percent individuals.

¹ "Diagnostic group" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015 and 2020 reporting periods. Data as of 03.03.2022.

Table 7.12. Individuals aged 18 years and older served in consecutive reporting periods, by mental health diagnosis and employment status: Number, 2015–2020

<i>Employment status</i>	<i>Diagnostic group¹</i>	<i>Reporting periods</i>				
		2015	2015–2017	2015–2018	2015–2019	2015–2020
Employed	Total	366,318	57,153	37,456	24,768	18,328
	Bipolar disorders	83,873	9,983	6,301	3,953	2,768
	Depressive disorders	158,635	18,029	11,159	6,689	4,705
	Schizophrenia and other psychotic disorders	36,130	8,974	6,608	4,858	3,652
	Alcohol or substance-related disorders	13,014	331	156	77	49
Unemployed	Total	516,727	50,663	29,510	16,497	10,863
	Bipolar disorders	139,588	9,005	5,008	2,715	1,693
	Depressive disorders	214,586	13,600	7,243	3,909	2,456
	Schizophrenia and other psychotic disorders	88,481	10,469	6,794	4,144	2,848
	Alcohol or substance-related disorders	24,313	804	358	201	134
Not in labor force	Total	881,698	263,419	191,088	139,039	108,564
	Bipolar disorders	212,081	44,794	30,459	21,007	15,611
	Depressive disorders	335,428	71,627	48,945	32,928	24,698
	Schizophrenia and other psychotic disorders	221,962	83,477	63,948	48,399	38,208
	Alcohol or substance-related disorders	26,950	1,004	508	307	234

¹ “Diagnostic group” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 7.13a. Individuals aged 18 years and older served in both 2015 and 2017, by mental health diagnosis in 2015 and employment status in 2015 and 2017: Number

<i>Employment status in 2015</i>		<i>Diagnostic group in 2015¹</i>	<i>Total</i>	<i>Employment status in 2017</i>		
				<i>Employed</i>	<i>Unemployed</i>	<i>Not in labor force</i>
Employed		Total	120,052	85,148	13,081	21,823
		Bipolar disorders	31,236	21,239	3,930	6,067
		Depressive disorders	50,514	36,350	5,185	8,979
		Schizophrenia and other psychotic disorders	20,545	13,949	1,878	4,718
		Alcohol or substance-related disorders	2,533	1,874	386	273
Unemployed		Total	132,530	16,701	78,765	37,064
		Bipolar disorders	36,431	4,745	21,930	9,756
		Depressive disorders	48,719	6,757	29,079	12,883
		Schizophrenia and other psychotic disorders	33,262	2,616	18,978	11,668
		Alcohol or substance-related disorders	5,496	533	4,135	828
Not in labor force		Total	392,119	24,019	26,945	341,155
		Bipolar disorders	96,469	6,677	7,324	82,468
		Depressive disorders	139,777	9,680	9,284	120,813
		Schizophrenia and other psychotic disorders	130,211	4,968	7,751	117,492
		Alcohol or substance-related disorders	7,588	393	967	6,228

¹ "Diagnostic group" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015 and 2017 reporting periods. Data as of 03.03.2022.

Table 7.13b. Individuals aged 18 years and older served in both 2015 and 2017, by mental health diagnosis in 2015 and employment status in 2015 and 2017: Row percent distribution

<i>Employment status in 2015</i>			<i>Employment status in 2017</i>			
<i>Diagnostic group in 2015¹</i>		<i>Total</i>	<i>Employed</i>	<i>Unemployed</i>	<i>Not in labor force</i>	
Employed	Total	100.0	70.9	10.9	18.2	
	Bipolar disorders	100.0	68.0	12.6	19.4	
	Depressive disorders	100.0	72.0	10.3	17.8	
	Schizophrenia and other psychotic disorders	100.0	67.9	9.1	23.0	
	Alcohol or substance-related disorders	100.0	74.0	15.2	10.8	
Unemployed	Total	100.0	12.6	59.4	28.0	
	Bipolar disorders	100.0	13.0	60.2	26.8	
	Depressive disorders	100.0	13.9	59.7	26.4	
	Schizophrenia and other psychotic disorders	100.0	7.9	57.1	35.1	
	Alcohol or substance-related disorders	100.0	9.7	75.2	15.1	
Not in labor force	Total	100.0	6.1	6.9	87.0	
	Bipolar disorders	100.0	6.9	7.6	85.5	
	Depressive disorders	100.0	6.9	6.6	86.4	
	Schizophrenia and other psychotic disorders	100.0	3.8	6.0	90.2	
	Alcohol or substance-related disorders	100.0	5.2	12.7	82.1	

¹ "Diagnostic group" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015 and 2017 reporting periods. Data as of 03.03.2022.

Table 7.14a. Individuals aged 18 years and older served in both 2015 and 2018 by mental health diagnosis in 2015 and employment status in 2015 and 2018: Number

<i>Employment status in 2015</i>		<i>Diagnostic group in 2015¹</i>	<i>Total</i>	<i>Employment status in 2018</i>		
				<i>Employed</i>	<i>Unemployed</i>	<i>Not in labor force</i>
Employed	Total		104,339	69,135	12,877	22,327
		Bipolar disorders	27,120	17,186	3,795	6,139
		Depressive disorders	43,419	29,201	5,076	9,142
		Schizophrenia and other psychotic disorders	19,142	12,167	1,942	5,033
		Alcohol or substance-related disorders	2,132	1,444	387	301
Unemployed	Total		113,533	15,196	63,835	34,502
		Bipolar disorders	30,742	4,370	17,253	9,119
		Depressive disorders	40,632	6,061	22,483	12,088
		Schizophrenia and other psychotic disorders	30,626	2,414	17,416	10,796
		Alcohol or substance-related disorders	4,783	537	3,438	808
Not in labor force	Total		342,305	24,316	29,774	288,215
		Bipolar disorders	83,590	6,689	8,007	68,894
		Depressive disorders	119,429	9,670	9,692	100,067
		Schizophrenia and other psychotic disorders	118,191	5,196	9,404	103,591
		Alcohol or substance-related disorders	6,939	397	924	5,618

¹ "Diagnostic group" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015 and 2018 reporting periods. Data as of 03.03.2022.

Table 7.14b. Individuals aged 18 years and older served in both 2015 and 2018 by mental health diagnosis in 2015 and employment status in 2015 and 2018: Row percent distribution

		Employment status in 2018			
Employment status in 2015					
	Diagnostic group in 2015 ¹	Total	Employed	Unemployed	Not in labor force
Employed	Total	100.0	66.3	12.3	21.4
	Bipolar disorders	100.0	63.4	14.0	22.6
	Depressive disorders	100.0	67.3	11.7	21.1
	Schizophrenia and other psychotic disorders	100.0	63.6	10.1	26.3
	Alcohol or substance-related disorders	100.0	67.7	18.2	14.1
Unemployed	Total	100.0	13.4	56.2	30.4
	Bipolar disorders	100.0	14.2	56.1	29.7
	Depressive disorders	100.0	14.9	55.3	29.7
	Schizophrenia and other psychotic disorders	100.0	7.9	56.9	35.3
	Alcohol or substance-related disorders	100.0	11.2	71.9	16.9
Not in labor force	Total	100.0	7.1	8.7	84.2
	Bipolar disorders	100.0	8.0	9.6	82.4
	Depressive disorders	100.0	8.1	8.1	83.8
	Schizophrenia and other psychotic disorders	100.0	4.4	8.0	87.6
	Alcohol or substance-related disorders	100.0	5.7	13.3	81.0

¹ "Diagnostic group" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015 and 2018 reporting periods. Data as of 03.03.2022.

Table 7.15a. Individuals aged 18 years and older served in both 2015 and 2019 by mental health diagnosis in 2015 and employment status in 2015 and 2019: Number

<i>Employment status in 2015</i>		<i>Diagnostic group in 2015¹</i>	<i>Total</i>	<i>Employment status in 2019</i>		
				<i>Employed</i>	<i>Unemployed</i>	<i>Not in labor force</i>
Employed	Total		84,067	52,955	9,820	21,292
	Bipolar disorders		21,679	13,023	2,929	5,727
	Depressive disorders		34,583	22,073	3,849	8,661
	Schizophrenia and other psychotic disorders		16,422	9,961	1,505	4,956
	Alcohol or substance-related disorders		1,774	1,146	355	273
Unemployed	Total		83,634	11,851	41,546	30,237
	Bipolar disorders		22,745	3,450	11,415	7,880
	Depressive disorders		29,578	4,672	14,586	10,320
	Schizophrenia and other psychotic disorders		23,310	2,001	11,223	10,086
	Alcohol or substance-related disorders		4,016	499	2,639	878
Not in labor force	Total		279,874	21,739	23,887	234,248
	Bipolar disorders		68,307	5,977	6,284	56,046
	Depressive disorders		95,920	8,688	7,529	79,703
	Schizophrenia and other psychotic disorders		99,663	4,773	7,875	87,015
	Alcohol or substance-related disorders		5,942	371	853	4,718

¹ "Diagnostic group" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015 and 2019 reporting periods. Data as of 03.03.2022.

Table 7.15b. Individuals aged 18 years and older served in both 2015 and 2019 by mental health diagnosis in 2015 and employment status in 2015 and 2019: Row percent distribution

<i>Employment status in 2015</i>			<i>Employment status in 2019</i>			
<i>Diagnostic group in 2015¹</i>		<i>Total</i>	<i>Employed</i>	<i>Unemployed</i>	<i>Not in labor force</i>	
Employed	Total	100.0	63.0	11.7	25.3	
	Bipolar disorders	100.0	60.1	13.5	26.4	
	Depressive disorders	100.0	63.8	11.1	25.0	
	Schizophrenia and other psychotic disorders	100.0	60.7	9.2	30.2	
	Alcohol or substance-related disorders	100.0	64.6	20.0	15.4	
Unemployed	Total	100.0	14.2	49.7	36.2	
	Bipolar disorders	100.0	15.2	50.2	34.6	
	Depressive disorders	100.0	15.8	49.3	34.9	
	Schizophrenia and other psychotic disorders	100.0	8.6	48.1	43.3	
	Alcohol or substance-related disorders	100.0	12.4	65.7	21.9	
Not in labor force	Total	100.0	7.8	8.5	83.7	
	Bipolar disorders	100.0	8.8	9.2	82.1	
	Depressive disorders	100.0	9.1	7.8	83.1	
	Schizophrenia and other psychotic disorders	100.0	4.8	7.9	87.3	
	Alcohol or substance-related disorders	100.0	6.2	14.4	79.4	

¹ "Diagnostic group" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015 and 2019 reporting periods. Data as of 03.03.2022.

Table 7.16a. Individuals aged 18 years and older served in both 2015 and 2020, by mental health diagnosis in 2015 and employment status in 2015 and 2020: Number

<i>Employment status in 2015</i>		<i>Diagnostic group in 2015¹</i>	<i>Total</i>	<i>Employment status in 2020</i>		
				<i>Employed</i>	<i>Unemployed</i>	<i>Not in labor force</i>
Employed	Total		75,016	45,829	8,420	20,767
	Bipolar disorders		19,201	11,190	2,432	5,579
	Depressive disorders		32,065	19,829	3,514	8,722
	Schizophrenia and other psychotic disorders		14,561	8,372	1,251	4,938
	Alcohol or substance-related disorders		1,421	897	281	243
Unemployed	Total		66,049	9,531	31,029	25,489
	Bipolar disorders		17,804	2,808	8,454	6,542
	Depressive disorders		23,960	3,831	11,245	8,884
	Schizophrenia and other psychotic disorders		18,385	1,609	8,113	8,663
	Alcohol or substance-related disorders		3,211	392	2,089	730
Not in labor force	Total		243,149	19,789	19,169	204,191
	Bipolar disorders		58,768	5,515	5,039	48,214
	Depressive disorders		86,459	8,183	6,533	71,743
	Schizophrenia and other psychotic disorders		85,257	4,120	5,949	75,188
	Alcohol or substance-related disorders		5,145	329	756	4,060

¹ "Diagnostic group" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015 and 2020 reporting periods. Data as of 03.03.2022.

Table 7.16b. Individuals aged 18 years and older served in both 2015 and 2020, by mental health diagnosis in 2015 and employment status in 2015 and 2020: Row percent distribution

		Employment status in 2020			
Employment status in 2015					
	Diagnostic group in 2015 ¹	Total	Employed	Unemployed	Not in labor force
Employed	Total	100.0	61.1	11.2	27.7
	Bipolar disorders	100.0	58.3	12.7	29.1
	Depressive disorders	100.0	61.8	11.0	27.2
	Schizophrenia and other psychotic disorders	100.0	57.5	8.6	33.9
	Alcohol or substance-related disorders	100.0	63.1	19.8	17.1
Unemployed	Total	100.0	14.4	47.0	38.6
	Bipolar disorders	100.0	15.8	47.5	36.7
	Depressive disorders	100.0	16.0	46.9	37.1
	Schizophrenia and other psychotic disorders	100.0	8.8	44.1	47.1
	Alcohol or substance-related disorders	100.0	12.2	65.1	22.7
Not in labor force	Total	100.0	8.1	7.9	84.0
	Bipolar disorders	100.0	9.4	8.6	82.0
	Depressive disorders	100.0	9.5	7.6	83.0
	Schizophrenia and other psychotic disorders	100.0	4.8	7.0	88.2
	Alcohol or substance-related disorders	100.0	6.4	14.7	78.9

¹ "Diagnostic group" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015 and 2020 reporting periods. Data as of 03.03.2022.

Table 8.1. Individuals, by mental health diagnosis, Census region, Census division, and state or jurisdiction: Number, 2020

Census region, Census division, and state or jurisdiction	Mental health diagnosis ¹								
	Total	Trauma- and stressor- related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ²
					Bipolar disorders	Depressive disorders			
Total	5,928,553	1,295,221	1,404,997	654,495	713,775	1,834,097	161,378	765,575	1,595,452
Census region³									
Northeast	988,116	197,772	168,852	109,798	109,860	271,137	27,848	117,531	299,956
Midwest	1,484,942	356,071	454,750	202,787	180,165	497,002	61,918	165,760	342,633
South	1,634,014	327,693	336,541	185,315	266,065	561,091	44,525	261,910	379,550
West	1,818,097	413,549	444,532	156,375	157,427	504,229	27,052	218,854	572,384
Census division³									
New England	102,829	26,235	27,088	8,928	14,367	34,444	4,513	19,296	22,129
Middle Atlantic	885,287	171,537	141,764	100,870	95,493	236,693	23,335	98,235	277,827
East North Central	882,610	195,574	229,799	120,455	113,388	263,057	38,441	105,884	223,663
West North Central	602,332	160,497	224,951	82,332	66,777	233,945	23,477	59,876	118,970
South Atlantic	748,688	156,690	137,265	73,418	122,818	258,095	22,836	133,092	175,820
East South Central	415,607	76,717	88,115	60,154	48,780	131,136	9,140	51,853	135,408
West South Central	469,719	94,286	111,161	51,743	94,467	171,860	12,549	76,965	68,322
Mountain	815,676	185,364	203,286	70,538	70,209	197,279	11,366	62,595	280,059
Pacific	1,002,421	228,185	241,246	85,837	87,218	306,950	15,686	156,259	292,325
State or jurisdiction									
Alabama	97,473	13,116	16,620	15,798	12,058	31,105	2,135	17,851	16,441
Alaska	10,202	3,329	1,521	512	527	2,613	97	648	2,203
Arizona	350,300	69,162	75,608	32,715	30,140	71,665	3,430	27,491	136,118
Arkansas	73,611	12,099	9,586	2,795	6,851	19,271	390	8,543	14,148
California	626,774	108,332	102,964	42,047	53,238	182,518	8,024	117,846	175,338
Colorado	126,315	40,976	26,738	9,644	14,828	39,422	2,604	12,549	25,857
Connecticut	41,625	9,682	10,148	2,130	7,356	14,395	2,057	9,827	6,468
Delaware	10,559	1,388	1,356	636	2,007	3,383	368	2,083	5,517
District of Columbia	35,161	3,871	1,525	2,157	5,839	13,701	73	6,372	3,893
Florida	244,240	44,036	21,148	13,991	32,579	76,895	952	35,609	27,475
Georgia	140,657	25,596	31,929	12,127	25,989	45,549	3,432	29,419	55,686
Hawaii	6,083	1,120	501	512	784	1,352	148	2,354	1,221
Idaho	7,082	805	1,237	691	1,440	2,389	514	1,993	1,306
Illinois	39,267	8,298	8,079	4,198	6,111	11,329	1,239	7,682	8,625

Continued. See notes at end of table.

Table 8.1. Individuals, by mental health diagnosis, Census region, Census division, and state or jurisdiction: Number, 2020 (continued)

Census region, Census division, and state or jurisdiction	Mental health diagnosis ¹								
	Total	Trauma- and stressor- related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ²
					Bipolar disorders	Depressive disorders			
Indiana	137,905	31,334	50,737	27,231	13,981	51,612	6,161	14,278	36,630
Iowa	56,738	4,714	22,784	7,147	6,501	20,852	1,157	7,155	9,473
Kansas	139,114	30,712	33,232	15,678	15,838	47,536	2,328	8,746	34,069
Kentucky	153,854	35,466	37,282	19,408	12,056	43,158	3,599	11,274	73,239
Louisiana	17,526	1,907	3,904	1,879	3,228	6,656	288	3,934	2,174
Maine	‡	‡	‡	‡	‡	‡	‡	‡	‡
Maryland	‡	‡	‡	‡	‡	‡	‡	‡	‡
Massachusetts	6,898	1,138	1,413	481	1,501	1,644	206	3,156	939
Michigan	207,770	40,535	45,722	28,347	32,606	61,284	15,136	34,612	90,889
Minnesota	284,601	107,500	147,707	48,532	24,968	126,039	17,267	25,217	58,153
Mississippi	101,461	14,273	17,736	20,989	12,179	29,870	2,231	14,574	38,370
Missouri	75,768	6,638	10,943	7,338	13,219	25,147	1,111	13,252	5,827
Montana	73,240	14,170	29,733	7,241	6,423	23,201	860	4,003	17,276
Nebraska	21,981	4,265	4,709	1,203	3,828	7,126	626	2,671	7,011
Nevada	13,036	1,930	2,244	669	1,123	2,395	409	2,321	4,183
New Hampshire	‡	‡	‡	‡	‡	‡	‡	‡	‡
New Jersey	315,988	61,405	35,098	25,562	31,660	79,778	9,266	37,502	176,753
New Mexico	179,605	43,583	41,569	13,116	9,932	35,556	565	7,082	81,353
New York ⁴	54,840	14,035	7,606	3,779	6,359	11,289	9,177	19,827	12,875
North Carolina	99,827	24,649	22,486	5,430	19,466	41,645	3,499	16,122	14,627
North Dakota	9,341	2,547	3,880	1,308	1,073	3,624	937	1,490	2,616
Ohio	429,022	107,550	110,594	56,771	54,712	126,003	7,226	39,576	73,300
Oklahoma	94,809	20,896	16,707	4,459	10,594	37,554	397	10,429	9,080
Oregon	145,349	54,561	57,606	16,169	12,514	45,310	3,299	13,150	30,378
Pennsylvania	514,459	96,097	99,060	71,529	57,474	145,626	4,892	40,906	88,199
Puerto Rico	2,950	88	291	192	226	601	29	1,379	792

Continued. See notes at end of table.

Table 8.1. Individuals, by mental health diagnosis, Census region, Census division, and state or jurisdiction: Number, 2020 (continued)

Census region, Census division, and state or jurisdiction	Mental health diagnosis ¹								
	Total	Trauma- and stressor- related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ²
					Bipolar disorders	Depressive disorders			
Rhode Island	29,366	5,568	10,635	3,724	4,176	13,251	1,487	4,466	5,434
South Carolina	95,263	25,244	22,886	16,762	12,157	29,930	4,420	17,323	34,166
South Dakota	14,789	4,121	1,696	1,126	1,350	3,621	51	1,345	1,821
Tennessee*	62,819	13,862	16,477	3,959	12,487	27,003	1,175	8,154	7,358
Texas	283,773	59,384	80,964	42,610	73,794	108,379	11,474	54,059	42,920
Utah	50,333	10,875	21,525	5,139	4,652	16,790	2,356	6,068	11,239
Vermont	24,940	9,847	4,892	2,593	1,334	5,154	763	1,847	9,288
Virginia	122,981	31,906	35,935	22,315	24,781	46,992	10,092	26,164	34,456
Washington	214,013	60,843	78,654	26,597	20,155	75,157	4,118	22,261	83,185
West Virginia	‡	‡	‡	‡	‡	‡	‡	‡	‡
Wisconsin	68,646	7,857	14,667	3,908	5,978	12,829	8,679	9,736	14,219
Wyoming	15,765	3,863	4,632	1,323	1,671	5,861	628	1,088	2,727
Other jurisdictions ⁵	434	48	31	28	32	37	6	141	137

‡ No data submitted

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes substance use diagnoses. See Appendix E.

³ The Commonwealth of the Northern Mariana Islands, the Republic of Palau, and the Commonwealth of Puerto Rico are not included in any Census region or division.

⁴ New York and Tennessee report only a subset of the total individuals served through their SMHAs.

⁵ Other jurisdictions include the Commonwealth of the Northern Mariana Islands and the Republic of Palau.

Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 8.2. Individuals per 100,000 population (2020 estimate), by mental health diagnosis, Census region, Census division, and state or jurisdiction: Number, 2020

<i>Census region, Census division, and state or jurisdiction</i>	<i>Mental health diagnosis¹</i>								
	<i>Total</i>				<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders²</i>
		<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>ADHD</i>	<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Total	1,859	406	441	205	224	575	51	240	500
Census region³									
Northeast	1,860	372	318	207	207	510	52	221	565
Midwest	2,174	521	666	297	264	728	91	243	502
South	1,375	276	283	156	224	472	37	220	319
West	2,311	526	565	199	200	641	34	278	728
Census division³									
New England	848	216	223	74	118	284	37	159	182
Middle Atlantic	2,159	418	346	246	233	577	57	240	678
East North Central	1,885	418	491	257	242	562	82	226	478
West North Central	2,804	747	1,047	383	311	1,089	109	279	554
South Atlantic	1,279	268	234	125	210	441	39	227	300
East South Central	2,159	398	458	312	253	681	47	269	703
West South Central	1,145	230	271	126	230	419	31	188	167
Mountain	3,235	735	806	280	278	782	45	248	1,111
Pacific	1,876	427	451	161	163	574	29	292	547
State or jurisdiction									
Alabama	1,981	267	338	321	245	632	43	363	334
Alaska	1,395	455	208	70	72	357	13	89	301
Arizona	4,720	932	1,019	441	406	966	46	370	1,834
Arkansas	2,429	399	316	92	226	636	13	282	467
California	1,592	275	262	107	135	464	20	299	445
Colorado	2,175	706	460	166	255	679	45	216	445
Connecticut	1,170	272	285	60	207	405	58	276	182
Delaware	1,070	141	137	64	203	343	37	211	559
District of Columbia	4,933	543	214	303	819	1,922	10	894	546
Florida	1,124	203	97	64	150	354	4	164	126
Georgia	1,313	239	298	113	243	425	32	275	520
Hawaii	432	80	36	36	56	96	11	167	87
Idaho	388	44	68	38	79	131	28	109	71
Illinois	312	66	64	33	49	90	10	61	69

Continued. See notes at end of table.

Table 8.2. Individuals per 100,000 population (2020 estimate), by mental health diagnosis, Census region, Census division, and state or jurisdiction: Number, 2020 (continued)

Census region, Census division, and state or jurisdiction	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ²
					Bipolar disorders	Depressive disorders			
Indiana	2,042	464	751	403	207	764	91	211	542
Iowa	1,793	149	720	226	206	659	37	226	299
Kansas	4,774	1,054	1,141	538	544	1,631	80	300	1,169
Kentucky	3,436	792	833	433	269	964	80	252	1,636
Louisiana	377	41	84	40	69	143	6	85	47
Maine	‡	‡	‡	‡	‡	‡	‡	‡	‡
Maryland	‡	‡	‡	‡	‡	‡	‡	‡	‡
Massachusetts	100	17	21	7	22	24	3	46	14
Michigan	2,085	407	459	284	327	615	152	347	912
Minnesota	5,031	1,900	2,611	858	441	2,228	305	446	1,028
Mississippi	3,420	481	598	707	411	1,007	75	491	1,293
Missouri	1,232	108	178	119	215	409	18	215	95
Montana	6,778	1,311	2,752	670	594	2,147	80	370	1,599
Nebraska	1,134	220	243	62	198	368	32	138	362
Nevada	415	62	72	21	36	76	13	74	133
New Hampshire	‡	‡	‡	‡	‡	‡	‡	‡	‡
New Jersey	3,557	691	395	288	356	898	104	422	1,990
New Mexico	8,527	2,069	1,974	623	472	1,688	27	336	3,862
New York ⁴	284	73	39	20	33	58	47	103	67
North Carolina	942	233	212	51	184	393	33	152	138
North Dakota	1,221	333	507	171	140	474	122	195	342
Ohio	3,669	920	946	486	468	1,078	62	338	627
Oklahoma	2,382	525	420	112	266	943	10	262	228
Oregon	3,427	1,286	1,358	381	295	1,068	78	310	716
Pennsylvania	4,024	752	775	560	450	1,139	38	320	690
Puerto Rico	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a

Continued. See notes at end of table.

Table 8.2. Individuals per 100,000 population (2020 estimate), by mental health diagnosis, Census region, Census division, and state or jurisdiction: Number, 2020 (continued)

<i>Census region, Census division, and state or jurisdiction</i>	<i>Mental health diagnosis¹</i>								
	<i>Total</i>	<i>Trauma- and stressor-related disorders</i>			<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders²</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Rhode Island	2,778	527	1,006	352	395	1,253	141	422	514
South Carolina	1,826	484	439	321	233	574	85	332	655
South Dakota	1,657	462	190	126	151	406	6	151	204
Tennessee ⁴	912	201	239	57	181	392	17	118	107
Texas	967	202	276	145	251	369	39	184	146
Utah	1,549	335	662	158	143	517	73	187	346
Vermont	4,001	1,580	785	416	214	827	122	296	1,490
Virginia	1,432	371	418	260	288	547	117	305	401
Washington	2,782	791	1,022	346	262	977	54	289	1,081
West Virginia	‡	‡	‡	‡	‡	‡	‡	‡	‡
Wisconsin	1,177	135	251	67	102	220	149	167	244
Wyoming	2,707	663	795	227	287	1,006	108	187	468
Other jurisdictions ⁵	<i>n/a</i>	<i>n/a</i>	<i>n/a</i>	<i>n/a</i>	<i>n/a</i>	<i>n/a</i>	<i>n/a</i>	<i>n/a</i>	<i>n/a</i>

‡ No data reported; † Less than 1 per 100,000 population; *n/a* Population detail required for adjustment was not available.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes substance use diagnoses. See Appendix E.

³ The Commonwealth of the Northern Mariana Islands, the Republic of Palau, and the Commonwealth of Puerto Rico are not included in any Census region or division.

⁴ New York and Tennessee report only a subset of the total individuals served through their SMHAs.

⁵ Other jurisdictions include the Commonwealth of the Northern Mariana Islands and the Republic of Palau.

NOTE: States that submitted insufficient data for inclusion in this report, as well as their reference populations, are excluded from calculations presented in this table.

SOURCES: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022; Population: U.S. Census Bureau, Population Estimates Program, SC-EST2020-alldata6: Annual State Resident Population Estimates for 6 Race Groups (5 Race Alone Groups and Two or More Races) by Age, Sex, and Hispanic Origin: April 1, 2010 to July 1, 2019; April 1, 2020; and July 1, 2020. <https://www2.census.gov/programs-surveys/popest/datasets/2010-2020/state/asrh/SC-EST2020-ALLDATA6.csv>.

**Table 8.3. Individuals per 100,000 population, by mental health diagnosis, Census region, Census division, and state or jurisdiction:
Number, 2020, adjusted to 2010 population estimate for age, gender, and race**

Census region, Census division, and state or jurisdiction	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ²
					Bipolar disorders	Depressive disorders			
Total	1,721	381	415	201	216	535	49	216	455
Census region³									
Northeast	1,736	356	309	215	198	466	48	189	491
Midwest	2,094	499	626	287	258	695	89	242	491
South	1,321	273	287	153	222	460	37	191	305
West	2,031	477	498	186	185	563	32	257	629
Census division³									
New England	766	205	206	73	110	256	35	150	150
Middle Atlantic	2,020	398	342	255	225	529	52	199	588
East North Central	1,834	412	479	260	234	542	81	217	477
West North Central	2,672	696	955	339	313	1,048	107	305	515
South Atlantic	1,214	266	240	125	205	423	38	184	282
East South Central	2,093	402	467	302	252	664	47	228	691
West South Central	1,100	221	267	120	231	410	31	180	155
Mountain	2,658	621	676	235	241	661	40	217	859
Pacific	1,660	397	401	156	156	497	28	268	495
State or jurisdiction									
Alabama	1,879	272	347	313	255	610	44	277	319
Alaska	1,042	319	152	54	74	261	11	74	215
Arizona	3,392	661	772	344	308	711	35	260	1,269
Arkansas	1,961	303	246	75	204	505	11	242	376
California	1,313	235	203	97	123	365	19	262	375
Colorado	2,156	708	437	168	250	662	43	231	439
Connecticut	1,105	266	274	60	199	378	55	258	172
Delaware	984	135	134	65	199	306	34	172	540
District of Columbia	1,664	185	78	120	284	612	5	322	184
Florida	1,168	217	103	74	154	364	4	153	135
Georgia	1,157	224	310	92	241	395	32	188	465
Hawaii	440	70	46	39	80	104	14	160	88
Idaho	403	42	66	37	80	125	26	130	74
Illinois	286	61	60	32	46	82	10	55	62

Continued. See notes at end of table.

**Table 8.3. Individuals per 100,000 population, by mental health diagnosis, Census region, Census division, and state or jurisdiction:
Number, 2020, adjusted to 2010 population estimate for age, gender, and race (continued)**

Census region, Census division, and state or jurisdiction	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ²
					Bipolar disorders	Depressive disorders			
Indiana	2,088	476	759	412	213	775	93	226	553
Iowa	1,611	144	643	174	203	598	35	240	251
Kansas	4,775	1,067	1,125	534	561	1,623	81	341	1,167
Kentucky	3,480	814	823	448	269	960	80	262	1,657
Louisiana	316	35	79	31	66	128	6	54	35
Maine	‡	‡	‡	‡	‡	‡	‡	‡	‡
Maryland	‡	‡	‡	‡	‡	‡	‡	‡	‡
Massachusetts	84	14	16	6	18	18	3	41	11
Michigan	2,092	420	466	295	329	616	154	330	919
Minnesota	4,632	1,739	2,385	720	440	2,130	300	487	917
Mississippi	2,972	474	644	575	424	912	76	321	1,082
Missouri	1,233	108	177	121	216	407	18	219	95
Montana	6,927	1,384	2,752	733	617	2,100	83	417	1,680
Nebraska	1,232	231	251	62	211	386	34	168	402
Nevada	343	53	56	18	31	64	11	63	110
New Hampshire	‡	‡	‡	‡	‡	‡	‡	‡	‡
New Jersey	2,786	548	324	237	307	682	89	311	1,553
New Mexico	7,723	1,967	1,868	566	456	1,529	25	310	3,322
New York ⁴	252	66	39	21	31	54	42	83	64
North Carolina	921	233	220	51	184	389	34	128	136
North Dakota	1,193	332	491	161	143	467	121	212	329
Ohio	3,412	872	884	466	427	992	55	304	592
Oklahoma	2,541	547	443	116	287	1,008	11	295	238
Oregon	3,416	1,306	1,312	419	280	1,039	71	314	721
Pennsylvania	4,249	813	808	619	466	1,178	39	318	737
Puerto Rico	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a

Continued. See notes at end of table.

Table 8.3. Individuals per 100,000 population, by mental health diagnosis, Census region, Census division, and state or jurisdiction: Number, 2020, adjusted to 2010 population estimate for age, gender, and race (continued)

Mental health diagnosis ¹									
Census region, Census division, and state or jurisdiction	Total	Trauma- and stressor-related disorders Anxiety disorders ADHD			Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ²
					Bipolar disorders	Depressive disorders			
		Rhode Island	2,505	468	903	338			
South Carolina	1,687	478	452	302	236	549	88	243	604
South Dakota	1,857	546	203	139	170	432	7	177	225
Tennessee*	880	195	236	54	178	382	17	108	102
Texas	971	204	279	141	256	373	40	185	143
Utah	1,525	330	634	143	154	516	77	229	313
Vermont	3,757	1,773	799	545	186	770	110	296	1,109
Virginia	1,388	370	432	267	286	538	114	260	385
Washington	2,894	854	1,056	373	277	1,019	56	312	1,109
West Virginia	-	-	-	‡	‡	‡	‡	‡	‡
Wisconsin	1,212	141	245	71	105	211	183	184	236
Wyoming	2,755	694	791	223	276	1,029	115	232	492
Other jurisdictions ⁵	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a

‡ No data reported; † Less than 1 per 100,000 population; *n/a* Population detail required for adjustment was not available.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes substance use diagnoses. See Appendix E.

³ The Commonwealth of the Northern Mariana Islands, the Republic of Palau, and the Commonwealth of Puerto Rico are not included in any Census region or division.

⁴ New York and Tennessee report only a subset of the total individuals served through their SMHAs.

⁵ Other jurisdictions include the Commonwealth of the Northern Mariana Islands and the Republic of Palau.

NOTE: States that submitted insufficient data for inclusion in this report, as well as their reference populations, are excluded from calculations presented in this table.

SOURCES: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022; Population: U.S. Census Bureau, Population Estimates Program, SC-EST2020-alldata6: Annual State Resident Population Estimates for 6 Race Groups (5 Race Alone Groups and Two or More Races) by Age, Sex, and Hispanic Origin: April 1, 2010 to July 1, 2019; April 1, 2020; and July 1, 2020. <https://www2.census.gov/programs-surveys/popest/datasets/2010-2020/state/asrh/SC-EST2020-ALLDATA6.csv>.

Table 8.4. Individuals with diagnosis of trauma- and stressor-related disorders, by Census region, Census division, and state or jurisdiction: Number, 2015–2020

<i>Census region, Census division, and state or jurisdiction</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals with diagnosis of trauma- and stressor-related disorders	853,482	878,639	1,035,177	1,161,339	1,253,914	1,295,221
Census region¹						
Northeast	82,433	112,486	190,011	194,991	196,595	197,772
Midwest	241,570	225,838	267,783	340,881	394,369	356,071
South	241,042	236,608	270,025	295,793	313,025	327,693
West	288,425	303,689	307,207	329,514	349,730	413,549
Census division¹						
New England	31,432	30,996	43,998	36,390	26,756	26,235
Middle Atlantic	51,001	81,490	146,013	158,601	169,839	171,537
East North Central	130,497	114,610	167,884	186,343	231,935	195,574
West North Central	111,073	111,228	99,899	154,538	162,434	160,497
South Atlantic	135,397	113,574	139,111	147,460	152,373	156,690
East South Central	61,256	60,623	60,177	68,804	72,759	76,717
West South Central	44,389	62,411	70,737	79,529	87,893	94,286
Mountain	122,146	127,021	136,247	134,868	122,011	185,364
Pacific	166,279	176,668	170,960	194,646	227,719	228,185
State or jurisdiction						
Alabama	11,618	10,585	11,259	12,038	11,232	13,116
Alaska	‡	‡	‡	‡	3,260	3,329
Arizona	38,779	40,497	21,792	23,938	‡	69,162
Arkansas	9,651	11,610	12,413	11,469	11,739	12,099
California	111,808	105,242	95,732	110,480	119,485	108,332
Colorado	24,329	24,783	41,317	46,293	42,209	40,976
Connecticut	14,329	14,831	20,716	21,254	10,158	9,682
Delaware	1,736	1,069	1,237	1,202	1,422	1,388
District of Columbia	2,555	2,546	3,173	3,856	3,764	3,871
Florida	21,425	24,093	34,988	30,829	28,781	44,036
Georgia	‡	‡	‡	‡	‡	25,596
Hawaii	2,653	2,309	2,567	3,044	2,542	1,120
Idaho	259	585	846	863	865	805
Illinois	21,692	2,411	12,469	12,353	8,886	8,298

Continued. See notes at end of table.

Table 8.4. Individuals with diagnosis of trauma- and stressor-related disorders, by Census region, Census division, and state or jurisdiction: Number, 2015–2020 (continued)

<i>Census region, Census division, and state or jurisdiction</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Indiana	26,088	24,724	25,841	27,140	29,311	31,334
Iowa	27,949	25,028	9,787	38,113	45,939	4,714
Kansas	‡	‡	‡	‡	‡	30,712
Kentucky	30,402	30,164	31,081	33,793	36,056	35,466
Louisiana	2,274	1,520	2,429	1,934	1,600	1,907
Maine	‡	‡	5,275	‡	‡	‡
Maryland	30,403	30,946	34,607	38,031	41,022	‡
Massachusetts	6,273	2,266	2,877	1,593	1,467	1,138
Michigan	‡	‡	23,546	35,541	43,451	40,535
Minnesota	65,164	72,479	75,755	101,004	99,344	107,500
Mississippi	7,316	8,322	8,007	10,823	11,473	14,273
Missouri	7,604	4,517	5,631	6,157	7,087	6,638
Montana	10,270	7,738	11,729	15,953	14,438	14,170
Nebraska	4,418	3,882	4,070	4,381	4,063	4,265
Nevada	4,132	1,823	2,021	1,931	2,029	1,930
New Hampshire	‡	‡	‡	‡	‡	‡
New Jersey	42,742	4,081	55,169	60,233	61,655	61,405
New Mexico	30,002	37,557	44,352	31,909	48,106	43,583
New York ²	8,259	11,587	11,625	12,490	13,272	14,035
North Carolina	43,282	18,906	25,579	21,506	24,196	24,649
North Dakota	2,138	1,491	1,099	940	1,918	2,547
Ohio	72,858	77,725	98,584	103,492	142,476	107,550
Oklahoma	10,558	13,892	15,257	18,458	19,507	20,896
Oregon	36,211	48,867	48,971	51,852	58,645	54,561
Pennsylvania	‡	65,822	79,219	85,878	94,912	96,097
Puerto Rico	7	7	130	152	160	88

Continued. See notes at end of table.

Table 8.4. Individuals with diagnosis of trauma- and stressor-related disorders, by Census region, Census division, and state or jurisdiction: Number, 2015–2020 (continued)

<i>Census region, Census division, and state or jurisdiction</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Rhode Island	4,278	6,011	6,526	3,958	5,305	5,568
South Carolina	13,786	14,128	15,406	18,177	22,828	25,244
South Dakota	3,800	3,831	3,557	3,943	4,083	4,121
Tennessee ²	11,920	11,552	9,830	12,150	13,998	13,862
Texas	21,906	35,389	40,638	47,668	55,047	59,384
Utah	10,093	9,543	10,017	10,092	10,340	10,875
Vermont	6,552	7,888	8,604	9,585	9,826	9,847
Virginia	22,210	21,886	24,121	27,840	30,360	31,906
Washington	15,607	20,250	23,690	29,270	43,787	60,843
West Virginia	‡	‡	‡	6,019	‡	‡
Wisconsin	9,859	9,750	7,444	7,817	7,811	7,857
Wyoming	4,282	4,495	4,173	3,889	4,024	3,863
Other jurisdictions ³	5	11	21	8	35	48

‡ No data submitted

¹ The Commonwealth of the Northern Mariana Islands, the Republic of Palau, and the Commonwealth of Puerto Rico are not included in any Census region or division.

² New York and Tennessee report only a subset of the total individuals served through their SMHAs.

³ Other jurisdictions include the Commonwealth of the Northern Mariana Islands and the Republic of Palau.

NOTE: This table refers to all mental health diagnoses, which is up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022

**Table 8.5. Individuals with diagnosis of anxiety disorders, by Census region, Census division, and state or jurisdiction:
Number, 2015–2020**

<i>Census region, Census division, and state or jurisdiction</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals with diagnosis of anxiety disorders	755,391	837,961	1,052,774	1,239,939	1,339,890	1,404,997
Census region¹						
Northeast	53,382	89,763	149,156	160,318	165,975	168,852
Midwest	228,197	240,110	324,443	414,077	500,694	454,750
South	251,198	263,647	279,628	320,252	325,004	336,541
West	222,607	244,424	299,382	344,956	347,845	444,532
Census division¹						
New England	20,465	24,425	34,127	32,517	28,204	27,088
Middle Atlantic	32,917	65,338	115,029	127,801	137,771	141,764
East North Central	120,552	109,128	190,510	215,865	254,841	229,799
West North Central	107,645	130,982	133,933	198,212	245,853	224,951
South Atlantic	125,440	109,395	124,135	148,315	140,466	137,265
East South Central	75,327	83,317	75,145	82,716	85,737	88,115
West South Central	50,431	70,935	80,348	89,221	98,801	111,161
Mountain	108,865	121,662	150,061	167,595	126,418	203,286
Pacific	113,742	122,762	149,321	177,361	221,427	241,246
State or jurisdiction						
Alabama	15,723	13,406	13,682	14,134	14,903	16,620
Alaska	‡	‡	‡	‡	1,574	1,521
Arizona	38,314	47,724	49,419	58,436	‡	75,608
Arkansas	7,721	9,601	10,588	8,839	9,171	9,586
California	82,469	81,057	89,222	109,661	114,960	102,964
Colorado	20,976	18,118	28,851	29,738	26,919	26,738
Connecticut	10,099	13,154	16,612	17,611	11,008	10,148
Delaware	787	498	670	829	1,172	1,356
District of Columbia	874	834	1,602	1,830	1,648	1,525
Florida	11,283	12,988	14,640	15,808	14,334	21,148
Georgia	‡	‡	‡	‡	‡	31,929
Hawaii	507	606	690	649	561	501
Idaho	216	1,322	1,434	1,447	1,399	1,237
Illinois	26,838	2,824	13,966	13,247	8,753	8,079

Continued. See notes at end of table.

**Table 8.5. Individuals with diagnosis of anxiety disorders, by Census region, Census division, and state or jurisdiction:
Number, 2015–2020 (continued)**

<i>Census region, Census division, and state or jurisdiction</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Indiana	31,280	40,948	42,251	43,577	46,769	50,737
Iowa	17,742	26,162	10,640	51,640	88,718	22,784
Kansas	‡	‡	‡	‡	‡	33,232
Kentucky	35,645	37,079	37,719	38,815	38,891	37,282
Louisiana	3,950	2,726	4,270	3,973	2,924	3,904
Maine	‡	‡	4,573	‡	‡	‡
Maryland	27,854	31,961	36,749	41,593	45,782	‡
Massachusetts	3,459	1,370	2,110	2,078	1,924	1,413
Michigan	‡	‡	33,824	47,125	50,718	45,722
Minnesota	74,419	86,835	103,951	126,735	134,815	147,707
Mississippi	6,384	17,296	8,909	12,974	14,299	17,736
Missouri	6,241	9,745	11,100	11,386	12,552	10,943
Montana	9,181	7,390	13,169	24,100	27,331	29,733
Nebraska	3,694	3,760	4,271	4,653	4,586	4,709
Nevada	4,799	2,021	2,062	1,981	2,200	2,244
New Hampshire	‡	‡	‡	‡	‡	‡
New Jersey	26,608	1,754	32,462	34,776	35,492	35,098
New Mexico	16,799	24,034	32,518	28,366	43,906	41,569
New York ²	6,309	9,568	7,291	7,442	7,527	7,606
North Carolina	44,636	20,609	25,413	21,286	22,646	22,486
North Dakota	4,327	3,220	2,566	2,188	3,548	3,880
Ohio	54,438	55,566	89,315	99,452	135,471	110,594
Oklahoma	9,883	13,483	14,883	16,688	15,674	16,707
Oregon	20,622	25,018	38,670	39,605	55,019	57,606
Pennsylvania	‡	54,016	75,276	85,583	94,752	99,060
Puerto Rico	5	5	156	326	352	291

Continued. See notes at end of table.

**Table 8.5. Individuals with diagnosis of anxiety disorders, by Census region, Census division, and state or jurisdiction:
Number, 2015–2020 (continued)**

<i>Census region, Census division, and state or jurisdiction</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Rhode Island	3,868	6,198	6,811	8,443	10,551	10,635
South Carolina	14,136	15,265	16,217	17,956	21,158	22,886
South Dakota	1,222	1,260	1,405	1,610	1,634	1,696
Tennessee ²	17,575	15,536	14,835	16,793	17,644	16,477
Texas	28,877	45,125	50,607	59,721	71,032	80,964
Utah	14,777	16,998	18,502	19,378	19,899	21,525
Vermont	3,039	3,703	4,021	4,385	4,721	4,892
Virginia	25,870	27,240	28,844	31,451	33,726	35,935
Washington	10,144	16,081	20,739	27,446	49,313	78,654
West Virginia	‡	‡	‡	17,562	‡	‡
Wisconsin	7,996	9,790	11,154	12,464	13,130	14,667
Wyoming	3,803	4,055	4,106	4,149	4,764	4,632
Other jurisdictions ³	2	12	9	10	20	31

‡ No data submitted

¹ The Commonwealth of the Northern Mariana Islands, the Republic of Palau, and the Commonwealth of Puerto Rico are not included in any Census region or division.

² New York and Tennessee report only a subset of the total individuals served through their SMHAs.

³ Other jurisdictions include the Commonwealth of the Northern Mariana Islands and the Republic of Palau.

NOTE: This table refers to all mental health diagnoses, which is up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022

Table 8.6. Individuals with diagnosis of attention deficit/hyperactivity disorder, by Census region, Census division, and state or jurisdiction: Number, 2015–2020

<i>Census region, Census division, and state or jurisdiction</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals with diagnosis of attention deficit/hyperactivity disorder	510,737	560,910	621,999	696,888	672,592	654,495
Census region¹						
Northeast	29,850	87,680	118,537	120,411	112,355	109,798
Midwest	162,851	152,877	183,578	223,281	244,014	202,787
South	198,390	193,375	197,171	208,109	201,611	185,315
West	119,578	126,900	122,332	144,738	114,166	156,375
Census division¹						
New England	12,323	11,297	18,198	17,732	9,140	8,928
Middle Atlantic	17,527	76,383	100,339	102,679	103,215	100,870
East North Central	97,931	85,233	124,433	130,864	143,315	120,455
West North Central	64,920	67,644	59,145	92,417	100,699	82,332
South Atlantic	104,653	87,046	94,324	99,129	94,070	73,418
East South Central	60,443	58,900	53,052	58,764	56,448	60,154
West South Central	33,294	47,429	49,795	50,216	51,093	51,743
Mountain	60,508	62,066	68,907	73,392	39,242	70,538
Pacific	59,070	64,834	53,425	71,346	74,924	85,837
State or jurisdiction						
Alabama	16,563	15,079	15,614	16,071	13,214	15,798
Alaska	‡	‡	‡	‡	620	512
Arizona	29,211	32,167	32,196	34,892	‡	32,715
Arkansas	3,755	3,575	2,974	1,371	1,588	2,795
California	51,492	47,160	46,373	49,613	48,633	42,047
Colorado	7,916	7,349	11,035	11,125	9,974	9,644
Connecticut	6,396	5,660	11,230	11,544	2,361	2,130
Delaware	538	523	652	559	736	636
District of Columbia	1,608	1,699	2,113	2,622	2,394	2,157
Florida	12,731	12,562	15,317	14,455	10,488	13,991
Georgia	‡	‡	‡	‡	‡	12,127
Hawaii	658	477	526	596	461	512
Idaho	114	620	656	697	657	691
Illinois	17,030	1,550	8,393	7,453	4,783	4,198

Continued. See notes at end of table.

Table 8.6. Individuals with diagnosis of attention deficit/hyperactivity disorder, by Census region, Census division, and state or jurisdiction: Number, 2015–2020 (continued)

<i>Census region, Census division, and state or jurisdiction</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Indiana	26,262	30,131	28,456	27,616	27,829	27,231
Iowa	19,859	19,460	7,373	34,304	41,398	7,147
Kansas	‡	‡	‡	‡	‡	15,678
Kentucky	22,807	21,199	20,688	20,670	20,600	19,408
Louisiana	3,749	2,324	2,729	2,456	2,120	1,879
Maine	‡	‡	795	‡	‡	‡
Maryland	27,608	29,601	31,481	33,333	34,894	‡
Massachusetts	2,086	541	882	753	728	481
Michigan	‡	‡	21,955	29,330	30,749	28,347
Minnesota	34,335	37,537	40,416	46,677	47,028	48,532
Mississippi	14,049	15,771	13,009	17,071	17,706	20,989
Missouri	6,291	6,779	7,779	7,828	8,517	7,338
Montana	5,164	4,033	5,594	7,909	7,370	7,241
Nebraska	1,309	1,171	1,221	1,383	1,272	1,203
Nevada	1,011	647	647	572	639	669
New Hampshire	‡	‡	‡	‡	‡	‡
New Jersey	14,165	4,408	21,691	25,880	25,891	25,562
New Mexico	9,169	9,783	11,305	10,875	13,674	13,116
New York ²	3,362	3,622	3,503	3,720	3,842	3,779
North Carolina	26,565	6,897	8,993	6,325	6,133	5,430
North Dakota	1,813	1,435	1,133	982	1,321	1,308
Ohio	50,933	49,789	62,445	62,844	76,029	56,771
Oklahoma	3,058	4,224	4,347	4,448	4,400	4,459
Oregon	3,555	12,168	1	12,383	7,693	16,169
Pennsylvania	‡	68,353	75,145	73,079	73,482	71,529
Puerto Rico	65	64	374	337	404	192

Continued. See notes at end of table.

Table 8.6. Individuals with diagnosis of attention deficit/hyperactivity disorder, by Census region, Census division, and state or jurisdiction: Number, 2015–2020 (continued)

<i>Census region, Census division, and state or jurisdiction</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Rhode Island	1,962	2,774	2,907	2,971	3,568	3,724
South Carolina	15,027	15,024	15,020	15,295	16,842	16,762
South Dakota	1,313	1,262	1,223	1,243	1,163	1,126
Tennessee ²	7,024	6,851	3,741	4,952	4,928	3,959
Texas	22,732	37,306	39,745	41,941	42,985	42,610
Utah	6,210	5,875	5,977	5,930	5,541	5,139
Vermont	1,879	2,322	2,384	2,464	2,483	2,593
Virginia	20,576	20,740	20,748	21,641	22,583	22,315
Washington	3,365	5,029	6,525	8,754	17,517	26,597
West Virginia	‡	‡	‡	4,899	‡	‡
Wisconsin	3,706	3,763	3,184	3,621	3,925	3,908
Wyoming	1,713	1,592	1,497	1,392	1,387	1,323
Other jurisdictions ³	3	14	7	12	42	28

‡ No data submitted

¹ The Commonwealth of the Northern Mariana Islands, the Republic of Palau, and the Commonwealth of Puerto Rico are not included in any Census region or division.

² New York and Tennessee report only a subset of the total individuals served through their SMHAs.

³ Other jurisdictions include the Commonwealth of the Northern Mariana Islands and the Republic of Palau.

NOTE: This table refers to all mental health diagnoses, which is up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 8.7. Individuals with diagnosis of bipolar disorders, by Census region, Census division, and state or jurisdiction: Number, 2015–2020

<i>Census region, Census division, and state or jurisdiction</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals with diagnosis of bipolar disorders	855,430	747,805	724,193	734,667	704,177	713,775
Census region¹						
Northeast	63,702	110,538	135,679	113,593	111,658	109,860
Midwest	196,543	158,783	169,536	191,783	195,820	180,165
South	363,198	278,714	268,298	269,324	259,268	266,065
West	231,976	199,746	150,535	159,727	137,128	157,427
Census division¹						
New England	27,288	17,892	21,335	15,965	15,616	14,367
Middle Atlantic	36,414	92,646	114,344	97,628	96,042	95,493
East North Central	124,530	85,110	113,232	123,770	133,065	113,388
West North Central	72,013	73,673	56,304	68,013	62,755	66,777
South Atlantic	181,516	121,068	125,598	127,016	117,198	122,818
East South Central	88,411	62,956	49,074	51,208	47,804	48,780
West South Central	93,271	94,690	93,626	91,100	94,266	94,467
Mountain	93,424	74,311	73,479	73,162	43,823	70,209
Pacific	138,552	125,435	77,056	86,565	93,305	87,218
State or jurisdiction						
Alabama	18,557	12,680	12,532	12,188	10,822	12,058
Alaska	‡	‡	‡	‡	625	527
Arizona	40,967	35,077	29,026	31,374	‡	30,140
Arkansas	13,589	8,137	7,924	6,781	6,619	6,851
California	112,395	99,014	51,714	60,765	62,226	53,238
Colorado	14,234	12,526	17,557	17,471	15,839	14,828
Connecticut	13,279	11,369	9,712	9,046	8,305	7,356
Delaware	2,205	1,419	1,450	1,834	1,907	2,007
District of Columbia	4,421	4,429	5,664	6,750	6,375	5,839
Florida	39,892	32,627	29,115	29,437	23,363	32,579
Georgia	‡	‡	‡	‡	‡	25,989
Hawaii	1,427	1,655	1,572	1,400	1,158	784
Idaho	1,179	1,660	1,687	1,665	1,579	1,440
Illinois	30,723	2,507	11,999	10,717	7,435	6,111

Continued. See notes at end of table.

Table 8.7. Individuals with diagnosis of bipolar disorders, by Census region, Census division, and state or jurisdiction: Number, 2015–2020 (continued)

<i>Census region, Census division, and state or jurisdiction</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Indiana	19,560	15,488	15,036	14,419	14,281	13,981
Iowa	13,498	23,501	11,537	16,977	18,500	6,501
Kansas	‡	‡	‡	‡	‡	15,838
Kentucky	30,853	23,087	13,616	13,793	13,114	12,056
Louisiana	9,606	4,742	4,577	3,858	3,407	3,228
Maine	‡	‡	4,246	‡	‡	‡
Maryland	31,771	24,814	26,024	27,614	28,830	‡
Massachusetts	8,671	1,710	2,352	1,930	1,756	1,501
Michigan	‡	‡	27,954	35,786	36,497	32,606
Minnesota	30,328	27,502	21,819	29,132	22,374	24,968
Mississippi	11,502	10,985	7,814	10,398	10,527	12,179
Missouri	19,231	15,713	16,097	15,299	15,630	13,219
Montana	6,954	5,038	4,990	5,854	6,697	6,423
Nebraska	5,198	3,890	4,091	3,998	3,771	3,828
Nevada	6,901	1,162	1,166	1,043	996	1,123
New Hampshire	‡	‡	‡	‡	‡	‡
New Jersey	27,725	638	32,789	33,131	32,535	31,660
New Mexico	12,885	12,115	12,942	9,593	12,516	9,932
New York ²	8,689	10,653	7,300	7,066	6,730	6,359
North Carolina	58,037	23,681	30,220	19,582	20,044	19,466
North Dakota	1,856	1,482	1,273	1,107	1,039	1,073
Ohio	62,299	56,299	51,921	56,634	68,642	54,712
Oklahoma	14,002	12,895	11,881	11,313	10,895	10,594
Oregon	11,150	11,025	11,061	11,584	13,083	12,514
Pennsylvania	‡	81,355	74,255	57,431	56,777	57,474
Puerto Rico	4	4	130	235	285	226

Continued. See notes at end of table.

Table 8.7. Individuals with diagnosis of bipolar disorders, by Census region, Census division, and state or jurisdiction: Number, 2015–2020 (continued)

<i>Census region, Census division, and state or jurisdiction</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Rhode Island	3,699	3,548	3,672	3,607	4,144	4,176
South Carolina	13,949	11,526	10,863	11,405	12,413	12,157
South Dakota	1,902	1,585	1,487	1,500	1,441	1,350
Tennessee ²	27,499	16,204	15,112	14,829	13,341	12,487
Texas	56,074	68,916	69,244	69,148	73,345	73,794
Utah	7,816	4,559	4,404	4,490	4,477	4,652
Vermont	1,639	1,265	1,353	1,382	1,411	1,334
Virginia	31,241	22,572	22,262	23,456	24,266	24,781
Washington	13,580	13,741	12,709	12,816	16,213	20,155
West Virginia	‡	‡	‡	6,938	‡	‡
Wisconsin	11,948	10,816	6,322	6,214	6,210	5,978
Wyoming	2,488	2,174	1,707	1,672	1,719	1,671
Other jurisdictions ³	7	20	15	5	18	32

‡ No data submitted

¹ The Commonwealth of the Northern Mariana Islands, the Republic of Palau, and the Commonwealth of Puerto Rico are not included in any Census region or division.

² New York and Tennessee report only a subset of the total individuals served through their SMHAs.

³ Other jurisdictions include the Commonwealth of the Northern Mariana Islands and the Republic of Palau.

NOTE: This table refers to all mental health diagnoses, which is up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 8.8. Individuals with diagnosis of primary depressive disorders, by Census region, Census division, and state or jurisdiction: Number, 2015–2020

<i>Census region, Census division, and state or jurisdiction</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals with diagnosis of depressive disorders	1,419,968	1,429,859	1,597,103	1,756,277	1,824,785	1,834,097
Census region¹						
Northeast	113,667	183,937	285,242	274,882	273,362	271,137
Midwest	373,003	361,627	401,871	495,746	548,234	497,002
South	528,881	502,258	525,479	550,919	564,185	561,091
West	404,390	381,998	384,023	433,863	438,092	504,229
Census division¹						
New England	36,596	37,035	48,540	41,192	36,606	34,444
Middle Atlantic	77,071	146,902	236,702	233,690	236,756	236,693
East North Central	212,521	170,042	240,316	276,524	314,714	263,057
West North Central	160,482	191,585	161,555	219,222	233,520	233,945
South Atlantic	272,348	224,209	244,119	258,864	252,945	258,095
East South Central	137,045	128,340	121,335	129,561	143,449	131,136
West South Central	119,488	149,709	160,025	162,494	167,791	171,860
Mountain	145,714	142,583	167,254	173,863	132,078	197,279
Pacific	258,676	239,415	216,769	260,000	306,014	306,950
State or jurisdiction						
Alabama	30,670	30,570	29,659	29,521	43,137	31,105
Alaska	‡	‡	‡	‡	2,691	2,613
Arizona	41,244	46,706	45,783	52,898	‡	71,665
Arkansas	18,984	22,164	22,195	20,001	19,301	19,271
California	199,702	173,986	146,360	183,683	200,795	182,518
Colorado	35,070	31,559	45,507	46,695	41,846	39,422
Connecticut	20,105	20,285	22,527	22,976	15,972	14,395
Delaware	2,072	1,418	1,606	2,283	2,985	3,383
District of Columbia	11,197	10,840	10,914	13,662	13,381	13,701
Florida	53,391	55,140	55,032	57,163	52,772	76,895
Georgia	‡	‡	‡	‡	‡	45,549
Hawaii	2,421	2,546	2,449	2,170	2,134	1,352
Idaho	766	2,282	2,542	2,601	2,571	2,389
Illinois	45,237	3,065	22,522	20,906	13,345	11,329

Continued. See notes at end of table.

Table 8.8. Individuals with diagnosis of primary depressive disorders, by Census region, Census division, and state or jurisdiction: Number, 2015–2020 (continued)

<i>Census region, Census division, and state or jurisdiction</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Indiana	47,319	46,103	48,194	48,787	50,195	51,612
Iowa	27,419	54,947	25,693	58,863	78,284	20,852
Kansas	‡	‡	‡	‡	‡	47,536
Kentucky	47,462	45,619	45,893	47,462	46,032	43,158
Louisiana	15,011	7,771	9,910	9,289	7,626	6,656
Maine	‡	‡	7,156	‡	‡	‡
Maryland	54,014	52,257	57,315	61,451	66,739	‡
Massachusetts	6,308	2,321	3,404	2,414	2,189	1,644
Michigan	‡	‡	47,208	66,185	71,111	61,284
Minnesota	94,482	100,686	97,879	121,450	113,168	126,039
Mississippi	22,294	21,171	17,868	24,488	25,560	29,870
Missouri	23,335	22,784	24,992	25,766	27,660	25,147
Montana	14,357	9,914	14,843	20,749	23,283	23,201
Nebraska	6,351	5,878	6,577	6,802	7,124	7,126
Nevada	7,829	2,246	2,385	2,183	2,054	2,395
New Hampshire	‡	‡	‡	‡	‡	‡
New Jersey	66,071	2,087	79,007	80,706	80,764	79,778
New Mexico	25,520	29,454	35,654	27,680	40,543	35,556
New York ²	11,000	13,745	11,727	11,861	11,753	11,289
North Carolina	87,236	41,804	54,745	39,032	41,679	41,645
North Dakota	4,972	3,763	2,892	2,493	3,422	3,624
Ohio	103,349	104,864	110,392	128,448	167,914	126,003
Oklahoma	22,847	30,838	32,951	34,584	36,134	37,554
Oregon	30,178	30,596	34,193	37,475	46,377	45,310
Pennsylvania	‡	131,070	145,968	141,123	144,239	145,626
Puerto Rico	17	17	468	848	876	601

Continued. See notes at end of table.

Table 8.8. Individuals with diagnosis of primary depressive disorders, by Census region, Census division, and state or jurisdiction: Number, 2015–2020 (continued)

<i>Census region, Census division, and state or jurisdiction</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Rhode Island	5,941	9,911	10,649	10,675	13,156	13,251
South Carolina	24,882	24,543	24,410	25,660	29,819	29,930
South Dakota	3,923	3,527	3,522	3,848	3,862	3,621
Tennessee ²	36,619	30,980	27,915	28,090	28,720	27,003
Texas	62,646	88,936	94,969	98,620	104,730	108,379
Utah	14,895	14,484	15,027	15,429	15,903	16,790
Vermont	4,242	4,518	4,804	5,127	5,289	5,154
Virginia	39,556	38,207	40,097	42,715	45,570	46,992
Washington	26,375	32,287	33,767	36,672	54,017	75,157
West Virginia	‡	‡	‡	16,898	‡	‡
Wisconsin	16,616	16,010	12,000	12,198	12,149	12,829
Wyoming	6,033	5,938	5,513	5,628	5,878	5,861
Other jurisdictions ³	10	22	20	19	36	37

‡ No data submitted

¹ The Commonwealth of the Northern Mariana Islands, the Republic of Palau, and the Commonwealth of Puerto Rico are not included in any Census region or division.

² New York and Tennessee report only a subset of the total individuals served through their SMHAs.

³ Other jurisdictions include the Commonwealth of the Northern Mariana Islands and the Republic of Palau.

NOTE: This table refers to all mental health diagnoses, which is up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 8.9. Individuals with diagnosis of primary schizophrenia and other psychotic disorders, by Census region, Census division, and state or jurisdiction: Number, 2015–2020

<i>Census region, Census division, and state or jurisdiction</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals with diagnosis of schizophrenia and other psychotic disorders	640,161	622,833	728,180	770,554	742,888	765,575
Census region¹						
Northeast	76,649	80,754	125,143	121,555	120,072	117,531
Midwest	122,593	112,195	152,522	170,607	174,492	165,760
South	231,053	228,992	240,744	246,243	234,628	261,910
West	209,844	200,786	209,380	231,331	212,285	218,854
Census division¹						
New England	25,995	20,204	22,697	20,707	20,555	19,296
Middle Atlantic	50,654	60,550	102,446	100,848	99,517	98,235
East North Central	74,967	60,906	105,045	113,583	118,140	105,884
West North Central	47,626	51,289	47,477	57,024	56,352	59,876
South Atlantic	120,280	104,920	116,527	120,789	111,275	133,092
East South Central	53,944	49,053	46,871	50,281	47,236	51,853
West South Central	56,829	75,019	77,346	75,173	76,117	76,965
Mountain	52,711	51,071	57,928	59,908	37,063	62,595
Pacific	157,133	149,715	151,452	171,423	175,222	156,259
State or jurisdiction						
Alabama	18,143	17,754	18,042	18,187	15,520	17,851
Alaska	‡	‡	‡	‡	940	648
Arizona	20,123	20,473	21,382	23,949	‡	27,491
Arkansas	8,624	8,943	9,508	7,890	7,235	8,543
California	126,499	116,143	116,847	137,633	137,727	117,846
Colorado	9,804	10,759	13,882	13,760	12,590	12,549
Connecticut	11,402	11,430	11,610	11,208	10,780	9,827
Delaware	1,928	1,299	1,238	1,819	1,943	2,083
District of Columbia	5,418	5,256	6,170	7,104	6,621	6,372
Florida	30,180	29,188	32,843	35,278	26,300	35,609
Georgia	‡	‡	‡	‡	‡	29,419
Hawaii	3,001	3,287	3,166	3,153	2,784	2,354
Idaho	799	1,909	2,045	2,071	1,999	1,993
Illinois	18,661	3,406	11,553	10,736	9,012	7,682

Continued. See notes at end of table.

Table 8.9. Individuals with diagnosis of primary schizophrenia and other psychotic disorders, by Census region, Census division, and state or jurisdiction: Number, 2015–2020 (continued)

<i>Census region, Census division, and state or jurisdiction</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Indiana	15,477	16,345	16,119	15,535	15,281	14,278
Iowa	6,909	11,490	6,378	13,254	14,200	7,155
Kansas	‡	‡	‡	‡	‡	8,746
Kentucky	12,521	11,495	11,935	12,214	11,955	11,274
Louisiana	8,132	5,086	5,154	4,805	4,459	3,934
Maine	‡	‡	1,342	‡	‡	‡
Maryland	15,498	15,870	16,326	16,785	16,968	‡
Massachusetts	11,621	3,013	3,951	3,567	3,452	3,156
Michigan	‡	‡	29,582	35,736	35,993	34,612
Minnesota	20,594	20,650	21,342	24,127	21,995	25,217
Mississippi	12,091	10,822	8,000	10,995	11,310	14,574
Missouri	14,233	13,986	14,530	14,403	14,748	13,252
Montana	3,906	2,877	3,299	3,983	4,036	4,003
Nebraska	2,369	2,121	2,361	2,442	2,567	2,671
Nevada	5,907	2,015	2,077	2,034	2,046	2,321
New Hampshire	‡	‡	‡	‡	‡	‡
New Jersey	30,550	158	40,494	40,021	39,098	37,502
New Mexico	5,554	6,458	8,587	7,264	9,416	7,082
New York ²	20,104	20,849	20,562	20,506	20,131	19,827
North Carolina	27,900	14,130	20,444	16,038	16,268	16,122
North Dakota	1,717	1,564	1,448	1,348	1,457	1,490
Ohio	29,908	30,286	38,926	42,146	48,110	39,576
Oklahoma	7,646	9,932	10,254	10,425	10,443	10,429
Oregon	10,843	11,677	12,962	12,772	13,380	13,150
Pennsylvania	‡	39,543	41,390	40,321	40,288	40,906
Puerto Rico	8	7	304	702	1,316	1,379

Continued. See notes at end of table.

Table 8.9. Individuals with diagnosis of primary schizophrenia and other psychotic disorders, by Census region, Census division, and state or jurisdiction: Number, 2015–2020 (continued)

<i>Census region, Census division, and state or jurisdiction</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Rhode Island	1,754	3,975	3,971	4,069	4,422	4,466
South Carolina	15,763	15,418	15,373	16,485	17,340	17,323
South Dakota	1,804	1,478	1,418	1,450	1,385	1,345
Tennessee ²	11,189	8,982	8,894	8,885	8,451	8,154
Texas	32,427	51,058	52,430	52,053	53,980	54,059
Utah	5,571	5,543	5,623	5,813	5,937	6,068
Vermont	1,218	1,786	1,823	1,863	1,901	1,847
Virginia	23,593	23,759	24,133	24,788	25,835	26,164
Washington	16,790	18,608	18,477	17,865	20,391	22,261
West Virginia	‡	‡	‡	2,492	‡	‡
Wisconsin	10,921	10,869	8,865	9,430	9,744	9,736
Wyoming	1,047	1,037	1,033	1,034	1,039	1,088
Other jurisdictions ³	14	99	87	116	95	141

‡ No data submitted

¹ The Commonwealth of the Northern Mariana Islands, the Republic of Palau, and the Commonwealth of Puerto Rico are not included in any Census region or division.

² New York and Tennessee report only a subset of the total individuals served through their SMHAs.

³ Other jurisdictions include the Commonwealth of the Northern Mariana Islands and the Republic of Palau.

NOTE: This table refers to all mental health diagnoses, which is up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 8.10. Individuals with co-occurring mental health and substance use disorders, by Census region, Census division, and state or jurisdiction: Number, 2015–2020

<i>Census region, Census division, and state or jurisdiction</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals with co-occurring mental health and substance use disorders¹	750,256	685,036	927,904	1,073,395	1,318,814	1,783,845
Census region²						
Northeast	104,013	85,924	105,008	126,209	132,335	388,399
Midwest	131,655	138,084	338,960	414,245	466,030	428,134
South	254,470	215,117	242,562	278,200	481,022	717,551
West	259,974	245,695	240,123	253,563	236,952	247,317
Census division²						
New England	52,026	36,106	34,630	43,486	48,732	69,467
Middle Atlantic	51,987	49,818	70,378	82,723	83,603	318,932
East North Central	42,057	43,318	246,038	277,745	318,050	285,908
West North Central	89,598	94,766	92,922	136,500	147,980	142,226
South Atlantic	141,423	99,408	120,550	147,112	298,313	451,200
East South Central	57,565	48,201	50,357	53,284	50,688	100,891
West South Central	55,482	67,508	71,655	77,804	132,021	165,460
Mountain	71,564	60,236	72,786	81,732	47,479	79,345
Pacific	188,410	185,459	167,337	171,831	189,473	167,972
State or jurisdiction						
Alabama	11,794	12,117	14,451	14,708	11,926	14,352
Alaska	‡	‡	‡	‡	10,078	9,230
Arizona	35,698	38,503	38,189	44,101	‡	29,155
Arkansas	852	1,452	1,317	962	686	4,656
California	147,563	133,851	108,284	113,754	110,290	85,997
Colorado	14,881	1,529	385	379	548	6,708
Connecticut	36,951	26,826	24,742	25,036	30,202	27,890
Delaware	353	3,807	3,571	5,610	8,160	8,393
District of Columbia	3,099	3,915	2,612	4,142	3,087	3,452
Florida	28,574	26,963	26,409	26,620	173,459	243,272
Georgia	‡	‡	‡	‡	‡	135,911
Hawaii	2,230	2,281	2,784	3,049	2,935	2,471
Idaho	522	2,225	2,359	2,386	2,257	2,141
Illinois	17,039	1,595	5,495	5,095	3,439	2,602

Continued. See notes at end of table.

Table 8.10. Individuals with co-occurring mental health and substance use disorders, by Census region, Census division, and state or jurisdiction: Number, 2015–2020 (continued)

<i>Census region, Census division, and state or jurisdiction</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Indiana	18,239	22,491	23,079	22,814	21,275	25,059
Iowa	1,345	1,209	755	18,606	22,724	11,846
Kansas	‡	‡	‡	‡	‡	‡
Kentucky	25,222	20,397	22,347	23,299	23,493	22,691
Louisiana	967	1,361	3,189	3,121	11,879	13,524
Maine	‡	‡	6	‡	‡	‡
Maryland	15,463	18,060	40,080	52,562	55,718	‡
Massachusetts	7,575	2,409	2,943	9,206	7,514	5,954
Michigan	‡	‡	97,533	163,392	187,283	171,530
Minnesota	40,787	44,999	46,763	80,291	66,436	71,429
Mississippi	‡	‡	‡	‡	‡	49,152
Missouri	30,060	33,036	33,518	31,183	32,123	30,483
Montana	211	3,935	9,467	18,506	22,280	23,826
Nebraska	9,166	8,340	5,494	81	20,155	21,421
Nevada	1,955	680	653	583	603	961
New Hampshire	‡	‡	‡	‡	‡	‡
New Jersey	29,384	3	25,159	25,064	24,782	24,421
New Mexico	7,315	10,731	15,324	9,190	15,193	9,944
New York ³	22,603	25,664	26,144	26,036	25,752	25,429
North Carolina	46,741	‡	‡	‡	91	104
North Dakota	5,430	4,317	3,586	3,424	3,731	4,141
Ohio	1,749	14,581	117,487	83,419	103,006	83,608
Oklahoma	7,028	8,668	8,688	9,500	48,305	73,299
Oregon	32,184	41,286	45,967	43,314	48,819	46,369
Pennsylvania	‡	24,151	19,075	31,623	33,069	269,082
Puerto Rico	133	131	1,218	1,148	2,383	2,391

Continued. See notes at end of table.

Table 8.10. Individuals with co-occurring mental health and substance use disorders, by Census region, Census division, and state or jurisdiction: Number, 2015–2020 (continued)

<i>Census region, Census division, and state or jurisdiction</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Rhode Island	4,322	6,871	6,939	9,244	11,016	11,185
South Carolina	11,396	10,963	10,910	12,911	15,482	15,756
South Dakota	2,810	2,865	2,806	2,915	2,811	2,906
Tennessee ³	20,549	15,687	13,559	15,277	15,269	14,696
Texas	46,635	56,027	58,461	64,221	71,151	73,981
Utah	5,017	637	4,676	4,872	4,977	5,096
Vermont	3,178	‡	‡	‡	‡	24,438
Virginia	35,797	35,700	36,968	40,188	42,316	44,312
Washington	6,433	8,041	10,302	11,714	17,351	23,905
West Virginia	‡	‡	‡	5,079	‡	‡
Wisconsin	5,030	4,651	2,444	3,025	3,047	3,109
Wyoming	5,965	1,996	1,733	1,715	1,621	1,514
Other jurisdictions ⁴	11	85	33	30	92	53

‡ No data submitted

¹ Total individuals with co-occurring mental health and substance use disorders who had at least one valid mental health diagnosis other than an alcohol- or substance-related disorder. Mental health diagnostic category definitions can be found in Appendix E.

² The Commonwealth of the Northern Mariana Islands, the Republic of Palau, and the Commonwealth of Puerto Rico are not included in any Census region or division.

³ New York and Tennessee report only a subset of the total individuals served through their SMHAs.

⁴ Other jurisdictions include the Commonwealth of the Northern Mariana Islands and the Republic of Palau.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 9.1a. Alabama – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	101,582	104,320	103,959	102,424	103,203	97,527
Mental health diagnosis¹						
Trauma- and stressor-related disorders	11,618	10,585	11,259	12,038	11,232	13,116
Anxiety disorders	15,723	13,406	13,682	14,134	14,903	16,620
Attention deficit/hyperactivity disorder	16,563	15,079	15,614	16,071	13,214	15,798
Mood disorders	48,923	42,985	41,916	41,495	53,680	42,841
Bipolar disorders	18,557	12,680	12,532	12,188	10,822	12,058
Depressive disorders	30,670	30,570	29,659	29,521	43,137	31,105
Personality disorders	6,673	2,935	2,559	2,362	2,195	2,135
Schizophrenia and other psychotic disorders	18,143	17,754	18,042	18,187	15,520	17,851
Alcohol or substance-related disorders	11,404	3,153	361	114	22	15
Other mental disorders ²	87,828	15,962	19,542	21,430	18,149	16,430
Total	101,305	94,533	98,315	101,695	102,998	97,473
Gender						
Male	47,601	48,854	48,896	48,528	48,628	45,706
Female	53,748	55,338	55,045	53,888	54,575	51,818
Total	101,349	104,192	103,941	102,416	103,203	97,524
Age³						
0 to 11 years	15,025	15,239	15,661	16,028	16,221	14,466
12 to 14 years	7,520	7,663	7,907	8,418	8,904	8,769
15 to 17 years	8,478	8,535	8,802	8,727	9,069	8,696
18 to 20 years	3,998	4,234	4,392	4,459	4,544	4,413
21 to 24 years	5,051	5,040	4,753	4,554	4,577	4,562
25 to 29 years	7,024	7,467	7,417	7,163	7,134	6,773
30 to 34 years	7,358	7,543	7,304	7,140	7,235	7,048
35 to 39 years	7,246	7,831	7,672	7,531	7,360	6,840
40 to 44 years	7,121	7,154	6,901	6,579	6,634	6,466
45 to 49 years	7,385	7,490	7,233	6,867	6,595	6,031
50 to 54 years	7,955	8,111	7,710	7,109	6,767	6,159
55 to 59 years	7,008	7,210	7,150	6,952	6,740	6,265
60 to 64 years	4,517	4,770	5,014	5,113	5,281	5,078
65 years and older	5,669	5,841	5,724	5,776	6,136	5,959
Total	101,355	104,128	103,640	102,416	103,197	97,525

Continued. See notes at end of table.

Table 9.1a. Alabama – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	306	289	294	294	313	310
Asian ⁴	178	190	220	212	213	216
Black or African American	36,647	38,863	39,054	38,046	39,025	36,600
Native Hawaiian or Other Pacific Islander	30	39	49	59	58	51
White	59,409	59,891	59,477	56,220	56,703	53,768
Other	3,862	4,356	4,848	6,265	6,890	6,581
Total	100,432	103,628	103,942	101,096	103,202	97,526
Ethnicity						
Not of Hispanic or Latino origin	99,178	101,322	101,142	99,689	100,136	93,160
Hispanic or Latino origin	1,868	2,664	2,764	2,727	3,067	4,367
Total	101,046	103,986	103,906	102,416	103,203	97,527
Specific Hispanic or Latino origin						
Mexican	739	1,302	1,150	1,099	1,153	1,046
Puerto Rican	213	223	264	367	364	296
Cuban	36	35	32	46	53	49
Other specific	863	1,078	1,298	1,198	1,479	2,957
Total	1,851	2,638	2,744	2,710	3,049	4,348
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	1,208	1,002	998	884
Not a veteran	<i>n/c</i>	<i>n/c</i>	62,341	54,256	55,108	51,018
Total	<i>n/c</i>	<i>n/c</i>	63,549	55,258	56,106	51,902

n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.1b. Alabama – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	11.5	11.2	11.5	11.8	10.9	13.5
Anxiety disorders	15.5	14.2	13.9	13.9	14.5	17.1
Attention deficit/hyperactivity disorder	16.3	16.0	15.9	15.8	12.8	16.2
Mood disorders	48.3	45.5	42.6	40.8	52.1	44.0
Bipolar disorders	18.3	13.4	12.7	12.0	10.5	12.4
Depressive disorders	30.3	32.3	30.2	29.0	41.9	31.9
Personality disorders	6.6	3.1	2.6	2.3	2.1	2.2
Schizophrenia and other psychotic disorders	17.9	18.8	18.4	17.9	15.1	18.3
Alcohol or substance-related disorders	11.3	3.3	†	†	†	†
Other mental disorders ²	86.7	16.9	19.9	21.1	17.6	16.9
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	47.0	46.9	47.0	47.4	47.1	46.9
Female	53.0	53.1	53.0	52.6	52.9	53.1
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	14.8	14.6	15.1	15.6	15.7	14.8
12 to 14 years	7.4	7.4	7.6	8.2	8.6	9.0
15 to 17 years	8.4	8.2	8.5	8.5	8.8	8.9
18 to 20 years	3.9	4.1	4.2	4.4	4.4	4.5
21 to 24 years	5.0	4.8	4.6	4.4	4.4	4.7
25 to 29 years	6.9	7.2	7.2	7.0	6.9	6.9
30 to 34 years	7.3	7.2	7.0	7.0	7.0	7.2
35 to 39 years	7.1	7.5	7.4	7.4	7.1	7.0
40 to 44 years	7.0	6.9	6.7	6.4	6.4	6.6
45 to 49 years	7.3	7.2	7.0	6.7	6.4	6.2
50 to 54 years	7.8	7.8	7.4	6.9	6.6	6.3
55 to 59 years	6.9	6.9	6.9	6.8	6.5	6.4
60 to 64 years	4.5	4.6	4.8	5.0	5.1	5.2
65 years and older	5.6	5.6	5.5	5.6	5.9	6.1
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.1b. Alabama – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	†	†	†	†	†	†
Asian ⁴	†	†	†	†	†	†
Black or African American	36.5	37.5	37.6	37.6	37.8	37.5
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	59.2	57.8	57.2	55.6	54.9	55.1
Other	3.8	4.2	4.7	6.2	6.7	6.7
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	98.2	97.4	97.3	97.3	97.0	95.5
Hispanic or Latino origin	1.8	2.6	2.7	2.7	3.0	4.5
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	39.9	49.4	41.9	40.6	37.8	24.1
Puerto Rican	11.5	8.5	9.6	13.5	11.9	6.8
Cuban	1.9	1.3	1.2	1.7	1.7	1.1
Other specific	46.6	40.9	47.3	44.2	48.5	68.0
Total	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	1.9	1.8	1.8	1.7
Not a veteran	<i>n/c</i>	<i>n/c</i>	98.1	98.2	98.2	98.3
Total	<i>n/c</i>	<i>n/c</i>	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals; n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.2a. Alaska – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	‡	‡	‡	‡	11,082	10,337
Mental health diagnosis¹						
Trauma- and stressor-related disorders	‡	‡	‡	‡	3,260	3,329
Anxiety disorders	‡	‡	‡	‡	1,574	1,521
Attention deficit/hyperactivity disorder	‡	‡	‡	‡	620	512
Mood disorders	‡	‡	‡	‡	3,309	3,137
Bipolar disorders	‡	‡	‡	‡	625	527
Depressive disorders	‡	‡	‡	‡	2,691	2,613
Personality disorders	‡	‡	‡	‡	135	97
Schizophrenia and other psychotic disorders	‡	‡	‡	‡	940	648
Alcohol or substance-related disorders	‡	‡	‡	‡	1,197	1,042
Other mental disorders ²	‡	‡	‡	‡	1,445	1,220
Total	‡	‡	‡	‡	10,850	10,202
Gender						
Male	‡	‡	‡	‡	4,878	4,474
Female	‡	‡	‡	‡	6,201	5,855
Total	‡	‡	‡	‡	11,079	10,329
Age³						
0 to 11 years	‡	‡	‡	‡	1,722	1,455
12 to 14 years	‡	‡	‡	‡	1,071	999
15 to 17 years	‡	‡	‡	‡	1,199	1,177
18 to 20 years	‡	‡	‡	‡	587	566
21 to 24 years	‡	‡	‡	‡	651	632
25 to 29 years	‡	‡	‡	‡	988	992
30 to 34 years	‡	‡	‡	‡	1,026	1,070
35 to 39 years	‡	‡	‡	‡	868	824
40 to 44 years	‡	‡	‡	‡	590	581
45 to 49 years	‡	‡	‡	‡	564	436
50 to 54 years	‡	‡	‡	‡	517	461
55 to 59 years	‡	‡	‡	‡	576	469
60 to 64 years	‡	‡	‡	‡	382	297
65 years and older	‡	‡	‡	‡	341	378
Total	‡	‡	‡	‡	11,082	10,337

Continued. See notes at end of table.

Table 9.2a. Alaska – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	‡	‡	‡	‡	2,890	2,896
Asian ⁴	‡	‡	‡	‡	144	126
Black or African American	‡	‡	‡	‡	316	327
Native Hawaiian or Other Pacific Islander	‡	‡	‡	‡	84	84
White	‡	‡	‡	‡	4,649	4,367
Other	‡	‡	‡	‡	2,841	2,341
Total	‡	‡	‡	‡	10,924	10,141
Ethnicity						
Not of Hispanic or Latino origin	‡	‡	‡	‡	10,047	9,297
Hispanic or Latino origin	‡	‡	‡	‡	696	632
Total	‡	‡	‡	‡	10,743	9,929
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	145	119
Puerto Rican	‡	‡	‡	‡	68	61
Cuban	‡	‡	‡	‡	15	11
Other specific	‡	‡	‡	‡	280	263
Total	‡	‡	‡	‡	508	454
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	‡	‡	246	242
Not a veteran	<i>n/c</i>	<i>n/c</i>	‡	‡	6,458	5,955
Total	<i>n/c</i>	<i>n/c</i>	‡	‡	6,704	6,197

‡ No data submitted; n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.2b. Alaska – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	‡	‡	‡	‡	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	‡	‡	‡	‡	30.0	32.6
Anxiety disorders	‡	‡	‡	‡	14.5	14.9
Attention deficit/hyperactivity disorder	‡	‡	‡	‡	5.7	5.0
Mood disorders	‡	‡	‡	‡	30.5	30.7
Bipolar disorders	‡	‡	‡	‡	5.8	5.2
Depressive disorders	‡	‡	‡	‡	24.8	25.6
Personality disorders	‡	‡	‡	‡	1.2	†
Schizophrenia and other psychotic disorders	‡	‡	‡	‡	8.7	6.4
Alcohol or substance-related disorders	‡	‡	‡	‡	11.0	10.2
Other mental disorders ²	‡	‡	‡	‡	13.3	12.0
Total	‡	‡	‡	‡	100.0	100.0
Gender						
Male	‡	‡	‡	‡	44.0	43.3
Female	‡	‡	‡	‡	56.0	56.7
Total	‡	‡	‡	‡	100.0	100.0
Age³						
0 to 11 years	‡	‡	‡	‡	15.5	14.1
12 to 14 years	‡	‡	‡	‡	9.7	9.7
15 to 17 years	‡	‡	‡	‡	10.8	11.4
18 to 20 years	‡	‡	‡	‡	5.3	5.5
21 to 24 years	‡	‡	‡	‡	5.9	6.1
25 to 29 years	‡	‡	‡	‡	8.9	9.6
30 to 34 years	‡	‡	‡	‡	9.3	10.4
35 to 39 years	‡	‡	‡	‡	7.8	8.0
40 to 44 years	‡	‡	‡	‡	5.3	5.6
45 to 49 years	‡	‡	‡	‡	5.1	4.2
50 to 54 years	‡	‡	‡	‡	4.7	4.5
55 to 59 years	‡	‡	‡	‡	5.2	4.5
60 to 64 years	‡	‡	‡	‡	3.4	2.9
65 years and older	‡	‡	‡	‡	3.1	3.7
Total	‡	‡	‡	‡	100.0	100.0

Continued. See notes at end of table.

Table 9.2b. Alaska – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	‡	‡	‡	‡	26.5	28.6
Asian ⁴	‡	‡	‡	‡	1.3	1.2
Black or African American	‡	‡	‡	‡	2.9	3.2
Native Hawaiian or Other Pacific Islander	‡	‡	‡	‡	†	†
White	‡	‡	‡	‡	42.6	43.1
Other	‡	‡	‡	‡	26.0	23.1
Total	‡	‡	‡	‡	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	‡	‡	‡	‡	93.5	93.6
Hispanic or Latino origin	‡	‡	‡	‡	6.5	6.4
Total	‡	‡	‡	‡	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	28.5	26.2
Puerto Rican	‡	‡	‡	‡	13.4	13.4
Cuban	‡	‡	‡	‡	3.0	2.4
Other specific	‡	‡	‡	‡	55.1	57.9
Total	‡	‡	‡	‡	100.0	100.0
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	‡	‡	3.7	3.9
Not a veteran	<i>n/c</i>	<i>n/c</i>	‡	‡	96.3	96.1
Total	<i>n/c</i>	<i>n/c</i>	‡	‡	100.0	100.0

† Less than 1 percent of individuals; ‡ No data submitted; n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.3a. Arizona – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	155,122	187,972	196,212	224,773	‡	384,334
Mental health diagnosis¹						
Trauma- and stressor-related disorders	38,779	40,497	21,792	23,938	‡	69,162
Anxiety disorders	38,314	47,724	49,419	58,436	‡	75,608
Attention deficit/hyperactivity disorder	29,211	32,167	32,196	34,892	‡	32,715
Mood disorders	80,829	80,656	73,718	82,807	‡	100,850
Bipolar disorders	40,967	35,077	29,026	31,374	‡	30,140
Depressive disorders	41,244	46,706	45,783	52,898	‡	71,665
Personality disorders	8,985	8,599	8,371	8,577	‡	3,430
Schizophrenia and other psychotic disorders	20,123	20,473	21,382	23,949	‡	27,491
Alcohol or substance-related disorders	894	2,693	97	59	‡	64,070
Other mental disorders ²	70,470	44,568	23,720	24,032	‡	77,060
Total	155,100	175,840	150,419	169,862	‡	350,300
Gender						
Male	73,212	88,531	93,786	108,015	‡	183,138
Female	81,910	99,441	102,417	116,758	‡	201,196
Total	155,122	187,972	196,203	224,773	‡	384,334
Age³						
0 to 11 years	26,839	37,380	42,808	47,296	‡	68,038
12 to 14 years	12,687	14,960	16,081	18,759	‡	27,569
15 to 17 years	12,090	14,598	15,582	18,146	‡	26,236
18 to 20 years	5,720	7,952	7,771	9,325	‡	16,634
21 to 24 years	7,492	9,179	10,014	11,774	‡	22,514
25 to 29 years	11,154	13,946	15,362	18,442	‡	34,633
30 to 34 years	11,602	14,022	14,868	17,629	‡	33,264
35 to 39 years	10,866	13,118	13,840	16,013	‡	29,330
40 to 44 years	10,639	11,843	11,809	13,314	‡	23,328
45 to 49 years	11,001	12,125	12,059	13,521	‡	21,705
50 to 54 years	12,252	13,172	12,476	13,715	‡	21,556
55 to 59 years	10,460	12,024	11,604	13,029	‡	21,828
60 to 64 years	6,900	7,870	7,731	8,944	‡	16,596
65 years and older	5,417	5,782	4,205	4,864	‡	21,100
Total	155,119	187,971	196,210	224,771	‡	384,331

Continued. See notes at end of table.

Table 9.3a. Arizona – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	5,474	7,242	6,166	6,620	‡	25,005
Asian ⁴	1,054	1,330	1,355	1,649	‡	3,311
Black or African American	10,648	13,071	14,193	17,456	‡	27,457
Native Hawaiian or Other Pacific Islander	378	433	468	580	‡	665
White	128,933	157,571	165,711	193,391	‡	192,467
Other	2,275	3,353	4,586	4,551	‡	7
Total	148,762	183,000	192,479	224,247	‡	248,912
Ethnicity						
Not of Hispanic or Latino origin	104,917	132,342	135,955	155,395	‡	--
Hispanic or Latino origin	43,736	50,421	56,459	68,771	‡	254,250
Total	148,653	182,763	192,414	224,166	‡	254,250
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	2,205	175	‡	1,953
Not a veteran	<i>n/c</i>	<i>n/c</i>	106,661	8,467	‡	76,977
Total	<i>n/c</i>	<i>n/c</i>	108,866	8,642	‡	78,930

-- Quantity is zero; ‡ No data submitted; n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.3b. Arizona – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	100.0	100.0	100.0	100.0	‡	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	25.0	23.0	14.5	14.1	‡	19.7
Anxiety disorders	24.7	27.1	32.9	34.4	‡	21.6
Attention deficit/hyperactivity disorder	18.8	18.3	21.4	20.5	‡	9.3
Mood disorders	52.1	45.9	49.0	48.7	‡	28.8
Bipolar disorders	26.4	19.9	19.3	18.5	‡	8.6
Depressive disorders	26.6	26.6	30.4	31.1	‡	20.5
Personality disorders	5.8	4.9	5.6	5.0	‡	†
Schizophrenia and other psychotic disorders	13.0	11.6	14.2	14.1	‡	7.8
Alcohol or substance-related disorders	†	1.5	†	†	‡	18.3
Other mental disorders ²	45.4	25.3	15.8	14.1	‡	22.0
Total	100.0	100.0	100.0	100.0	‡	100.0
Gender						
Male	47.2	47.1	47.8	48.1	‡	47.7
Female	52.8	52.9	52.2	51.9	‡	52.3
Total	100.0	100.0	100.0	100.0	‡	100.0
Age³						
0 to 11 years	17.3	19.9	21.8	21.0	‡	17.7
12 to 14 years	8.2	8.0	8.2	8.3	‡	7.2
15 to 17 years	7.8	7.8	7.9	8.1	‡	6.8
18 to 20 years	3.7	4.2	4.0	4.1	‡	4.3
21 to 24 years	4.8	4.9	5.1	5.2	‡	5.9
25 to 29 years	7.2	7.4	7.8	8.2	‡	9.0
30 to 34 years	7.5	7.5	7.6	7.8	‡	8.7
35 to 39 years	7.0	7.0	7.1	7.1	‡	7.6
40 to 44 years	6.9	6.3	6.0	5.9	‡	6.1
45 to 49 years	7.1	6.5	6.1	6.0	‡	5.6
50 to 54 years	7.9	7.0	6.4	6.1	‡	5.6
55 to 59 years	6.7	6.4	5.9	5.8	‡	5.7
60 to 64 years	4.4	4.2	3.9	4.0	‡	4.3
65 years and older	3.5	3.1	2.1	2.2	‡	5.5
Total	100.0	100.0	100.0	100.0	‡	100.0

Continued. See notes at end of table.

Table 9.3b. Arizona – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	3.7	4.0	3.2	3.0	‡	10.0
Asian ⁴	†	†	†	†	‡	1.3
Black or African American	7.2	7.1	7.4	7.8	‡	11.0
Native Hawaiian or Other Pacific Islander	†	†	†	†	‡	†
White	86.7	86.1	86.1	86.2	‡	77.3
Other	1.5	1.8	2.4	2.0	‡	†
Total	100.0	100.0	100.0	100.0	‡	100.0
Ethnicity						
Not of Hispanic or Latino origin	70.6	72.4	70.7	69.3	‡	--
Hispanic or Latino origin	29.4	27.6	29.3	30.7	‡	100.0
Total	100.0	100.0	100.0	100.0	‡	100.0
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	2.0	2.0	‡	2.5
Not a veteran	<i>n/c</i>	<i>n/c</i>	98.0	98.0	‡	97.5
Total	<i>n/c</i>	<i>n/c</i>	100.0	100.0	‡	100.0

† Less than 1 percent of individuals; -- Quantity is zero; ‡ No data submitted; *n/c* Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.4a. Arkansas – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	69,398	74,458	74,985	74,888	76,139	77,552
Mental health diagnosis¹						
Trauma- and stressor-related disorders	9,651	11,610	12,413	11,469	11,739	12,099
Anxiety disorders	7,721	9,601	10,588	8,839	9,171	9,586
Attention deficit/hyperactivity disorder	3,755	3,575	2,974	1,371	1,588	2,795
Mood disorders	31,640	29,677	29,499	26,168	25,443	26,122
Bipolar disorders	13,589	8,137	7,924	6,781	6,619	6,851
Depressive disorders	18,984	22,164	22,195	20,001	19,301	19,271
Personality disorders	273	475	612	566	541	390
Schizophrenia and other psychotic disorders	8,624	8,943	9,508	7,890	7,235	8,543
Alcohol or substance-related disorders	11	--	--	1	--	4,177
Other mental disorders ²	9,585	13,117	11,731	10,058	10,062	9,978
Total	65,555	67,965	69,236	59,602	60,109	73,611
Gender						
Male	32,721	35,054	35,107	35,099	36,015	37,319
Female	36,622	39,372	39,792	39,690	40,045	40,183
Total	69,343	74,426	74,899	74,789	76,060	77,502
Age³						
0 to 11 years	10,597	10,902	6,300	10,582	10,845	10,527
12 to 14 years	4,648	4,849	3,271	5,223	5,523	5,994
15 to 17 years	5,055	5,549	3,557	5,898	6,160	6,213
18 to 20 years	3,192	3,700	2,490	4,250	4,441	4,650
21 to 24 years	3,875	4,092	2,244	3,954	4,071	4,351
25 to 29 years	5,663	6,185	3,599	6,054	5,862	6,100
30 to 34 years	5,984	6,433	4,007	6,169	6,313	6,625
35 to 39 years	5,766	6,364	4,204	6,285	6,209	6,201
40 to 44 years	5,219	5,572	3,683	5,377	5,593	5,701
45 to 49 years	5,003	5,356	3,695	5,152	5,124	5,129
50 to 54 years	5,304	5,471	3,936	4,985	4,799	4,734
55 to 59 years	4,235	4,489	3,497	4,714	4,769	4,628
60 to 64 years	2,537	2,833	2,233	3,113	3,153	3,275
65 years and older	2,319	2,662	28,269	3,130	3,277	3,422
Total	69,397	74,457	74,985	74,886	76,139	77,550

Continued. See notes at end of table.

Table 9.4a. Arkansas – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	205	249	235	273	305	306
Asian ⁴	157	202	200	220	227	195
Black or African American	13,875	14,575	14,997	14,738	15,240	14,556
Native Hawaiian or Other Pacific Islander	66	67	61	66	87	87
White	43,962	46,997	49,323	47,261	45,856	45,215
Other	715	815	874	847	720	683
Total	58,980	62,905	65,690	63,405	62,435	61,042
Ethnicity						
Not of Hispanic or Latino origin	56,816	63,211	65,835	64,739	64,796	64,010
Hispanic or Latino origin	2,217	2,433	2,689	2,939	3,201	3,496
Total	59,033	65,644	68,524	67,678	67,997	67,506
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡
Not a veteran	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡
Total	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡

- - Quantity is zero; ‡ No data submitted; n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.4b. Arkansas – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	14.7	17.1	17.9	19.2	19.5	16.4
Anxiety disorders	11.8	14.1	15.3	14.8	15.3	13.0
Attention deficit/hyperactivity disorder	5.7	5.3	4.3	2.3	2.6	3.8
Mood disorders	48.3	43.7	42.6	43.9	42.3	35.5
Bipolar disorders	20.7	12.0	11.4	11.4	11.0	9.3
Depressive disorders	29.0	32.6	32.1	33.6	32.1	26.2
Personality disorders	†	†	†	†	†	†
psychotic disorders	13.2	13.2	13.7	13.2	12.0	11.6
Alcohol or substance-related disorders	†	--	--	†	--	5.7
Other mental disorders ²	14.6	19.3	16.9	16.9	16.7	13.6
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	47.2	47.1	46.9	46.9	47.4	48.2
Female	52.8	52.9	53.1	53.1	52.6	51.8
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	15.3	14.6	8.4	14.1	14.2	13.6
12 to 14 years	6.7	6.5	4.4	7.0	7.3	7.7
15 to 17 years	7.3	7.5	4.7	7.9	8.1	8.0
18 to 20 years	4.6	5.0	3.3	5.7	5.8	6.0
21 to 24 years	5.6	5.5	3.0	5.3	5.3	5.6
25 to 29 years	8.2	8.3	4.8	8.1	7.7	7.9
30 to 34 years	8.6	8.6	5.3	8.2	8.3	8.5
35 to 39 years	8.3	8.5	5.6	8.4	8.2	8.0
40 to 44 years	7.5	7.5	4.9	7.2	7.3	7.4
45 to 49 years	7.2	7.2	4.9	6.9	6.7	6.6
50 to 54 years	7.6	7.3	5.2	6.7	6.3	6.1
55 to 59 years	6.1	6.0	4.7	6.3	6.3	6.0
60 to 64 years	3.7	3.8	3.0	4.2	4.1	4.2
65 years and older	3.3	3.6	37.7	4.2	4.3	4.4
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.4b. Arkansas – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	†	†	†	†	†	†
Asian ⁴	†	†	†	†	†	†
Black or African American	23.5	23.2	22.8	23.2	24.4	23.8
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	74.5	74.7	75.1	74.5	73.4	74.1
Other	1.2	1.3	1.3	1.3	1.2	1.1
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	96.2	96.3	96.1	95.7	95.3	94.8
Hispanic or Latino origin	3.8	3.7	3.9	4.3	4.7	5.2
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡
Not a veteran	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡
Total	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡

† Less than 1 percent of individuals; - - Quantity is zero; ‡ No data submitted; *n/c* Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.5a. California – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	695,815	647,328	638,394	708,521	742,725	665,415
Mental health diagnosis¹						
Trauma- and stressor-related disorders	111,808	105,242	95,732	110,480	119,485	108,332
Anxiety disorders	82,469	81,057	89,222	109,661	114,960	102,964
Attention deficit/hyperactivity disorder	51,492	47,160	46,373	49,613	48,633	42,047
Mood disorders	310,003	271,064	196,962	243,089	261,433	235,052
Bipolar disorders	112,395	99,014	51,714	60,765	62,226	53,238
Depressive disorders	199,702	173,986	146,360	183,683	200,795	182,518
Personality disorders	23,815	21,992	14,559	12,659	10,512	8,024
Schizophrenia and other psychotic disorders	126,499	116,143	116,847	137,633	137,727	117,846
Alcohol or substance-related disorders	112,754	92,519	64,364	70,791	68,327	63,517
Other mental disorders ²	530,389	476,307	140,147	148,940	140,016	119,714
Total	691,155	642,186	577,266	673,734	698,176	626,774
Gender						
Male	367,499	347,320	339,302	370,630	385,482	341,739
Female	327,415	298,860	295,889	337,061	356,402	322,855
Total	694,914	646,180	635,191	707,691	741,884	664,594
Age³						
0 to 11 years	116,816	111,276	110,841	122,971	130,173	113,384
12 to 14 years	54,600	51,705	51,264	57,823	65,336	61,237
15 to 17 years	66,476	63,618	61,802	67,676	73,818	67,142
18 to 20 years	34,028	31,679	30,981	32,588	34,738	32,356
21 to 24 years	37,136	35,503	34,896	37,915	39,159	35,192
25 to 29 years	49,289	47,168	48,631	55,141	58,398	53,821
30 to 34 years	50,319	47,341	46,861	51,488	53,585	49,267
35 to 39 years	43,110	41,000	41,936	47,430	49,327	44,742
40 to 44 years	43,944	38,270	35,903	39,362	40,260	36,269
45 to 49 years	49,354	43,069	40,458	43,642	42,437	36,049
50 to 54 years	53,561	47,056	43,769	47,641	46,278	38,993
55 to 59 years	43,738	39,814	38,472	44,888	45,131	39,394
60 to 64 years	28,514	26,530	26,096	30,883	32,408	29,317
65 years and older	22,221	20,625	21,797	26,632	29,008	27,687
Total	693,106	644,654	633,707	706,080	740,056	664,850

Continued. See notes at end of table.

Table 9.5a. California – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	4,813	4,740	4,638	5,356	5,416	5,455
Asian ⁴	25,027	23,460	24,584	27,225	28,721	24,065
Black or African American	86,605	81,275	81,262	95,170	97,171	89,224
Native Hawaiian or Other Pacific Islander	1,909	2,099	2,463	3,261	4,496	2,145
White	213,081	211,125	201,216	215,584	215,046	189,344
Other	222,727	214,944	225,108	248,431	264,130	250,385
Total	554,162	537,643	539,271	595,027	614,980	560,618
Ethnicity						
Not of Hispanic or Latino origin	294,488	280,859	274,896	319,203	323,810	282,989
Hispanic or Latino origin	235,804	226,006	232,016	255,670	273,804	257,757
Total	530,292	506,865	506,912	574,873	597,614	540,746
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	456	429	397	429
Not a veteran	<i>n/c</i>	<i>n/c</i>	6,116	6,299	5,147	5,118
Total	<i>n/c</i>	<i>n/c</i>	6,572	6,728	5,544	5,547

‡ No data submitted; n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.5b. California – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	16.2	16.4	16.6	16.4	17.1	17.3
Anxiety disorders	11.9	12.6	15.5	16.3	16.5	16.4
Attention deficit/hyperactivity disorder	7.5	7.3	8.0	7.4	7.0	6.7
Mood disorders	44.9	42.2	34.1	36.1	37.4	37.5
Bipolar disorders	16.3	15.4	9.0	9.0	8.9	8.5
Depressive disorders	28.9	27.1	25.4	27.3	28.8	29.1
Personality disorders	3.4	3.4	2.5	1.9	1.5	1.3
Schizophrenia and other psychotic disorders	18.3	18.1	20.2	20.4	19.7	18.8
Alcohol or substance-related disorders	16.3	14.4	11.1	10.5	9.8	10.1
Other mental disorders ²	76.7	74.2	24.3	22.1	20.1	19.1
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	52.9	53.7	53.4	52.4	52.0	51.4
Female	47.1	46.3	46.6	47.6	48.0	48.6
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	16.9	17.3	17.5	17.4	17.6	17.1
12 to 14 years	7.9	8.0	8.1	8.2	8.8	9.2
15 to 17 years	9.6	9.9	9.8	9.6	10.0	10.1
18 to 20 years	4.9	4.9	4.9	4.6	4.7	4.9
21 to 24 years	5.4	5.5	5.5	5.4	5.3	5.3
25 to 29 years	7.1	7.3	7.7	7.8	7.9	8.1
30 to 34 years	7.3	7.3	7.4	7.3	7.2	7.4
35 to 39 years	6.2	6.4	6.6	6.7	6.7	6.7
40 to 44 years	6.3	5.9	5.7	5.6	5.4	5.5
45 to 49 years	7.1	6.7	6.4	6.2	5.7	5.4
50 to 54 years	7.7	7.3	6.9	6.7	6.3	5.9
55 to 59 years	6.3	6.2	6.1	6.4	6.1	5.9
60 to 64 years	4.1	4.1	4.1	4.4	4.4	4.4
65 years and older	3.2	3.2	3.4	3.8	3.9	4.2
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.5b. California – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	†	†	†	†	†	†
Asian ⁴	4.5	4.4	4.6	4.6	4.7	4.3
Black or African American	15.6	15.1	15.1	16.0	15.8	15.9
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	38.5	39.3	37.3	36.2	35.0	33.8
Other	40.2	40.0	41.7	41.8	42.9	44.7
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	55.5	55.4	54.2	55.5	54.2	52.3
Hispanic or Latino origin	44.5	44.6	45.8	44.5	45.8	47.7
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	6.9	6.4	7.2	7.7
Not a veteran	<i>n/c</i>	<i>n/c</i>	93.1	93.6	92.8	92.3
Total	<i>n/c</i>	<i>n/c</i>	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals; ‡ No data submitted; n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.6a. Colorado – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	130,537	124,011	147,250	150,547	137,279	127,969
Mental health diagnosis¹						
Trauma- and stressor-related disorders	24,329	24,783	41,317	46,293	42,209	40,976
Anxiety disorders	20,976	18,118	28,851	29,738	26,919	26,738
Attention deficit/hyperactivity disorder	7,916	7,349	11,035	11,125	9,974	9,644
Mood disorders	49,155	43,981	62,741	63,758	57,358	53,914
Bipolar disorders	14,234	12,526	17,557	17,471	15,839	14,828
Depressive disorders	35,070	31,559	45,507	46,695	41,846	39,422
Personality disorders	3,411	1,988	3,632	3,846	2,932	2,604
Schizophrenia and other psychotic disorders	9,804	10,759	13,882	13,760	12,590	12,549
Alcohol or substance-related disorders	6,932	9,828	17,063	17,431	14,500	13,709
Other mental disorders ²	25,398	16,247	18,349	18,157	15,111	12,944
Total	107,948	109,885	145,379	148,869	135,408	126,315
Gender						
Male	60,361	57,146	67,344	69,153	63,329	59,178
Female	70,176	66,865	79,906	81,394	73,950	68,791
Total	130,537	124,011	147,250	150,547	137,279	127,969
Age³						
0 to 11 years	16,678	14,237	18,618	19,142	16,027	13,909
12 to 14 years	10,582	9,296	12,025	12,188	10,909	10,237
15 to 17 years	11,514	10,456	13,543	13,673	12,498	11,857
18 to 20 years	5,781	5,560	6,682	7,203	7,036	6,731
21 to 24 years	7,901	7,649	8,580	8,662	7,790	7,516
25 to 29 years	11,993	12,007	13,984	14,519	12,969	12,223
30 to 34 years	12,401	12,003	13,905	13,940	12,915	12,208
35 to 39 years	10,349	10,285	12,591	13,006	11,812	11,130
40 to 44 years	9,371	8,809	9,617	9,955	9,276	8,883
45 to 49 years	8,819	8,575	9,364	9,363	8,438	7,677
50 to 54 years	9,242	8,669	9,394	8,947	7,990	7,367
55 to 59 years	7,335	7,362	8,347	8,404	7,924	7,124
60 to 64 years	4,417	4,741	5,500	5,829	5,551	5,246
65 years and older	4,154	4,361	5,099	5,716	6,144	5,861
Total	130,537	124,010	147,249	150,547	137,279	127,969

Continued. See notes at end of table.

Table 9.6a. Colorado – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	2,948	2,622	3,279	3,389	3,167	2,947
Asian ⁴	1,214	1,193	1,493	1,543	1,563	1,358
Black or African American	8,632	7,747	9,087	8,865	7,805	7,735
Native Hawaiian or Other Pacific Islander	342	340	422	426	376	438
White	100,665	96,917	112,418	103,474	100,884	92,238
Other	4,466	4,601	6,379	15,385	4,576	23,238
Total	118,267	113,420	133,078	133,082	118,371	127,954
Ethnicity						
Not of Hispanic or Latino origin	95,911	89,680	101,186	101,984	90,542	82,404
Hispanic or Latino origin	30,078	28,488	36,662	37,886	35,697	33,020
Total	125,989	118,168	137,848	139,870	126,239	115,424
Specific Hispanic or Latino origin						
Mexican	21,740	18,310	23,380	23,897	22,710	20,941
Puerto Rican	410	670	947	889	837	829
Cuban	128	304	474	400	419	376
Other specific	7,697	9,098	11,747	12,564	11,547	10,723
Total	29,975	28,382	36,548	37,750	35,513	32,869
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	3,124	3,181	3,128	2,709
Not a veteran	<i>n/c</i>	<i>n/c</i>	99,608	101,923	93,913	88,522
Total	<i>n/c</i>	<i>n/c</i>	102,732	105,104	97,041	91,231

n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.6b. Colorado – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	22.5	22.6	28.4	31.1	31.2	32.4
Anxiety disorders	19.4	16.5	19.8	20.0	19.9	21.2
Attention deficit/hyperactivity disorder	7.3	6.7	7.6	7.5	7.4	7.6
Mood disorders	45.5	40.0	43.2	42.8	42.4	42.7
Bipolar disorders	13.2	11.4	12.1	11.7	11.7	11.7
Depressive disorders	32.5	28.7	31.3	31.4	30.9	31.2
Personality disorders	3.2	1.8	2.5	2.6	2.2	2.1
Schizophrenia and other psychotic disorders	9.1	9.8	9.5	9.2	9.3	9.9
Alcohol or substance-related disorders	6.4	8.9	11.7	11.7	10.7	10.9
Other mental disorders ²	23.5	14.8	12.6	12.2	11.2	10.2
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	46.2	46.1	45.7	45.9	46.1	46.2
Female	53.8	53.9	54.3	54.1	53.9	53.8
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	12.8	11.5	12.6	12.7	11.7	10.9
12 to 14 years	8.1	7.5	8.2	8.1	7.9	8.0
15 to 17 years	8.8	8.4	9.2	9.1	9.1	9.3
18 to 20 years	4.4	4.5	4.5	4.8	5.1	5.3
21 to 24 years	6.1	6.2	5.8	5.8	5.7	5.9
25 to 29 years	9.2	9.7	9.5	9.6	9.4	9.6
30 to 34 years	9.5	9.7	9.4	9.3	9.4	9.5
35 to 39 years	7.9	8.3	8.6	8.6	8.6	8.7
40 to 44 years	7.2	7.1	6.5	6.6	6.8	6.9
45 to 49 years	6.8	6.9	6.4	6.2	6.1	6.0
50 to 54 years	7.1	7.0	6.4	5.9	5.8	5.8
55 to 59 years	5.6	5.9	5.7	5.6	5.8	5.6
60 to 64 years	3.4	3.8	3.7	3.9	4.0	4.1
65 years and older	3.2	3.5	3.5	3.8	4.5	4.6
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.6b. Colorado – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	2.5	2.3	2.5	2.5	2.7	2.3
Asian ⁴	1.0	1.1	1.1	1.2	1.3	1.1
Black or African American	7.3	6.8	6.8	6.7	6.6	6.0
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	85.1	85.4	84.5	77.8	85.2	72.1
Other	3.8	4.1	4.8	11.6	3.9	18.2
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	76.1	75.9	73.4	72.9	71.7	71.4
Hispanic or Latino origin	23.9	24.1	26.6	27.1	28.3	28.6
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	72.5	64.5	64.0	63.3	63.9	63.7
Puerto Rican	1.4	2.4	2.6	2.4	2.4	2.5
Cuban	†	1.1	1.3	1.1	1.2	1.1
Other specific	25.7	32.1	32.1	33.3	32.5	32.6
Total	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	3.0	3.0	3.2	3.0
Not a veteran	<i>n/c</i>	<i>n/c</i>	97.0	97.0	96.8	97.0
Total	<i>n/c</i>	<i>n/c</i>	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals; n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.7a. Connecticut – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	90,524	94,541	97,130	93,947	94,199	86,915
Mental health diagnosis¹						
Trauma- and stressor-related disorders	14,329	14,831	20,716	21,254	10,158	9,682
Anxiety disorders	10,099	13,154	16,612	17,611	11,008	10,148
Attention deficit/hyperactivity disorder	6,396	5,660	11,230	11,544	2,361	2,130
Mood disorders	32,846	31,308	31,977	31,653	24,014	21,540
Bipolar disorders	13,279	11,369	9,712	9,046	8,305	7,356
Depressive disorders	20,105	20,285	22,527	22,976	15,972	14,395
Personality disorders	4,269	3,149	3,126	3,516	2,726	2,057
Schizophrenia and other psychotic disorders	11,402	11,430	11,610	11,208	10,780	9,827
Alcohol or substance-related disorders	536	274	2,740	2,489	271	187
Other mental disorders ²	16,609	12,411	17,953	19,910	7,622	6,284
Total	62,569	64,000	78,543	77,885	45,729	41,625
Gender						
Male	46,134	48,903	50,586	49,096	49,277	45,319
Female	44,375	45,579	46,492	44,781	44,860	41,523
Total	90,509	94,482	97,078	93,877	94,137	86,842
Age³						
0 to 11 years	14,684	15,287	16,374	16,608	16,971	15,387
12 to 14 years	8,825	8,701	9,118	9,298	9,900	8,694
15 to 17 years	9,659	9,748	9,889	9,832	9,817	8,152
18 to 20 years	3,910	4,132	4,181	4,218	4,135	3,147
21 to 24 years	4,066	4,227	4,240	3,922	3,771	3,717
25 to 29 years	5,200	5,615	5,815	5,519	5,542	5,261
30 to 34 years	5,312	5,593	5,830	5,548	5,628	5,532
35 to 39 years	4,733	5,093	5,372	5,186	5,226	5,269
40 to 44 years	5,023	5,019	4,922	4,443	4,390	4,533
45 to 49 years	6,497	6,637	6,313	5,531	5,207	4,603
50 to 54 years	7,427	7,720	7,621	6,893	6,391	5,836
55 to 59 years	6,545	7,093	7,234	6,786	6,505	6,304
60 to 64 years	4,366	4,840	5,137	5,030	5,176	5,015
65 years and older	4,252	4,830	5,081	5,131	5,536	5,464
Total	90,499	94,535	97,127	93,945	94,195	86,914

Continued. See notes at end of table.

Table 9.7a. Connecticut – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	479	581	575	448	451	457
Asian ⁴	758	828	894	933	1,064	902
Black or African American	15,542	16,964	17,695	16,919	17,125	15,756
Native Hawaiian or Other Pacific Islander	238	220	228	230	230	266
White	54,746	57,434	58,801	55,336	55,144	50,560
Other	14,716	14,442	13,985	14,583	14,574	13,469
Total	86,479	90,469	92,178	88,449	88,588	81,410
Ethnicity						
Not of Hispanic or Latino origin	62,986	65,717	66,950	63,632	63,612	58,474
Hispanic or Latino origin	23,271	24,549	25,881	25,279	26,267	22,915
Total	86,257	90,266	92,831	88,911	89,879	81,389
Specific Hispanic or Latino origin						
Mexican	794	822	570	389	263	322
Puerto Rican	11,629	12,030	10,154	8,361	6,424	6,574
Cuban	147	162	170	110	93	92
Other specific	4,439	5,108	5,092	4,939	4,697	4,598
Total	17,009	18,122	15,986	13,799	11,477	11,586
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	‡	2,107	2,094	1,915
Not a veteran	<i>n/c</i>	<i>n/c</i>	‡	46,297	47,175	46,334
Total	<i>n/c</i>	<i>n/c</i>	‡	48,404	49,269	48,249

‡ No data submitted; n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.7b. Connecticut – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	22.9	23.2	26.4	27.3	22.2	23.3
Anxiety disorders	16.1	20.6	21.2	22.6	24.1	24.4
Attention deficit/hyperactivity disorder	10.2	8.8	14.3	14.8	5.2	5.1
Mood disorders	52.5	48.9	40.7	40.6	52.5	51.7
Bipolar disorders	21.2	17.8	12.4	11.6	18.2	17.7
Depressive disorders	32.1	31.7	28.7	29.5	34.9	34.6
Personality disorders	6.8	4.9	4.0	4.5	6.0	4.9
Schizophrenia and other psychotic disorders	18.2	17.9	14.8	14.4	23.6	23.6
Alcohol or substance-related disorders	†	†	3.5	3.2	†	†
Other mental disorders ²	26.5	19.4	22.9	25.6	16.7	15.1
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	51.0	51.8	52.1	52.3	52.3	52.2
Female	49.0	48.2	47.9	47.7	47.7	47.8
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	16.2	16.2	16.9	17.7	18.0	17.7
12 to 14 years	9.8	9.2	9.4	9.9	10.5	10.0
15 to 17 years	10.7	10.3	10.2	10.5	10.4	9.4
18 to 20 years	4.3	4.4	4.3	4.5	4.4	3.6
21 to 24 years	4.5	4.5	4.4	4.2	4.0	4.3
25 to 29 years	5.7	5.9	6.0	5.9	5.9	6.1
30 to 34 years	5.9	5.9	6.0	5.9	6.0	6.4
35 to 39 years	5.2	5.4	5.5	5.5	5.5	6.1
40 to 44 years	5.6	5.3	5.1	4.7	4.7	5.2
45 to 49 years	7.2	7.0	6.5	5.9	5.5	5.3
50 to 54 years	8.2	8.2	7.8	7.3	6.8	6.7
55 to 59 years	7.2	7.5	7.4	7.2	6.9	7.3
60 to 64 years	4.8	5.1	5.3	5.4	5.5	5.8
65 years and older	4.7	5.1	5.2	5.5	5.9	6.3
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.7b. Connecticut – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	†	†	†	†	†	†
Asian ⁴	†	†	†	1.1	1.2	1.1
Black or African American	18.0	18.8	19.2	19.1	19.3	19.4
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	63.3	63.5	63.8	62.6	62.2	62.1
Other	17.0	16.0	15.2	16.5	16.5	16.5
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	73.0	72.8	72.1	71.6	70.8	71.8
Hispanic or Latino origin	27.0	27.2	27.9	28.4	29.2	28.2
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	4.7	4.5	3.6	2.8	2.3	2.8
Puerto Rican	68.4	66.4	63.5	60.6	56.0	56.7
Cuban	†	†	1.1	†	†	†
Other specific	26.1	28.2	31.9	35.8	40.9	39.7
Total	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	‡	4.4	4.3	4.0
Not a veteran	<i>n/c</i>	<i>n/c</i>	‡	95.6	95.7	96.0
Total	<i>n/c</i>	<i>n/c</i>	‡	100.0	100.0	100.0

† Less than 1 percent of individuals; ‡ No data submitted; *n/c* Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.8a. Delaware – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	9,680	7,456	9,578	9,889	14,032	12,230
Mental health diagnosis¹						
Trauma- and stressor-related disorders	1,736	1,069	1,237	1,202	1,422	1,388
Anxiety disorders	787	498	670	829	1,172	1,356
Attention deficit/hyperactivity disorder	538	523	652	559	736	636
Mood disorders	4,037	2,803	3,022	4,013	4,784	5,227
Bipolar disorders	2,205	1,419	1,450	1,834	1,907	2,007
Depressive disorders	2,072	1,418	1,606	2,283	2,985	3,383
Personality disorders	253	274	120	316	346	368
Schizophrenia and other psychotic disorders	1,928	1,299	1,238	1,819	1,943	2,083
Alcohol or substance-related disorders	2,077	2,399	2,532	3,568	3,556	3,728
Other mental disorders ²	1,784	1,493	1,276	1,556	1,968	2,104
Total	8,590	6,479	7,625	8,802	10,467	10,559
Gender						
Male	5,252	4,017	5,042	5,334	7,424	6,585
Female	4,408	3,370	4,343	4,399	6,495	5,528
Total	9,660	7,387	9,385	9,733	13,919	12,113
Age³						
0 to 11 years	882	410	879	313	1,292	337
12 to 14 years	855	416	835	397	1,246	350
15 to 17 years	1,124	487	839	456	1,196	452
18 to 20 years	253	332	386	478	671	661
21 to 24 years	479	604	647	785	863	921
25 to 29 years	731	853	959	1,178	1,362	1,430
30 to 34 years	703	714	860	1,025	1,239	1,360
35 to 39 years	618	639	800	1,062	1,238	1,238
40 to 44 years	575	520	611	791	964	1,079
45 to 49 years	697	567	639	814	903	945
50 to 54 years	858	655	678	795	841	953
55 to 59 years	765	570	618	768	945	993
60 to 64 years	551	308	363	498	612	740
65 years and older	589	378	459	526	660	770
Total	9,680	7,453	9,573	9,886	14,032	12,229

Continued. See notes at end of table.

Table 9.8a. Delaware – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	40	50	61	69	90	81
Asian ⁴	69	53	55	65	101	89
Black or African American	3,240	2,278	3,046	3,140	4,486	3,740
Native Hawaiian or Other Pacific Islander	53	27	38	59	65	64
White	5,986	4,751	5,999	6,105	7,944	7,068
Other	65	84	93	129	295	214
Total	9,453	7,243	9,292	9,567	12,981	11,256
Ethnicity						
Not of Hispanic or Latino origin	8,463	6,679	8,442	8,644	11,289	9,796
Hispanic or Latino origin	715	401	603	566	966	646
Total	9,178	7,080	9,045	9,210	12,255	10,442
Specific Hispanic or Latino origin						
Mexican	89	49	33	58	67	64
Puerto Rican	89	60	72	120	168	161
Cuban	13	2	3	5	15	10
Other specific	114	85	105	150	180	204
Total	305	196	213	333	430	439
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	‡	292	361	400
Not a veteran	<i>n/c</i>	<i>n/c</i>	‡	6,182	7,675	8,189
Total	<i>n/c</i>	<i>n/c</i>	‡	6,474	8,036	8,589

‡ No data submitted; n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.8b. Delaware – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	20.2	16.5	16.2	13.7	13.6	13.1
Anxiety disorders	9.2	7.7	8.8	9.4	11.2	12.8
Attention deficit/hyperactivity disorder	6.3	8.1	8.6	6.4	7.0	6.0
Mood disorders	47.0	43.3	39.6	45.6	45.7	49.5
Bipolar disorders	25.7	21.9	19.0	20.8	18.2	19.0
Depressive disorders	24.1	21.9	21.1	25.9	28.5	32.0
Personality disorders	2.9	4.2	1.6	3.6	3.3	3.5
Schizophrenia and other psychotic disorders	22.4	20.0	16.2	20.7	18.6	19.7
Alcohol or substance-related disorders	24.2	37.0	33.2	40.5	34.0	35.3
Other mental disorders ²	20.8	23.0	16.7	17.7	18.8	19.9
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	54.4	54.4	53.7	54.8	53.3	54.4
Female	45.6	45.6	46.3	45.2	46.7	45.6
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	9.1	5.5	9.2	3.2	9.2	2.8
12 to 14 years	8.8	5.6	8.7	4.0	8.9	2.9
15 to 17 years	11.6	6.5	8.8	4.6	8.5	3.7
18 to 20 years	2.6	4.5	4.0	4.8	4.8	5.4
21 to 24 years	4.9	8.1	6.8	7.9	6.2	7.5
25 to 29 years	7.6	11.4	10.0	11.9	9.7	11.7
30 to 34 years	7.3	9.6	9.0	10.4	8.8	11.1
35 to 39 years	6.4	8.6	8.4	10.7	8.8	10.1
40 to 44 years	5.9	7.0	6.4	8.0	6.9	8.8
45 to 49 years	7.2	7.6	6.7	8.2	6.4	7.7
50 to 54 years	8.9	8.8	7.1	8.0	6.0	7.8
55 to 59 years	7.9	7.6	6.5	7.8	6.7	8.1
60 to 64 years	5.7	4.1	3.8	5.0	4.4	6.1
65 years and older	6.1	5.1	4.8	5.3	4.7	6.3
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.8b. Delaware – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	†	†	†	†	†	†
Asian ⁴	†	†	†	†	†	†
Black or African American	34.3	31.5	32.8	32.8	34.6	33.2
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	63.3	65.6	64.6	63.8	61.2	62.8
Other	†	1.2	1.0	1.3	2.3	1.9
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	92.2	94.3	93.3	93.9	92.1	93.8
Hispanic or Latino origin	7.8	5.7	6.7	6.1	7.9	6.2
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	29.2	25.0	15.5	17.4	15.6	14.6
Puerto Rican	29.2	30.6	33.8	36.0	39.1	36.7
Cuban	4.3	1.0	1.4	1.5	3.5	2.3
Other specific	37.4	43.4	49.3	45.0	41.9	46.5
Total	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	‡	4.5	4.5	4.7
Not a veteran	<i>n/c</i>	<i>n/c</i>	‡	95.5	95.5	95.3
Total	<i>n/c</i>	<i>n/c</i>	‡	100.0	100.0	100.0

† Less than 1 percent of individuals; ‡ No data submitted; *n/c* Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

**Table 9.9a. District of Columbia – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Number, 2015–2020**

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	28,531	35,028	36,668	45,560	45,189	51,073
Mental health diagnosis¹						
Trauma- and stressor-related disorders	2,555	2,546	3,173	3,856	3,764	3,871
Anxiety disorders	874	834	1,602	1,830	1,648	1,525
Attention deficit/hyperactivity disorder	1,608	1,699	2,113	2,622	2,394	2,157
Mood disorders	15,497	15,178	16,113	19,918	19,601	19,383
Bipolar disorders	4,421	4,429	5,664	6,750	6,375	5,839
Depressive disorders	11,197	10,840	10,914	13,662	13,381	13,701
Personality disorders	56	47	69	83	65	73
Schizophrenia and other psychotic disorders	5,418	5,256	6,170	7,104	6,621	6,372
Alcohol or substance-related disorders	1,208	1,061	511	735	1,622	1,369
Other mental disorders ²	1,191	1,259	2,777	3,419	2,929	2,529
Total	26,420	26,197	29,625	35,830	36,588	35,161
Gender						
Male	14,986	18,306	18,923	23,509	23,342	26,357
Female	13,466	16,629	17,626	21,925	21,746	24,600
Total	28,452	34,935	36,549	45,434	45,088	50,957
Age³						
0 to 11 years	1,997	2,581	2,334	2,746	1,586	1,503
12 to 14 years	1,462	1,564	1,378	1,706	1,529	1,759
15 to 17 years	1,546	1,900	1,798	2,158	1,800	1,847
18 to 20 years	1,166	1,581	1,534	2,032	1,994	2,361
21 to 24 years	1,647	2,063	2,188	2,635	2,429	2,897
25 to 29 years	2,295	3,157	3,440	4,292	3,944	4,471
30 to 34 years	2,229	2,866	3,095	3,877	3,818	4,680
35 to 39 years	2,040	2,635	2,936	3,726	3,792	4,278
40 to 44 years	1,830	2,183	2,319	3,019	3,294	3,867
45 to 49 years	2,802	2,961	2,995	3,385	3,205	3,383
50 to 54 years	3,526	4,004	4,025	4,684	4,755	4,873
55 to 59 years	2,952	3,557	4,009	4,978	5,416	5,943
60 to 64 years	1,835	2,338	2,710	3,574	4,185	4,870
65 years and older	1,204	1,638	1,907	2,748	3,442	4,341
Total	28,531	35,028	36,668	45,560	45,189	51,073

Continued. See notes at end of table.

**Table 9.9a. District of Columbia – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Number, 2015–2020 (continued)**

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	61	90	76	112	116	114
Asian ⁴	98	130	117	156	158	157
Black or African American	25,415	31,261	32,911	40,747	40,793	45,449
Native Hawaiian or Other Pacific Islander	4	2	3	3	3	2
White	1,123	1,366	1,312	1,641	1,647	1,666
Other	28	59	85	114	124	114
Total	26,729	32,908	34,504	42,773	42,841	47,502
Ethnicity						
Not of Hispanic or Latino origin	2,574	9,834	12,968	17,374	18,828	21,359
Hispanic or Latino origin	376	686	747	971	1,039	1,159
Total	2,950	10,520	13,715	18,345	19,867	22,518
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡
Not a veteran	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡
Total	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡

‡ No data submitted; n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

**Table 9.9b. District of Columbia – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Column percent distribution, 2015–2020**

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	9.7	9.7	10.7	10.8	10.3	11.0
Anxiety disorders	3.3	3.2	5.4	5.1	4.5	4.3
Attention deficit/hyperactivity disorder	6.1	6.5	7.1	7.3	6.5	6.1
Mood disorders	58.7	57.9	54.4	55.6	53.6	55.1
Bipolar disorders	16.7	16.9	19.1	18.8	17.4	16.6
Depressive disorders	42.4	41.4	36.8	38.1	36.6	39.0
Personality disorders	†	†	†	†	†	†
Schizophrenia and other psychotic disorders	20.5	20.1	20.8	19.8	18.1	18.1
Alcohol or substance-related disorders	4.6	4.1	1.7	2.1	4.4	3.9
Other mental disorders ²	4.5	4.8	9.4	9.5	8.0	7.2
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	52.7	52.4	51.8	51.7	51.8	51.7
Female	47.3	47.6	48.2	48.3	48.2	48.3
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	7.0	7.4	6.4	6.0	3.5	2.9
12 to 14 years	5.1	4.5	3.8	3.7	3.4	3.4
15 to 17 years	5.4	5.4	4.9	4.7	4.0	3.6
18 to 20 years	4.1	4.5	4.2	4.5	4.4	4.6
21 to 24 years	5.8	5.9	6.0	5.8	5.4	5.7
25 to 29 years	8.0	9.0	9.4	9.4	8.7	8.8
30 to 34 years	7.8	8.2	8.4	8.5	8.4	9.2
35 to 39 years	7.2	7.5	8.0	8.2	8.4	8.4
40 to 44 years	6.4	6.2	6.3	6.6	7.3	7.6
45 to 49 years	9.8	8.5	8.2	7.4	7.1	6.6
50 to 54 years	12.4	11.4	11.0	10.3	10.5	9.5
55 to 59 years	10.3	10.2	10.9	10.9	12.0	11.6
60 to 64 years	6.4	6.7	7.4	7.8	9.3	9.5
65 years and older	4.2	4.7	5.2	6.0	7.6	8.5
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

**Table 9.9b. District of Columbia – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Column percent distribution, 2015–2020 (continued)**

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	†	†	†	†	†	†
Asian ⁴	†	†	†	†	†	†
Black or African American	95.1	95.0	95.4	95.3	95.2	95.7
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	4.2	4.2	3.8	3.8	3.8	3.5
Other	†	†	†	†	†	†
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	87.3	93.5	94.6	94.7	94.8	94.9
Hispanic or Latino origin	12.7	6.5	5.4	5.3	5.2	5.1
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡
Not a veteran	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡
Total	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡

† Less than 1 percent of individuals; ‡ No data submitted; *n/c* Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.10a. Florida – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	230,652	214,257	213,758	218,368	204,076	282,228
Mental health diagnosis¹						
Trauma- and stressor-related disorders	21,425	24,093	34,988	30,829	28,781	44,036
Anxiety disorders	11,283	12,988	14,640	15,808	14,334	21,148
Attention deficit/hyperactivity disorder	12,731	12,562	15,317	14,455	10,488	13,991
Mood disorders	93,283	87,762	82,067	84,501	75,983	108,593
Bipolar disorders	39,892	32,627	29,115	29,437	23,363	32,579
Depressive disorders	53,391	55,140	55,032	57,163	52,772	76,895
Personality disorders	243	509	644	693	659	952
Schizophrenia and other psychotic disorders	30,180	29,188	32,843	35,278	26,300	35,609
Alcohol or substance-related disorders	789	1,378	1,008	1,472	1,291	661
Other mental disorders ²	7,690	10,289	18,807	21,090	18,977	26,840
Total	177,417	178,478	181,039	185,759	174,330	244,240
Gender						
Male	112,591	105,359	106,001	109,510	99,993	138,815
Female	118,014	108,868	107,757	108,857	104,083	143,413
Total	230,605	214,227	213,758	218,367	204,076	282,228
Age³						
0 to 11 years	16,554	15,201	14,621	13,903	15,836	19,546
12 to 14 years	10,513	9,926	9,927	10,386	11,500	15,455
15 to 17 years	14,864	13,360	12,864	12,637	12,452	17,737
18 to 20 years	10,429	9,733	9,888	10,004	8,584	13,464
21 to 24 years	13,818	12,690	12,661	13,014	12,311	18,105
25 to 29 years	20,907	20,034	20,630	20,768	19,764	27,952
30 to 34 years	21,150	20,068	20,427	21,319	19,615	28,493
35 to 39 years	18,843	18,393	19,079	20,044	18,252	26,344
40 to 44 years	17,848	16,204	16,159	16,548	15,119	21,472
45 to 49 years	19,710	17,971	17,459	17,279	15,918	20,665
50 to 54 years	22,820	20,506	19,192	18,570	16,735	21,321
55 to 59 years	19,617	18,455	18,484	18,778	16,704	21,907
60 to 64 years	12,728	12,134	12,323	13,155	11,503	15,333
65 years and older	10,850	9,580	10,043	11,961	9,780	14,434
Total	230,651	214,255	213,757	218,366	204,073	282,228

Continued. See notes at end of table.

Table 9.10a. Florida – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	913	831	723	786	678	1,016
Asian ⁴	1,284	1,419	1,346	1,276	1,238	1,650
Black or African American	51,235	47,286	46,896	47,914	43,960	59,494
Native Hawaiian or Other Pacific Islander	448	411	377	419	363	543
White	158,821	145,967	144,095	146,366	134,851	186,639
Other	17,033	14,781	13,531	14,196	22,960	32,877
Total	229,734	210,695	206,968	210,957	204,050	282,219
Ethnicity						
Not of Hispanic or Latino origin	195,154	179,459	177,707	178,641	163,109	229,423
Hispanic or Latino origin	35,445	34,749	36,022	39,693	38,970	50,471
Total	230,599	214,208	213,729	218,334	202,079	279,894
Specific Hispanic or Latino origin						
Mexican	3,037	2,693	2,878	3,063	2,864	3,993
Puerto Rican	6,986	7,251	8,017	8,146	8,035	10,603
Cuban	7,292	6,776	6,273	7,280	7,021	8,618
Other specific	14,481	14,439	14,632	16,392	5,414	7,538
Total	31,796	31,159	31,800	34,881	23,334	30,752
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	9,230	7,199	33,516	53,721
Not a veteran	<i>n/c</i>	<i>n/c</i>	144,543	151,304	124,896	168,193
Total	<i>n/c</i>	<i>n/c</i>	153,773	158,503	158,412	221,914

n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.10b. Florida – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	12.1	13.5	19.3	16.6	16.5	18.0
Anxiety disorders	6.4	7.3	8.1	8.5	8.2	8.7
Attention deficit/hyperactivity disorder	7.2	7.0	8.5	7.8	6.0	5.7
Mood disorders	52.6	49.2	45.3	45.5	43.6	44.5
Bipolar disorders	22.5	18.3	16.1	15.8	13.4	13.3
Depressive disorders	30.1	30.9	30.4	30.8	30.3	31.5
Personality disorders	†	†	†	†	†	†
Schizophrenia and other psychotic disorders	17.0	16.4	18.1	19.0	15.1	14.6
Alcohol or substance-related disorders	†	†	†	†	†	†
Other mental disorders ²	4.3	5.8	10.4	11.4	10.9	11.0
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	48.8	49.2	49.6	50.1	49.0	49.2
Female	51.2	50.8	50.4	49.9	51.0	50.8
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	7.2	7.1	6.8	6.4	7.8	6.9
12 to 14 years	4.6	4.6	4.6	4.8	5.6	5.5
15 to 17 years	6.4	6.2	6.0	5.8	6.1	6.3
18 to 20 years	4.5	4.5	4.6	4.6	4.2	4.8
21 to 24 years	6.0	5.9	5.9	6.0	6.0	6.4
25 to 29 years	9.1	9.4	9.7	9.5	9.7	9.9
30 to 34 years	9.2	9.4	9.6	9.8	9.6	10.1
35 to 39 years	8.2	8.6	8.9	9.2	8.9	9.3
40 to 44 years	7.7	7.6	7.6	7.6	7.4	7.6
45 to 49 years	8.5	8.4	8.2	7.9	7.8	7.3
50 to 54 years	9.9	9.6	9.0	8.5	8.2	7.6
55 to 59 years	8.5	8.6	8.6	8.6	8.2	7.8
60 to 64 years	5.5	5.7	5.8	6.0	5.6	5.4
65 years and older	4.7	4.5	4.7	5.5	4.8	5.1
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.10b. Florida – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	†	†	†	†	†	†
Asian ⁴	†	†	†	†	†	†
Black or African American	22.3	22.4	22.7	22.7	21.5	21.1
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	69.1	69.3	69.6	69.4	66.1	66.1
Other	7.4	7.0	6.5	6.7	11.3	11.6
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	84.6	83.8	83.1	81.8	80.7	82.0
Hispanic or Latino origin	15.4	16.2	16.9	18.2	19.3	18.0
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	9.6	8.6	9.1	8.8	12.3	13.0
Puerto Rican	22.0	23.3	25.2	23.4	34.4	34.5
Cuban	22.9	21.7	19.7	20.9	30.1	28.0
Other specific	45.5	46.3	46.0	47.0	23.2	24.5
Total	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	6.0	4.5	21.2	24.2
Not a veteran	<i>n/c</i>	<i>n/c</i>	94.0	95.5	78.8	75.8
Total	<i>n/c</i>	<i>n/c</i>	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals; n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.11a. Georgia – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020

<i>Selected characteristics</i>	2015	2016	2017	2018	2019	2020
Total individuals	‡	‡	‡	‡	‡	140,661
Mental health diagnosis¹						
Trauma- and stressor-related disorders	‡	‡	‡	‡	‡	25,596
Anxiety disorders	‡	‡	‡	‡	‡	31,929
Attention deficit/hyperactivity disorder	‡	‡	‡	‡	‡	12,127
Mood disorders	‡	‡	‡	‡	‡	70,212
Bipolar disorders	‡	‡	‡	‡	‡	25,989
Depressive disorders	‡	‡	‡	‡	‡	45,549
Personality disorders	‡	‡	‡	‡	‡	3,432
Schizophrenia and other psychotic disorders	‡	‡	‡	‡	‡	29,419
Alcohol or substance-related disorders	‡	‡	‡	‡	‡	32,553
Other mental disorders ²	‡	‡	‡	‡	‡	27,874
Total	‡	‡	‡	‡	‡	140,657
Gender						
Male	‡	‡	‡	‡	‡	69,316
Female	‡	‡	‡	‡	‡	71,071
Total	‡	‡	‡	‡	‡	140,387
Age³						
0 to 11 years	‡	‡	‡	‡	‡	4,892
12 to 14 years	‡	‡	‡	‡	‡	4,024
15 to 17 years	‡	‡	‡	‡	‡	4,364
18 to 20 years	‡	‡	‡	‡	‡	5,983
21 to 24 years	‡	‡	‡	‡	‡	9,597
25 to 29 years	‡	‡	‡	‡	‡	15,575
30 to 34 years	‡	‡	‡	‡	‡	15,798
35 to 39 years	‡	‡	‡	‡	‡	14,378
40 to 44 years	‡	‡	‡	‡	‡	12,432
45 to 49 years	‡	‡	‡	‡	‡	12,342
50 to 54 years	‡	‡	‡	‡	‡	12,940
55 to 59 years	‡	‡	‡	‡	‡	13,104
60 to 64 years	‡	‡	‡	‡	‡	8,723
65 years and older	‡	‡	‡	‡	‡	6,507
Total	‡	‡	‡	‡	‡	140,659

Continued. See notes at end of table.

Table 9.11a. Georgia – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	‡	‡	‡	‡	‡	263
Asian ⁴	‡	‡	‡	‡	‡	1,686
Black or African American	‡	‡	‡	‡	‡	59,357
Native Hawaiian or Other Pacific Islander	‡	‡	‡	‡	‡	162
White	‡	‡	‡	‡	‡	68,846
Other	‡	‡	‡	‡	‡	5,737
Total	‡	‡	‡	‡	‡	136,051
Ethnicity						
Not of Hispanic or Latino origin	‡	‡	‡	‡	‡	125,795
Hispanic or Latino origin	‡	‡	‡	‡	‡	5,804
Total	‡	‡	‡	‡	‡	131,599
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	2,458
Not a veteran	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	105,286
Total	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	107,744

‡ No data submitted; n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.11b. Georgia – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	‡	‡	‡	‡	‡	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	‡	‡	‡	‡	‡	18.2
Anxiety disorders	‡	‡	‡	‡	‡	22.7
Attention deficit/hyperactivity disorder	‡	‡	‡	‡	‡	8.6
Mood disorders	‡	‡	‡	‡	‡	49.9
Bipolar disorders	‡	‡	‡	‡	‡	18.5
Depressive disorders	‡	‡	‡	‡	‡	32.4
Personality disorders	‡	‡	‡	‡	‡	2.4
Schizophrenia and other psychotic disorders	‡	‡	‡	‡	‡	20.9
Alcohol or substance-related disorders	‡	‡	‡	‡	‡	23.1
Other mental disorders ²	‡	‡	‡	‡	‡	19.8
Total	‡	‡	‡	‡	‡	100.0
Gender						
Male	‡	‡	‡	‡	‡	49.4
Female	‡	‡	‡	‡	‡	50.6
Total	‡	‡	‡	‡	‡	100.0
Age³						
0 to 11 years	‡	‡	‡	‡	‡	3.5
12 to 14 years	‡	‡	‡	‡	‡	2.9
15 to 17 years	‡	‡	‡	‡	‡	3.1
18 to 20 years	‡	‡	‡	‡	‡	4.3
21 to 24 years	‡	‡	‡	‡	‡	6.8
25 to 29 years	‡	‡	‡	‡	‡	11.1
30 to 34 years	‡	‡	‡	‡	‡	11.2
35 to 39 years	‡	‡	‡	‡	‡	10.2
40 to 44 years	‡	‡	‡	‡	‡	8.8
45 to 49 years	‡	‡	‡	‡	‡	8.8
50 to 54 years	‡	‡	‡	‡	‡	9.2
55 to 59 years	‡	‡	‡	‡	‡	9.3
60 to 64 years	‡	‡	‡	‡	‡	6.2
65 years and older	‡	‡	‡	‡	‡	4.6
Total	‡	‡	‡	‡	‡	100.0

Continued. See notes at end of table.

Table 9.11b. Georgia – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	‡	‡	‡	‡	‡	†
Asian ⁴	‡	‡	‡	‡	‡	1.2
Black or African American	‡	‡	‡	‡	‡	43.6
Native Hawaiian or Other Pacific Islander	‡	‡	‡	‡	‡	†
White	‡	‡	‡	‡	‡	50.6
Other	‡	‡	‡	‡	‡	4.2
Total	‡	‡	‡	‡	‡	100.0
Ethnicity						
Not of Hispanic or Latino origin	‡	‡	‡	‡	‡	95.6
Hispanic or Latino origin	‡	‡	‡	‡	‡	4.4
Total	‡	‡	‡	‡	‡	100.0
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	2.3
Not a veteran	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	97.7
Total	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	100.0

† Less than 1 percent of individuals; ‡ No data submitted; n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.12a. Hawaii – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	10,645	10,117	9,915	10,006	9,905	10,014
Mental health diagnosis¹						
Trauma- and stressor-related disorders	2,653	2,309	2,567	3,044	2,542	1,120
Anxiety disorders	507	606	690	649	561	501
Attention deficit/hyperactivity disorder	658	477	526	596	461	512
Mood disorders	3,790	3,881	3,769	3,393	3,154	2,051
Bipolar disorders	1,427	1,655	1,572	1,400	1,158	784
Depressive disorders	2,421	2,546	2,449	2,170	2,134	1,352
Personality disorders	316	475	339	589	457	148
Schizophrenia and other psychotic disorders	3,001	3,287	3,166	3,153	2,784	2,354
Alcohol or substance-related disorders	346	91	146	162	45	103
Other mental disorders ²	2,158	1,906	1,841	2,105	1,178	1,134
Total	9,880	9,394	9,099	9,097	8,602	6,083
Gender						
Male	6,045	5,840	5,708	5,823	5,838	5,909
Female	4,392	4,253	4,204	4,180	4,052	3,942
Total	10,437	10,093	9,912	10,003	9,890	9,851
Age³						
0 to 11 years	712	805	820	797	879	597
12 to 14 years	559	549	582	595	703	498
15 to 17 years	893	892	822	831	986	750
18 to 20 years	384	403	337	317	415	302
21 to 24 years	388	299	323	334	326	352
25 to 29 years	591	545	560	601	555	594
30 to 34 years	714	662	662	680	609	776
35 to 39 years	672	632	619	683	679	766
40 to 44 years	756	660	622	632	579	736
45 to 49 years	960	828	738	759	681	745
50 to 54 years	1,121	1,073	967	901	780	890
55 to 59 years	1,164	1,131	1,087	1,032	885	996
60 to 64 years	834	784	814	872	799	872
65 years and older	895	852	954	968	1,024	1,110
Total	10,643	10,115	9,907	10,002	9,900	9,984

Continued. See notes at end of table.

Table 9.12a. Hawaii – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	59	57	54	69	53	72
Asian ⁴	1,510	1,503	1,453	1,395	1,339	1,384
Black or African American	154	155	176	217	188	179
Native Hawaiian or Other Pacific Islander	964	999	937	973	1,002	1,141
White	2,034	2,088	2,099	2,149	2,115	2,158
Other	2,363	2,078	1,850	1,604	2,134	1,992
Total	7,084	6,880	6,569	6,407	6,831	6,926
Ethnicity						
Not of Hispanic or Latino origin	6,543	7,960	5,776	5,699	6,332	5,828
Hispanic or Latino origin	809	830	689	529	630	677
Total	7,352	8,790	6,465	6,228	6,962	6,505
Specific Hispanic or Latino origin						
Mexican	33	39	40	30	83	120
Puerto Rican	120	152	139	145	258	304
Cuban	6	6	5	4	2	4
Other specific	117	153	138	116	111	103
Total	276	350	322	295	454	531
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡
Not a veteran	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡
Total	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡

‡ No data submitted; n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.12b. Hawaii – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	26.9	24.6	28.2	33.5	29.6	18.4
Anxiety disorders	5.1	6.5	7.6	7.1	6.5	8.2
Attention deficit/hyperactivity disorder	6.7	5.1	5.8	6.6	5.4	8.4
Mood disorders	38.4	41.3	41.4	37.3	36.7	33.7
Bipolar disorders	14.4	17.6	17.3	15.4	13.5	12.9
Depressive disorders	24.5	27.1	26.9	23.9	24.8	22.2
Personality disorders	3.2	5.1	3.7	6.5	5.3	2.4
Schizophrenia and other psychotic disorders	30.4	35.0	34.8	34.7	32.4	38.7
Alcohol or substance-related disorders	3.5	†	1.6	1.8	†	1.7
Other mental disorders ²	21.8	20.3	20.2	23.1	13.7	18.6
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	57.9	57.9	57.6	58.2	59.0	60.0
Female	42.1	42.1	42.4	41.8	41.0	40.0
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	6.7	8.0	8.3	8.0	8.9	6.0
12 to 14 years	5.3	5.4	5.9	5.9	7.1	5.0
15 to 17 years	8.4	8.8	8.3	8.3	10.0	7.5
18 to 20 years	3.6	4.0	3.4	3.2	4.2	3.0
21 to 24 years	3.6	3.0	3.3	3.3	3.3	3.5
25 to 29 years	5.6	5.4	5.7	6.0	5.6	5.9
30 to 34 years	6.7	6.5	6.7	6.8	6.2	7.8
35 to 39 years	6.3	6.2	6.2	6.8	6.9	7.7
40 to 44 years	7.1	6.5	6.3	6.3	5.8	7.4
45 to 49 years	9.0	8.2	7.4	7.6	6.9	7.5
50 to 54 years	10.5	10.6	9.8	9.0	7.9	8.9
55 to 59 years	10.9	11.2	11.0	10.3	8.9	10.0
60 to 64 years	7.8	7.8	8.2	8.7	8.1	8.7
65 years and older	8.4	8.4	9.6	9.7	10.3	11.1
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.12b. Hawaii – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	†	†	†	1.1	†	1.0
Asian ⁴	21.3	21.8	22.1	21.8	19.6	20.0
Black or African American	2.2	2.3	2.7	3.4	2.8	2.6
Native Hawaiian or Other Pacific Islander	13.6	14.5	14.3	15.2	14.7	16.5
White	28.7	30.3	32.0	33.5	31.0	31.2
Other	33.4	30.2	28.2	25.0	31.2	28.8
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	89.0	90.6	89.3	91.5	91.0	89.6
Hispanic or Latino origin	11.0	9.4	10.7	8.5	9.0	10.4
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	12.0	11.1	12.4	10.2	18.3	22.6
Puerto Rican	43.5	43.4	43.2	49.2	56.8	57.3
Cuban	2.2	1.7	1.6	1.4	†	†
Other specific	42.4	43.7	42.9	39.3	24.4	19.4
Total	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡
Not a veteran	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡
Total	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡

† Less than 1 percent of individuals; ‡ No data submitted; n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.13a. Idaho – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020

<i>Selected characteristics</i>	2015	2016	2017	2018	2019	2020
Total individuals	14,425	15,320	15,424	16,718	17,465	16,364
Mental health diagnosis¹						
Trauma- and stressor-related disorders	259	585	846	863	865	805
Anxiety disorders	216	1,322	1,434	1,447	1,399	1,237
Attention deficit/hyperactivity disorder	114	620	656	697	657	691
Mood disorders	1,876	3,874	4,152	4,192	4,078	3,773
Bipolar disorders	1,179	1,660	1,687	1,665	1,579	1,440
Depressive disorders	766	2,282	2,542	2,601	2,571	2,389
Personality disorders	1,018	544	684	686	618	514
Schizophrenia and other psychotic disorders	799	1,909	2,045	2,071	1,999	1,993
Alcohol or substance-related disorders	562	8	17	2	- -	4
Other mental disorders ²	1,459	1,557	1,457	1,439	1,304	1,302
Total	3,276	7,278	7,661	7,667	7,421	7,082
Gender						
Male	8,581	9,125	9,191	9,925	10,345	9,663
Female	5,838	6,191	6,224	6,782	7,108	6,680
Total	14,419	15,316	15,415	16,707	17,453	16,343
Age³						
0 to 11 years	177	230	357	727	1,147	988
12 to 14 years	547	516	484	707	920	933
15 to 17 years	1,002	975	1,013	1,058	1,107	1,261
18 to 20 years	1,090	1,096	1,063	1,138	1,125	1,122
21 to 24 years	1,494	1,594	1,544	1,557	1,500	1,333
25 to 29 years	1,871	1,995	2,061	2,177	2,109	1,869
30 to 34 years	1,730	1,914	1,897	1,987	1,982	1,791
35 to 39 years	1,547	1,717	1,704	1,830	1,864	1,693
40 to 44 years	1,166	1,238	1,260	1,346	1,383	1,314
45 to 49 years	1,110	1,201	1,233	1,222	1,164	1,049
50 to 54 years	1,069	1,137	1,005	989	1,052	958
55 to 59 years	768	804	824	914	932	863
60 to 64 years	445	455	469	508	567	516
65 years and older	399	421	484	527	582	657
Total	14,415	15,293	15,398	16,687	17,434	16,347

Continued. See notes at end of table.

Table 9.13a. Idaho – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	331	396	368	375	533	362
Asian ⁴	74	80	75	95	103	87
Black or African American	268	272	287	293	458	337
Native Hawaiian or Other Pacific Islander	36	43	38	54	69	63
White	11,372	11,785	11,776	12,659	13,371	12,176
Other	1,204	1,378	1,437	1,761	1,398	1,597
Total	13,285	13,954	13,981	15,237	15,932	14,622
Ethnicity						
Not of Hispanic or Latino origin	11,463	11,900	11,896	12,795	13,611	12,276
Hispanic or Latino origin	1,410	1,574	1,552	1,753	1,830	1,816
Total	12,873	13,474	13,448	14,548	15,441	14,092
Specific Hispanic or Latino origin						
Mexican	294	252	206	213	225	226
Puerto Rican	26	22	22	22	24	23
Cuban	6	9	9	5	5	3
Other specific	325	220	157	160	148	153
Total	651	503	394	400	402	405
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	497	470	474	413
Not a veteran	<i>n/c</i>	<i>n/c</i>	10,253	10,724	10,697	9,709
Total	<i>n/c</i>	<i>n/c</i>	10,750	11,194	11,171	10,122

- - Quantity is zero; n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.13b. Idaho – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	7.9	8.0	11.0	11.3	11.7	11.4
Anxiety disorders	6.6	18.2	18.7	18.9	18.9	17.5
Attention deficit/hyperactivity disorder	3.5	8.5	8.6	9.1	8.9	9.8
Mood disorders	57.3	53.2	54.2	54.7	55.0	53.3
Bipolar disorders	36.0	22.8	22.0	21.7	21.3	20.3
Depressive disorders	23.4	31.4	33.2	33.9	34.6	33.7
Personality disorders	31.1	7.5	8.9	8.9	8.3	7.3
Schizophrenia and other psychotic disorders	24.4	26.2	26.7	27.0	26.9	28.1
Alcohol or substance-related disorders	17.2	†	†	†	--	†
Other mental disorders ²	44.5	21.4	19.0	18.8	17.6	18.4
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	59.5	59.6	59.6	59.4	59.3	59.1
Female	40.5	40.4	40.4	40.6	40.7	40.9
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	1.2	1.5	2.3	4.4	6.6	6.0
12 to 14 years	3.8	3.4	3.1	4.2	5.3	5.7
15 to 17 years	7.0	6.4	6.6	6.3	6.3	7.7
18 to 20 years	7.6	7.2	6.9	6.8	6.5	6.9
21 to 24 years	10.4	10.4	10.0	9.3	8.6	8.2
25 to 29 years	13.0	13.0	13.4	13.0	12.1	11.4
30 to 34 years	12.0	12.5	12.3	11.9	11.4	11.0
35 to 39 years	10.7	11.2	11.1	11.0	10.7	10.4
40 to 44 years	8.1	8.1	8.2	8.1	7.9	8.0
45 to 49 years	7.7	7.9	8.0	7.3	6.7	6.4
50 to 54 years	7.4	7.4	6.5	5.9	6.0	5.9
55 to 59 years	5.3	5.3	5.4	5.5	5.3	5.3
60 to 64 years	3.1	3.0	3.0	3.0	3.3	3.2
65 years and older	2.8	2.8	3.1	3.2	3.3	4.0
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.13b. Idaho – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	2.5	2.8	2.6	2.5	3.3	2.5
Asian ⁴	†	†	†	†	†	†
Black or African American	2.0	1.9	2.1	1.9	2.9	2.3
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	85.6	84.5	84.2	83.1	83.9	83.3
Other	9.1	9.9	10.3	11.6	8.8	10.9
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	89.0	88.3	88.5	88.0	88.1	87.1
Hispanic or Latino origin	11.0	11.7	11.5	12.0	11.9	12.9
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	45.2	50.1	52.3	53.3	56.0	55.8
Puerto Rican	4.0	4.4	5.6	5.5	6.0	5.7
Cuban	†	1.8	2.3	1.3	1.2	†
Other specific	49.9	43.7	39.8	40.0	36.8	37.8
Total	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	4.6	4.2	4.2	4.1
Not a veteran	<i>n/c</i>	<i>n/c</i>	95.4	95.8	95.8	95.9
Total	<i>n/c</i>	<i>n/c</i>	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals; - - Quantity is zero; n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.14a. Illinois – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	130,002	78,703	71,195	66,656	45,670	39,500
Mental health diagnosis¹						
Trauma- and stressor-related disorders	21,692	2,411	12,469	12,353	8,886	8,298
Anxiety disorders	26,838	2,824	13,966	13,247	8,753	8,079
Attention deficit/hyperactivity disorder	17,030	1,550	8,393	7,453	4,783	4,198
Mood disorders	75,102	5,541	34,296	31,386	20,608	17,252
Bipolar disorders	30,723	2,507	11,999	10,717	7,435	6,111
Depressive disorders	45,237	3,065	22,522	20,906	13,345	11,329
Personality disorders	6,118	814	2,772	2,356	1,571	1,239
Schizophrenia and other psychotic disorders	18,661	3,406	11,553	10,736	9,012	7,682
Alcohol or substance-related disorders	23	2,276	2,574	2,645	2,304	2,011
Other mental disorders ²	103,786	5,430	12,165	11,309	7,876	6,719
Total	129,996	17,324	70,802	66,237	45,442	39,267
Gender						
Male	62,453	38,763	34,141	32,109	22,780	19,886
Female	67,547	39,938	37,054	34,547	22,890	19,614
Total	130,000	78,701	71,195	66,656	45,670	39,500
Age³						
0 to 11 years	15,204	9,085	8,205	7,745	4,842	4,480
12 to 14 years	9,459	5,513	4,871	4,732	3,311	3,133
15 to 17 years	10,196	5,881	5,276	5,038	3,365	3,261
18 to 20 years	6,650	3,818	3,499	3,218	2,496	2,240
21 to 24 years	8,402	4,726	4,369	3,967	2,407	2,126
25 to 29 years	10,901	6,635	6,191	5,876	3,578	3,059
30 to 34 years	10,964	6,414	5,866	5,482	3,392	2,884
35 to 39 years	10,121	6,113	5,730	5,358	3,384	2,802
40 to 44 years	9,733	5,750	5,078	4,594	3,000	2,673
45 to 49 years	9,627	5,982	5,087	4,730	3,143	2,556
50 to 54 years	10,661	6,591	5,632	4,954	3,384	2,710
55 to 59 years	8,396	5,730	5,207	4,961	3,718	3,039
60 to 64 years	5,020	3,466	3,217	3,208	2,763	2,224
65 years and older	4,666	2,996	2,966	2,793	2,887	2,313
Total	130,000	78,700	71,194	66,656	45,670	39,500

Continued. See notes at end of table.

Table 9.14a. Illinois – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	330	235	194	212	153	149
Asian ⁴	1,655	924	776	675	609	542
Black or African American	25,586	15,461	12,415	11,911	9,910	8,583
Native Hawaiian or Other Pacific Islander	134	66	59	51	40	33
White	90,658	55,367	51,775	48,401	29,982	25,782
Other	1,323	1,002	997	916	459	388
Total	119,686	73,055	66,216	62,166	41,153	35,477
Ethnicity						
Not of Hispanic or Latino origin	107,168	67,380	62,032	57,957	37,158	31,321
Hispanic or Latino origin	17,273	8,090	5,706	5,262	4,484	4,223
Total	124,441	75,470	67,738	63,219	41,642	35,544
Specific Hispanic or Latino origin						
Mexican	11,314	4,887	3,286	2,980	2,564	2,507
Puerto Rican	2,330	1,243	927	822	532	415
Cuban	191	85	70	54	45	32
Other specific	3,129	1,497	1,206	1,252	1,235	1,177
Total	16,964	7,712	5,489	5,108	4,376	4,131
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	1,266	1,161	767	590
Not a veteran	<i>n/c</i>	<i>n/c</i>	47,992	44,161	29,934	24,891
Total	<i>n/c</i>	<i>n/c</i>	49,258	45,322	30,701	25,481

n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.14b. Illinois – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	16.7	13.9	17.6	18.6	19.6	21.1
Anxiety disorders	20.6	16.3	19.7	20.0	19.3	20.6
Attention deficit/hyperactivity disorder	13.1	8.9	11.9	11.3	10.5	10.7
Mood disorders	57.8	32.0	48.4	47.4	45.4	43.9
Bipolar disorders	23.6	14.5	16.9	16.2	16.4	15.6
Depressive disorders	34.8	17.7	31.8	31.6	29.4	28.9
Personality disorders	4.7	4.7	3.9	3.6	3.5	3.2
Schizophrenia and other psychotic disorders	14.4	19.7	16.3	16.2	19.8	19.6
Alcohol or substance-related disorders	†	13.1	3.6	4.0	5.1	5.1
Other mental disorders ²	79.8	31.3	17.2	17.1	17.3	17.1
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	48.0	49.3	48.0	48.2	49.9	50.3
Female	52.0	50.7	52.0	51.8	50.1	49.7
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	11.7	11.5	11.5	11.6	10.6	11.3
12 to 14 years	7.3	7.0	6.8	7.1	7.2	7.9
15 to 17 years	7.8	7.5	7.4	7.6	7.4	8.3
18 to 20 years	5.1	4.9	4.9	4.8	5.5	5.7
21 to 24 years	6.5	6.0	6.1	6.0	5.3	5.4
25 to 29 years	8.4	8.4	8.7	8.8	7.8	7.7
30 to 34 years	8.4	8.1	8.2	8.2	7.4	7.3
35 to 39 years	7.8	7.8	8.0	8.0	7.4	7.1
40 to 44 years	7.5	7.3	7.1	6.9	6.6	6.8
45 to 49 years	7.4	7.6	7.1	7.1	6.9	6.5
50 to 54 years	8.2	8.4	7.9	7.4	7.4	6.9
55 to 59 years	6.5	7.3	7.3	7.4	8.1	7.7
60 to 64 years	3.9	4.4	4.5	4.8	6.0	5.6
65 years and older	3.6	3.8	4.2	4.2	6.3	5.9
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.14b. Illinois – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	†	†	†	†	†	†
Asian ⁴	1.4	1.3	1.2	1.1	1.5	1.5
Black or African American	21.4	21.2	18.7	19.2	24.1	24.2
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	75.7	75.8	78.2	77.9	72.9	72.7
Other	1.1	1.4	1.5	1.5	1.1	1.1
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	86.1	89.3	91.6	91.7	89.2	88.1
Hispanic or Latino origin	13.9	10.7	8.4	8.3	10.8	11.9
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	66.7	63.4	59.9	58.3	58.6	60.7
Puerto Rican	13.7	16.1	16.9	16.1	12.2	10.0
Cuban	1.1	1.1	1.3	1.1	1.0	†
Other specific	18.4	19.4	22.0	24.5	28.2	28.5
Total	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	2.6	2.6	2.5	2.3
Not a veteran	<i>n/c</i>	<i>n/c</i>	97.4	97.4	97.5	97.7
Total	<i>n/c</i>	<i>n/c</i>	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals; n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.15a. Indiana – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	134,121	135,384	137,628	137,357	139,126	138,697
Mental health diagnosis¹						
Trauma- and stressor-related disorders	26,088	24,724	25,841	27,140	29,311	31,334
Anxiety disorders	31,280	40,948	42,251	43,577	46,769	50,737
Attention deficit/hyperactivity disorder	26,262	30,131	28,456	27,616	27,829	27,231
Mood disorders	66,173	61,043	62,620	62,559	63,695	64,715
Bipolar disorders	19,560	15,488	15,036	14,419	14,281	13,981
Depressive disorders	47,319	46,103	48,194	48,787	50,195	51,612
Personality disorders	7,512	8,353	7,592	7,223	6,324	6,161
Schizophrenia and other psychotic disorders	15,477	16,345	16,119	15,535	15,281	14,278
Alcohol or substance-related disorders	2	21	10	15	6,719	10
Other mental disorders ²	32,120	39,063	37,517	36,838	40,722	36,621
Total	133,292	135,062	137,591	137,337	138,836	137,905
Gender						
Male	63,170	63,921	64,835	65,048	65,837	65,398
Female	70,951	71,463	72,793	72,309	73,289	73,299
Total	134,121	135,384	137,628	137,357	139,126	138,697
Age³						
0 to 11 years	28,517	28,617	29,157	29,661	30,493	29,303
12 to 14 years	13,093	12,840	13,041	13,684	14,718	14,488
15 to 17 years	11,968	12,588	12,458	12,416	12,633	12,362
18 to 20 years	5,331	5,365	5,744	5,881	6,172	6,337
21 to 24 years	6,591	6,467	6,642	6,657	6,740	7,118
25 to 29 years	8,772	8,933	9,365	9,441	9,499	10,083
30 to 34 years	9,250	9,078	9,185	8,837	9,049	9,561
35 to 39 years	8,759	8,805	9,144	8,867	8,781	9,205
40 to 44 years	8,400	8,302	8,143	7,929	7,840	7,931
45 to 49 years	8,421	8,387	8,327	8,042	7,815	7,566
50 to 54 years	9,185	9,253	8,897	8,314	7,778	7,328
55 to 59 years	7,379	7,703	7,968	7,859	7,628	7,402
60 to 64 years	4,426	4,760	5,100	5,211	5,291	5,227
65 years and older	4,028	4,285	4,455	4,558	4,688	4,785
Total	134,120	135,383	137,626	137,357	139,125	138,696

Continued. See notes at end of table.

Table 9.15a. Indiana – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	674	642	575	505	451	424
Asian ⁴	432	435	519	521	555	563
Black or African American	19,380	19,523	20,136	20,163	20,031	19,212
Native Hawaiian or Other Pacific Islander	101	105	98	127	110	110
White	103,503	104,601	105,898	106,692	107,128	106,481
Other	10,029	10,077	10,400	9,349	10,849	11,903
Total	134,119	135,383	137,626	137,357	139,124	138,693
Ethnicity						
Not of Hispanic or Latino origin	121,098	122,498	124,840	125,498	126,776	126,950
Hispanic or Latino origin	8,134	8,416	8,727	9,617	9,476	9,855
Total	129,232	130,914	133,567	135,115	136,252	136,805
Specific Hispanic or Latino origin						
Mexican	3,149	3,232	3,262	3,413	3,637	3,233
Puerto Rican	602	616	614	737	689	602
Cuban	52	54	51	69	67	87
Other specific	4,293	4,468	4,760	5,366	5,047	5,899
Total	8,096	8,370	8,687	9,585	9,440	9,821
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	2,678	2,628	2,336	2,146
Not a veteran	<i>n/c</i>	<i>n/c</i>	611	651	77,623	78,940
Total	<i>n/c</i>	<i>n/c</i>	3,289	3,279	79,959	81,086

n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.15b. Indiana – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	19.6	18.3	18.8	19.8	21.1	22.7
Anxiety disorders	23.5	30.3	30.7	31.7	33.7	36.8
Attention deficit/hyperactivity disorder	19.7	22.3	20.7	20.1	20.0	19.7
Mood disorders	49.6	45.2	45.5	45.6	45.9	46.9
Bipolar disorders	14.7	11.5	10.9	10.5	10.3	10.1
Depressive disorders	35.5	34.1	35.0	35.5	36.2	37.4
Personality disorders	5.6	6.2	5.5	5.3	4.6	4.5
Schizophrenia and other psychotic disorders	11.6	12.1	11.7	11.3	11.0	10.4
Alcohol or substance-related disorders	†	†	†	†	4.8	†
Other mental disorders ²	24.1	28.9	27.3	26.8	29.3	26.6
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	47.1	47.2	47.1	47.4	47.3	47.2
Female	52.9	52.8	52.9	52.6	52.7	52.8
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	21.3	21.1	21.2	21.6	21.9	21.1
12 to 14 years	9.8	9.5	9.5	10.0	10.6	10.4
15 to 17 years	8.9	9.3	9.1	9.0	9.1	8.9
18 to 20 years	4.0	4.0	4.2	4.3	4.4	4.6
21 to 24 years	4.9	4.8	4.8	4.8	4.8	5.1
25 to 29 years	6.5	6.6	6.8	6.9	6.8	7.3
30 to 34 years	6.9	6.7	6.7	6.4	6.5	6.9
35 to 39 years	6.5	6.5	6.6	6.5	6.3	6.6
40 to 44 years	6.3	6.1	5.9	5.8	5.6	5.7
45 to 49 years	6.3	6.2	6.1	5.9	5.6	5.5
50 to 54 years	6.8	6.8	6.5	6.1	5.6	5.3
55 to 59 years	5.5	5.7	5.8	5.7	5.5	5.3
60 to 64 years	3.3	3.5	3.7	3.8	3.8	3.8
65 years and older	3.0	3.2	3.2	3.3	3.4	3.4
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.15b. Indiana – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	†	†	†	†	†	†
Asian ⁴	†	†	†	†	†	†
Black or African American	14.4	14.4	14.6	14.7	14.4	13.9
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	77.2	77.3	76.9	77.7	77.0	76.8
Other	7.5	7.4	7.6	6.8	7.8	8.6
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	93.7	93.6	93.5	92.9	93.0	92.8
Hispanic or Latino origin	6.3	6.4	6.5	7.1	7.0	7.2
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	38.9	38.6	37.6	35.6	38.5	32.9
Puerto Rican	7.4	7.4	7.1	7.7	7.3	6.1
Cuban	†	†	†	†	†	†
Other specific	53.0	53.4	54.8	56.0	53.5	60.1
Total	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	81.4	80.1	2.9	2.6
Not a veteran	<i>n/c</i>	<i>n/c</i>	18.6	19.9	97.1	97.4
Total	<i>n/c</i>	<i>n/c</i>	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals; n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.16a. Iowa – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020

<i>Selected characteristics</i>	2015	2016	2017	2018	2019	2020
Total individuals	116,443	115,628	101,763	189,347	263,112	236,041
Mental health diagnosis¹						
Trauma- and stressor-related disorders	27,949	25,028	9,787	38,113	45,939	4,714
Anxiety disorders	17,742	26,162	10,640	51,640	88,718	22,784
Attention deficit/hyperactivity disorder	19,859	19,460	7,373	34,304	41,398	7,147
Mood disorders	37,861	70,023	34,313	71,637	91,982	26,097
Bipolar disorders	13,498	23,501	11,537	16,977	18,500	6,501
Depressive disorders	27,419	54,947	25,693	58,863	78,284	20,852
Personality disorders	684	1,525	589	1,985	3,924	1,157
Schizophrenia and other psychotic disorders	6,909	11,490	6,378	13,254	14,200	7,155
Alcohol or substance-related disorders	1,491	2,363	853	--	--	--
Other mental disorders ²	14,265	19,176	8,003	29,266	40,618	9,473
Total	96,692	105,796	55,800	174,011	203,180	56,738
Gender						
Male	55,013	51,623	43,115	84,182	111,755	103,123
Female	61,430	64,004	58,460	105,026	151,184	132,733
Total	116,443	115,627	101,575	189,208	262,939	235,856
Age³						
0 to 11 years	24,238	15,664	7,496	33,235	39,349	36,361
12 to 14 years	10,109	7,670	5,147	14,784	19,639	18,835
15 to 17 years	9,896	8,687	7,106	14,456	19,040	18,044
18 to 20 years	5,601	5,568	5,270	9,940	14,679	13,403
21 to 24 years	7,187	7,149	6,788	11,397	16,314	14,895
25 to 29 years	9,831	10,280	9,736	17,152	24,651	21,567
30 to 34 years	9,674	10,085	9,585	15,522	22,737	20,521
35 to 39 years	8,260	9,258	9,006	14,455	20,822	18,528
40 to 44 years	6,932	7,834	7,425	11,290	16,526	14,470
45 to 49 years	6,591	7,616	7,514	10,694	15,069	12,283
50 to 54 years	7,177	8,339	8,043	10,660	15,150	12,031
55 to 59 years	5,574	7,306	7,549	10,507	16,126	12,869
60 to 64 years	3,119	4,766	5,032	7,076	12,012	9,954
65 years and older	2,252	5,404	6,060	8,178	10,897	12,266
Total	116,441	115,626	101,757	189,346	263,011	236,027

Continued. See notes at end of table.

Table 9.16a. Iowa – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	1,413	1,337	1,248	2,527	3,995	3,641
Asian ⁴	511	492	451	916	2,946	2,425
Black or African American	7,247	6,740	5,881	10,399	15,412	14,104
Native Hawaiian or Other Pacific Islander	426	387	293	751	--	8
White	78,119	78,211	68,565	121,638	168,703	149,084
Other	3,572	3,146	2,414	6,231	4,344	7,899
Total	91,288	90,313	78,852	142,462	195,400	177,161
Ethnicity						
Not of Hispanic or Latino origin	403	461	466	480	217	171,772
Hispanic or Latino origin	34	44	48	41	13,767	12,850
Total	437	505	514	521	13,984	184,622
Specific Hispanic or Latino origin						
Mexican	3	1	1	‡	‡	‡
Puerto Rican	--	--	--	‡	‡	‡
Cuban	--	--	--	‡	‡	‡
Other specific	--	--	--	‡	‡	‡
Total	3	1	1	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡
Not a veteran	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡
Total	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡

-- Quantity is zero; ‡ No data submitted; n/c Not collected.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.16b. Iowa – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	28.9	23.7	17.5	21.9	22.6	8.3
Anxiety disorders	18.3	24.7	19.1	29.7	43.7	40.2
Attention deficit/hyperactivity disorder	20.5	18.4	13.2	19.7	20.4	12.6
Mood disorders	39.2	66.2	61.5	41.2	45.3	46.0
Bipolar disorders	14.0	22.2	20.7	9.8	9.1	11.5
Depressive disorders	28.4	51.9	46.0	33.8	38.5	36.8
Personality disorders	†	1.4	1.1	1.1	1.9	2.0
Schizophrenia and other psychotic disorders	7.1	10.9	11.4	7.6	7.0	12.6
Alcohol or substance-related disorders	1.5	2.2	1.5	--	--	--
Other mental disorders ²	14.8	18.1	14.3	16.8	20.0	16.7
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	47.2	44.6	42.4	44.5	42.5	43.7
Female	52.8	55.4	57.6	55.5	57.5	56.3
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	20.8	13.5	7.4	17.6	15.0	15.4
12 to 14 years	8.7	6.6	5.1	7.8	7.5	8.0
15 to 17 years	8.5	7.5	7.0	7.6	7.2	7.6
18 to 20 years	4.8	4.8	5.2	5.2	5.6	5.7
21 to 24 years	6.2	6.2	6.7	6.0	6.2	6.3
25 to 29 years	8.4	8.9	9.6	9.1	9.4	9.1
30 to 34 years	8.3	8.7	9.4	8.2	8.6	8.7
35 to 39 years	7.1	8.0	8.9	7.6	7.9	7.8
40 to 44 years	6.0	6.8	7.3	6.0	6.3	6.1
45 to 49 years	5.7	6.6	7.4	5.6	5.7	5.2
50 to 54 years	6.2	7.2	7.9	5.6	5.8	5.1
55 to 59 years	4.8	6.3	7.4	5.5	6.1	5.5
60 to 64 years	2.7	4.1	4.9	3.7	4.6	4.2
65 years and older	1.9	4.7	6.0	4.3	4.1	5.2
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.16b. Iowa – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	1.5	1.5	1.6	1.8	2.0	2.1
Asian ⁴	†	†	†	†	1.5	1.4
Black or African American	7.9	7.5	7.5	7.3	7.9	8.0
Native Hawaiian or Other Pacific Islander	†	†	†	†	--	†
White	85.6	86.6	87.0	85.4	86.3	84.2
Other	3.9	3.5	3.1	4.4	2.2	4.5
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	92.2	91.3	90.7	92.1	1.6	93.0
Hispanic or Latino origin	7.8	8.7	9.3	7.9	98.4	7.0
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	100.0	100.0	100.0	‡	‡	‡
Puerto Rican	--	--	--	‡	‡	‡
Cuban	--	--	--	‡	‡	‡
Other specific	--	--	--	‡	‡	‡
Total	100.0	100.0	100.0	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡
Not a veteran	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡
Total	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡

† Less than 1 percent of individuals; -- Quantity is zero; ‡ No data submitted; *n/c* Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.17a. Kansas – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020

<i>Selected characteristics</i>	2015	2016	2017	2018	2019	2020
Total individuals	‡	‡	‡	‡	‡	146,152
Mental health diagnosis¹						
Trauma- and stressor-related disorders	‡	‡	‡	‡	‡	30,712
Anxiety disorders	‡	‡	‡	‡	‡	33,232
Attention deficit/hyperactivity disorder	‡	‡	‡	‡	‡	15,678
Mood disorders	‡	‡	‡	‡	‡	63,049
Bipolar disorders	‡	‡	‡	‡	‡	15,838
Depressive disorders	‡	‡	‡	‡	‡	47,536
Personality disorders	‡	‡	‡	‡	‡	2,328
Schizophrenia and other psychotic disorders	‡	‡	‡	‡	‡	8,746
Alcohol or substance-related disorders	‡	‡	‡	‡	‡	8,287
Other mental disorders ²	‡	‡	‡	‡	‡	26,105
Total	‡	‡	‡	‡	‡	139,114
Gender						
Male	‡	‡	‡	‡	‡	68,591
Female	‡	‡	‡	‡	‡	76,991
Total	‡	‡	‡	‡	‡	145,582
Age³						
0 to 11 years	‡	‡	‡	‡	‡	17,451
12 to 14 years	‡	‡	‡	‡	‡	10,724
15 to 17 years	‡	‡	‡	‡	‡	11,525
18 to 20 years	‡	‡	‡	‡	‡	9,126
21 to 24 years	‡	‡	‡	‡	‡	10,556
25 to 29 years	‡	‡	‡	‡	‡	13,090
30 to 34 years	‡	‡	‡	‡	‡	12,528
35 to 39 years	‡	‡	‡	‡	‡	11,783
40 to 44 years	‡	‡	‡	‡	‡	9,900
45 to 49 years	‡	‡	‡	‡	‡	8,291
50 to 54 years	‡	‡	‡	‡	‡	7,936
55 to 59 years	‡	‡	‡	‡	‡	7,941
60 to 64 years	‡	‡	‡	‡	‡	6,171
65 years and older	‡	‡	‡	‡	‡	9,130
Total	‡	‡	‡	‡	‡	146,152

Continued. See notes at end of table.

Table 9.17a. Kansas – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	‡	‡	‡	‡	‡	3,670
Asian ⁴	‡	‡	‡	‡	‡	1,486
Black or African American	‡	‡	‡	‡	‡	11,230
Native Hawaiian or Other Pacific Islander	‡	‡	‡	‡	‡	261
White	‡	‡	‡	‡	‡	110,177
Other	‡	‡	‡	‡	‡	18,183
Total	‡	‡	‡	‡	‡	145,007
Ethnicity						
Not of Hispanic or Latino origin	‡	‡	‡	‡	‡	133,056
Hispanic or Latino origin	‡	‡	‡	‡	‡	10,843
Total	‡	‡	‡	‡	‡	143,899
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	1,813
Not a veteran	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	55,355
Total	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	57,168

‡ No data submitted; n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.17b. Kansas – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	‡	‡	‡	‡	‡	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	‡	‡	‡	‡	‡	22.1
Anxiety disorders	‡	‡	‡	‡	‡	23.9
Attention deficit/hyperactivity disorder	‡	‡	‡	‡	‡	11.3
Mood disorders	‡	‡	‡	‡	‡	45.3
Bipolar disorders	‡	‡	‡	‡	‡	11.4
Depressive disorders	‡	‡	‡	‡	‡	34.2
Personality disorders	‡	‡	‡	‡	‡	1.7
Schizophrenia and other psychotic disorders	‡	‡	‡	‡	‡	6.3
Alcohol or substance-related disorders	‡	‡	‡	‡	‡	6.0
Other mental disorders ²	‡	‡	‡	‡	‡	18.8
Total	‡	‡	‡	‡	‡	100.0
Gender						
Male	‡	‡	‡	‡	‡	47.1
Female	‡	‡	‡	‡	‡	52.9
Total	‡	‡	‡	‡	‡	100.0
Age³						
0 to 11 years	‡	‡	‡	‡	‡	11.9
12 to 14 years	‡	‡	‡	‡	‡	7.3
15 to 17 years	‡	‡	‡	‡	‡	7.9
18 to 20 years	‡	‡	‡	‡	‡	6.2
21 to 24 years	‡	‡	‡	‡	‡	7.2
25 to 29 years	‡	‡	‡	‡	‡	9.0
30 to 34 years	‡	‡	‡	‡	‡	8.6
35 to 39 years	‡	‡	‡	‡	‡	8.1
40 to 44 years	‡	‡	‡	‡	‡	6.8
45 to 49 years	‡	‡	‡	‡	‡	5.7
50 to 54 years	‡	‡	‡	‡	‡	5.4
55 to 59 years	‡	‡	‡	‡	‡	5.4
60 to 64 years	‡	‡	‡	‡	‡	4.2
65 years and older	‡	‡	‡	‡	‡	6.2
Total	‡	‡	‡	‡	‡	100.0

Continued. See notes at end of table.

Table 9.17b. Kansas – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	‡	‡	‡	‡	‡	2.5
Asian ⁴	‡	‡	‡	‡	‡	1.0
Black or African American	‡	‡	‡	‡	‡	7.7
Native Hawaiian or Other Pacific Islander	‡	‡	‡	‡	‡	†
White	‡	‡	‡	‡	‡	76.0
Other	‡	‡	‡	‡	‡	12.5
Total	‡	‡	‡	‡	‡	100.0
Ethnicity						
Not of Hispanic or Latino origin	‡	‡	‡	‡	‡	92.5
Hispanic or Latino origin	‡	‡	‡	‡	‡	7.5
Total	‡	‡	‡	‡	‡	100.0
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	3.2
Not a veteran	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	96.8
Total	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	100.0

† Less than 1 percent of individuals; ‡ No data submitted; n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.18a. Kentucky – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	160,096	155,014	155,192	157,732	165,999	156,223
Mental health diagnosis¹						
Trauma- and stressor-related disorders	30,402	30,164	31,081	33,793	36,056	35,466
Anxiety disorders	35,645	37,079	37,719	38,815	38,891	37,282
Attention deficit/hyperactivity disorder	22,807	21,199	20,688	20,670	20,600	19,408
Mood disorders	77,056	67,748	58,818	60,365	58,154	54,201
Bipolar disorders	30,853	23,087	13,616	13,793	13,114	12,056
Depressive disorders	47,462	45,619	45,893	47,462	46,032	43,158
Personality disorders	6,034	4,826	4,370	4,175	3,981	3,599
Schizophrenia and other psychotic disorders	12,521	11,495	11,935	12,214	11,955	11,274
Alcohol or substance-related disorders	5,430	7,378	28,473	29,531	38,855	36,572
Other mental disorders ²	31,981	26,349	41,100	40,144	41,469	38,345
Total	155,648	148,396	153,342	156,225	164,785	153,854
Gender						
Male	77,167	74,017	73,958	75,774	81,897	77,696
Female	82,905	80,947	81,209	81,914	83,547	77,704
Total	160,072	154,964	155,167	157,688	165,444	155,400
Age³						
0 to 11 years	29,922	26,022	25,882	26,645	27,479	25,235
12 to 14 years	13,526	12,490	12,461	12,710	13,435	13,195
15 to 17 years	13,071	12,717	12,822	12,936	12,993	12,346
18 to 20 years	6,740	6,732	6,840	7,338	7,916	7,621
21 to 24 years	9,200	9,180	9,018	9,133	9,744	9,042
25 to 29 years	12,022	12,539	12,818	13,282	14,929	13,462
30 to 34 years	12,339	12,025	12,102	12,244	13,844	13,032
35 to 39 years	11,506	11,800	11,810	11,753	13,318	12,414
40 to 44 years	10,661	10,310	9,916	9,977	11,171	10,711
45 to 49 years	10,716	10,544	10,287	9,887	10,521	9,751
50 to 54 years	11,046	10,871	10,611	10,030	10,007	9,309
55 to 59 years	8,505	8,658	8,581	8,654	8,895	8,496
60 to 64 years	5,147	5,330	5,601	5,728	5,874	5,800
65 years and older	5,687	5,746	6,407	7,375	5,830	5,791
Total	160,088	154,964	155,156	157,692	165,956	156,205

Continued. See notes at end of table.

Table 9.18a. Kentucky – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	312	310	334	306	329	286
Asian ⁴	259	379	419	426	435	407
Black or African American	16,572	15,623	15,554	15,046	16,110	14,834
Native Hawaiian or Other Pacific Islander	103	114	136	152	167	133
White	132,094	126,191	125,397	125,432	130,441	123,068
Other	10,683	10,114	10,539	13,187	15,514	15,194
Total	160,023	152,731	152,379	154,549	162,996	153,922
Ethnicity						
Not of Hispanic or Latino origin	154,186	148,372	148,174	150,198	157,486	148,252
Hispanic or Latino origin	3,150	3,319	3,483	3,453	3,963	3,881
Total	157,336	151,691	151,657	153,651	161,449	152,133
Specific Hispanic or Latino origin						
Mexican	1,396	1,346	1,398	1,392	1,566	1,481
Puerto Rican	551	496	493	506	531	479
Cuban	264	266	288	287	315	299
Other specific	920	1,184	1,259	1,241	1,520	1,591
Total	3,131	3,292	3,438	3,426	3,932	3,850
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	2,352	1,159	2,814	2,400
Not a veteran	<i>n/c</i>	<i>n/c</i>	97,031	59,376	101,935	99,446
Total	<i>n/c</i>	<i>n/c</i>	99,383	60,535	104,749	101,846

n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.18b. Kentucky – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	19.5	20.3	20.3	21.6	21.9	23.1
Anxiety disorders	22.9	25.0	24.6	24.8	23.6	24.2
Attention deficit/hyperactivity disorder	14.7	14.3	13.5	13.2	12.5	12.6
Mood disorders	49.5	45.7	38.4	38.6	35.3	35.2
Bipolar disorders	19.8	15.6	8.9	8.8	8.0	7.8
Depressive disorders	30.5	30.7	29.9	30.4	27.9	28.1
Personality disorders	3.9	3.3	2.8	2.7	2.4	2.3
Schizophrenia and other psychotic disorders	8.0	7.7	7.8	7.8	7.3	7.3
Alcohol or substance-related disorders	3.5	5.0	18.6	18.9	23.6	23.8
Other mental disorders ²	20.5	17.8	26.8	25.7	25.2	24.9
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	48.2	47.8	47.7	48.1	49.5	50.0
Female	51.8	52.2	52.3	51.9	50.5	50.0
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	18.7	16.8	16.7	16.9	16.6	16.2
12 to 14 years	8.4	8.1	8.0	8.1	8.1	8.4
15 to 17 years	8.2	8.2	8.3	8.2	7.8	7.9
18 to 20 years	4.2	4.3	4.4	4.7	4.8	4.9
21 to 24 years	5.7	5.9	5.8	5.8	5.9	5.8
25 to 29 years	7.5	8.1	8.3	8.4	9.0	8.6
30 to 34 years	7.7	7.8	7.8	7.8	8.3	8.3
35 to 39 years	7.2	7.6	7.6	7.5	8.0	7.9
40 to 44 years	6.7	6.7	6.4	6.3	6.7	6.9
45 to 49 years	6.7	6.8	6.6	6.3	6.3	6.2
50 to 54 years	6.9	7.0	6.8	6.4	6.0	6.0
55 to 59 years	5.3	5.6	5.5	5.5	5.4	5.4
60 to 64 years	3.2	3.4	3.6	3.6	3.5	3.7
65 years and older	3.6	3.7	4.1	4.7	3.5	3.7
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.18b. Kentucky – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	†	†	†	†	†	†
Asian ⁴	†	†	†	†	†	†
Black or African American	10.4	10.2	10.2	9.7	9.9	9.6
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	82.5	82.6	82.3	81.2	80.0	80.0
Other	6.7	6.6	6.9	8.5	9.5	9.9
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	98.0	97.8	97.7	97.8	97.5	97.4
Hispanic or Latino origin	2.0	2.2	2.3	2.2	2.5	2.6
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	44.6	40.9	40.7	40.6	39.8	38.5
Puerto Rican	17.6	15.1	14.3	14.8	13.5	12.4
Cuban	8.4	8.1	8.4	8.4	8.0	7.8
Other specific	29.4	36.0	36.6	36.2	38.7	41.3
Total	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	2.4	1.9	2.7	2.4
Not a veteran	<i>n/c</i>	<i>n/c</i>	97.6	98.1	97.3	97.6
Total	<i>n/c</i>	<i>n/c</i>	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals; n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.19a. Louisiana – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	43,831	34,508	49,302	39,413	36,271	35,891
Mental health diagnosis¹						
Trauma- and stressor-related disorders	2,274	1,520	2,429	1,934	1,600	1,907
Anxiety disorders	3,950	2,726	4,270	3,973	2,924	3,904
Attention deficit/hyperactivity disorder	3,749	2,324	2,729	2,456	2,120	1,879
Mood disorders	24,351	12,370	14,285	12,973	10,935	9,702
Bipolar disorders	9,606	4,742	4,577	3,858	3,407	3,228
Depressive disorders	15,011	7,771	9,910	9,289	7,626	6,656
Personality disorders	1,651	594	428	379	235	288
Schizophrenia and other psychotic disorders	8,132	5,086	5,154	4,805	4,459	3,934
Alcohol or substance-related disorders	--	--	--	--	1	1
Other mental disorders ²	10,081	5,104	3,328	3,060	2,100	2,174
Total	40,983	23,569	25,560	23,002	20,333	17,526
Gender						
Male	19,128	15,482	22,354	17,869	16,222	15,925
Female	23,927	18,747	26,641	21,438	19,954	19,882
Total	43,055	34,229	48,995	39,307	36,176	35,807
Age³						
0 to 11 years	2,996	2,601	3,208	2,318	2,245	2,316
12 to 14 years	1,946	1,551	2,145	1,552	1,533	1,656
15 to 17 years	1,972	1,434	2,230	1,644	1,584	1,672
18 to 20 years	1,585	1,215	1,854	1,377	1,314	1,287
21 to 24 years	2,537	2,030	2,975	2,198	1,853	1,876
25 to 29 years	3,875	3,081	4,606	3,664	3,182	3,031
30 to 34 years	4,107	3,311	4,950	3,957	3,480	3,353
35 to 39 years	3,753	3,116	4,743	3,921	3,439	3,395
40 to 44 years	3,496	2,713	3,868	3,226	2,967	2,912
45 to 49 years	4,110	2,995	4,169	3,347	2,920	2,774
50 to 54 years	4,926	3,692	4,923	4,025	3,573	3,268
55 to 59 years	4,242	3,355	4,668	3,861	3,558	3,553
60 to 64 years	2,521	2,023	2,900	2,564	2,636	2,668
65 years and older	1,764	1,391	2,063	1,759	1,987	2,129
Total	43,830	34,508	49,302	39,413	36,271	35,890

Continued. See notes at end of table.

Table 9.19a. Louisiana – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	243	218	367	351	312	286
Asian ⁴	147	121	138	117	125	140
Black or African American	19,156	15,056	20,907	16,553	15,180	15,048
Native Hawaiian or Other Pacific Islander	63	31	55	42	63	91
White	18,976	15,302	22,603	18,067	16,156	17,200
Other	567	417	659	1,917	569	632
Total	39,152	31,145	44,729	37,047	32,405	33,397
Ethnicity						
Not of Hispanic or Latino origin	36,375	30,469	44,420	36,270	32,703	32,647
Hispanic or Latino origin	786	842	1,244	1,035	997	1,089
Total	37,161	31,311	45,664	37,305	33,700	33,736
Specific Hispanic or Latino origin						
Mexican	19	38	70	44	32	40
Puerto Rican	12	20	30	23	19	22
Cuban	13	15	17	14	9	13
Other specific	18	35	56	45	45	52
Total	62	108	173	126	105	127
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	‡	459	324	1,855
Not a veteran	<i>n/c</i>	<i>n/c</i>	‡	28,170	24,027	23,363
Total	<i>n/c</i>	<i>n/c</i>	‡	28,629	24,351	25,218

- - Quantity is zero; ‡ No data submitted; n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.19b. Louisiana – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	5.5	6.4	9.5	8.4	7.9	10.9
Anxiety disorders	9.6	11.6	16.7	17.3	14.4	22.3
Attention deficit/hyperactivity disorder	9.1	9.9	10.7	10.7	10.4	10.7
Mood disorders	59.4	52.5	55.9	56.4	53.8	55.4
Bipolar disorders	23.4	20.1	17.9	16.8	16.8	18.4
Depressive disorders	36.6	33.0	38.8	40.4	37.5	38.0
Personality disorders	4.0	2.5	1.7	1.6	1.2	1.6
Schizophrenia and other psychotic disorders	19.8	21.6	20.2	20.9	21.9	22.4
Alcohol or substance-related disorders	--	--	--	--	†	†
Other mental disorders ²	24.6	21.7	13.0	13.3	10.3	12.4
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	44.4	45.2	45.6	45.5	44.8	44.5
Female	55.6	54.8	54.4	54.5	55.2	55.5
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	6.8	7.5	6.5	5.9	6.2	6.5
12 to 14 years	4.4	4.5	4.4	3.9	4.2	4.6
15 to 17 years	4.5	4.2	4.5	4.2	4.4	4.7
18 to 20 years	3.6	3.5	3.8	3.5	3.6	3.6
21 to 24 years	5.8	5.9	6.0	5.6	5.1	5.2
25 to 29 years	8.8	8.9	9.3	9.3	8.8	8.4
30 to 34 years	9.4	9.6	10.0	10.0	9.6	9.3
35 to 39 years	8.6	9.0	9.6	9.9	9.5	9.5
40 to 44 years	8.0	7.9	7.8	8.2	8.2	8.1
45 to 49 years	9.4	8.7	8.5	8.5	8.1	7.7
50 to 54 years	11.2	10.7	10.0	10.2	9.9	9.1
55 to 59 years	9.7	9.7	9.5	9.8	9.8	9.9
60 to 64 years	5.8	5.9	5.9	6.5	7.3	7.4
65 years and older	4.0	4.0	4.2	4.5	5.5	5.9
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.19b. Louisiana – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	†	†	†	†	†	†
Asian ⁴	†	†	†	†	†	†
Black or African American	48.9	48.3	46.7	44.7	46.8	45.1
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	48.5	49.1	50.5	48.8	49.9	51.5
Other	1.4	1.3	1.5	5.2	1.8	1.9
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	97.9	97.3	97.3	97.2	97.0	96.8
Hispanic or Latino origin	2.1	2.7	2.7	2.8	3.0	3.2
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	30.6	35.2	40.5	34.9	30.5	31.5
Puerto Rican	19.4	18.5	17.3	18.3	18.1	17.3
Cuban	21.0	13.9	9.8	11.1	8.6	10.2
Other specific	29.0	32.4	32.4	35.7	42.9	40.9
Total	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	‡	1.6	1.3	7.4
Not a veteran	<i>n/c</i>	<i>n/c</i>	‡	98.4	98.7	92.6
Total	<i>n/c</i>	<i>n/c</i>	‡	100.0	100.0	100.0

† Less than 1 percent of individuals; - - Quantity is zero; ‡ No data submitted; *n/c* Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.20a. Maine – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	‡	‡	49,502	‡	‡	‡
Mental health diagnosis¹						
Trauma- and stressor-related disorders	‡	‡	5,275	‡	‡	‡
Anxiety disorders	‡	‡	4,573	‡	‡	‡
Attention deficit/hyperactivity disorder	‡	‡	795	‡	‡	‡
Mood disorders	‡	‡	11,300	‡	‡	‡
Bipolar disorders	‡	‡	4,246	‡	‡	‡
Depressive disorders	‡	‡	7,156	‡	‡	‡
Personality disorders	‡	‡	20	‡	‡	‡
Schizophrenia and other psychotic disorders	‡	‡	1,342	‡	‡	‡
Alcohol or substance-related disorders	‡	‡	1,672	‡	‡	‡
Other mental disorders ²	‡	‡	512	‡	‡	‡
Total	‡	‡	18,855	‡	‡	‡
Gender						
Male	‡	‡	17,394	‡	‡	‡
Female	‡	‡	25,677	‡	‡	‡
Total	‡	‡	43,071	‡	‡	‡
Age³						
0 to 11 years	‡	‡	3	‡	‡	‡
12 to 14 years	‡	‡	--	‡	‡	‡
15 to 17 years	‡	‡	--	‡	‡	‡
18 to 20 years	‡	‡	382	‡	‡	‡
21 to 24 years	‡	‡	2,210	‡	‡	‡
25 to 29 years	‡	‡	4,815	‡	‡	‡
30 to 34 years	‡	‡	5,651	‡	‡	‡
35 to 39 years	‡	‡	5,800	‡	‡	‡
40 to 44 years	‡	‡	5,184	‡	‡	‡
45 to 49 years	‡	‡	5,209	‡	‡	‡
50 to 54 years	‡	‡	5,455	‡	‡	‡
55 to 59 years	‡	‡	5,285	‡	‡	‡
60 to 64 years	‡	‡	4,015	‡	‡	‡
65 years and older	‡	‡	5,489	‡	‡	‡
Total	‡	‡	49,498	‡	‡	‡

Continued. See notes at end of table.

Table 9.20a. Maine – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	‡	‡	15	‡	‡	‡
Asian ⁴	‡	‡	3	‡	‡	‡
Black or African American	‡	‡	10	‡	‡	‡
Native Hawaiian or Other Pacific Islander	‡	‡	--	‡	‡	‡
White	‡	‡	1,070	‡	‡	‡
Other	‡	‡	4	‡	‡	‡
Total	‡	‡	1,102	‡	‡	‡
Ethnicity						
Not of Hispanic or Latino origin	‡	‡	26	‡	‡	‡
Hispanic or Latino origin	‡	‡	8	‡	‡	‡
Total	‡	‡	34	‡	‡	‡
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡
Not a veteran	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡
Total	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡

-- Quantity is zero; ‡ No data submitted; n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.20b. Maine – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	‡	‡	100.0	‡	‡	‡
Mental health diagnosis¹						
Trauma- and stressor-related disorders	‡	‡	28.0	‡	‡	‡
Anxiety disorders	‡	‡	24.3	‡	‡	‡
Attention deficit/hyperactivity disorder	‡	‡	4.2	‡	‡	‡
Mood disorders	‡	‡	59.9	‡	‡	‡
Bipolar disorders	‡	‡	22.5	‡	‡	‡
Depressive disorders	‡	‡	38.0	‡	‡	‡
Personality disorders	‡	‡	†	‡	‡	‡
Schizophrenia and other psychotic disorders	‡	‡	7.1	‡	‡	‡
Alcohol or substance-related disorders	‡	‡	8.9	‡	‡	‡
Other mental disorders ²	‡	‡	2.7	‡	‡	‡
Total	‡	‡	100.0	‡	‡	‡
Gender						
Male	‡	‡	40.4	‡	‡	‡
Female	‡	‡	59.6	‡	‡	‡
Total	‡	‡	100.0	‡	‡	‡
Age³						
0 to 11 years	‡	‡	†	‡	‡	‡
12 to 14 years	‡	‡	--	‡	‡	‡
15 to 17 years	‡	‡	--	‡	‡	‡
18 to 20 years	‡	‡	†	‡	‡	‡
21 to 24 years	‡	‡	4.5	‡	‡	‡
25 to 29 years	‡	‡	9.7	‡	‡	‡
30 to 34 years	‡	‡	11.4	‡	‡	‡
35 to 39 years	‡	‡	11.7	‡	‡	‡
40 to 44 years	‡	‡	10.5	‡	‡	‡
45 to 49 years	‡	‡	10.5	‡	‡	‡
50 to 54 years	‡	‡	11.0	‡	‡	‡
55 to 59 years	‡	‡	10.7	‡	‡	‡
60 to 64 years	‡	‡	8.1	‡	‡	‡
65 years and older	‡	‡	11.1	‡	‡	‡
Total	‡	‡	100.0	‡	‡	‡

Continued. See notes at end of table.

Table 9.20b. Maine – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	‡	‡	1.4	‡	‡	‡
Asian ⁴	‡	‡	†	‡	‡	‡
Black or African American	‡	‡	†	‡	‡	‡
Native Hawaiian or Other Pacific Islander	‡	‡	--	‡	‡	‡
White	‡	‡	97.1	‡	‡	‡
Other	‡	‡	†	‡	‡	‡
Total	‡	‡	100.0	‡	‡	‡
Ethnicity						
Not of Hispanic or Latino origin	‡	‡	76.5	‡	‡	‡
Hispanic or Latino origin	‡	‡	23.5	‡	‡	‡
Total	‡	‡	100.0	‡	‡	‡
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡
Not a veteran	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡
Total	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡

† Less than 1 percent of individuals; -- Quantity is zero; ‡ No data submitted; *n/c* Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.21a. Maryland – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	185,306	202,972	201,718	213,605	226,003	‡
Mental health diagnosis¹						
Trauma- and stressor-related disorders	30,403	30,946	34,607	38,031	41,022	‡
Anxiety disorders	27,854	31,961	36,749	41,593	45,782	‡
Attention deficit/hyperactivity disorder	27,608	29,601	31,481	33,333	34,894	‡
Mood disorders	85,256	76,706	82,958	88,592	94,927	‡
Bipolar disorders	31,771	24,814	26,024	27,614	28,830	‡
Depressive disorders	54,014	52,257	57,315	61,451	66,739	‡
Personality disorders	1,165	904	933	1,081	1,136	‡
Schizophrenia and other psychotic disorders	15,498	15,870	16,326	16,785	16,968	‡
Alcohol or substance-related disorders	2,372	54	3,294	7	3	‡
Other mental disorders ²	18,576	25,479	27,076	27,827	28,643	‡
Total	180,276	184,895	200,329	211,910	223,895	‡
Gender						
Male	85,362	91,880	92,962	98,289	103,027	‡
Female	99,938	111,078	108,740	115,299	122,963	‡
Total	185,300	202,958	201,702	213,588	225,990	‡
Age³						
0 to 11 years	31,162	32,406	33,965	37,514	39,137	‡
12 to 14 years	15,696	15,987	16,479	17,837	19,563	‡
15 to 17 years	15,033	15,844	16,261	16,696	17,856	‡
18 to 20 years	8,715	9,567	9,813	10,545	11,430	‡
21 to 24 years	10,746	12,189	11,502	12,061	12,662	‡
25 to 29 years	17,359	19,999	19,580	20,952	22,145	‡
30 to 34 years	16,286	18,166	17,950	19,212	20,978	‡
35 to 39 years	12,827	14,667	14,931	15,903	17,316	‡
40 to 44 years	11,650	12,244	11,528	12,030	12,588	‡
45 to 49 years	13,184	14,201	13,036	13,051	12,805	‡
50 to 54 years	14,119	15,672	14,493	14,470	14,368	‡
55 to 59 years	10,430	12,418	12,059	12,466	13,226	‡
60 to 64 years	5,937	7,123	7,282	7,687	8,387	‡
65 years and older	2,160	2,487	2,838	3,178	3,538	‡
Total	185,304	202,970	201,717	213,602	225,999	‡

Continued. See notes at end of table.

Table 9.21a. Maryland – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	739	843	823	948	1,066	‡
Asian ⁴	2,044	2,445	2,508	2,864	3,080	‡
Black or African American	86,719	93,642	92,424	98,374	105,933	‡
Native Hawaiian or Other Pacific Islander	332	413	436	586	667	‡
White	87,817	95,431	94,891	100,152	103,470	‡
Other	3,094	3,594	4,015	4,464	4,704	‡
Total	180,745	196,368	195,097	207,388	218,920	‡
Ethnicity						
Not of Hispanic or Latino origin	157,031	166,800	169,940	180,624	187,886	‡
Hispanic or Latino origin	16,340	18,272	19,971	22,700	22,100	‡
Total	173,371	185,072	189,911	203,324	209,986	‡
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	3,858	4,067	4,152	‡
Not a veteran	<i>n/c</i>	<i>n/c</i>	109,295	129,180	137,011	‡
Total	<i>n/c</i>	<i>n/c</i>	113,153	133,247	141,163	‡

‡ No data submitted; n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.21b. Maryland – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	100.0	100.0	100.0	100.0	100.0	‡
Mental health diagnosis¹						
Trauma- and stressor-related disorders	16.9	16.7	17.3	17.9	18.3	‡
Anxiety disorders	15.5	17.3	18.3	19.6	20.4	‡
Attention deficit/hyperactivity disorder	15.3	16.0	15.7	15.7	15.6	‡
Mood disorders	47.3	41.5	41.4	41.8	42.4	‡
Bipolar disorders	17.6	13.4	13.0	13.0	12.9	‡
Depressive disorders	30.0	28.3	28.6	29.0	29.8	‡
Personality disorders	†	†	†	†	†	‡
Schizophrenia and other psychotic disorders	8.6	8.6	8.1	7.9	7.6	‡
Alcohol or substance-related disorders	1.3	†	1.6	†	†	‡
Other mental disorders ²	10.3	13.8	13.5	13.1	12.8	‡
Total	100.0	100.0	100.0	100.0	100.0	‡
Gender						
Male	46.1	45.3	46.1	46.0	45.6	‡
Female	53.9	54.7	53.9	54.0	54.4	‡
Total	100.0	100.0	100.0	100.0	100.0	‡
Age³						
0 to 11 years	16.8	16.0	16.8	17.6	17.3	‡
12 to 14 years	8.5	7.9	8.2	8.4	8.7	‡
15 to 17 years	8.1	7.8	8.1	7.8	7.9	‡
18 to 20 years	4.7	4.7	4.9	4.9	5.1	‡
21 to 24 years	5.8	6.0	5.7	5.6	5.6	‡
25 to 29 years	9.4	9.9	9.7	9.8	9.8	‡
30 to 34 years	8.8	9.0	8.9	9.0	9.3	‡
35 to 39 years	6.9	7.2	7.4	7.4	7.7	‡
40 to 44 years	6.3	6.0	5.7	5.6	5.6	‡
45 to 49 years	7.1	7.0	6.5	6.1	5.7	‡
50 to 54 years	7.6	7.7	7.2	6.8	6.4	‡
55 to 59 years	5.6	6.1	6.0	5.8	5.9	‡
60 to 64 years	3.2	3.5	3.6	3.6	3.7	‡
65 years and older	1.2	1.2	1.4	1.5	1.6	‡
Total	100.0	100.0	100.0	100.0	100.0	‡

Continued. See notes at end of table.

Table 9.21b. Maryland – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	†	†	†	†	†	‡
Asian ⁴	1.1	1.2	1.3	1.4	1.4	‡
Black or African American	48.0	47.7	47.4	47.4	48.4	‡
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	‡
White	48.6	48.6	48.6	48.3	47.3	‡
Other	1.7	1.8	2.1	2.2	2.1	‡
Total	100.0	100.0	100.0	100.0	100.0	‡
Ethnicity						
Not of Hispanic or Latino origin	90.6	90.1	89.5	88.8	89.5	‡
Hispanic or Latino origin	9.4	9.9	10.5	11.2	10.5	‡
Total	100.0	100.0	100.0	100.0	100.0	‡
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	3.4	3.1	2.9	‡
Not a veteran	<i>n/c</i>	<i>n/c</i>	96.6	96.9	97.1	‡
Total	<i>n/c</i>	<i>n/c</i>	100.0	100.0	100.0	‡

† Less than 1 percent of individuals; ‡ No data submitted; n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

**Table 9.22a. Massachusetts – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Number, 2015–2020**

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	33,556	33,921	31,659	26,383	26,275	24,755
Mental health diagnosis¹						
Trauma- and stressor-related disorders	6,273	2,266	2,877	1,593	1,467	1,138
Anxiety disorders	3,459	1,370	2,110	2,078	1,924	1,413
Attention deficit/hyperactivity disorder	2,086	541	882	753	728	481
Mood disorders	14,726	3,982	5,655	4,224	3,847	3,061
Bipolar disorders	8,671	1,710	2,352	1,930	1,756	1,501
Depressive disorders	6,308	2,321	3,404	2,414	2,189	1,644
Personality disorders	2,031	3	12	228	237	206
Schizophrenia and other psychotic disorders	11,621	3,013	3,951	3,567	3,452	3,156
Alcohol or substance-related disorders	1	--	1	--	--	--
Other mental disorders ²	3,043	1,728	2,083	1,245	1,210	939
Total	31,022	10,056	13,173	9,249	8,557	6,898
Gender						
Male	18,712	18,948	17,680	14,760	14,832	13,923
Female	14,843	14,973	13,979	11,623	11,443	10,832
Total	33,555	33,921	31,659	26,383	26,275	24,755
Age³						
0 to 11 years	1,967	1,924	1,681	1,266	1,178	646
12 to 14 years	1,471	1,618	1,393	985	881	604
15 to 17 years	1,861	1,822	1,621	1,009	1,006	853
18 to 20 years	1,432	1,512	1,380	1,027	1,022	869
21 to 24 years	2,245	2,136	1,962	1,544	1,489	1,436
25 to 29 years	2,994	3,156	2,848	2,179	2,250	2,163
30 to 34 years	2,798	2,873	2,676	2,147	2,185	2,160
35 to 39 years	2,420	2,497	2,437	2,010	2,051	2,085
40 to 44 years	2,384	2,277	2,060	1,764	1,837	1,816
45 to 49 years	2,940	2,858	2,603	2,126	2,045	1,877
50 to 54 years	3,707	3,588	3,325	2,768	2,586	2,424
55 to 59 years	3,408	3,484	3,325	3,092	3,053	2,995
60 to 64 years	2,239	2,356	2,423	2,418	2,532	2,531
65 years and older	1,690	1,820	1,925	2,046	2,159	2,295
Total	33,556	33,921	31,659	26,381	26,274	24,754

Continued. See notes at end of table.

**Table 9.22a. Massachusetts – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Number, 2015–2020 (continued)**

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	108	110	100	69	72	76
Asian ⁴	633	619	615	593	599	594
Black or African American	4,392	4,576	4,401	3,757	3,844	3,635
Native Hawaiian or Other Pacific Islander	20	15	14	12	19	13
White	22,554	22,362	20,699	15,958	16,011	15,782
Other	591	604	520	361	405	403
Total	28,298	28,286	26,349	20,750	20,950	20,503
Ethnicity						
Not of Hispanic or Latino origin	14,993	24,502	22,389	17,139	18,111	17,800
Hispanic or Latino origin	3,165	2,942	2,248	2,042	2,059	1,942
Total	18,158	27,444	24,637	19,181	20,170	19,742
Specific Hispanic or Latino origin						
Mexican	19	17	19	18	15	17
Puerto Rican	1,027	889	991	810	821	782
Cuban	--	--	--	--	--	--
Other specific	449	1,377	422	407	402	357
Total	1,495	2,283	1,432	1,235	1,238	1,156
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	678	570	564	521
Not a veteran	<i>n/c</i>	<i>n/c</i>	21,364	19,263	19,465	19,219
Total	<i>n/c</i>	<i>n/c</i>	22,042	19,833	20,029	19,740

-- Quantity is zero; n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

**Table 9.22b. Massachusetts – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Column percent distribution, 2015–2020**

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	20.2	22.5	21.8	17.2	17.1	16.5
Anxiety disorders	11.2	13.6	16.0	22.5	22.5	20.5
Attention deficit/hyperactivity disorder	6.7	5.4	6.7	8.1	8.5	7.0
Mood disorders	47.5	39.6	42.9	45.7	45.0	44.4
Bipolar disorders	28.0	17.0	17.9	20.9	20.5	21.8
Depressive disorders	20.3	23.1	25.8	26.1	25.6	23.8
Personality disorders	6.5	†	†	2.5	2.8	3.0
Schizophrenia and other psychotic disorders	37.5	30.0	30.0	38.6	40.3	45.8
Alcohol or substance-related disorders	†	--	†	--	--	--
Other mental disorders ²	9.8	17.2	15.8	13.5	14.1	13.6
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	55.8	55.9	55.8	55.9	56.4	56.2
Female	44.2	44.1	44.2	44.1	43.6	43.8
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	5.9	5.7	5.3	4.8	4.5	2.6
12 to 14 years	4.4	4.8	4.4	3.7	3.4	2.4
15 to 17 years	5.5	5.4	5.1	3.8	3.8	3.4
18 to 20 years	4.3	4.5	4.4	3.9	3.9	3.5
21 to 24 years	6.7	6.3	6.2	5.9	5.7	5.8
25 to 29 years	8.9	9.3	9.0	8.3	8.6	8.7
30 to 34 years	8.3	8.5	8.5	8.1	8.3	8.7
35 to 39 years	7.2	7.4	7.7	7.6	7.8	8.4
40 to 44 years	7.1	6.7	6.5	6.7	7.0	7.3
45 to 49 years	8.8	8.4	8.2	8.1	7.8	7.6
50 to 54 years	11.0	10.6	10.5	10.5	9.8	9.8
55 to 59 years	10.2	10.3	10.5	11.7	11.6	12.1
60 to 64 years	6.7	6.9	7.7	9.2	9.6	10.2
65 years and older	5.0	5.4	6.1	7.8	8.2	9.3
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

**Table 9.22b. Massachusetts – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Column percent distribution, 2015–2020 (continued)**

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	†	†	†	†	†	†
Asian ⁴	2.2	2.2	2.3	2.9	2.9	2.9
Black or African American	15.5	16.2	16.7	18.1	18.3	17.7
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	79.7	79.1	78.6	76.9	76.4	77.0
Other	2.1	2.1	2.0	1.7	1.9	2.0
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	82.6	89.3	90.9	89.4	89.8	90.2
Hispanic or Latino origin	17.4	10.7	9.1	10.6	10.2	9.8
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	1.3	†	1.3	1.5	1.2	1.5
Puerto Rican	68.7	38.9	69.2	65.6	66.3	67.6
Cuban	--	--	--	--	--	--
Other specific	30.0	60.3	29.5	33.0	32.5	30.9
Total	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	3.1	2.9	2.8	2.6
Not a veteran	<i>n/c</i>	<i>n/c</i>	96.9	97.1	97.2	97.4
Total	<i>n/c</i>	<i>n/c</i>	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals; -- Quantity is zero; n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.23a. Michigan – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020

<i>Selected characteristics</i>	2015	2016	2017	2018	2019	2020
Total individuals	‡	‡	177,289	231,953	244,168	224,250
Mental health diagnosis¹						
Trauma- and stressor-related disorders	‡	‡	23,546	35,541	43,451	40,535
Anxiety disorders	‡	‡	33,824	47,125	50,718	45,722
Attention deficit/hyperactivity disorder	‡	‡	21,955	29,330	30,749	28,347
Mood disorders	‡	‡	72,706	98,338	103,469	90,025
Bipolar disorders	‡	‡	27,954	35,786	36,497	32,606
Depressive disorders	‡	‡	47,208	66,185	71,111	61,284
Personality disorders	‡	‡	11,879	15,587	16,609	15,136
Schizophrenia and other psychotic disorders	‡	‡	29,582	35,736	35,993	34,612
Alcohol or substance-related disorders	‡	‡	9,591	13,849	11,642	7,259
Other mental disorders ²	‡	‡	72,836	92,937	93,343	84,571
Total	‡	‡	170,998	224,703	233,394	207,770
Gender						
Male	‡	‡	92,096	119,664	126,010	116,961
Female	‡	‡	85,193	112,289	118,158	107,289
Total	‡	‡	177,289	231,953	244,168	224,250
Age³						
0 to 11 years	‡	‡	21,897	30,567	32,982	30,969
12 to 14 years	‡	‡	9,996	13,291	14,044	13,307
15 to 17 years	‡	‡	10,613	14,839	15,925	13,778
18 to 20 years	‡	‡	7,748	10,664	12,049	10,157
21 to 24 years	‡	‡	11,695	14,608	14,906	13,578
25 to 29 years	‡	‡	15,876	21,425	22,898	20,940
30 to 34 years	‡	‡	14,596	19,184	20,306	18,996
35 to 39 years	‡	‡	13,426	17,646	18,753	17,170
40 to 44 years	‡	‡	12,339	15,441	16,025	14,987
45 to 49 years	‡	‡	13,865	17,176	17,449	15,435
50 to 54 years	‡	‡	15,191	18,412	18,131	16,138
55 to 59 years	‡	‡	13,541	17,075	17,449	16,181
60 to 64 years	‡	‡	8,650	11,209	11,864	11,729
65 years and older	‡	‡	7,854	10,416	11,386	10,883
Total	‡	‡	177,287	231,953	244,167	224,248

Continued. See notes at end of table.

Table 9.23a. Michigan – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	‡	‡	1,123	1,418	1,530	1,322
Asian ⁴	‡	‡	754	974	1,111	1,139
Black or African American	‡	‡	34,398	46,258	52,620	51,695
Native Hawaiian or Other Pacific Islander	‡	‡	92	139	160	151
White	‡	‡	107,077	137,522	144,985	135,648
Other	‡	‡	32,534	43,805	41,104	32,798
Total	‡	‡	175,978	230,116	241,510	222,753
Ethnicity						
Not of Hispanic or Latino origin	‡	‡	151,252	198,528	206,993	191,552
Hispanic or Latino origin	‡	‡	9,326	13,494	15,339	13,792
Total	‡	‡	160,578	212,022	222,332	205,344
Specific Hispanic or Latino origin						
Mexican	‡	‡	2,088	3,351	3,815	3,678
Puerto Rican	‡	‡	662	993	1,181	1,113
Cuban	‡	‡	98	147	182	155
Other specific	‡	‡	1,085	1,625	1,917	2,527
Total	‡	‡	3,933	6,116	7,095	7,473
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	‡	3,858	4,101	3,833
Not a veteran	<i>n/c</i>	<i>n/c</i>	‡	159,936	164,616	151,917
Total	<i>n/c</i>	<i>n/c</i>	‡	163,794	168,717	155,750

‡ No data submitted; n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.23b. Michigan – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	‡	‡	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	‡	‡	13.8	15.8	18.6	19.5
Anxiety disorders	‡	‡	19.8	21.0	21.7	22.0
Attention deficit/hyperactivity disorder	‡	‡	12.8	13.1	13.2	13.6
Mood disorders	‡	‡	42.5	43.8	44.3	43.3
Bipolar disorders	‡	‡	16.3	15.9	15.6	15.7
Depressive disorders	‡	‡	27.6	29.5	30.5	29.5
Personality disorders	‡	‡	6.9	6.9	7.1	7.3
Schizophrenia and other psychotic disorders	‡	‡	17.3	15.9	15.4	16.7
Alcohol or substance-related disorders	‡	‡	5.6	6.2	5.0	3.5
Other mental disorders ²	‡	‡	42.6	41.4	40.0	40.7
Total	‡	‡	100.0	100.0	100.0	100.0
Gender						
Male	‡	‡	51.9	51.6	51.6	52.2
Female	‡	‡	48.1	48.4	48.4	47.8
Total	‡	‡	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	‡	‡	12.4	13.2	13.5	13.8
12 to 14 years	‡	‡	5.6	5.7	5.8	5.9
15 to 17 years	‡	‡	6.0	6.4	6.5	6.1
18 to 20 years	‡	‡	4.4	4.6	4.9	4.5
21 to 24 years	‡	‡	6.6	6.3	6.1	6.1
25 to 29 years	‡	‡	9.0	9.2	9.4	9.3
30 to 34 years	‡	‡	8.2	8.3	8.3	8.5
35 to 39 years	‡	‡	7.6	7.6	7.7	7.7
40 to 44 years	‡	‡	7.0	6.7	6.6	6.7
45 to 49 years	‡	‡	7.8	7.4	7.1	6.9
50 to 54 years	‡	‡	8.6	7.9	7.4	7.2
55 to 59 years	‡	‡	7.6	7.4	7.1	7.2
60 to 64 years	‡	‡	4.9	4.8	4.9	5.2
65 years and older	‡	‡	4.4	4.5	4.7	4.9
Total	‡	‡	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.23b. Michigan – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	‡	‡	†	†	†	†
Asian ⁴	‡	‡	†	†	†	†
Black or African American	‡	‡	19.5	20.1	21.8	23.2
Native Hawaiian or Other Pacific Islander	‡	‡	†	†	†	†
White	‡	‡	60.8	59.8	60.0	60.9
Other	‡	‡	18.5	19.0	17.0	14.7
Total	‡	‡	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	‡	‡	94.2	93.6	93.1	93.3
Hispanic or Latino origin	‡	‡	5.8	6.4	6.9	6.7
Total	‡	‡	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	‡	‡	53.1	54.8	53.8	49.2
Puerto Rican	‡	‡	16.8	16.2	16.6	14.9
Cuban	‡	‡	2.5	2.4	2.6	2.1
Other specific	‡	‡	27.6	26.6	27.0	33.8
Total	‡	‡	100.0	100.0	100.0	100.0
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	‡	2.4	2.4	2.5
Not a veteran	<i>n/c</i>	<i>n/c</i>	‡	97.6	97.6	97.5
Total	<i>n/c</i>	<i>n/c</i>	‡	100.0	100.0	100.0

† Less than 1 percent of individuals; ‡ No data submitted; n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.24a. Minnesota – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	225,786	259,024	270,651	287,962	287,670	318,448
Mental health diagnosis¹						
Trauma- and stressor-related disorders	65,164	72,479	75,755	101,004	99,344	107,500
Anxiety disorders	74,419	86,835	103,951	126,735	134,815	147,707
Attention deficit/hyperactivity disorder	34,335	37,537	40,416	46,677	47,028	48,532
Mood disorders	118,381	123,482	116,241	143,475	130,883	146,212
Bipolar disorders	30,328	27,502	21,819	29,132	22,374	24,968
Depressive disorders	94,482	100,686	97,879	121,450	113,168	126,039
Personality disorders	12,147	12,507	14,494	14,516	14,802	17,267
Schizophrenia and other psychotic disorders	20,594	20,650	21,342	24,127	21,995	25,217
Alcohol or substance-related disorders	1,277	1,206	139	488	44	120
Other mental disorders ²	45,161	53,481	49,850	63,333	58,465	58,049
Total	215,348	238,763	245,172	269,422	261,644	284,601
Gender						
Male	102,433	117,156	121,791	129,777	129,708	142,328
Female	123,353	141,868	148,860	158,184	157,962	175,961
Total	225,786	259,024	270,651	287,961	287,670	318,289
Age³						
0 to 11 years	31,865	40,307	41,832	45,172	49,379	51,594
12 to 14 years	13,824	16,738	17,271	18,702	20,490	22,135
15 to 17 years	13,623	16,550	17,468	18,931	19,770	20,967
18 to 20 years	9,273	10,503	11,332	12,472	13,361	14,372
21 to 24 years	13,241	14,918	15,081	15,965	15,674	16,536
25 to 29 years	20,810	23,946	25,190	26,864	26,560	28,357
30 to 34 years	20,752	23,380	24,316	25,603	24,932	27,683
35 to 39 years	17,271	19,790	21,340	22,861	22,625	25,336
40 to 44 years	15,906	17,019	17,400	17,992	17,559	20,243
45 to 49 years	16,266	17,361	17,881	18,120	16,793	18,534
50 to 54 years	17,999	19,226	19,090	19,102	17,061	18,708
55 to 59 years	14,423	16,163	17,298	18,339	17,198	19,766
60 to 64 years	9,498	10,937	11,833	12,921	12,098	15,152
65 years and older	11,016	12,133	13,279	14,896	14,158	19,062
Total	225,767	258,971	270,611	287,940	287,658	318,445

Continued. See notes at end of table.

Table 9.24a. Minnesota – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	8,986	9,699	9,854	10,579	10,994	12,075
Asian ⁴	6,492	7,334	7,894	8,552	8,806	9,862
Black or African American	29,332	32,493	33,616	35,146	35,978	39,400
Native Hawaiian or Other Pacific Islander	211	251	278	295	335	356
White	145,660	161,658	165,516	172,627	173,131	184,898
Other	8,375	10,232	11,311	12,623	13,375	14,545
Total	199,056	221,667	228,469	239,822	242,619	261,136
Ethnicity						
Not of Hispanic or Latino origin	209,901	239,409	249,560	248,890	247,561	257,296
Hispanic or Latino origin	12,956	15,375	16,232	16,627	17,554	37,001
Total	222,857	254,784	265,792	265,517	265,115	294,297
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡
Not a veteran	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡
Total	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡

‡ No data submitted; n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.24b. Minnesota – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	30.3	30.4	30.9	37.5	38.0	37.8
Anxiety disorders	34.6	36.4	42.4	47.0	51.5	51.9
Attention deficit/hyperactivity disorder	15.9	15.7	16.5	17.3	18.0	17.1
Mood disorders	55.0	51.7	47.4	53.3	50.0	51.4
Bipolar disorders	14.1	11.5	8.9	10.8	8.6	8.8
Depressive disorders	43.9	42.2	39.9	45.1	43.3	44.3
Personality disorders	5.6	5.2	5.9	5.4	5.7	6.1
Schizophrenia and other psychotic disorders	9.6	8.6	8.7	9.0	8.4	8.9
Alcohol or substance-related disorders	†	†	†	†	†	†
Other mental disorders ²	21.0	22.4	20.3	23.5	22.3	20.4
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	45.4	45.2	45.0	45.1	45.1	44.7
Female	54.6	54.8	55.0	54.9	54.9	55.3
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	14.1	15.6	15.5	15.7	17.2	16.2
12 to 14 years	6.1	6.5	6.4	6.5	7.1	7.0
15 to 17 years	6.0	6.4	6.5	6.6	6.9	6.6
18 to 20 years	4.1	4.1	4.2	4.3	4.6	4.5
21 to 24 years	5.9	5.8	5.6	5.5	5.4	5.2
25 to 29 years	9.2	9.2	9.3	9.3	9.2	8.9
30 to 34 years	9.2	9.0	9.0	8.9	8.7	8.7
35 to 39 years	7.6	7.6	7.9	7.9	7.9	8.0
40 to 44 years	7.0	6.6	6.4	6.2	6.1	6.4
45 to 49 years	7.2	6.7	6.6	6.3	5.8	5.8
50 to 54 years	8.0	7.4	7.1	6.6	5.9	5.9
55 to 59 years	6.4	6.2	6.4	6.4	6.0	6.2
60 to 64 years	4.2	4.2	4.4	4.5	4.2	4.8
65 years and older	4.9	4.7	4.9	5.2	4.9	6.0
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.24b. Minnesota – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	4.5	4.4	4.3	4.4	4.5	4.6
Asian ⁴	3.3	3.3	3.5	3.6	3.6	3.8
Black or African American	14.7	14.7	14.7	14.7	14.8	15.1
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	73.2	72.9	72.4	72.0	71.4	70.8
Other	4.2	4.6	5.0	5.3	5.5	5.6
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	94.2	94.0	93.9	93.7	93.4	87.4
Hispanic or Latino origin	5.8	6.0	6.1	6.3	6.6	12.6
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡
Not a veteran	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡
Total	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡

† Less than 1 percent of individuals; ‡ No data submitted; *n/c* Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.25a. Mississippi – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	90,279	83,432	73,867	85,449	87,406	102,061
Mental health diagnosis¹						
Trauma- and stressor-related disorders	7,316	8,322	8,007	10,823	11,473	14,273
Anxiety disorders	6,384	17,296	8,909	12,974	14,299	17,736
Attention deficit/hyperactivity disorder	14,049	15,771	13,009	17,071	17,706	20,989
Mood disorders	33,749	32,001	25,457	34,538	35,765	41,652
Bipolar disorders	11,502	10,985	7,814	10,398	10,527	12,179
Depressive disorders	22,294	21,171	17,868	24,488	25,560	29,870
Personality disorders	3,433	1,960	1,374	1,535	1,548	2,231
Schizophrenia and other psychotic disorders	12,091	10,822	8,000	10,995	11,310	14,574
Alcohol or substance-related disorders	4,284	7,269	7,288	12,545	13,237	14,904
Other mental disorders ²	14,197	13,871	14,886	18,070	19,064	24,120
Total	83,119	81,128	61,880	84,923	86,884	101,461
Gender						
Male	42,319	35,434	35,345	41,517	42,454	49,164
Female	47,922	40,366	38,479	43,874	44,932	52,852
Total	90,241	75,800	73,824	85,391	87,386	102,016
Age³						
0 to 11 years	14,145	13,408	13,901	15,514	14,926	15,561
12 to 14 years	7,560	6,969	6,627	7,571	8,056	9,593
15 to 17 years	7,832	7,248	6,448	7,049	6,996	8,201
18 to 20 years	4,655	4,485	3,363	3,998	4,303	5,233
21 to 24 years	4,311	3,905	3,221	4,008	4,107	4,946
25 to 29 years	5,863	5,261	4,760	6,250	6,420	7,485
30 to 34 years	6,343	5,660	4,857	6,223	6,496	7,468
35 to 39 years	6,015	5,616	4,938	6,232	6,557	7,492
40 to 44 years	5,636	5,121	4,185	5,098	5,347	6,512
45 to 49 years	6,012	5,395	4,592	5,141	5,219	5,998
50 to 54 years	6,970	6,250	4,972	5,394	5,295	6,210
55 to 59 years	6,253	5,787	4,871	5,313	5,411	6,597
60 to 64 years	4,005	3,861	3,382	3,736	4,015	5,147
65 years and older	4,679	4,466	3,750	3,921	4,258	5,617
Total	90,279	83,432	73,867	85,448	87,406	102,060

Continued. See notes at end of table.

Table 9.25a. Mississippi – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	177	132	142	212	226	227
Asian ⁴	135	122	112	145	164	168
Black or African American	46,392	37,102	38,963	42,640	43,412	51,007
Native Hawaiian or Other Pacific Islander	8	12	7	12	13	21
White	42,192	37,027	33,050	40,255	41,134	47,612
Other	933	839	912	1,215	1,321	1,739
Total	89,837	75,234	73,186	84,479	86,270	100,774
Ethnicity						
Not of Hispanic or Latino origin	88,811	73,444	69,909	79,512	80,612	95,029
Hispanic or Latino origin	840	874	1,099	1,336	1,489	1,821
Total	89,651	74,318	71,008	80,848	82,101	96,850
Specific Hispanic or Latino origin						
Mexican	395	395	413	537	608	711
Puerto Rican	86	80	75	85	93	115
Cuban	38	26	19	27	32	45
Other specific	--	--	--	--	--	210
Total	519	501	507	649	733	1,081
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	‡	1,172	1,227	1,473
Not a veteran	<i>n/c</i>	<i>n/c</i>	‡	52,422	54,055	64,864
Total	<i>n/c</i>	<i>n/c</i>	‡	53,594	55,282	66,337

-- Quantity is zero; ‡ No data submitted; n/c Not collected.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.25b. Mississippi – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	8.8	10.3	12.9	12.7	13.2	14.1
Anxiety disorders	7.7	21.3	14.4	15.3	16.5	17.5
Attention deficit/hyperactivity disorder	16.9	19.4	21.0	20.1	20.4	20.7
Mood disorders	40.6	39.4	41.1	40.7	41.2	41.1
Bipolar disorders	13.8	13.5	12.6	12.2	12.1	12.0
Depressive disorders	26.8	26.1	28.9	28.8	29.4	29.4
Personality disorders	4.1	2.4	2.2	1.8	1.8	2.2
Schizophrenia and other psychotic disorders	14.5	13.3	12.9	12.9	13.0	14.4
Alcohol or substance-related disorders	5.2	9.0	11.8	14.8	15.2	14.7
Other mental disorders ²	17.1	17.1	24.1	21.3	21.9	23.8
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	46.9	46.7	47.9	48.6	48.6	48.2
Female	53.1	53.3	52.1	51.4	51.4	51.8
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	15.7	16.1	18.8	18.2	17.1	15.2
12 to 14 years	8.4	8.4	9.0	8.9	9.2	9.4
15 to 17 years	8.7	8.7	8.7	8.2	8.0	8.0
18 to 20 years	5.2	5.4	4.6	4.7	4.9	5.1
21 to 24 years	4.8	4.7	4.4	4.7	4.7	4.8
25 to 29 years	6.5	6.3	6.4	7.3	7.3	7.3
30 to 34 years	7.0	6.8	6.6	7.3	7.4	7.3
35 to 39 years	6.7	6.7	6.7	7.3	7.5	7.3
40 to 44 years	6.2	6.1	5.7	6.0	6.1	6.4
45 to 49 years	6.7	6.5	6.2	6.0	6.0	5.9
50 to 54 years	7.7	7.5	6.7	6.3	6.1	6.1
55 to 59 years	6.9	6.9	6.6	6.2	6.2	6.5
60 to 64 years	4.4	4.6	4.6	4.4	4.6	5.0
65 years and older	5.2	5.4	5.1	4.6	4.9	5.5
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.25b. Mississippi – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	†	†	†	†	†	†
Asian ⁴	†	†	†	†	†	†
Black or African American	51.6	49.3	53.2	50.5	50.3	50.6
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	47.0	49.2	45.2	47.7	47.7	47.2
Other	1.0	1.1	1.2	1.4	1.5	1.7
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	99.1	98.8	98.5	98.3	98.2	98.1
Hispanic or Latino origin	†	1.2	1.5	1.7	1.8	1.9
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	76.1	78.8	81.5	82.7	82.9	65.8
Puerto Rican	16.6	16.0	14.8	13.1	12.7	10.6
Cuban	7.3	5.2	3.7	4.2	4.4	4.2
Other specific	--	--	--	--	--	19.4
Total	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	‡	2.2	2.2	2.2
Not a veteran	<i>n/c</i>	<i>n/c</i>	‡	97.8	97.8	97.8
Total	<i>n/c</i>	<i>n/c</i>	‡	100.0	100.0	100.0

† Less than 1 percent of individuals; -- Quantity is zero; ‡ No data submitted; *n/c* Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.26a. Missouri – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	75,472	76,268	77,497	76,409	80,771	82,117
Mental health diagnosis¹						
Trauma- and stressor-related disorders	7,604	4,517	5,631	6,157	7,087	6,638
Anxiety disorders	6,241	9,745	11,100	11,386	12,552	10,943
Attention deficit/hyperactivity disorder	6,291	6,779	7,779	7,828	8,517	7,338
Mood disorders	40,831	36,858	39,341	38,955	40,783	37,480
Bipolar disorders	19,231	15,713	16,097	15,299	15,630	13,219
Depressive disorders	23,335	22,784	24,992	25,766	27,660	25,147
Personality disorders	547	700	970	1,077	1,337	1,111
Schizophrenia and other psychotic disorders	14,233	13,986	14,530	14,403	14,748	13,252
Alcohol or substance-related disorders	5	--	--	--	--	--
Other mental disorders ²	3,748	6,074	6,056	6,359	7,062	5,827
Total	72,319	68,807	74,021	73,128	76,963	75,768
Gender						
Male	35,439	36,036	36,780	36,584	38,936	39,385
Female	40,033	40,232	40,717	39,825	41,834	42,729
Total	75,472	76,268	77,497	76,409	80,770	82,114
Age³						
0 to 11 years	7,146	7,600	8,053	8,138	9,114	9,039
12 to 14 years	4,444	4,592	4,661	4,872	5,763	5,846
15 to 17 years	4,262	4,499	4,596	4,290	4,865	4,922
18 to 20 years	2,385	2,494	2,650	2,657	2,948	2,991
21 to 24 years	3,516	3,564	3,526	3,385	3,648	3,889
25 to 29 years	5,787	5,678	5,687	5,584	6,006	6,289
30 to 34 years	6,639	6,453	6,538	6,136	6,406	6,748
35 to 39 years	6,490	6,629	6,820	6,537	6,824	6,923
40 to 44 years	6,550	6,156	6,045	5,869	6,139	6,388
45 to 49 years	7,041	6,937	6,870	6,643	6,559	6,526
50 to 54 years	8,496	8,281	8,005	7,450	7,060	6,775
55 to 59 years	6,568	6,794	6,956	7,240	7,342	7,295
60 to 64 years	3,691	3,955	4,270	4,620	4,935	5,048
65 years and older	2,456	2,636	2,820	2,987	3,161	3,438
Total	75,471	76,268	77,497	76,408	80,770	82,117

Continued. See notes at end of table.

Table 9.26a. Missouri – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	181	196	196	174	169	165
Asian ⁴	183	181	204	225	223	235
Black or African American	15,627	15,503	16,041	16,353	17,422	17,617
Native Hawaiian or Other Pacific Islander	30	37	40	32	37	55
White	55,430	56,106	56,369	54,875	57,613	58,062
Other	2,118	2,406	2,763	2,857	3,170	3,230
Total	73,569	74,429	75,613	74,516	78,634	79,364
Ethnicity						
Not of Hispanic or Latino origin	73,780	74,506	75,621	74,515	78,596	79,874
Hispanic or Latino origin	1,679	1,753	1,863	1,893	2,166	2,239
Total	75,459	76,259	77,484	76,408	80,762	82,113
Specific Hispanic or Latino origin						
Mexican	909	991	1,051	1,039	1,078	1,112
Puerto Rican	197	213	193	170	193	193
Cuban	55	55	51	55	129	92
Other specific	499	480	551	606	755	830
Total	1,660	1,739	1,846	1,870	2,155	2,227
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡
Not a veteran	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡
Total	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡

- - Quantity is zero; ‡ No data submitted; n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.26b. Missouri – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	10.5	6.6	7.6	8.4	9.2	8.8
Anxiety disorders	8.6	14.2	15.0	15.6	16.3	14.4
Attention deficit/hyperactivity disorder	8.7	9.9	10.5	10.7	11.1	9.7
Mood disorders	56.5	53.6	53.1	53.3	53.0	49.5
Bipolar disorders	26.6	22.8	21.7	20.9	20.3	17.4
Depressive disorders	32.3	33.1	33.8	35.2	35.9	33.2
Personality disorders	†	1.0	1.3	1.5	1.7	1.5
Schizophrenia and other psychotic disorders	19.7	20.3	19.6	19.7	19.2	17.5
Alcohol or substance-related disorders	†	--	--	--	--	--
Other mental disorders ²	5.2	8.8	8.2	8.7	9.2	7.7
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	47.0	47.2	47.5	47.9	48.2	48.0
Female	53.0	52.8	52.5	52.1	51.8	52.0
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	9.5	10.0	10.4	10.7	11.3	11.0
12 to 14 years	5.9	6.0	6.0	6.4	7.1	7.1
15 to 17 years	5.6	5.9	5.9	5.6	6.0	6.0
18 to 20 years	3.2	3.3	3.4	3.5	3.6	3.6
21 to 24 years	4.7	4.7	4.5	4.4	4.5	4.7
25 to 29 years	7.7	7.4	7.3	7.3	7.4	7.7
30 to 34 years	8.8	8.5	8.4	8.0	7.9	8.2
35 to 39 years	8.6	8.7	8.8	8.6	8.4	8.4
40 to 44 years	8.7	8.1	7.8	7.7	7.6	7.8
45 to 49 years	9.3	9.1	8.9	8.7	8.1	7.9
50 to 54 years	11.3	10.9	10.3	9.8	8.7	8.3
55 to 59 years	8.7	8.9	9.0	9.5	9.1	8.9
60 to 64 years	4.9	5.2	5.5	6.0	6.1	6.1
65 years and older	3.3	3.5	3.6	3.9	3.9	4.2
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.26b. Missouri – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	†	†	†	†	†	†
Asian ⁴	†	†	†	†	†	†
Black or African American	21.2	20.8	21.2	21.9	22.2	22.2
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	75.3	75.4	74.5	73.6	73.3	73.2
Other	2.9	3.2	3.7	3.8	4.0	4.1
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	97.8	97.7	97.6	97.5	97.3	97.3
Hispanic or Latino origin	2.2	2.3	2.4	2.5	2.7	2.7
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	54.8	57.0	56.9	55.6	50.0	49.9
Puerto Rican	11.9	12.2	10.5	9.1	9.0	8.7
Cuban	3.3	3.2	2.8	2.9	6.0	4.1
Other specific	30.1	27.6	29.8	32.4	35.0	37.3
Total	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡
Not a veteran	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡
Total	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡

† Less than 1 percent of individuals; - - Quantity is zero; ‡ No data submitted; *n/c* Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.27a. Montana – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	46,288	38,476	50,515	63,050	71,770	74,413
Mental health diagnosis¹						
Trauma- and stressor-related disorders	10,270	7,738	11,729	15,953	14,438	14,170
Anxiety disorders	9,181	7,390	13,169	24,100	27,331	29,733
Attention deficit/hyperactivity disorder	5,164	4,033	5,594	7,909	7,370	7,241
Mood disorders	19,824	14,091	19,078	25,495	28,640	28,417
Bipolar disorders	6,954	5,038	4,990	5,854	6,697	6,423
Depressive disorders	14,357	9,914	14,843	20,749	23,283	23,201
Personality disorders	1,385	1,039	711	1,414	987	860
Schizophrenia and other psychotic disorders	3,906	2,877	3,299	3,983	4,036	4,003
Alcohol or substance-related disorders	2,729	1,687	443	96	28	123
Other mental disorders ²	9,429	7,033	14,791	19,572	17,743	17,159
Total	40,759	31,106	49,274	61,239	70,761	73,240
Gender						
Male	20,936	17,426	23,310	28,091	30,572	31,092
Female	25,352	21,049	27,205	34,933	41,198	43,321
Total	46,288	38,475	50,515	63,024	71,770	74,413
Age³						
0 to 11 years	11,095	9,114	13,588	13,320	12,943	12,344
12 to 14 years	4,416	3,814	5,223	5,661	5,935	6,018
15 to 17 years	4,084	3,637	4,871	5,470	5,769	5,831
18 to 20 years	1,916	1,634	2,098	3,076	3,669	3,774
21 to 24 years	1,955	1,517	2,159	3,371	4,256	4,522
25 to 29 years	3,095	2,506	3,470	5,371	6,640	6,962
30 to 34 years	3,117	2,625	3,484	5,251	6,506	6,929
35 to 39 years	2,741	2,352	3,145	4,613	5,874	6,288
40 to 44 years	2,313	1,815	2,390	3,464	4,306	4,897
45 to 49 years	2,183	1,893	2,334	3,155	3,653	3,888
50 to 54 years	2,524	2,006	2,378	3,017	3,476	3,448
55 to 59 years	2,349	2,015	2,320	2,921	3,612	3,808
60 to 64 years	1,584	1,361	1,542	2,135	2,626	2,907
65 years and older	2,916	2,186	1,510	2,024	2,503	2,797
Total	46,288	38,475	50,512	62,849	71,768	74,413

Continued. See notes at end of table.

Table 9.27a. Montana – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	5,368	4,479	5,801	7,315	8,288	8,897
Asian ⁴	123	86	154	176	210	230
Black or African American	432	394	551	668	710	775
Native Hawaiian or Other Pacific Islander	27	19	32	47	57	67
White	36,950	30,496	38,968	47,929	54,413	56,389
Other	923	839	1,127	1,497	1,687	1,825
Total	43,823	36,313	46,633	57,632	65,365	68,183
Ethnicity						
Not of Hispanic or Latino origin	44,647	37,052	48,332	58,036	64,387	71,492
Hispanic or Latino origin	1,610	1,395	1,899	2,408	2,775	2,916
Total	46,257	38,447	50,231	60,444	67,162	74,408
Specific Hispanic or Latino origin						
Mexican	2	4	‡	‡	‡	‡
Puerto Rican	--	--	‡	‡	‡	‡
Cuban	--	--	‡	‡	‡	‡
Other specific	--	1	‡	‡	‡	‡
Total	2	5	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	200	174	171	161
Not a veteran	<i>n/c</i>	<i>n/c</i>	8,138	8,220	9,224	15,008
Total	<i>n/c</i>	<i>n/c</i>	8,338	8,394	9,395	15,169

-- Quantity is zero; ‡ No data submitted; n/c Not collected.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.27b. Montana – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	25.2	24.9	23.8	26.1	20.4	19.3
Anxiety disorders	22.5	23.8	26.7	39.4	38.6	40.6
Attention deficit/hyperactivity disorder	12.7	13.0	11.4	12.9	10.4	9.9
Mood disorders	48.6	45.3	38.7	41.6	40.5	38.8
Bipolar disorders	17.1	16.2	10.1	9.6	9.5	8.8
Depressive disorders	35.2	31.9	30.1	33.9	32.9	31.7
Personality disorders	3.4	3.3	1.4	2.3	1.4	1.2
Schizophrenia and other psychotic disorders	9.6	9.2	6.7	6.5	5.7	5.5
Alcohol or substance-related disorders	6.7	5.4	†	†	†	†
Other mental disorders ²	23.1	22.6	30.0	32.0	25.1	23.4
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	45.2	45.3	46.1	44.6	42.6	41.8
Female	54.8	54.7	53.9	55.4	57.4	58.2
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	24.0	23.7	26.9	21.2	18.0	16.6
12 to 14 years	9.5	9.9	10.3	9.0	8.3	8.1
15 to 17 years	8.8	9.5	9.6	8.7	8.0	7.8
18 to 20 years	4.1	4.2	4.2	4.9	5.1	5.1
21 to 24 years	4.2	3.9	4.3	5.4	5.9	6.1
25 to 29 years	6.7	6.5	6.9	8.5	9.3	9.4
30 to 34 years	6.7	6.8	6.9	8.4	9.1	9.3
35 to 39 years	5.9	6.1	6.2	7.3	8.2	8.5
40 to 44 years	5.0	4.7	4.7	5.5	6.0	6.6
45 to 49 years	4.7	4.9	4.6	5.0	5.1	5.2
50 to 54 years	5.5	5.2	4.7	4.8	4.8	4.6
55 to 59 years	5.1	5.2	4.6	4.6	5.0	5.1
60 to 64 years	3.4	3.5	3.1	3.4	3.7	3.9
65 years and older	6.3	5.7	3.0	3.2	3.5	3.8
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.27b. Montana – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	12.2	12.3	12.4	12.7	12.7	13.0
Asian ⁴	†	†	†	†	†	†
Black or African American	†	1.1	1.2	1.2	1.1	1.1
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	84.3	84.0	83.6	83.2	83.2	82.7
Other	2.1	2.3	2.4	2.6	2.6	2.7
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	96.5	96.4	96.2	96.0	95.9	96.1
Hispanic or Latino origin	3.5	3.6	3.8	4.0	4.1	3.9
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	100.0	80.0	‡	‡	‡	‡
Puerto Rican	--	--	‡	‡	‡	‡
Cuban	--	--	‡	‡	‡	‡
Other specific	--	20.0	‡	‡	‡	‡
Total	100.0	100.0	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	2.4	2.1	1.8	1.1
Not a veteran	<i>n/c</i>	<i>n/c</i>	97.6	97.9	98.2	98.9
Total	<i>n/c</i>	<i>n/c</i>	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals; -- Quantity is zero; ‡ No data submitted; *n/c* Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.28a. Nebraska – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	19,441	19,739	21,927	24,983	26,280	29,728
Mental health diagnosis¹						
Trauma- and stressor-related disorders	4,418	3,882	4,070	4,381	4,063	4,265
Anxiety disorders	3,694	3,760	4,271	4,653	4,586	4,709
Attention deficit/hyperactivity disorder	1,309	1,171	1,221	1,383	1,272	1,203
Mood disorders	11,339	9,634	10,515	10,583	10,699	10,773
Bipolar disorders	5,198	3,890	4,091	3,998	3,771	3,828
Depressive disorders	6,351	5,878	6,577	6,802	7,124	7,126
Personality disorders	2,013	409	601	688	587	626
Schizophrenia and other psychotic disorders	2,369	2,121	2,361	2,442	2,567	2,671
Alcohol or substance-related disorders	--	2	3,017	3,741	2,769	2,981
Other mental disorders ²	6,008	2,346	2,822	3,618	3,560	4,262
Total	18,011	16,920	19,218	20,013	20,628	21,981
Gender						
Male	9,636	9,866	10,994	12,339	12,791	15,571
Female	9,802	9,840	10,753	11,766	12,254	13,307
Total	19,438	19,706	21,747	24,105	25,045	28,878
Age³						
0 to 11 years	692	715	757	904	958	866
12 to 14 years	482	501	690	933	1,044	1,021
15 to 17 years	644	645	956	1,548	1,512	1,542
18 to 20 years	927	984	1,110	1,414	1,409	1,496
21 to 24 years	1,796	1,859	1,908	2,148	2,183	2,254
25 to 29 years	2,370	2,434	2,678	3,045	3,119	3,536
30 to 34 years	2,495	2,508	2,797	2,915	3,004	3,397
35 to 39 years	2,009	2,130	2,315	2,581	2,818	3,275
40 to 44 years	1,770	1,755	1,910	2,007	2,184	2,648
45 to 49 years	1,882	1,820	1,944	2,029	2,040	2,383
50 to 54 years	1,962	1,839	1,909	1,969	1,973	2,178
55 to 59 years	1,329	1,374	1,449	1,614	1,759	2,103
60 to 64 years	704	735	831	965	1,028	1,357
65 years and older	378	415	672	910	1,248	1,672
Total	19,440	19,714	21,926	24,982	26,279	29,728

Continued. See notes at end of table.

Table 9.28a. Nebraska – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	477	505	661	615	684	681
Asian ⁴	144	131	140	161	184	219
Black or African American	1,670	1,732	1,982	2,176	2,385	2,728
Native Hawaiian or Other Pacific Islander	125	126	64	93	111	116
White	16,731	16,499	17,545	18,378	18,896	20,922
Other	252	583	770	1,368	1,383	3,313
Total	19,399	19,576	21,162	22,791	23,643	27,979
Ethnicity						
Not of Hispanic or Latino origin	17,157	16,029	17,536	18,393	19,046	19,171
Hispanic or Latino origin	1,635	1,604	1,750	2,118	2,367	2,468
Total	18,792	17,633	19,286	20,511	21,413	21,639
Specific Hispanic or Latino origin						
Mexican	535	‡	‡	‡	‡	‡
Puerto Rican	44	‡	‡	‡	‡	‡
Cuban	24	‡	‡	‡	‡	‡
Other specific	196	‡	‡	‡	‡	‡
Total	799	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	‡	398	325	386
Not a veteran	<i>n/c</i>	<i>n/c</i>	‡	17,643	17,814	21,344
Total	<i>n/c</i>	<i>n/c</i>	‡	18,041	18,139	21,730

- - Quantity is zero; ‡ No data submitted; n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.28b. Nebraska – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	24.5	22.9	21.2	21.9	19.7	19.4
Anxiety disorders	20.5	22.2	22.2	23.2	22.2	21.4
Attention deficit/hyperactivity disorder	7.3	6.9	6.4	6.9	6.2	5.5
Mood disorders	63.0	56.9	54.7	52.9	51.9	49.0
Bipolar disorders	28.9	23.0	21.3	20.0	18.3	17.4
Depressive disorders	35.3	34.7	34.2	34.0	34.5	32.4
Personality disorders	11.2	2.4	3.1	3.4	2.8	2.8
Schizophrenia and other psychotic disorders	13.2	12.5	12.3	12.2	12.4	12.2
Alcohol or substance-related disorders	--	†	15.7	18.7	13.4	13.6
Other mental disorders ²	33.4	13.9	14.7	18.1	17.3	19.4
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	49.6	50.1	50.6	51.2	51.1	53.9
Female	50.4	49.9	49.4	48.8	48.9	46.1
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	3.6	3.6	3.5	3.6	3.6	2.9
12 to 14 years	2.5	2.5	3.1	3.7	4.0	3.4
15 to 17 years	3.3	3.3	4.4	6.2	5.8	5.2
18 to 20 years	4.8	5.0	5.1	5.7	5.4	5.0
21 to 24 years	9.2	9.4	8.7	8.6	8.3	7.6
25 to 29 years	12.2	12.3	12.2	12.2	11.9	11.9
30 to 34 years	12.8	12.7	12.8	11.7	11.4	11.4
35 to 39 years	10.3	10.8	10.6	10.3	10.7	11.0
40 to 44 years	9.1	8.9	8.7	8.0	8.3	8.9
45 to 49 years	9.7	9.2	8.9	8.1	7.8	8.0
50 to 54 years	10.1	9.3	8.7	7.9	7.5	7.3
55 to 59 years	6.8	7.0	6.6	6.5	6.7	7.1
60 to 64 years	3.6	3.7	3.8	3.9	3.9	4.6
65 years and older	1.9	2.1	3.1	3.6	4.7	5.6
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.28b. Nebraska – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	2.5	2.6	3.1	2.7	2.9	2.4
Asian ⁴	†	†	†	†	†	†
Black or African American	8.6	8.8	9.4	9.5	10.1	9.8
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	86.2	84.3	82.9	80.6	79.9	74.8
Other	1.3	3.0	3.6	6.0	5.8	11.8
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	91.3	90.9	90.9	89.7	88.9	88.6
Hispanic or Latino origin	8.7	9.1	9.1	10.3	11.1	11.4
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	67.0	‡	‡	‡	‡	‡
Puerto Rican	5.5	‡	‡	‡	‡	‡
Cuban	3.0	‡	‡	‡	‡	‡
Other specific	24.5	‡	‡	‡	‡	‡
Total	100.0	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	‡	2.2	1.8	1.8
Not a veteran	<i>n/c</i>	<i>n/c</i>	‡	97.8	98.2	98.2
Total	<i>n/c</i>	<i>n/c</i>	‡	100.0	100.0	100.0

† Less than 1 percent of individuals; - - Quantity is zero; ‡ No data submitted; *n/c* Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.29a. Nevada – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	28,589	23,945	20,535	19,635	18,937	16,416
Mental health diagnosis¹						
Trauma- and stressor-related disorders	4,132	1,823	2,021	1,931	2,029	1,930
Anxiety disorders	4,799	2,021	2,062	1,981	2,200	2,244
Attention deficit/hyperactivity disorder	1,011	647	647	572	639	669
Mood disorders	14,591	3,402	3,547	3,223	3,047	3,506
Bipolar disorders	6,901	1,162	1,166	1,043	996	1,123
Depressive disorders	7,829	2,246	2,385	2,183	2,054	2,395
Personality disorders	48	647	600	517	445	409
Schizophrenia and other psychotic disorders	5,907	2,015	2,077	2,034	2,046	2,321
Alcohol or substance-related disorders	7,813	3,246	3,243	2,829	2,538	1,810
Other mental disorders ²	2,194	2,147	2,161	2,066	1,982	2,459
Total	25,005	13,861	14,249	13,263	12,942	13,036
Gender						
Male	13,760	11,739	10,135	9,827	9,569	8,402
Female	14,676	12,151	10,383	9,792	9,349	7,998
Total	28,436	23,890	20,518	19,619	18,918	16,400
Age³						
0 to 11 years	1,832	1,727	1,851	1,924	1,741	1,460
12 to 14 years	942	1,011	1,202	1,236	1,345	1,155
15 to 17 years	1,154	1,349	1,477	1,528	1,500	1,366
18 to 20 years	921	849	864	793	788	675
21 to 24 years	1,801	1,452	1,259	1,279	1,144	968
25 to 29 years	2,823	2,316	1,881	1,774	1,720	1,460
30 to 34 years	2,960	2,356	1,810	1,741	1,666	1,424
35 to 39 years	2,615	2,151	1,684	1,572	1,484	1,302
40 to 44 years	2,575	1,923	1,516	1,440	1,362	1,157
45 to 49 years	2,942	2,255	1,686	1,534	1,483	1,237
50 to 54 years	3,167	2,373	1,898	1,691	1,532	1,286
55 to 59 years	2,502	2,007	1,588	1,438	1,384	1,263
60 to 64 years	1,474	1,242	1,001	919	948	900
65 years and older	857	923	807	727	788	735
Total	28,565	23,934	20,524	19,596	18,885	16,388

Continued. See notes at end of table.

Table 9.29a. Nevada – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	351	292	344	366	339	297
Asian ⁴	514	469	423	471	439	393
Black or African American	3,684	3,008	2,585	2,538	2,432	2,148
Native Hawaiian or Other Pacific Islander	142	136	121	121	126	112
White	16,591	13,257	10,948	10,412	9,746	8,586
Other	3,097	2,460	1,978	1,890	1,765	1,411
Total	24,379	19,622	16,399	15,798	14,847	12,947
Ethnicity						
Not of Hispanic or Latino origin	18,471	15,050	12,166	11,948	11,244	9,958
Hispanic or Latino origin	3,774	3,369	3,039	2,996	2,929	2,770
Total	22,245	18,419	15,205	14,944	14,173	12,728
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	14	17	9	37
Not a veteran	<i>n/c</i>	<i>n/c</i>	8,863	4,868	197	1,201
Total	<i>n/c</i>	<i>n/c</i>	8,877	4,885	206	1,238

‡ No data submitted; n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.29b. Nevada – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	16.5	13.2	14.2	14.6	15.7	14.8
Anxiety disorders	19.2	14.6	14.5	14.9	17.0	17.2
Attention deficit/hyperactivity disorder	4.0	4.7	4.5	4.3	4.9	5.1
Mood disorders	58.4	24.5	24.9	24.3	23.5	26.9
Bipolar disorders	27.6	8.4	8.2	7.9	7.7	8.6
Depressive disorders	31.3	16.2	16.7	16.5	15.9	18.4
Personality disorders	†	4.7	4.2	3.9	3.4	3.1
Schizophrenia and other psychotic disorders	23.6	14.5	14.6	15.3	15.8	17.8
Alcohol or substance-related disorders	31.2	23.4	22.8	21.3	19.6	13.9
Other mental disorders ²	8.8	15.5	15.2	15.6	15.3	18.9
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	48.4	49.1	49.4	50.1	50.6	51.2
Female	51.6	50.9	50.6	49.9	49.4	48.8
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	6.4	7.2	9.0	9.8	9.2	8.9
12 to 14 years	3.3	4.2	5.9	6.3	7.1	7.0
15 to 17 years	4.0	5.6	7.2	7.8	7.9	8.3
18 to 20 years	3.2	3.5	4.2	4.0	4.2	4.1
21 to 24 years	6.3	6.1	6.1	6.5	6.1	5.9
25 to 29 years	9.9	9.7	9.2	9.1	9.1	8.9
30 to 34 years	10.4	9.8	8.8	8.9	8.8	8.7
35 to 39 years	9.2	9.0	8.2	8.0	7.9	7.9
40 to 44 years	9.0	8.0	7.4	7.3	7.2	7.1
45 to 49 years	10.3	9.4	8.2	7.8	7.9	7.5
50 to 54 years	11.1	9.9	9.2	8.6	8.1	7.8
55 to 59 years	8.8	8.4	7.7	7.3	7.3	7.7
60 to 64 years	5.2	5.2	4.9	4.7	5.0	5.5
65 years and older	3.0	3.9	3.9	3.7	4.2	4.5
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.29b. Nevada – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	1.4	1.5	2.1	2.3	2.3	2.3
Asian ⁴	2.1	2.4	2.6	3.0	3.0	3.0
Black or African American	15.1	15.3	15.8	16.1	16.4	16.6
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	68.1	67.6	66.8	65.9	65.6	66.3
Other	12.7	12.5	12.1	12.0	11.9	10.9
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	83.0	81.7	80.0	80.0	79.3	78.2
Hispanic or Latino origin	17.0	18.3	20.0	20.0	20.7	21.8
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	†	†	4.4	3.0
Not a veteran	<i>n/c</i>	<i>n/c</i>	99.8	99.7	95.6	97.0
Total	<i>n/c</i>	<i>n/c</i>	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals; ‡ No data submitted; *n/c* Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.30a. New Jersey – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	325,102	323,392	402,469	421,376	427,465	415,832
Mental health diagnosis¹						
Trauma- and stressor-related disorders	42,742	4,081	55,169	60,233	61,655	61,405
Anxiety disorders	26,608	1,754	32,462	34,776	35,492	35,098
Attention deficit/hyperactivity disorder	14,165	4,408	21,691	25,880	25,891	25,562
Mood disorders	93,375	2,704	111,273	113,319	112,774	110,934
Bipolar disorders	27,725	638	32,789	33,131	32,535	31,660
Depressive disorders	66,071	2,087	79,007	80,706	80,764	79,778
Personality disorders	7,621	73	9,311	9,362	9,280	9,266
Schizophrenia and other psychotic disorders	30,550	158	40,494	40,021	39,098	37,502
Alcohol or substance-related disorders	31,905	1,152	39,458	39,796	39,334	38,827
Other mental disorders ²	107,073	7,789	139,642	146,256	145,805	144,397
Total	243,422	16,772	308,859	320,958	320,679	315,988
Gender						
Male	154,484	153,428	191,045	201,665	204,563	197,957
Female	170,002	169,258	210,562	218,887	222,043	217,077
Total	324,486	322,686	401,607	420,552	426,606	415,034
Age³						
0 to 11 years	22,951	22,990	21,704	25,639	25,731	22,311
12 to 14 years	16,862	15,886	15,175	18,040	19,151	17,058
15 to 17 years	20,757	20,080	19,999	22,540	22,975	20,998
18 to 20 years	12,746	11,766	16,440	18,494	18,060	16,691
21 to 24 years	15,043	14,643	19,300	18,892	17,905	16,318
25 to 29 years	20,529	20,155	27,079	27,063	26,685	25,447
30 to 34 years	22,038	21,437	27,371	27,554	27,981	27,457
35 to 39 years	20,864	21,349	27,874	28,684	28,846	28,052
40 to 44 years	22,109	20,697	25,579	25,771	26,174	26,404
45 to 49 years	25,852	25,070	31,458	30,659	29,802	27,893
50 to 54 years	29,841	28,847	36,597	35,798	34,599	32,953
55 to 59 years	27,743	28,252	37,640	38,322	38,784	37,871
60 to 64 years	20,720	21,898	30,139	32,126	33,932	34,937
65 years and older	46,690	50,254	65,897	71,571	76,620	81,234
Total	324,745	323,324	402,252	421,153	427,245	415,624

Continued. See notes at end of table.

Table 9.30a. New Jersey – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	2,556	2,516	3,551	3,533	3,535	3,462
Asian ⁴	4,757	4,823	6,227	6,623	6,723	6,501
Black or African American	63,691	63,074	79,298	82,021	81,366	79,147
Native Hawaiian or Other Pacific Islander	44	29	62	62	51	37
White	170,132	168,291	207,226	214,241	213,466	209,782
Other	23,136	21,919	29,027	30,060	29,448	27,912
Total	264,316	260,652	325,391	336,540	334,589	326,841
Ethnicity						
Not of Hispanic or Latino origin	‡	‡	305,416	312,257	310,716	304,857
Hispanic or Latino origin	‡	‡	70,218	73,931	75,855	73,420
Total	‡	‡	375,634	386,188	386,571	378,277
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	5,156	- -	6,971	7,210
Not a veteran	<i>n/c</i>	<i>n/c</i>	294,504	345,469	294,789	290,796
Total	<i>n/c</i>	<i>n/c</i>	299,660	345,469	301,760	298,006

- - Quantity is zero; ‡ No data submitted; n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.30b. New Jersey – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	17.6	24.3	17.9	18.8	19.2	19.4
Anxiety disorders	10.9	10.5	10.5	10.8	11.1	11.1
Attention deficit/hyperactivity disorder	5.8	26.3	7.0	8.1	8.1	8.1
Mood disorders	38.4	16.1	36.0	35.3	35.2	35.1
Bipolar disorders	11.4	3.8	10.6	10.3	10.1	10.0
Depressive disorders	27.1	12.4	25.6	25.1	25.2	25.2
Personality disorders	3.1	†	3.0	2.9	2.9	2.9
Schizophrenia and other psychotic disorders	12.6	†	13.1	12.5	12.2	11.9
Alcohol or substance-related disorders	13.1	6.9	12.8	12.4	12.3	12.3
Other mental disorders ²	44.0	46.4	45.2	45.6	45.5	45.7
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	47.6	47.5	47.6	48.0	48.0	47.7
Female	52.4	52.5	52.4	52.0	52.0	52.3
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	7.1	7.1	5.4	6.1	6.0	5.4
12 to 14 years	5.2	4.9	3.8	4.3	4.5	4.1
15 to 17 years	6.4	6.2	5.0	5.4	5.4	5.1
18 to 20 years	3.9	3.6	4.1	4.4	4.2	4.0
21 to 24 years	4.6	4.5	4.8	4.5	4.2	3.9
25 to 29 years	6.3	6.2	6.7	6.4	6.2	6.1
30 to 34 years	6.8	6.6	6.8	6.5	6.5	6.6
35 to 39 years	6.4	6.6	6.9	6.8	6.8	6.7
40 to 44 years	6.8	6.4	6.4	6.1	6.1	6.4
45 to 49 years	8.0	7.8	7.8	7.3	7.0	6.7
50 to 54 years	9.2	8.9	9.1	8.5	8.1	7.9
55 to 59 years	8.5	8.7	9.4	9.1	9.1	9.1
60 to 64 years	6.4	6.8	7.5	7.6	7.9	8.4
65 years and older	14.4	15.5	16.4	17.0	17.9	19.5
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.30b. New Jersey – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	†	†	1.1	1.0	1.1	1.1
Asian ⁴	1.8	1.9	1.9	2.0	2.0	2.0
Black or African American	24.1	24.2	24.4	24.4	24.3	24.2
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	64.4	64.6	63.7	63.7	63.8	64.2
Other	8.8	8.4	8.9	8.9	8.8	8.5
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	‡	‡	81.3	80.9	80.4	80.6
Hispanic or Latino origin	‡	‡	18.7	19.1	19.6	19.4
Total	‡	‡	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	1.7	--	2.3	2.4
Not a veteran	<i>n/c</i>	<i>n/c</i>	98.3	100.0	97.7	97.6
Total	<i>n/c</i>	<i>n/c</i>	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals; -- Quantity is zero; ‡ No data submitted; *n/c* Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.31a. New Mexico – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	121,603	131,220	147,669	149,503	203,037	179,627
Mental health diagnosis¹						
Trauma- and stressor-related disorders	30,002	37,557	44,352	31,909	48,106	43,583
Anxiety disorders	16,799	24,034	32,518	28,366	43,906	41,569
Attention deficit/hyperactivity disorder	9,169	9,783	11,305	10,875	13,674	13,116
Mood disorders	36,367	39,804	46,518	36,053	51,048	43,891
Bipolar disorders	12,885	12,115	12,942	9,593	12,516	9,932
Depressive disorders	25,520	29,454	35,654	27,680	40,543	35,556
Personality disorders	614	714	890	541	697	565
Schizophrenia and other psychotic disorders	5,554	6,458	8,587	7,264	9,416	7,082
Alcohol or substance-related disorders	18,635	22,934	32,261	30,636	44,994	32,278
Other mental disorders ²	22,549	27,396	23,497	42,387	54,232	50,226
Total	105,156	126,002	144,055	149,500	203,017	179,605
Gender						
Male	59,253	63,080	70,053	73,683	97,935	85,978
Female	62,178	68,062	77,554	75,602	104,578	93,538
Total	121,431	131,142	147,607	149,285	202,513	179,516
Age³						
0 to 11 years	29,081	30,205	25,268	33,847	45,314	40,939
12 to 14 years	9,158	9,886	10,105	10,109	15,226	13,275
15 to 17 years	9,025	10,142	10,471	9,859	14,299	12,587
18 to 20 years	4,786	5,332	5,988	5,738	9,458	8,032
21 to 24 years	6,513	7,249	8,799	7,395	11,388	9,651
25 to 29 years	10,126	11,285	14,294	12,833	18,614	15,410
30 to 34 years	10,047	11,230	13,846	13,041	17,780	15,427
35 to 39 years	8,381	9,737	12,168	11,876	15,851	13,826
40 to 44 years	7,144	7,614	9,379	9,110	11,993	10,862
45 to 49 years	6,858	7,171	9,018	8,449	10,224	9,007
50 to 54 years	7,306	7,435	9,213	8,342	9,800	8,416
55 to 59 years	5,785	6,089	8,032	7,672	9,489	8,706
60 to 64 years	3,894	3,975	5,328	5,449	6,564	6,488
65 years and older	3,490	3,867	5,755	5,779	7,035	6,999
Total	121,594	131,217	147,664	149,499	203,035	179,625

Continued. See notes at end of table.

Table 9.31a. New Mexico – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	4,543	4,678	7,773	11,849	19,048	17,312
Asian ⁴	713	693	3,804	1,602	1,847	1,757
Black or African American	2,498	2,653	3,305	3,013	4,373	3,805
Native Hawaiian or Other Pacific Islander	133	80	112	144	242	229
White	75,050	81,802	101,876	104,088	145,340	129,964
Other	3,593	3,245	3,023	3,359	4,404	4,261
Total	86,530	93,151	119,893	124,055	175,254	157,328
Ethnicity						
Not of Hispanic or Latino origin	39,669	41,091	62,187	57,549	81,757	73,734
Hispanic or Latino origin	49,259	54,695	54,446	66,607	94,696	84,090
Total	88,928	95,786	116,633	124,156	176,453	157,824
Specific Hispanic or Latino origin						
Mexican	1,377	1,248	1,040	835	853	684
Puerto Rican	121	107	86	64	87	66
Cuban	62	52	56	43	43	28
Other specific	667	601	537	495	496	371
Total	2,227	2,008	1,719	1,437	1,479	1,149
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	199	697	772	871
Not a veteran	<i>n/c</i>	<i>n/c</i>	4,930	92,393	126,059	108,047
Total	<i>n/c</i>	<i>n/c</i>	5,129	93,090	126,831	108,918

n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.31b. New Mexico – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	28.5	29.8	30.8	21.3	23.7	24.3
Anxiety disorders	16.0	19.1	22.6	19.0	21.6	23.1
Attention deficit/hyperactivity disorder	8.7	7.8	7.8	7.3	6.7	7.3
Mood disorders	34.6	31.6	32.3	24.1	25.1	24.4
Bipolar disorders	12.3	9.6	9.0	6.4	6.2	5.5
Depressive disorders	24.3	23.4	24.8	18.5	20.0	19.8
Personality disorders	†	†	†	†	†	†
Schizophrenia and other psychotic disorders	5.3	5.1	6.0	4.9	4.6	3.9
Alcohol or substance-related disorders	17.7	18.2	22.4	20.5	22.2	18.0
Other mental disorders ²	21.4	21.7	16.3	28.4	26.7	28.0
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	48.8	48.1	47.5	49.4	48.4	47.9
Female	51.2	51.9	52.5	50.6	51.6	52.1
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	23.9	23.0	17.1	22.6	22.3	22.8
12 to 14 years	7.5	7.5	6.8	6.8	7.5	7.4
15 to 17 years	7.4	7.7	7.1	6.6	7.0	7.0
18 to 20 years	3.9	4.1	4.1	3.8	4.7	4.5
21 to 24 years	5.4	5.5	6.0	4.9	5.6	5.4
25 to 29 years	8.3	8.6	9.7	8.6	9.2	8.6
30 to 34 years	8.3	8.6	9.4	8.7	8.8	8.6
35 to 39 years	6.9	7.4	8.2	7.9	7.8	7.7
40 to 44 years	5.9	5.8	6.4	6.1	5.9	6.0
45 to 49 years	5.6	5.5	6.1	5.7	5.0	5.0
50 to 54 years	6.0	5.7	6.2	5.6	4.8	4.7
55 to 59 years	4.8	4.6	5.4	5.1	4.7	4.8
60 to 64 years	3.2	3.0	3.6	3.6	3.2	3.6
65 years and older	2.9	2.9	3.9	3.9	3.5	3.9
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.31b. New Mexico – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	5.3	5.0	6.5	9.6	10.9	11.0
Asian ⁴	†	†	3.2	1.3	1.1	1.1
Black or African American	2.9	2.8	2.8	2.4	2.5	2.4
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	86.7	87.8	85.0	83.9	82.9	82.6
Other	4.2	3.5	2.5	2.7	2.5	2.7
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	44.6	42.9	53.3	46.4	46.3	46.7
Hispanic or Latino origin	55.4	57.1	46.7	53.6	53.7	53.3
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	61.8	62.2	60.5	58.1	57.7	59.5
Puerto Rican	5.4	5.3	5.0	4.5	5.9	5.7
Cuban	2.8	2.6	3.3	3.0	2.9	2.4
Other specific	30.0	29.9	31.2	34.4	33.5	32.3
Total	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	3.9	†	†	†
Not a veteran	<i>n/c</i>	<i>n/c</i>	96.1	99.3	99.4	99.2
Total	<i>n/c</i>	<i>n/c</i>	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals; n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.32a. New York – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	54,524	56,968	58,040	57,929	57,523	55,997
Mental health diagnosis¹						
Trauma- and stressor-related disorders	8,259	11,587	11,625	12,490	13,272	14,035
Anxiety disorders	6,309	9,568	7,291	7,442	7,527	7,606
Attention deficit/hyperactivity disorder	3,362	3,622	3,503	3,720	3,842	3,779
Mood disorders	19,464	22,418	18,199	18,157	17,737	16,886
Bipolar disorders	8,689	10,653	7,300	7,066	6,730	6,359
Depressive disorders	11,000	13,745	11,727	11,861	11,753	11,289
Personality disorders	9,845	11,645	9,316	9,477	9,359	9,177
Schizophrenia and other psychotic disorders	20,104	20,849	20,562	20,506	20,131	19,827
Alcohol or substance-related disorders	26	578	7	21	1	1
Other mental disorders ²	5,143	6,222	13,061	13,523	13,425	12,875
Total	52,873	55,837	56,581	56,534	56,182	54,840
Gender						
Male	35,414	36,941	37,976	37,755	37,530	36,719
Female	19,109	20,024	20,063	20,173	19,993	19,278
Total	54,523	56,965	58,039	57,928	57,523	55,997
Age³						
0 to 11 years	1,978	2,197	2,261	2,280	2,220	2,122
12 to 14 years	2,089	2,217	2,348	2,414	2,490	2,379
15 to 17 years	2,648	2,899	3,005	3,032	3,063	2,821
18 to 20 years	1,369	1,536	1,483	1,458	1,412	1,333
21 to 24 years	3,028	3,108	3,119	2,938	2,928	2,645
25 to 29 years	4,593	5,055	5,363	5,301	5,405	5,152
30 to 34 years	4,839	5,130	5,241	5,390	5,489	5,594
35 to 39 years	4,591	4,889	5,251	5,325	5,365	5,360
40 to 44 years	4,344	4,392	4,348	4,419	4,437	4,584
45 to 49 years	5,477	5,352	5,236	4,931	4,600	4,186
50 to 54 years	6,115	6,126	5,900	5,742	5,325	5,025
55 to 59 years	5,286	5,420	5,579	5,549	5,485	5,278
60 to 64 years	3,661	3,851	4,054	4,103	4,255	4,261
65 years and older	4,506	4,796	4,852	5,045	5,048	5,256
Total	54,524	56,968	58,040	57,927	57,522	55,996

Continued. See notes at end of table.

Table 9.32a. New York – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	266	352	367	374	360	358
Asian ⁴	775	873	939	968	1,060	1,022
Black or African American	15,585	16,054	16,494	16,434	16,234	16,074
Native Hawaiian or Other Pacific Islander	69	64	60	53	58	47
White	27,626	28,846	29,407	29,285	29,123	28,007
Other	4,613	5,502	6,429	6,833	6,741	6,636
Total	48,934	51,691	53,696	53,947	53,576	52,144
Ethnicity						
Not of Hispanic or Latino origin	38,180	41,136	42,793	42,986	42,832	41,685
Hispanic or Latino origin	8,534	8,600	8,802	8,643	8,462	8,200
Total	46,714	49,736	51,595	51,629	51,294	49,885
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	359	331	313	290
Not a veteran	<i>n/c</i>	<i>n/c</i>	13,547	12,680	11,759	10,948
Total	<i>n/c</i>	<i>n/c</i>	13,906	13,011	12,072	11,238

‡ No data submitted; n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.32b. New York – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	15.6	20.8	20.5	22.1	23.6	25.6
Anxiety disorders	11.9	17.1	12.9	13.2	13.4	13.9
Attention deficit/hyperactivity disorder	6.4	6.5	6.2	6.6	6.8	6.9
Mood disorders	36.8	40.1	32.2	32.1	31.6	30.8
Bipolar disorders	16.4	19.1	12.9	12.5	12.0	11.6
Depressive disorders	20.8	24.6	20.7	21.0	20.9	20.6
Personality disorders	18.6	20.9	16.5	16.8	16.7	16.7
Schizophrenia and other psychotic disorders	38.0	37.3	36.3	36.3	35.8	36.2
Alcohol or substance-related disorders	†	1.0	†	†	†	†
Other mental disorders ²	9.7	11.1	23.1	23.9	23.9	23.5
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	65.0	64.8	65.4	65.2	65.2	65.6
Female	35.0	35.2	34.6	34.8	34.8	34.4
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	3.6	3.9	3.9	3.9	3.9	3.8
12 to 14 years	3.8	3.9	4.0	4.2	4.3	4.2
15 to 17 years	4.9	5.1	5.2	5.2	5.3	5.0
18 to 20 years	2.5	2.7	2.6	2.5	2.5	2.4
21 to 24 years	5.6	5.5	5.4	5.1	5.1	4.7
25 to 29 years	8.4	8.9	9.2	9.2	9.4	9.2
30 to 34 years	8.9	9.0	9.0	9.3	9.5	10.0
35 to 39 years	8.4	8.6	9.0	9.2	9.3	9.6
40 to 44 years	8.0	7.7	7.5	7.6	7.7	8.2
45 to 49 years	10.0	9.4	9.0	8.5	8.0	7.5
50 to 54 years	11.2	10.8	10.2	9.9	9.3	9.0
55 to 59 years	9.7	9.5	9.6	9.6	9.5	9.4
60 to 64 years	6.7	6.8	7.0	7.1	7.4	7.6
65 years and older	8.3	8.4	8.4	8.7	8.8	9.4
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.32b. New York – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	†	†	†	†	†	†
Asian ⁴	1.6	1.7	1.7	1.8	2.0	2.0
Black or African American	31.8	31.1	30.7	30.5	30.3	30.8
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	56.5	55.8	54.8	54.3	54.4	53.7
Other	9.4	10.6	12.0	12.7	12.6	12.7
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	81.7	82.7	82.9	83.3	83.5	83.6
Hispanic or Latino origin	18.3	17.3	17.1	16.7	16.5	16.4
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	2.6	2.5	2.6	2.6
Not a veteran	<i>n/c</i>	<i>n/c</i>	97.4	97.5	97.4	97.4
Total	<i>n/c</i>	<i>n/c</i>	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals; ‡ No data submitted; n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

**Table 9.33a. North Carolina – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Number, 2015–2020**

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	225,452	138,079	142,665	101,046	107,091	109,011
Mental health diagnosis¹						
Trauma- and stressor-related disorders	43,282	18,906	25,579	21,506	24,196	24,649
Anxiety disorders	44,636	20,609	25,413	21,286	22,646	22,486
Attention deficit/hyperactivity disorder	26,565	6,897	8,993	6,325	6,133	5,430
Mood disorders	130,588	59,460	79,839	54,273	57,979	58,088
Bipolar disorders	58,037	23,681	30,220	19,582	20,044	19,466
Depressive disorders	87,236	41,804	54,745	39,032	41,679	41,645
Personality disorders	5,607	3,498	4,681	3,679	3,678	3,499
Schizophrenia and other psychotic disorders	27,900	14,130	20,444	16,038	16,268	16,122
Alcohol or substance-related disorders	--	2	1	1	7	--
Other mental disorders ²	30,340	14,229	20,442	15,342	15,183	14,627
Total	218,361	90,822	132,239	90,186	97,747	99,827
Gender						
Male	106,398	66,277	70,348	52,269	54,638	56,075
Female	119,054	71,767	72,278	48,760	52,441	52,924
Total	225,452	138,044	142,626	101,029	107,079	108,999
Age³						
0 to 11 years	25,244	4,800	4,073	3,569	3,414	2,936
12 to 14 years	14,410	5,109	4,516	3,470	3,712	3,563
15 to 17 years	15,834	7,122	6,554	4,218	4,257	3,970
18 to 20 years	11,132	6,027	6,946	4,565	4,885	5,002
21 to 24 years	16,115	10,863	11,800	8,939	9,323	9,905
25 to 29 years	22,170	15,609	17,284	13,375	14,762	15,074
30 to 34 years	20,773	14,761	15,869	11,861	12,960	14,018
35 to 39 years	18,482	13,674	14,777	10,603	11,295	11,871
40 to 44 years	17,599	12,812	12,890	9,239	9,792	10,025
45 to 49 years	17,546	13,664	13,738	9,542	9,757	9,686
50 to 54 years	18,106	14,005	13,527	8,908	9,038	8,785
55 to 59 years	13,325	10,331	10,720	6,775	7,226	7,360
60 to 64 years	7,485	5,612	5,814	3,725	4,088	4,218
65 years and older	7,231	3,687	4,157	2,257	2,581	2,598
Total	225,452	138,076	142,665	101,046	107,090	109,011

Continued. See notes at end of table.

**Table 9.33a. North Carolina – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Number, 2015–2020 (continued)**

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	5,480	2,271	2,465	1,765	1,778	1,785
Asian ⁴	840	620	705	559	539	539
Black or African American	71,740	42,124	44,329	32,235	33,841	34,139
Native Hawaiian or Other Pacific Islander	194	75	81	51	48	73
White	133,655	86,746	88,431	61,901	65,941	67,250
Other	11,926	5,625	6,031	4,230	4,556	4,892
Total	223,835	137,461	142,042	100,741	106,703	108,678
Ethnicity						
Not of Hispanic or Latino origin	213,132	112,014	115,526	96,332	102,036	103,462
Hispanic or Latino origin	10,609	4,819	5,422	4,402	4,709	5,320
Total	223,741	116,833	120,948	100,734	106,745	108,782
Specific Hispanic or Latino origin						
Mexican	4,509	1,852	1,913	1,350	1,390	1,525
Puerto Rican	907	467	506	384	427	451
Cuban	280	132	148	106	94	91
Other specific	4,890	2,315	2,805	2,511	2,736	3,185
Total	10,586	4,766	5,372	4,351	4,647	5,252
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	‡	1,680	1,785	1,650
Not a veteran	<i>n/c</i>	<i>n/c</i>	‡	87,458	93,233	96,087
Total	<i>n/c</i>	<i>n/c</i>	‡	89,138	95,018	97,737

- - Quantity is zero; ‡ No data submitted; n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

**Table 9.33b. North Carolina – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Column percent distribution, 2015–2020**

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	19.8	20.8	19.3	23.8	24.8	24.7
Anxiety disorders	20.4	22.7	19.2	23.6	23.2	22.5
Attention deficit/hyperactivity disorder	12.2	7.6	6.8	7.0	6.3	5.4
Mood disorders	59.8	65.5	60.4	60.2	59.3	58.2
Bipolar disorders	26.6	26.1	22.9	21.7	20.5	19.5
Depressive disorders	40.0	46.0	41.4	43.3	42.6	41.7
Personality disorders	2.6	3.9	3.5	4.1	3.8	3.5
Schizophrenia and other psychotic disorders	12.8	15.6	15.5	17.8	16.6	16.1
Alcohol or substance-related disorders	--	†	†	†	†	--
Other mental disorders ²	13.9	15.7	15.5	17.0	15.5	14.7
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	47.2	48.0	49.3	51.7	51.0	51.4
Female	52.8	52.0	50.7	48.3	49.0	48.6
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	11.2	3.5	2.9	3.5	3.2	2.7
12 to 14 years	6.4	3.7	3.2	3.4	3.5	3.3
15 to 17 years	7.0	5.2	4.6	4.2	4.0	3.6
18 to 20 years	4.9	4.4	4.9	4.5	4.6	4.6
21 to 24 years	7.1	7.9	8.3	8.8	8.7	9.1
25 to 29 years	9.8	11.3	12.1	13.2	13.8	13.8
30 to 34 years	9.2	10.7	11.1	11.7	12.1	12.9
35 to 39 years	8.2	9.9	10.4	10.5	10.5	10.9
40 to 44 years	7.8	9.3	9.0	9.1	9.1	9.2
45 to 49 years	7.8	9.9	9.6	9.4	9.1	8.9
50 to 54 years	8.0	10.1	9.5	8.8	8.4	8.1
55 to 59 years	5.9	7.5	7.5	6.7	6.7	6.8
60 to 64 years	3.3	4.1	4.1	3.7	3.8	3.9
65 years and older	3.2	2.7	2.9	2.2	2.4	2.4
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

**Table 9.33b. North Carolina – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Column percent distribution, 2015–2020 (continued)**

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	2.4	1.7	1.7	1.8	1.7	1.6
Asian ⁴	†	†	†	†	†	†
Black or African American	32.1	30.6	31.2	32.0	31.7	31.4
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	59.7	63.1	62.3	61.4	61.8	61.9
Other	5.3	4.1	4.2	4.2	4.3	4.5
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	95.3	95.9	95.5	95.6	95.6	95.1
Hispanic or Latino origin	4.7	4.1	4.5	4.4	4.4	4.9
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	42.6	38.9	35.6	31.0	29.9	29.0
Puerto Rican	8.6	9.8	9.4	8.8	9.2	8.6
Cuban	2.6	2.8	2.8	2.4	2.0	1.7
Other specific	46.2	48.6	52.2	57.7	58.9	60.6
Total	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	‡	1.9	1.9	1.7
Not a veteran	<i>n/c</i>	<i>n/c</i>	‡	98.1	98.1	98.3
Total	<i>n/c</i>	<i>n/c</i>	‡	100.0	100.0	100.0

† Less than 1 percent of individuals; - - Quantity is zero; ‡ No data submitted; *n/c* Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

**Table 9.34a. North Dakota – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Number, 2015–2020**

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	14,943	14,842	13,978	13,866	15,143	14,382
Mental health diagnosis¹						
Trauma- and stressor-related disorders	2,138	1,491	1,099	940	1,918	2,547
Anxiety disorders	4,327	3,220	2,566	2,188	3,548	3,880
Attention deficit/hyperactivity disorder	1,813	1,435	1,133	982	1,321	1,308
Mood disorders	6,766	5,200	4,127	3,570	4,424	4,649
Bipolar disorders	1,856	1,482	1,273	1,107	1,039	1,073
Depressive disorders	4,972	3,763	2,892	2,493	3,422	3,624
Personality disorders	1,381	1,161	999	885	956	937
Schizophrenia and other psychotic disorders	1,717	1,564	1,448	1,348	1,457	1,490
Alcohol or substance-related disorders	39	78	55	2	--	--
Other mental disorders ²	7,742	5,813	4,741	4,400	2,637	2,616
Total	12,868	9,854	8,016	7,182	8,939	9,341
Gender						
Male	7,291	7,259	6,885	7,044	7,937	7,721
Female	7,612	7,536	7,053	6,747	7,082	6,626
Total	14,903	14,795	13,938	13,791	15,019	14,347
Age³						
0 to 11 years	1,387	1,473	1,399	1,212	988	803
12 to 14 years	731	752	699	639	719	672
15 to 17 years	841	924	860	779	782	762
18 to 20 years	669	679	651	646	741	697
21 to 24 years	1,091	1,025	910	932	1,093	1,070
25 to 29 years	1,551	1,486	1,380	1,532	1,790	1,758
30 to 34 years	1,504	1,448	1,413	1,542	1,772	1,705
35 to 39 years	1,295	1,253	1,232	1,255	1,524	1,470
40 to 44 years	1,079	1,070	993	959	1,144	1,170
45 to 49 years	1,071	1,045	944	986	1,019	941
50 to 54 years	1,178	1,150	1,038	942	982	914
55 to 59 years	1,013	977	931	941	988	917
60 to 64 years	667	717	678	680	723	711
65 years and older	860	838	845	814	876	792
Total	14,937	14,837	13,973	13,859	15,141	14,382

Continued. See notes at end of table.

**Table 9.34a. North Dakota – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Number, 2015–2020 (continued)**

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	1,541	1,609	1,520	1,551	1,904	2,238
Asian ⁴	51	75	86	76	92	124
Black or African American	304	350	356	403	508	603
Native Hawaiian or Other Pacific Islander	15	18	14	10	8	- -
White	12,581	12,233	11,451	11,180	11,760	10,351
Other	422	499	504	572	759	915
Total	14,914	14,784	13,931	13,792	15,031	14,231
Ethnicity						
Not of Hispanic or Latino origin	14,557	14,409	13,545	13,449	14,661	13,848
Hispanic or Latino origin	347	380	388	347	449	510
Total	14,904	14,789	13,933	13,796	15,110	14,358
Specific Hispanic or Latino origin						
Mexican	83	92	75	47	24	‡
Puerto Rican	13	16	21	19	4	‡
Cuban	4	4	2	1	1	‡
Other specific	81	82	71	48	18	‡
Total	181	194	169	115	47	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	‡	‡	181	317
Not a veteran	<i>n/c</i>	<i>n/c</i>	‡	‡	6,804	10,422
Total	<i>n/c</i>	<i>n/c</i>	‡	‡	6,985	10,739

- - Quantity is zero; ‡ No data submitted; n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.34b. North Dakota – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Column percent distribution, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	16.6	15.1	13.7	13.1	21.5	27.3
Anxiety disorders	33.6	32.7	32.0	30.5	39.7	41.5
Attention deficit/hyperactivity disorder	14.1	14.6	14.1	13.7	14.8	14.0
Mood disorders	52.6	52.8	51.5	49.7	49.5	49.8
Bipolar disorders	14.4	15.0	15.9	15.4	11.6	11.5
Depressive disorders	38.6	38.2	36.1	34.7	38.3	38.8
Personality disorders	10.7	11.8	12.5	12.3	10.7	10.0
Schizophrenia and other psychotic disorders	13.3	15.9	18.1	18.8	16.3	16.0
Alcohol or substance-related disorders	†	†	†	†	--	--
Other mental disorders ²	60.2	59.0	59.1	61.3	29.5	28.0
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	48.9	49.1	49.4	51.1	52.8	53.8
Female	51.1	50.9	50.6	48.9	47.2	46.2
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	9.3	9.9	10.0	8.7	6.5	5.6
12 to 14 years	4.9	5.1	5.0	4.6	4.7	4.7
15 to 17 years	5.6	6.2	6.2	5.6	5.2	5.3
18 to 20 years	4.5	4.6	4.7	4.7	4.9	4.8
21 to 24 years	7.3	6.9	6.5	6.7	7.2	7.4
25 to 29 years	10.4	10.0	9.9	11.1	11.8	12.2
30 to 34 years	10.1	9.8	10.1	11.1	11.7	11.9
35 to 39 years	8.7	8.4	8.8	9.1	10.1	10.2
40 to 44 years	7.2	7.2	7.1	6.9	7.6	8.1
45 to 49 years	7.2	7.0	6.8	7.1	6.7	6.5
50 to 54 years	7.9	7.8	7.4	6.8	6.5	6.4
55 to 59 years	6.8	6.6	6.7	6.8	6.5	6.4
60 to 64 years	4.5	4.8	4.9	4.9	4.8	4.9
65 years and older	5.8	5.6	6.0	5.9	5.8	5.5
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

**Table 9.34b. North Dakota – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Column percent distribution, 2015–2020 (continued)**

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	10.3	10.9	10.9	11.2	12.7	15.7
Asian ⁴	†	†	†	†	†	†
Black or African American	2.0	2.4	2.6	2.9	3.4	4.2
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	--
White	84.4	82.7	82.2	81.1	78.2	72.7
Other	2.8	3.4	3.6	4.1	5.0	6.4
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	97.7	97.4	97.2	97.5	97.0	96.4
Hispanic or Latino origin	2.3	2.6	2.8	2.5	3.0	3.6
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	45.9	47.4	44.4	40.9	51.1	‡
Puerto Rican	7.2	8.2	12.4	16.5	8.5	‡
Cuban	2.2	2.1	1.2	†	2.1	‡
Other specific	44.8	42.3	42.0	41.7	38.3	‡
Total	100.0	100.0	100.0	100.0	100.0	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	‡	‡	2.6	3.0
Not a veteran	<i>n/c</i>	<i>n/c</i>	‡	‡	97.4	97.0
Total	<i>n/c</i>	<i>n/c</i>	‡	‡	100.0	100.0

† Less than 1 percent of individuals; -- Quantity is zero; ‡ No data submitted; *n/c* Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.35a. Ohio – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	415,094	430,145	531,703	482,098	579,276	515,398
Mental health diagnosis¹						
Trauma- and stressor-related disorders	72,858	77,725	98,584	103,492	142,476	107,550
Anxiety disorders	54,438	55,566	89,315	99,452	135,471	110,594
Attention deficit/hyperactivity disorder	50,933	49,789	62,445	62,844	76,029	56,771
Mood disorders	164,402	160,124	161,011	182,802	233,782	178,502
Bipolar disorders	62,299	56,299	51,921	56,634	68,642	54,712
Depressive disorders	103,349	104,864	110,392	128,448	167,914	126,003
Personality disorders	5,138	4,620	7,896	8,103	9,198	7,226
Schizophrenia and other psychotic disorders	29,908	30,286	38,926	42,146	48,110	39,576
Alcohol or substance-related disorders	36,343	38,650	810	431	404	458
Other mental disorders ²	80,841	61,399	77,922	86,011	104,586	72,867
Total	389,414	412,118	432,983	449,184	574,476	429,022
Gender						
Male	195,810	200,320	259,214	224,455	264,945	244,714
Female	214,836	222,720	261,983	247,798	302,560	258,968
Total	410,646	423,040	521,197	472,253	567,505	503,682
Age³						
0 to 11 years	75,135	77,980	82,572	79,137	105,925	86,272
12 to 14 years	35,909	36,246	38,024	37,601	48,119	37,026
15 to 17 years	35,220	37,337	41,498	37,954	46,999	30,322
18 to 20 years	18,124	19,397	24,390	21,409	26,938	19,959
21 to 24 years	23,552	24,459	33,851	26,736	33,194	30,009
25 to 29 years	32,487	34,527	52,422	42,728	49,379	50,524
30 to 34 years	33,042	34,346	50,106	41,758	46,632	50,537
35 to 39 years	29,538	31,516	44,653	38,721	43,247	46,022
40 to 44 years	27,478	27,421	34,859	30,444	34,802	35,596
45 to 49 years	27,140	27,489	33,948	30,575	34,734	32,524
50 to 54 years	29,380	28,932	33,706	30,309	34,148	31,035
55 to 59 years	23,943	24,696	29,866	28,700	32,419	28,960
60 to 64 years	13,533	14,591	18,169	19,243	22,106	19,096
65 years and older	10,611	11,198	13,635	16,776	20,624	17,510
Total	415,092	430,135	531,699	482,091	579,266	515,392

Continued. See notes at end of table.

Table 9.35a. Ohio – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	465	835	1,474	1,527	2,019	1,899
Asian ⁴	1,161	1,423	2,049	2,164	3,078	2,613
Black or African American	96,663	86,900	115,991	107,673	132,537	116,058
Native Hawaiian or Other Pacific Islander	145	158	382	414	561	500
White	279,089	286,230	347,742	316,050	380,114	341,033
Other	3,376	2,073	3,258	2,445	2,742	1,970
Total	380,899	377,619	470,896	430,273	521,051	464,073
Ethnicity						
Not of Hispanic or Latino origin	291,823	180,809	478,596	437,949	526,497	464,863
Hispanic or Latino origin	7,442	6,449	14,116	13,584	16,918	14,043
Total	299,265	187,258	492,712	451,533	543,415	478,906
Specific Hispanic or Latino origin						
Mexican	343	532	557	431	477	414
Puerto Rican	2,812	1,347	580	419	454	311
Cuban	23	30	25	23	23	24
Other specific	1,351	1,197	3,152	2,171	2,380	1,394
Total	4,529	3,106	4,314	3,044	3,334	2,143
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡
Not a veteran	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡
Total	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡

‡ No data submitted; n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.35b. Ohio – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	18.7	18.9	22.8	23.0	24.8	25.1
Anxiety disorders	14.0	13.5	20.6	22.1	23.6	25.8
Attention deficit/hyperactivity disorder	13.1	12.1	14.4	14.0	13.2	13.2
Mood disorders	42.2	38.9	37.2	40.7	40.7	41.6
Bipolar disorders	16.0	13.7	12.0	12.6	11.9	12.8
Depressive disorders	26.5	25.4	25.5	28.6	29.2	29.4
Personality disorders	1.3	1.1	1.8	1.8	1.6	1.7
Schizophrenia and other psychotic disorders	7.7	7.3	9.0	9.4	8.4	9.2
Alcohol or substance-related disorders	9.3	9.4	†	†	†	†
Other mental disorders ²	20.8	14.9	18.0	19.1	18.2	17.0
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	47.7	47.4	49.7	47.5	46.7	48.6
Female	52.3	52.6	50.3	52.5	53.3	51.4
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	18.1	18.1	15.5	16.4	18.3	16.7
12 to 14 years	8.7	8.4	7.2	7.8	8.3	7.2
15 to 17 years	8.5	8.7	7.8	7.9	8.1	5.9
18 to 20 years	4.4	4.5	4.6	4.4	4.7	3.9
21 to 24 years	5.7	5.7	6.4	5.5	5.7	5.8
25 to 29 years	7.8	8.0	9.9	8.9	8.5	9.8
30 to 34 years	8.0	8.0	9.4	8.7	8.1	9.8
35 to 39 years	7.1	7.3	8.4	8.0	7.5	8.9
40 to 44 years	6.6	6.4	6.6	6.3	6.0	6.9
45 to 49 years	6.5	6.4	6.4	6.3	6.0	6.3
50 to 54 years	7.1	6.7	6.3	6.3	5.9	6.0
55 to 59 years	5.8	5.7	5.6	6.0	5.6	5.6
60 to 64 years	3.3	3.4	3.4	4.0	3.8	3.7
65 years and older	2.6	2.6	2.6	3.5	3.6	3.4
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.35b. Ohio – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	†	†	†	†	†	†
Asian ⁴	†	†	†	†	†	†
Black or African American	25.4	23.0	24.6	25.0	25.4	25.0
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	73.3	75.8	73.8	73.5	73.0	73.5
Other	†	†	†	†	†	†
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	97.5	96.6	97.1	97.0	96.9	97.1
Hispanic or Latino origin	2.5	3.4	2.9	3.0	3.1	2.9
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	7.6	17.1	12.9	14.2	14.3	19.3
Puerto Rican	62.1	43.4	13.4	13.8	13.6	14.5
Cuban	†	†	†	†	†	1.1
Other specific	29.8	38.5	73.1	71.3	71.4	65.0
Total	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡
Not a veteran	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡
Total	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡

† Less than 1 percent of individuals; ‡ No data submitted; n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.36a. Oklahoma – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	74,836	92,310	96,368	101,831	106,638	109,073
Mental health diagnosis¹						
Trauma- and stressor-related disorders	10,558	13,892	15,257	18,458	19,507	20,896
Anxiety disorders	9,883	13,483	14,883	16,688	15,674	16,707
Attention deficit/hyperactivity disorder	3,058	4,224	4,347	4,448	4,400	4,459
Mood disorders	36,341	43,412	44,523	45,212	46,552	47,808
Bipolar disorders	14,002	12,895	11,881	11,313	10,895	10,594
Depressive disorders	22,847	30,838	32,951	34,584	36,134	37,554
Personality disorders	1,058	936	763	973	569	397
Schizophrenia and other psychotic disorders	7,646	9,932	10,254	10,425	10,443	10,429
Alcohol or substance-related disorders	--	3	7	--	1	7
Other mental disorders ²	10,043	10,037	8,015	8,459	9,055	9,075
Total	65,794	82,899	85,138	87,517	91,952	94,809
Gender						
Male	34,903	42,394	44,570	47,871	49,931	51,406
Female	39,933	49,916	51,798	53,960	56,707	57,667
Total	74,836	92,310	96,368	101,831	106,638	109,073
Age³						
0 to 11 years	9,009	11,174	11,704	12,844	14,311	14,524
12 to 14 years	4,655	5,351	5,543	6,243	6,993	7,488
15 to 17 years	4,998	5,962	6,090	6,601	6,794	7,179
18 to 20 years	3,463	4,175	4,555	5,009	5,241	5,472
21 to 24 years	5,337	6,253	6,296	6,623	6,844	7,210
25 to 29 years	7,253	8,888	9,570	10,260	10,689	10,694
30 to 34 years	7,677	9,049	9,601	9,969	10,269	10,605
35 to 39 years	6,525	8,247	8,930	9,463	9,919	9,969
40 to 44 years	6,009	7,221	7,427	7,578	7,901	8,378
45 to 49 years	5,813	7,218	7,428	7,527	7,488	7,354
50 to 54 years	6,095	7,639	7,352	7,185	6,964	6,800
55 to 59 years	4,490	6,076	6,263	6,487	6,628	6,473
60 to 64 years	2,231	3,174	3,451	3,711	3,942	4,091
65 years and older	1,233	1,882	2,157	2,330	2,655	2,836
Total	74,788	92,309	96,367	101,830	106,638	109,073

Continued. See notes at end of table.

Table 9.36a. Oklahoma – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	6,210	7,727	8,275	8,954	9,559	9,849
Asian ⁴	427	496	542	556	614	657
Black or African American	9,598	12,231	12,713	13,142	13,537	13,553
Native Hawaiian or Other Pacific Islander	225	354	358	403	453	464
White	54,651	66,966	70,017	72,982	75,984	78,030
Other	3,707	4,519	4,453	5,788	6,484	6,517
Total	74,818	92,293	96,358	101,825	106,631	109,070
Ethnicity						
Not of Hispanic or Latino origin	69,642	85,376	89,087	93,894	98,015	99,877
Hispanic or Latino origin	5,185	6,926	7,277	7,929	8,614	9,188
Total	74,827	92,302	96,364	101,823	106,629	109,065
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	‡	1,645	1,358	1,901
Not a veteran	<i>n/c</i>	<i>n/c</i>	‡	60,475	51,585	74,978
Total	<i>n/c</i>	<i>n/c</i>	‡	62,120	52,943	76,879

- - Quantity is zero; ‡ No data submitted; n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.36b. Oklahoma – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	16.0	16.8	17.9	21.1	21.2	22.0
Anxiety disorders	15.0	16.3	17.5	19.1	17.0	17.6
Attention deficit/hyperactivity disorder	4.6	5.1	5.1	5.1	4.8	4.7
Mood disorders	55.2	52.4	52.3	51.7	50.6	50.4
Bipolar disorders	21.3	15.6	14.0	12.9	11.8	11.2
Depressive disorders	34.7	37.2	38.7	39.5	39.3	39.6
Personality disorders	1.6	1.1	†	1.1	†	†
Schizophrenia and other psychotic disorders	11.6	12.0	12.0	11.9	11.4	11.0
Alcohol or substance-related disorders	--	†	†	--	†	†
Other mental disorders ²	15.3	12.1	9.4	9.7	9.8	9.6
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	46.6	45.9	46.2	47.0	46.8	47.1
Female	53.4	54.1	53.8	53.0	53.2	52.9
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	12.0	12.1	12.1	12.6	13.4	13.3
12 to 14 years	6.2	5.8	5.8	6.1	6.6	6.9
15 to 17 years	6.7	6.5	6.3	6.5	6.4	6.6
18 to 20 years	4.6	4.5	4.7	4.9	4.9	5.0
21 to 24 years	7.1	6.8	6.5	6.5	6.4	6.6
25 to 29 years	9.7	9.6	9.9	10.1	10.0	9.8
30 to 34 years	10.3	9.8	10.0	9.8	9.6	9.7
35 to 39 years	8.7	8.9	9.3	9.3	9.3	9.1
40 to 44 years	8.0	7.8	7.7	7.4	7.4	7.7
45 to 49 years	7.8	7.8	7.7	7.4	7.0	6.7
50 to 54 years	8.1	8.3	7.6	7.1	6.5	6.2
55 to 59 years	6.0	6.6	6.5	6.4	6.2	5.9
60 to 64 years	3.0	3.4	3.6	3.6	3.7	3.8
65 years and older	1.6	2.0	2.2	2.3	2.5	2.6
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.36b. Oklahoma – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	8.3	8.4	8.6	8.8	9.0	9.0
Asian ⁴	†	†	†	†	†	†
Black or African American	12.8	13.3	13.2	12.9	12.7	12.4
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	73.0	72.6	72.7	71.7	71.3	71.5
Other	5.0	4.9	4.6	5.7	6.1	6.0
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	93.1	92.5	92.4	92.2	91.9	91.6
Hispanic or Latino origin	6.9	7.5	7.6	7.8	8.1	8.4
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	‡	2.6	2.6	2.5
Not a veteran	<i>n/c</i>	<i>n/c</i>	‡	97.4	97.4	97.5
Total	<i>n/c</i>	<i>n/c</i>	‡	100.0	100.0	100.0

† Less than 1 percent of individuals; - - Quantity is zero; ‡ No data submitted; *n/c* Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.37a. Oregon – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	131,581	126,319	126,493	134,743	142,930	145,434
Mental health diagnosis¹						
Trauma- and stressor-related disorders	36,211	48,867	48,971	51,852	58,645	54,561
Anxiety disorders	20,622	25,018	38,670	39,605	55,019	57,606
Attention deficit/hyperactivity disorder	3,555	12,168	1	12,383	7,693	16,169
Mood disorders	40,650	41,302	44,719	48,290	57,680	56,203
Bipolar disorders	11,150	11,025	11,061	11,584	13,083	12,514
Depressive disorders	30,178	30,596	34,193	37,475	46,377	45,310
Personality disorders	1,460	4,116	2,604	2,227	3,050	3,299
Schizophrenia and other psychotic disorders	10,843	11,677	12,962	12,772	13,380	13,150
Alcohol or substance-related disorders	1	6,580	30	15	21	30
Other mental disorders ²	7,164	9,286	13,712	21,963	26,682	30,354
Total	95,464	126,020	126,410	134,706	141,300	145,349
Gender						
Male	56,178	53,825	51,211	57,131	56,824	59,862
Female	75,385	72,439	75,221	77,508	85,842	85,568
Total	131,563	126,264	126,432	134,639	142,666	145,430
Age³						
0 to 11 years	18,654	19,518	15,394	20,028	16,304	22,375
12 to 14 years	10,179	9,347	7,515	9,142	9,355	11,596
15 to 17 years	11,261	10,648	10,358	11,298	11,611	12,143
18 to 20 years	6,251	5,242	5,599	7,010	8,289	6,906
21 to 24 years	7,483	6,790	7,168	7,105	8,181	8,462
25 to 29 years	10,904	11,321	12,666	12,525	13,295	13,868
30 to 34 years	11,217	11,311	12,382	12,644	13,719	13,422
35 to 39 years	9,877	9,932	10,877	11,282	12,069	11,905
40 to 44 years	9,162	8,394	8,802	8,752	9,755	9,403
45 to 49 years	8,882	8,399	8,846	8,573	8,879	8,571
50 to 54 years	9,320	8,704	8,834	8,010	8,465	7,985
55 to 59 years	7,667	7,488	7,904	7,942	8,696	8,099
60 to 64 years	5,284	5,106	5,652	5,842	6,691	5,857
65 years and older	5,438	4,119	4,496	4,590	7,621	4,842
Total	131,579	126,319	126,493	134,743	142,930	145,434

Continued. See notes at end of table.

Table 9.37a. Oregon – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	4,775	5,680	4,935	5,544	6,463	5,781
Asian ⁴	2,033	6,572	3,588	1,935	1,961	1,908
Black or African American	4,971	4,140	3,934	4,261	4,303	5,723
Native Hawaiian or Other Pacific Islander	528	387	420	566	697	646
White	94,375	88,991	85,853	92,812	94,289	93,160
Other	9,281	7,940	15,366	20,353	30,418	25,527
Total	115,963	113,710	114,096	125,471	138,131	132,745
Ethnicity						
Not of Hispanic or Latino origin	92,349	107,097	101,350	104,228	117,106	111,195
Hispanic or Latino origin	15,179	17,934	18,249	18,008	22,096	20,891
Total	107,528	125,031	119,599	122,236	139,202	132,086
Specific Hispanic or Latino origin						
Mexican	2,139	2,660	6,108	5,280	7,830	7,140
Puerto Rican	205	232	421	412	582	511
Cuban	162	178	376	377	518	471
Other specific	983	1,272	2,809	2,226	2,920	2,680
Total	3,489	4,342	9,714	8,295	11,850	10,802
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	2,016	2,301	2,429	1,776
Not a veteran	<i>n/c</i>	<i>n/c</i>	48,651	62,517	68,327	10,555
Total	<i>n/c</i>	<i>n/c</i>	50,667	64,818	70,756	12,331

n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.37b. Oregon – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	37.9	38.8	38.7	38.5	41.5	37.5
Anxiety disorders	21.6	19.9	30.6	29.4	38.9	39.6
Attention deficit/hyperactivity disorder	3.7	9.7	†	9.2	5.4	11.1
Mood disorders	42.6	32.8	35.4	35.8	40.8	38.7
Bipolar disorders	11.7	8.7	8.8	8.6	9.3	8.6
Depressive disorders	31.6	24.3	27.0	27.8	32.8	31.2
Personality disorders	1.5	3.3	2.1	1.7	2.2	2.3
Schizophrenia and other psychotic disorders	11.4	9.3	10.3	9.5	9.5	9.0
Alcohol or substance-related disorders	†	5.2	†	†	†	†
Other mental disorders ²	7.5	7.4	10.8	16.3	18.9	20.9
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	42.7	42.6	40.5	42.4	39.8	41.2
Female	57.3	57.4	59.5	57.6	60.2	58.8
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	14.2	15.5	12.2	14.9	11.4	15.4
12 to 14 years	7.7	7.4	5.9	6.8	6.5	8.0
15 to 17 years	8.6	8.4	8.2	8.4	8.1	8.3
18 to 20 years	4.8	4.1	4.4	5.2	5.8	4.7
21 to 24 years	5.7	5.4	5.7	5.3	5.7	5.8
25 to 29 years	8.3	9.0	10.0	9.3	9.3	9.5
30 to 34 years	8.5	9.0	9.8	9.4	9.6	9.2
35 to 39 years	7.5	7.9	8.6	8.4	8.4	8.2
40 to 44 years	7.0	6.6	7.0	6.5	6.8	6.5
45 to 49 years	6.8	6.6	7.0	6.4	6.2	5.9
50 to 54 years	7.1	6.9	7.0	5.9	5.9	5.5
55 to 59 years	5.8	5.9	6.2	5.9	6.1	5.6
60 to 64 years	4.0	4.0	4.5	4.3	4.7	4.0
65 years and older	4.1	3.3	3.6	3.4	5.3	3.3
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.37b. Oregon – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	4.1	5.0	4.3	4.4	4.7	4.4
Asian ⁴	1.8	5.8	3.1	1.5	1.4	1.4
Black or African American	4.3	3.6	3.4	3.4	3.1	4.3
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	81.4	78.3	75.2	74.0	68.3	70.2
Other	8.0	7.0	13.5	16.2	22.0	19.2
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	85.9	85.7	84.7	85.3	84.1	84.2
Hispanic or Latino origin	14.1	14.3	15.3	14.7	15.9	15.8
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	61.3	61.3	62.9	63.7	66.1	66.1
Puerto Rican	5.9	5.3	4.3	5.0	4.9	4.7
Cuban	4.6	4.1	3.9	4.5	4.4	4.4
Other specific	28.2	29.3	28.9	26.8	24.6	24.8
Total	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	4.0	3.5	3.4	14.4
Not a veteran	<i>n/c</i>	<i>n/c</i>	96.0	96.5	96.6	85.6
Total	<i>n/c</i>	<i>n/c</i>	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals; n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

**Table 9.38a. Pennsylvania – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Number, 2015–2020**

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	‡	543,618	573,439	565,437	586,913	603,630
Mental health diagnosis¹						
Trauma- and stressor-related disorders	‡	65,822	79,219	85,878	94,912	96,097
Anxiety disorders	‡	54,016	75,276	85,583	94,752	99,060
Attention deficit/hyperactivity disorder	‡	68,353	75,145	73,079	73,482	71,529
Mood disorders	‡	210,735	218,244	196,943	199,485	202,122
Bipolar disorders	‡	81,355	74,255	57,431	56,777	57,474
Depressive disorders	‡	131,070	145,968	141,123	144,239	145,626
Personality disorders	‡	4,079	5,499	5,867	5,758	4,892
Schizophrenia and other psychotic disorders	‡	39,543	41,390	40,321	40,288	40,906
Alcohol or substance-related disorders	‡	--	28	23	21	8
Other mental disorders ²	‡	77,112	92,200	95,606	97,308	88,195
Total	‡	460,504	499,313	488,061	503,931	514,459
Gender						
Male	‡	255,646	275,196	272,293	281,727	287,521
Female	‡	287,972	298,243	293,143	305,185	316,109
Total	‡	543,618	573,439	565,436	586,912	603,630
Age³						
0 to 11 years	‡	96,368	98,114	95,588	98,880	100,561
12 to 14 years	‡	39,494	41,073	39,908	42,395	44,863
15 to 17 years	‡	39,163	40,897	39,755	41,062	42,531
18 to 20 years	‡	25,527	25,706	25,079	26,308	27,092
21 to 24 years	‡	31,497	31,834	30,964	31,204	31,336
25 to 29 years	‡	45,573	49,534	49,958	51,436	52,213
30 to 34 years	‡	42,999	46,701	47,120	49,479	51,451
35 to 39 years	‡	36,360	40,717	41,798	44,216	46,257
40 to 44 years	‡	33,704	35,093	34,136	34,989	36,258
45 to 49 years	‡	36,765	38,411	36,768	36,881	36,537
50 to 54 years	‡	39,546	41,697	39,184	39,111	38,996
55 to 59 years	‡	32,249	34,890	34,829	36,077	37,101
60 to 64 years	‡	20,816	23,360	23,970	26,086	27,627
65 years and older	‡	23,552	25,409	26,375	28,784	30,804
Total	‡	543,613	573,436	565,432	586,908	603,627

Continued. See notes at end of table.

**Table 9.38a. Pennsylvania – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Number, 2015–2020 (continued)**

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	‡	967	1,019	1,111	1,192	1,299
Asian ⁴	‡	6,223	6,897	7,730	8,345	8,767
Black or African American	‡	120,993	125,989	123,197	127,212	129,743
Native Hawaiian or Other Pacific Islander	‡	267	259	287	439	568
White	‡	348,977	369,793	366,963	378,580	388,623
Other	‡	62,692	63,900	58,714	62,002	63,934
Total	‡	540,119	567,857	558,002	577,770	592,934
Ethnicity						
Not of Hispanic or Latino origin	‡	469,290	497,817	494,553	511,507	525,726
Hispanic or Latino origin	‡	72,763	74,309	68,596	73,455	76,143
Total	‡	542,053	572,126	563,149	584,962	601,869
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡
Not a veteran	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡
Total	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡

- - Quantity is zero; ‡ No data submitted; n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.38b. Pennsylvania – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	‡	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	‡	14.3	15.9	17.6	18.8	18.7
Anxiety disorders	‡	11.7	15.1	17.5	18.8	19.3
Attention deficit/hyperactivity disorder	‡	14.8	15.0	15.0	14.6	13.9
Mood disorders	‡	45.8	43.7	40.4	39.6	39.3
Bipolar disorders	‡	17.7	14.9	11.8	11.3	11.2
Depressive disorders	‡	28.5	29.2	28.9	28.6	28.3
Personality disorders	‡	†	1.1	1.2	1.1	†
Schizophrenia and other psychotic disorders	‡	8.6	8.3	8.3	8.0	8.0
Alcohol or substance-related disorders	‡	--	†	†	†	†
Other mental disorders ²	‡	16.7	18.5	19.6	19.3	17.1
Total	‡	100.0	100.0	100.0	100.0	100.0
Gender						
Male	‡	47.0	48.0	48.2	48.0	47.6
Female	‡	53.0	52.0	51.8	52.0	52.4
Total	‡	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	‡	17.7	17.1	16.9	16.8	16.7
12 to 14 years	‡	7.3	7.2	7.1	7.2	7.4
15 to 17 years	‡	7.2	7.1	7.0	7.0	7.0
18 to 20 years	‡	4.7	4.5	4.4	4.5	4.5
21 to 24 years	‡	5.8	5.6	5.5	5.3	5.2
25 to 29 years	‡	8.4	8.6	8.8	8.8	8.6
30 to 34 years	‡	7.9	8.1	8.3	8.4	8.5
35 to 39 years	‡	6.7	7.1	7.4	7.5	7.7
40 to 44 years	‡	6.2	6.1	6.0	6.0	6.0
45 to 49 years	‡	6.8	6.7	6.5	6.3	6.1
50 to 54 years	‡	7.3	7.3	6.9	6.7	6.5
55 to 59 years	‡	5.9	6.1	6.2	6.1	6.1
60 to 64 years	‡	3.8	4.1	4.2	4.4	4.6
65 years and older	‡	4.3	4.4	4.7	4.9	5.1
Total	‡	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.38b. Pennsylvania – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	‡	†	†	†	†	†
Asian ⁴	‡	1.2	1.2	1.4	1.4	1.5
Black or African American	‡	22.4	22.2	22.1	22.0	21.9
Native Hawaiian or Other Pacific Islander	‡	†	†	†	†	†
White	‡	64.6	65.1	65.8	65.5	65.5
Other	‡	11.6	11.3	10.5	10.7	10.8
Total	‡	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	‡	86.6	87.0	87.8	87.4	87.3
Hispanic or Latino origin	‡	13.4	13.0	12.2	12.6	12.7
Total	‡	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡
Not a veteran	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡
Total	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡

† Less than 1 percent of individuals; - - Quantity is zero; ‡ No data submitted; n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.39a. Puerto Rico – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020

<i>Selected characteristics</i>	2015	2016	2017	2018	2019	2020
Total individuals	23,194	17,567	25,529	2,918	3,947	3,126
Mental health diagnosis¹						
Trauma- and stressor-related disorders	7	7	130	152	160	88
Anxiety disorders	5	5	156	326	352	291
Attention deficit/hyperactivity disorder	65	64	374	337	404	192
Mood disorders	21	21	585	1,063	1,137	814
Bipolar disorders	4	4	130	235	285	226
Depressive disorders	17	17	468	848	876	601
Personality disorders	--	--	5	33	37	29
Schizophrenia and other psychotic disorders	8	7	304	702	1,316	1,379
Alcohol or substance-related disorders	11	10	407	620	696	441
Other mental disorders ²	62	62	437	622	682	393
Total	141	139	1,667	2,669	3,582	2,950
Gender						
Male	12,054	9,098	13,490	1,700	2,320	1,921
Female	11,139	8,468	12,037	1,218	1,627	1,205
Total	23,193	17,566	25,527	2,918	3,947	3,126
Age³						
0 to 11 years	1,192	652	1,062	240	286	162
12 to 14 years	809	436	810	194	200	67
15 to 17 years	1,061	624	1,110	215	272	116
18 to 20 years	736	731	1,110	76	104	74
21 to 24 years	997	662	1,004	119	160	105
25 to 29 years	1,396	1,129	1,560	194	254	198
30 to 34 years	1,503	1,141	1,593	193	262	218
35 to 39 years	1,592	1,190	1,746	193	253	220
40 to 44 years	1,690	1,239	1,739	188	262	220
45 to 49 years	2,125	1,501	2,050	217	314	265
50 to 54 years	2,547	1,934	2,713	285	365	316
55 to 59 years	2,528	1,949	2,717	266	410	394
60 to 64 years	2,080	1,685	2,425	207	312	324
65 years and older	2,936	2,694	3,888	331	493	447
Total	23,192	17,567	25,527	2,918	3,947	3,126

Continued. See notes at end of table.

Table 9.39a. Puerto Rico – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	36	8	64	34	63	41
Asian ⁴	9	--	12	2	12	5
Black or African American	259	207	303	49	71	32
Native Hawaiian or Other Pacific Islander	--	--	4	6	4	--
White	1,166	955	1,685	439	639	422
Other	15,318	11,933	16,962	2,200	2,883	2,472
Total	16,788	13,103	19,030	2,730	3,672	2,972
Ethnicity						
Not of Hispanic or Latino origin	106	76	114	21	39	42
Hispanic or Latino origin	22,864	17,299	25,186	2,879	3,875	3,068
Total	22,970	17,375	25,300	2,900	3,914	3,110
Specific Hispanic or Latino origin						
Mexican	29	23	30	10	9	8
Puerto Rican	22,197	16,715	24,494	2,787	3,754	2,950
Cuban	44	37	49	6	9	10
Other specific	509	468	531	39	42	41
Total	22,779	17,243	25,104	2,842	3,814	3,009
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	‡	86	127	94
Not a veteran	<i>n/c</i>	<i>n/c</i>	‡	1,926	2,612	2,365
Total	<i>n/c</i>	<i>n/c</i>	‡	2,012	2,739	2,459

-- Quantity is zero; ‡ No data submitted; n/c Not collected.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.39b. Puerto Rico – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	5.0	5.0	7.8	5.7	4.5	3.0
Anxiety disorders	3.5	3.6	9.4	12.2	9.8	9.9
Attention deficit/hyperactivity disorder	46.1	46.0	22.4	12.6	11.3	6.5
Mood disorders	14.9	15.1	35.1	39.8	31.7	27.6
Bipolar disorders	2.8	2.9	7.8	8.8	8.0	7.7
Depressive disorders	12.1	12.2	28.1	31.8	24.5	20.4
Personality disorders	--	--	†	1.2	1.0	†
Schizophrenia and other psychotic disorders	5.7	5.0	18.2	26.3	36.7	46.7
Alcohol or substance-related disorders	7.8	7.2	24.4	23.2	19.4	14.9
Other mental disorders ²	44.0	44.6	26.2	23.3	19.0	13.3
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	52.0	51.8	52.8	58.3	58.8	61.5
Female	48.0	48.2	47.2	41.7	41.2	38.5
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	5.1	3.7	4.2	8.2	7.2	5.2
12 to 14 years	3.5	2.5	3.2	6.6	5.1	2.1
15 to 17 years	4.6	3.6	4.3	7.4	6.9	3.7
18 to 20 years	3.2	4.2	4.3	2.6	2.6	2.4
21 to 24 years	4.3	3.8	3.9	4.1	4.1	3.4
25 to 29 years	6.0	6.4	6.1	6.6	6.4	6.3
30 to 34 years	6.5	6.5	6.2	6.6	6.6	7.0
35 to 39 years	6.9	6.8	6.8	6.6	6.4	7.0
40 to 44 years	7.3	7.1	6.8	6.4	6.6	7.0
45 to 49 years	9.2	8.5	8.0	7.4	8.0	8.5
50 to 54 years	11.0	11.0	10.6	9.8	9.2	10.1
55 to 59 years	10.9	11.1	10.6	9.1	10.4	12.6
60 to 64 years	9.0	9.6	9.5	7.1	7.9	10.4
65 years and older	12.7	15.3	15.2	11.3	12.5	14.3
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.39b. Puerto Rico – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	†	†	†	1.2	1.7	1.4
Asian ⁴	†	--	†	†	†	†
Black or African American	1.5	1.6	1.6	1.8	1.9	1.1
Native Hawaiian or Other Pacific Islander	--	--	†	†	†	--
White	6.9	7.3	8.9	16.1	17.4	14.2
Other	91.2	91.1	89.1	80.6	78.5	83.2
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	†	†	†	†	†	1.4
Hispanic or Latino origin	99.5	99.6	99.5	99.3	99.0	98.6
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	†	†	†	†	†	†
Puerto Rican	97.4	96.9	97.6	98.1	98.4	98.0
Cuban	†	†	†	†	†	†
Other specific	2.2	2.7	2.1	1.4	1.1	1.4
Total	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	‡	4.3	4.6	3.8
Not a veteran	<i>n/c</i>	<i>n/c</i>	‡	95.7	95.4	96.2
Total	<i>n/c</i>	<i>n/c</i>	‡	100.0	100.0	100.0

† Less than 1 percent of individuals; -- Quantity is zero; ‡ No data submitted; *n/c* Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

**Table 9.40a. Rhode Island – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Number, 2015–2020**

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	32,061	31,643	31,249	30,383	32,563	32,645
Mental health diagnosis¹						
Trauma- and stressor-related disorders	4,278	6,011	6,526	3,958	5,305	5,568
Anxiety disorders	3,868	6,198	6,811	8,443	10,551	10,635
Attention deficit/hyperactivity disorder	1,962	2,774	2,907	2,971	3,568	3,724
Mood disorders	9,500	13,211	14,026	13,814	16,708	16,816
Bipolar disorders	3,699	3,548	3,672	3,607	4,144	4,176
Depressive disorders	5,941	9,911	10,649	10,675	13,156	13,251
Personality disorders	923	1,319	1,305	1,388	1,487	1,487
Schizophrenia and other psychotic disorders	1,754	3,975	3,971	4,069	4,422	4,466
Alcohol or substance-related disorders	110	3	1	4	5	8
Other mental disorders ²	2,203	4,177	4,220	4,584	5,458	5,427
Total	15,841	23,666	24,895	23,899	29,131	29,366
Gender						
Male	16,021	15,830	15,512	15,082	16,201	16,273
Female	16,040	15,813	15,737	15,284	16,347	16,370
Total	32,061	31,643	31,249	30,366	32,548	32,643
Age³						
0 to 11 years	3,200	3,170	3,567	3,090	3,215	2,888
12 to 14 years	1,801	1,729	1,742	1,751	2,006	1,984
15 to 17 years	2,239	2,108	1,999	2,005	2,121	2,055
18 to 20 years	1,480	1,548	1,527	1,489	1,694	1,810
21 to 24 years	1,755	1,779	1,722	1,658	1,858	1,879
25 to 29 years	2,367	2,423	2,472	2,335	2,498	2,449
30 to 34 years	2,389	2,323	2,286	2,337	2,528	2,601
35 to 39 years	2,247	2,207	2,244	2,254	2,398	2,500
40 to 44 years	2,396	2,210	2,003	1,921	2,079	2,168
45 to 49 years	2,937	2,754	2,604	2,460	2,471	2,349
50 to 54 years	3,163	3,169	2,853	2,790	2,867	2,733
55 to 59 years	2,599	2,705	2,584	2,623	2,753	2,852
60 to 64 years	1,607	1,669	1,718	1,853	2,012	2,121
65 years and older	1,880	1,849	1,928	1,816	2,063	2,256
Total	32,060	31,643	31,249	30,382	32,563	32,645

Continued. See notes at end of table.

**Table 9.40a. Rhode Island – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Number, 2015–2020 (continued)**

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	243	243	230	244	235	239
Asian ⁴	373	355	338	330	349	335
Black or African American	3,330	3,489	3,541	3,510	3,825	3,850
Native Hawaiian or Other Pacific Islander	40	43	49	55	75	74
White	22,356	22,138	21,738	20,961	22,515	22,235
Other	277	257	269	634	922	1,160
Total	26,619	26,525	26,165	25,734	27,921	27,893
Ethnicity						
Not of Hispanic or Latino origin	23,048	22,777	22,592	21,648	22,879	22,219
Hispanic or Latino origin	6,677	6,771	6,706	6,690	7,197	7,194
Total	29,725	29,548	29,298	28,338	30,076	29,413
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	678	582	586	561
Not a veteran	<i>n/c</i>	<i>n/c</i>	19,975	18,414	20,276	21,364
Total	<i>n/c</i>	<i>n/c</i>	20,653	18,996	20,862	21,925

‡ No data submitted; n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.40b. Rhode Island – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	27.0	25.4	26.2	16.6	18.2	19.0
Anxiety disorders	24.4	26.2	27.4	35.3	36.2	36.2
Attention deficit/hyperactivity disorder	12.4	11.7	11.7	12.4	12.2	12.7
Mood disorders	60.0	55.8	56.3	57.8	57.4	57.3
Bipolar disorders	23.4	15.0	14.7	15.1	14.2	14.2
Depressive disorders	37.5	41.9	42.8	44.7	45.2	45.1
Personality disorders	5.8	5.6	5.2	5.8	5.1	5.1
Schizophrenia and other psychotic disorders	11.1	16.8	16.0	17.0	15.2	15.2
Alcohol or substance-related disorders	†	†	†	†	†	†
Other mental disorders ²	13.9	17.6	17.0	19.2	18.7	18.5
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	50.0	50.0	49.6	49.7	49.8	49.9
Female	50.0	50.0	50.4	50.3	50.2	50.1
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	10.0	10.0	11.4	10.2	9.9	8.8
12 to 14 years	5.6	5.5	5.6	5.8	6.2	6.1
15 to 17 years	7.0	6.7	6.4	6.6	6.5	6.3
18 to 20 years	4.6	4.9	4.9	4.9	5.2	5.5
21 to 24 years	5.5	5.6	5.5	5.5	5.7	5.8
25 to 29 years	7.4	7.7	7.9	7.7	7.7	7.5
30 to 34 years	7.5	7.3	7.3	7.7	7.8	8.0
35 to 39 years	7.0	7.0	7.2	7.4	7.4	7.7
40 to 44 years	7.5	7.0	6.4	6.3	6.4	6.6
45 to 49 years	9.2	8.7	8.3	8.1	7.6	7.2
50 to 54 years	9.9	10.0	9.1	9.2	8.8	8.4
55 to 59 years	8.1	8.5	8.3	8.6	8.5	8.7
60 to 64 years	5.0	5.3	5.5	6.1	6.2	6.5
65 years and older	5.9	5.8	6.2	6.0	6.3	6.9
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.40b. Rhode Island – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	†	†	†	†	†	†
Asian ⁴	1.4	1.3	1.3	1.3	1.2	1.2
Black or African American	12.5	13.2	13.5	13.6	13.7	13.8
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	84.0	83.5	83.1	81.5	80.6	79.7
Other	1.0	†	1.0	2.5	3.3	4.2
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	77.5	77.1	77.1	76.4	76.1	75.5
Hispanic or Latino origin	22.5	22.9	22.9	23.6	23.9	24.5
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	3.3	3.1	2.8	2.6
Not a veteran	<i>n/c</i>	<i>n/c</i>	96.7	96.9	97.2	97.4
Total	<i>n/c</i>	<i>n/c</i>	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals; ‡ No data submitted; n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

**Table 9.41a. South Carolina – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Number, 2015–2020**

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	83,286	85,254	85,754	87,192	99,514	99,755
Mental health diagnosis¹						
Trauma- and stressor-related disorders	13,786	14,128	15,406	18,177	22,828	25,244
Anxiety disorders	14,136	15,265	16,217	17,956	21,158	22,886
Attention deficit/hyperactivity disorder	15,027	15,024	15,020	15,295	16,842	16,762
Mood disorders	38,595	35,855	35,041	36,752	41,894	41,740
Bipolar disorders	13,949	11,526	10,863	11,405	12,413	12,157
Depressive disorders	24,882	24,543	24,410	25,660	29,819	29,930
Personality disorders	3,812	3,596	3,545	3,897	4,492	4,420
Schizophrenia and other psychotic disorders	15,763	15,418	15,373	16,485	17,340	17,323
Alcohol or substance-related disorders	12,132	10,924	10,690	12,842	15,210	15,207
Other mental disorders ²	14,472	16,474	17,176	18,010	19,887	20,090
Total	80,113	79,913	80,489	84,342	94,854	95,263
Gender						
Male	38,692	39,471	39,724	40,711	46,392	46,697
Female	44,459	45,770	46,022	46,472	53,108	53,037
Total	83,151	85,241	85,746	87,183	99,500	99,734
Age³						
0 to 11 years	11,659	12,159	12,620	13,249	15,284	14,995
12 to 14 years	7,235	7,340	7,337	7,765	9,538	10,017
15 to 17 years	7,731	7,906	8,006	8,199	9,600	10,042
18 to 20 years	3,086	3,255	3,303	3,450	4,462	4,660
21 to 24 years	3,975	4,057	3,918	3,965	4,669	4,887
25 to 29 years	5,626	5,768	6,065	6,043	6,926	6,826
30 to 34 years	5,749	5,939	5,762	5,812	6,781	6,963
35 to 39 years	5,531	5,803	5,779	5,781	6,580	6,547
40 to 44 years	5,746	5,604	5,301	5,365	5,847	5,798
45 to 49 years	6,384	6,245	6,161	5,816	6,242	5,814
50 to 54 years	7,160	7,129	6,743	6,431	6,588	6,168
55 to 59 years	5,959	6,131	6,236	6,226	6,757	6,520
60 to 64 years	3,818	3,992	4,231	4,463	4,833	4,861
65 years and older	3,621	3,919	4,289	4,623	5,403	5,657
Total	83,280	85,247	85,751	87,188	99,510	99,755

Continued. See notes at end of table.

**Table 9.41a. South Carolina – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Number, 2015–2020 (continued)**

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	258	271	273	279	343	450
Asian ⁴	257	285	272	61	274	337
Black or African American	33,090	33,758	34,076	34,214	38,137	37,837
Native Hawaiian or Other Pacific Islander	11	18	18	16	18	14
White	45,170	46,045	45,728	45,681	51,566	50,777
Other	3,090	3,247	4,044	4,376	7,131	7,913
Total	81,876	83,624	84,411	84,627	97,469	97,328
Ethnicity						
Not of Hispanic or Latino origin	61,715	63,533	64,559	65,519	73,498	72,403
Hispanic or Latino origin	2,494	2,771	2,947	3,144	3,733	3,944
Total	64,209	66,304	67,506	68,663	77,231	76,347
Specific Hispanic or Latino origin						
Mexican	636	847	955	937	1,123	1,189
Puerto Rican	153	207	240	286	298	334
Cuban	578	508	417	368	388	362
Other specific	1,109	1,187	1,315	1,539	1,910	2,041
Total	2,476	2,749	2,927	3,130	3,719	3,926
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	905	836	930	930
Not a veteran	<i>n/c</i>	<i>n/c</i>	36,906	57,086	64,099	63,740
Total	<i>n/c</i>	<i>n/c</i>	37,811	57,922	65,029	64,670

n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

**Table 9.41b. South Carolina – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Column percent distribution, 2015–2020**

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	17.2	17.7	19.1	21.6	24.1	26.5
Anxiety disorders	17.6	19.1	20.1	21.3	22.3	24.0
Attention deficit/hyperactivity disorder	18.8	18.8	18.7	18.1	17.8	17.6
Mood disorders	48.2	44.9	43.5	43.6	44.2	43.8
Bipolar disorders	17.4	14.4	13.5	13.5	13.1	12.8
Depressive disorders	31.1	30.7	30.3	30.4	31.4	31.4
Personality disorders	4.8	4.5	4.4	4.6	4.7	4.6
Schizophrenia and other psychotic disorders	19.7	19.3	19.1	19.5	18.3	18.2
Alcohol or substance-related disorders	15.1	13.7	13.3	15.2	16.0	16.0
Other mental disorders ²	18.1	20.6	21.3	21.4	21.0	21.1
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	46.5	46.3	46.3	46.7	46.6	46.8
Female	53.5	53.7	53.7	53.3	53.4	53.2
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	14.0	14.3	14.7	15.2	15.4	15.0
12 to 14 years	8.7	8.6	8.6	8.9	9.6	10.0
15 to 17 years	9.3	9.3	9.3	9.4	9.6	10.1
18 to 20 years	3.7	3.8	3.9	4.0	4.5	4.7
21 to 24 years	4.8	4.8	4.6	4.5	4.7	4.9
25 to 29 years	6.8	6.8	7.1	6.9	7.0	6.8
30 to 34 years	6.9	7.0	6.7	6.7	6.8	7.0
35 to 39 years	6.6	6.8	6.7	6.6	6.6	6.6
40 to 44 years	6.9	6.6	6.2	6.2	5.9	5.8
45 to 49 years	7.7	7.3	7.2	6.7	6.3	5.8
50 to 54 years	8.6	8.4	7.9	7.4	6.6	6.2
55 to 59 years	7.2	7.2	7.3	7.1	6.8	6.5
60 to 64 years	4.6	4.7	4.9	5.1	4.9	4.9
65 years and older	4.3	4.6	5.0	5.3	5.4	5.7
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

**Table 9.41b. South Carolina – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Column percent distribution, 2015–2020 (continued)**

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	†	†	†	†	†	†
Asian ⁴	†	†	†	†	†	†
Black or African American	40.4	40.4	40.4	40.4	39.1	38.9
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	55.2	55.1	54.2	54.0	52.9	52.2
Other	3.8	3.9	4.8	5.2	7.3	8.1
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	96.1	95.8	95.6	95.4	95.2	94.8
Hispanic or Latino origin	3.9	4.2	4.4	4.6	4.8	5.2
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	25.7	30.8	32.6	29.9	30.2	30.3
Puerto Rican	6.2	7.5	8.2	9.1	8.0	8.5
Cuban	23.3	18.5	14.2	11.8	10.4	9.2
Other specific	44.8	43.2	44.9	49.2	51.4	52.0
Total	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	2.4	1.4	1.4	1.4
Not a veteran	<i>n/c</i>	<i>n/c</i>	97.6	98.6	98.6	98.6
Total	<i>n/c</i>	<i>n/c</i>	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals; n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

**Table 9.42a. South Dakota – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Number, 2015–2020**

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	15,928	16,055	15,726	16,278	16,207	15,806
Mental health diagnosis¹						
Trauma- and stressor-related disorders	3,800	3,831	3,557	3,943	4,083	4,121
Anxiety disorders	1,222	1,260	1,405	1,610	1,634	1,696
Attention deficit/hyperactivity disorder	1,313	1,262	1,223	1,243	1,163	1,126
Mood disorders	5,821	5,107	5,005	5,344	5,298	4,964
Bipolar disorders	1,902	1,585	1,487	1,500	1,441	1,350
Depressive disorders	3,923	3,527	3,522	3,848	3,862	3,621
Personality disorders	166	98	78	64	49	51
Schizophrenia and other psychotic disorders	1,804	1,478	1,418	1,450	1,385	1,345
Alcohol or substance-related disorders	115	65	52	56	56	46
Other mental disorders ²	2,200	2,486	2,327	1,938	1,787	1,778
Total	15,837	15,167	14,717	15,310	15,148	14,789
Gender						
Male	7,709	7,569	7,339	7,509	7,396	7,198
Female	8,219	8,486	8,387	8,769	8,811	8,608
Total	15,928	16,055	15,726	16,278	16,207	15,806
Age³						
0 to 11 years	2,545	2,582	2,389	2,508	2,495	2,320
12 to 14 years	1,259	1,290	1,216	1,481	1,563	1,592
15 to 17 years	1,414	1,414	1,427	1,445	1,545	1,533
18 to 20 years	689	790	731	817	743	731
21 to 24 years	913	949	930	934	889	902
25 to 29 years	1,179	1,210	1,219	1,158	1,182	1,179
30 to 34 years	1,241	1,253	1,191	1,181	1,213	1,133
35 to 39 years	1,086	1,131	1,140	1,218	1,165	1,162
40 to 44 years	995	901	888	895	886	898
45 to 49 years	917	870	901	931	919	850
50 to 54 years	1,178	1,079	959	930	846	798
55 to 59 years	918	977	1,013	1,022	1,015	942
60 to 64 years	656	642	673	696	701	725
65 years and older	937	966	1,049	1,061	1,045	1,041
Total	15,927	16,054	15,726	16,277	16,207	15,806

Continued. See notes at end of table.

**Table 9.42a. South Dakota – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Number, 2015–2020 (continued)**

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	297	293	228	218	277	281
Asian ⁴	112	90	116	113	117	107
Black or African American	431	445	478	479	500	489
Native Hawaiian or Other Pacific Islander	28	19	27	22	18	23
White	11,690	11,630	11,311	11,549	11,416	11,276
Other	3,322	3,551	3,543	3,844	3,837	3,592
Total	15,880	16,028	15,703	16,225	16,165	15,768
Ethnicity						
Not of Hispanic or Latino origin	15,104	15,349	15,012	15,440	15,414	14,999
Hispanic or Latino origin	646	642	669	771	728	745
Total	15,750	15,991	15,681	16,211	16,142	15,744
Specific Hispanic or Latino origin						
Mexican	212	210	204	249	246	239
Puerto Rican	50	40	68	64	53	62
Cuban	15	8	15	16	18	11
Other specific	106	89	99	120	129	128
Total	383	347	386	449	446	440
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡
Not a veteran	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡
Total	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡

‡ No data submitted; n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.42b. South Dakota – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Column percent distribution, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	24.0	25.3	24.2	25.8	27.0	27.9
Anxiety disorders	7.7	8.3	9.5	10.5	10.8	11.5
Attention deficit/hyperactivity disorder	8.3	8.3	8.3	8.1	7.7	7.6
Mood disorders	36.8	33.7	34.0	34.9	35.0	33.6
Bipolar disorders	12.0	10.5	10.1	9.8	9.5	9.1
Depressive disorders	24.8	23.3	23.9	25.1	25.5	24.5
Personality disorders	1.0	†	†	†	†	†
Schizophrenia and other psychotic disorders	11.4	9.7	9.6	9.5	9.1	9.1
Alcohol or substance-related disorders	†	†	†	†	†	†
Other mental disorders ²	13.9	16.4	15.8	12.7	11.8	12.0
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	48.4	47.1	46.7	46.1	45.6	45.5
Female	51.6	52.9	53.3	53.9	54.4	54.5
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	16.0	16.1	15.2	15.4	15.4	14.7
12 to 14 years	7.9	8.0	7.7	9.1	9.6	10.1
15 to 17 years	8.9	8.8	9.1	8.9	9.5	9.7
18 to 20 years	4.3	4.9	4.6	5.0	4.6	4.6
21 to 24 years	5.7	5.9	5.9	5.7	5.5	5.7
25 to 29 years	7.4	7.5	7.8	7.1	7.3	7.5
30 to 34 years	7.8	7.8	7.6	7.3	7.5	7.2
35 to 39 years	6.8	7.0	7.2	7.5	7.2	7.4
40 to 44 years	6.2	5.6	5.6	5.5	5.5	5.7
45 to 49 years	5.8	5.4	5.7	5.7	5.7	5.4
50 to 54 years	7.4	6.7	6.1	5.7	5.2	5.0
55 to 59 years	5.8	6.1	6.4	6.3	6.3	6.0
60 to 64 years	4.1	4.0	4.3	4.3	4.3	4.6
65 years and older	5.9	6.0	6.7	6.5	6.4	6.6
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

**Table 9.42b. South Dakota – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Column percent distribution, 2015–2020 (continued)**

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	1.9	1.8	1.5	1.3	1.7	1.8
Asian ⁴	†	†	†	†	†	†
Black or African American	2.7	2.8	3.0	3.0	3.1	3.1
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	73.6	72.6	72.0	71.2	70.6	71.5
Other	20.9	22.2	22.6	23.7	23.7	22.8
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	95.9	96.0	95.7	95.2	95.5	95.3
Hispanic or Latino origin	4.1	4.0	4.3	4.8	4.5	4.7
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	55.4	60.5	52.8	55.5	55.2	54.3
Puerto Rican	13.1	11.5	17.6	14.3	11.9	14.1
Cuban	3.9	2.3	3.9	3.6	4.0	2.5
Other specific	27.7	25.6	25.6	26.7	28.9	29.1
Total	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡
Not a veteran	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡
Total	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡

† Less than 1 percent of individuals; ‡ No data submitted; n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.43a. Tennessee – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	115,168	118,609	108,269	117,555	112,426	98,452
Mental health diagnosis¹						
Trauma- and stressor-related disorders	11,920	11,552	9,830	12,150	13,998	13,862
Anxiety disorders	17,575	15,536	14,835	16,793	17,644	16,477
Attention deficit/hyperactivity disorder	7,024	6,851	3,741	4,952	4,928	3,959
Mood disorders	60,981	45,537	41,452	41,488	40,667	38,453
Bipolar disorders	27,499	16,204	15,112	14,829	13,341	12,487
Depressive disorders	36,619	30,980	27,915	28,090	28,720	27,003
Personality disorders	1,932	1,315	1,146	1,522	1,263	1,175
Schizophrenia and other psychotic disorders	11,189	8,982	8,894	8,885	8,451	8,154
Alcohol or substance-related disorders	35	19	--	--	1	--
Other mental disorders ²	10,805	12,270	7,458	8,504	8,621	7,358
Total	87,823	76,068	63,388	64,039	65,194	62,819
Gender						
Male	53,099	54,977	50,047	56,091	53,841	48,718
Female	61,449	63,050	57,823	61,100	58,188	49,431
Total	114,548	118,027	107,870	117,191	112,029	98,149
Age³						
0 to 11 years	8,503	8,598	3,401	7,081	7,667	5,619
12 to 14 years	5,921	6,194	5,142	6,107	5,993	4,867
15 to 17 years	6,995	7,601	7,102	6,976	6,719	5,169
18 to 20 years	4,895	5,547	5,486	5,677	5,512	4,846
21 to 24 years	8,391	8,510	8,166	8,448	8,077	7,481
25 to 29 years	11,647	12,347	12,373	13,242	12,425	11,436
30 to 34 years	11,627	12,096	11,521	12,208	11,798	11,155
35 to 39 years	10,675	11,029	10,928	11,676	11,069	10,116
40 to 44 years	10,481	10,095	9,325	9,838	9,310	9,016
45 to 49 years	10,504	10,441	9,931	10,419	9,604	8,243
50 to 54 years	10,793	10,623	9,744	9,963	8,924	7,676
55 to 59 years	7,565	7,814	7,561	8,007	7,582	6,533
60 to 64 years	3,971	4,228	4,221	4,640	4,371	3,830
65 years and older	3,198	3,484	3,366	3,272	3,375	2,465
Total	115,166	118,607	108,267	117,554	112,426	98,452

Continued. See notes at end of table.

Table 9.43a. Tennessee – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	465	410	308	275	285	200
Asian ⁴	266	234	4,840	11,382	272	207
Black or African American	21,071	22,260	21,729	31,715	30,960	24,837
Native Hawaiian or Other Pacific Islander	66	69	53	64	107	121
White	78,123	78,987	67,287	65,379	72,468	65,320
Other	882	968	1,059	1,006	939	935
Total	100,873	102,928	95,276	109,821	105,031	91,620
Ethnicity						
Not of Hispanic or Latino origin	68,240	72,365	64,698	74,507	70,975	54,579
Hispanic or Latino origin	2,261	2,560	1,718	1,742	1,752	1,413
Total	70,501	74,925	66,416	76,249	72,727	55,992
Specific Hispanic or Latino origin						
Mexican	50	45	37	24	13	26
Puerto Rican	13	10	10	6	4	- -
Cuban	5	4	3	3	4	1
Other specific	49	70	49	46	31	57
Total	117	129	99	79	52	84
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	1,274	1,322	1,684	1,281
Not a veteran	<i>n/c</i>	<i>n/c</i>	31,680	36,018	45,216	41,901
Total	<i>n/c</i>	<i>n/c</i>	32,954	37,340	46,900	43,182

- - Quantity is zero; n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.43b. Tennessee – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	13.6	15.2	15.5	19.0	21.5	22.1
Anxiety disorders	20.0	20.4	23.4	26.2	27.1	26.2
Attention deficit/hyperactivity disorder	8.0	9.0	5.9	7.7	7.6	6.3
Mood disorders	69.4	59.9	65.4	64.8	62.4	61.2
Bipolar disorders	31.3	21.3	23.8	23.2	20.5	19.9
Depressive disorders	41.7	40.7	44.0	43.9	44.1	43.0
Personality disorders	2.2	1.7	1.8	2.4	1.9	1.9
Schizophrenia and other psychotic disorders	12.7	11.8	14.0	13.9	13.0	13.0
Alcohol or substance-related disorders	†	†	--	--	†	--
Other mental disorders ²	12.3	16.1	11.8	13.3	13.2	11.7
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	46.4	46.6	46.4	47.9	48.1	49.6
Female	53.6	53.4	53.6	52.1	51.9	50.4
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	7.4	7.2	3.1	6.0	6.8	5.7
12 to 14 years	5.1	5.2	4.7	5.2	5.3	4.9
15 to 17 years	6.1	6.4	6.6	5.9	6.0	5.3
18 to 20 years	4.3	4.7	5.1	4.8	4.9	4.9
21 to 24 years	7.3	7.2	7.5	7.2	7.2	7.6
25 to 29 years	10.1	10.4	11.4	11.3	11.1	11.6
30 to 34 years	10.1	10.2	10.6	10.4	10.5	11.3
35 to 39 years	9.3	9.3	10.1	9.9	9.8	10.3
40 to 44 years	9.1	8.5	8.6	8.4	8.3	9.2
45 to 49 years	9.1	8.8	9.2	8.9	8.5	8.4
50 to 54 years	9.4	9.0	9.0	8.5	7.9	7.8
55 to 59 years	6.6	6.6	7.0	6.8	6.7	6.6
60 to 64 years	3.4	3.6	3.9	3.9	3.9	3.9
65 years and older	2.8	2.9	3.1	2.8	3.0	2.5
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.43b. Tennessee – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	†	†	†	†	†	†
Asian ⁴	†	†	5.1	10.4	†	†
Black or African American	20.9	21.6	22.8	28.9	29.5	27.1
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	77.4	76.7	70.6	59.5	69.0	71.3
Other	†	†	1.1	†	†	1.0
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	96.8	96.6	97.4	97.7	97.6	97.5
Hispanic or Latino origin	3.2	3.4	2.6	2.3	2.4	2.5
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	42.7	34.9	37.4	30.4	25.0	31.0
Puerto Rican	11.1	7.8	10.1	7.6	7.7	--
Cuban	4.3	3.1	3.0	3.8	7.7	1.2
Other specific	41.9	54.3	49.5	58.2	59.6	67.9
Total	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	3.9	3.5	3.6	3.0
Not a veteran	<i>n/c</i>	<i>n/c</i>	96.1	96.5	96.4	97.0
Total	<i>n/c</i>	<i>n/c</i>	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals; -- Quantity is zero; n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.44a. Texas – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020

<i>Selected characteristics</i>	2015	2016	2017	2018	2019	2020
Total individuals	348,147	368,722	379,956	400,154	416,338	415,350
Mental health diagnosis¹						
Trauma- and stressor-related disorders	21,906	35,389	40,638	47,668	55,047	59,384
Anxiety disorders	28,877	45,125	50,607	59,721	71,032	80,964
Attention deficit/hyperactivity disorder	22,732	37,306	39,745	41,941	42,985	42,610
Mood disorders	117,377	157,070	162,768	165,857	176,508	180,428
Bipolar disorders	56,074	68,916	69,244	69,148	73,345	73,794
Depressive disorders	62,646	88,936	94,969	98,620	104,730	108,379
Personality disorders	14,680	13,866	13,196	13,179	12,223	11,474
Schizophrenia and other psychotic disorders	32,427	51,058	52,430	52,053	53,980	54,059
Alcohol or substance-related disorders	--	2	--	1	--	--
Other mental disorders ²	64,397	40,567	39,614	43,471	44,040	42,920
Total	175,633	255,712	264,246	268,766	281,114	283,773
Gender						
Male	173,549	185,317	190,881	200,987	208,722	207,692
Female	174,597	183,404	189,074	199,167	207,615	207,658
Total	348,146	368,721	379,955	400,154	416,337	415,350
Age³						
0 to 11 years	32,241	34,937	36,009	38,920	39,661	37,071
12 to 14 years	20,418	21,908	22,381	24,991	27,438	26,300
15 to 17 years	22,596	24,355	25,254	27,087	29,338	29,151
18 to 20 years	15,668	17,040	18,210	20,025	20,988	21,209
21 to 24 years	23,769	24,921	25,199	26,643	27,550	28,335
25 to 29 years	33,045	35,645	37,162	39,410	40,608	41,168
30 to 34 years	33,409	35,505	36,282	38,269	39,684	40,941
35 to 39 years	29,327	32,324	33,718	36,121	37,382	38,294
40 to 44 years	27,608	27,973	28,188	28,955	30,375	31,585
45 to 49 years	28,739	29,597	29,647	29,683	29,538	28,608
50 to 54 years	31,166	31,699	31,106	29,998	29,302	28,493
55 to 59 years	24,706	26,012	27,093	27,546	28,288	27,696
60 to 64 years	13,937	15,106	16,383	17,549	18,929	19,510
65 years and older	11,516	11,699	13,320	14,956	17,254	16,988
Total	348,145	368,721	379,952	400,153	416,335	415,349

Continued. See notes at end of table.

Table 9.44a. Texas – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	958	1,087	1,153	1,278	1,428	1,427
Asian ⁴	2,706	2,816	2,946	3,301	3,505	3,602
Black or African American	81,621	84,822	84,750	86,405	90,194	89,199
Native Hawaiian or Other Pacific Islander	154	194	246	311	355	423
White	250,914	266,236	274,240	291,325	303,574	300,537
Other	11,785	13,554	16,612	17,518	17,272	20,151
Total	348,138	368,709	379,947	400,138	416,328	415,339
Ethnicity						
Not of Hispanic or Latino origin	256,546	272,107	281,244	295,061	302,961	294,709
Hispanic or Latino origin	91,577	96,585	98,693	105,071	113,356	120,624
Total	348,123	368,692	379,937	400,132	416,317	415,333
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡
Not a veteran	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡
Total	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡

- - Quantity is zero; ‡ No data submitted; n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.44b. Texas – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	12.5	13.8	15.4	17.7	19.6	20.9
Anxiety disorders	16.4	17.6	19.2	22.2	25.3	28.5
Attention deficit/hyperactivity disorder	12.9	14.6	15.0	15.6	15.3	15.0
Mood disorders	66.8	61.4	61.6	61.7	62.8	63.6
Bipolar disorders	31.9	27.0	26.2	25.7	26.1	26.0
Depressive disorders	35.7	34.8	35.9	36.7	37.3	38.2
Personality disorders	8.4	5.4	5.0	4.9	4.3	4.0
Schizophrenia and other psychotic disorders	18.5	20.0	19.8	19.4	19.2	19.1
Alcohol or substance-related disorders	--	†	--	†	--	--
Other mental disorders ²	36.7	15.9	15.0	16.2	15.7	15.1
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	49.8	50.3	50.2	50.2	50.1	50.0
Female	50.2	49.7	49.8	49.8	49.9	50.0
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	9.3	9.5	9.5	9.7	9.5	8.9
12 to 14 years	5.9	5.9	5.9	6.2	6.6	6.3
15 to 17 years	6.5	6.6	6.6	6.8	7.0	7.0
18 to 20 years	4.5	4.6	4.8	5.0	5.0	5.1
21 to 24 years	6.8	6.8	6.6	6.7	6.6	6.8
25 to 29 years	9.5	9.7	9.8	9.8	9.8	9.9
30 to 34 years	9.6	9.6	9.5	9.6	9.5	9.9
35 to 39 years	8.4	8.8	8.9	9.0	9.0	9.2
40 to 44 years	7.9	7.6	7.4	7.2	7.3	7.6
45 to 49 years	8.3	8.0	7.8	7.4	7.1	6.9
50 to 54 years	9.0	8.6	8.2	7.5	7.0	6.9
55 to 59 years	7.1	7.1	7.1	6.9	6.8	6.7
60 to 64 years	4.0	4.1	4.3	4.4	4.5	4.7
65 years and older	3.3	3.2	3.5	3.7	4.1	4.1
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.44b. Texas – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	†	†	†	†	†	†
Asian ⁴	†	†	†	†	†	†
Black or African American	23.4	23.0	22.3	21.6	21.7	21.5
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	72.1	72.2	72.2	72.8	72.9	72.4
Other	3.4	3.7	4.4	4.4	4.1	4.9
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	73.7	73.8	74.0	73.7	72.8	71.0
Hispanic or Latino origin	26.3	26.2	26.0	26.3	27.2	29.0
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡
Not a veteran	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡
Total	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡

† Less than 1 percent of individuals; - - Quantity is zero; ‡ No data submitted; *n/c* Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.45a. Utah – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	51,080	53,708	55,525	56,095	56,224	57,456
Mental health diagnosis¹						
Trauma- and stressor-related disorders	10,093	9,543	10,017	10,092	10,340	10,875
Anxiety disorders	14,777	16,998	18,502	19,378	19,899	21,525
Attention deficit/hyperactivity disorder	6,210	5,875	5,977	5,930	5,541	5,139
Mood disorders	22,352	18,853	19,259	19,676	20,067	21,099
Bipolar disorders	7,816	4,559	4,404	4,490	4,477	4,652
Depressive disorders	14,895	14,484	15,027	15,429	15,903	16,790
Personality disorders	4,162	3,096	2,904	2,580	2,367	2,356
Schizophrenia and other psychotic disorders	5,571	5,543	5,623	5,813	5,937	6,068
Alcohol or substance-related disorders	454	4,678	156	23	160	158
Other mental disorders ²	9,837	11,797	11,411	11,162	11,329	11,118
Total	44,446	47,714	48,506	48,330	48,484	50,333
Gender						
Male	24,379	25,645	26,496	27,059	27,082	27,289
Female	26,701	28,063	29,029	29,036	29,142	30,163
Total	51,080	53,708	55,525	56,095	56,224	57,452
Age³						
0 to 11 years	11,705	12,157	12,430	12,032	11,611	10,929
12 to 14 years	4,760	5,179	5,368	5,583	5,405	5,428
15 to 17 years	3,671	4,080	4,322	4,510	4,323	4,415
18 to 20 years	1,697	1,743	1,862	2,059	2,173	2,370
21 to 24 years	2,742	2,928	2,973	2,909	2,964	3,238
25 to 29 years	4,082	4,292	4,487	4,502	4,595	4,966
30 to 34 years	4,366	4,435	4,642	4,546	4,651	4,855
35 to 39 years	3,763	4,008	4,360	4,410	4,523	4,749
40 to 44 years	3,061	3,254	3,206	3,313	3,527	3,859
45 to 49 years	2,672	2,724	2,833	2,948	2,989	3,092
50 to 54 years	2,853	2,843	2,676	2,667	2,632	2,675
55 to 59 years	2,213	2,326	2,443	2,470	2,568	2,606
60 to 64 years	1,385	1,500	1,590	1,709	1,743	1,837
65 years and older	2,107	2,238	2,332	2,435	2,518	2,437
Total	51,077	53,707	55,524	56,093	56,222	57,456

Continued. See notes at end of table.

Table 9.45a. Utah – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	983	1,034	978	1,346	1,094	1,115
Asian ⁴	467	475	394	387	370	373
Black or African American	1,438	1,515	1,281	1,222	1,229	1,175
Native Hawaiian or Other Pacific Islander	4,654	6,056	341	1,026	462	408
White	34,516	35,555	42,176	41,893	42,535	43,952
Other	4,613	4,968	5,860	5,669	5,569	5,811
Total	46,671	49,603	51,030	51,543	51,259	52,834
Ethnicity						
Not of Hispanic or Latino origin	40,189	42,184	43,098	42,970	42,910	44,031
Hispanic or Latino origin	7,040	7,835	8,045	8,078	7,787	7,874
Total	47,229	50,019	51,143	51,048	50,697	51,905
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	759	783	797	836
Not a veteran	<i>n/c</i>	<i>n/c</i>	28,498	28,462	28,633	30,874
Total	<i>n/c</i>	<i>n/c</i>	29,257	29,245	29,430	31,710

‡ No data submitted; n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.45b. Utah – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	22.7	20.0	20.7	20.9	21.3	21.6
Anxiety disorders	33.2	35.6	38.1	40.1	41.0	42.8
Attention deficit/hyperactivity disorder	14.0	12.3	12.3	12.3	11.4	10.2
Mood disorders	50.3	39.5	39.7	40.7	41.4	41.9
Bipolar disorders	17.6	9.6	9.1	9.3	9.2	9.2
Depressive disorders	33.5	30.4	31.0	31.9	32.8	33.4
Personality disorders	9.4	6.5	6.0	5.3	4.9	4.7
Schizophrenia and other psychotic disorders	12.5	11.6	11.6	12.0	12.2	12.1
Alcohol or substance-related disorders	1.0	9.8	†	†	†	†
Other mental disorders ²	22.1	24.7	23.5	23.1	23.4	22.1
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	47.7	47.7	47.7	48.2	48.2	47.5
Female	52.3	52.3	52.3	51.8	51.8	52.5
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	22.9	22.6	22.4	21.5	20.7	19.0
12 to 14 years	9.3	9.6	9.7	10.0	9.6	9.4
15 to 17 years	7.2	7.6	7.8	8.0	7.7	7.7
18 to 20 years	3.3	3.2	3.4	3.7	3.9	4.1
21 to 24 years	5.4	5.5	5.4	5.2	5.3	5.6
25 to 29 years	8.0	8.0	8.1	8.0	8.2	8.6
30 to 34 years	8.5	8.3	8.4	8.1	8.3	8.4
35 to 39 years	7.4	7.5	7.9	7.9	8.0	8.3
40 to 44 years	6.0	6.1	5.8	5.9	6.3	6.7
45 to 49 years	5.2	5.1	5.1	5.3	5.3	5.4
50 to 54 years	5.6	5.3	4.8	4.8	4.7	4.7
55 to 59 years	4.3	4.3	4.4	4.4	4.6	4.5
60 to 64 years	2.7	2.8	2.9	3.0	3.1	3.2
65 years and older	4.1	4.2	4.2	4.3	4.5	4.2
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.45b. Utah – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	2.1	2.1	1.9	2.6	2.1	2.1
Asian ⁴	1.0	†	†	†	†	†
Black or African American	3.1	3.1	2.5	2.4	2.4	2.2
Native Hawaiian or Other Pacific Islander	10.0	12.2	†	2.0	†	†
White	74.0	71.7	82.6	81.3	83.0	83.2
Other	9.9	10.0	11.5	11.0	10.9	11.0
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	85.1	84.3	84.3	84.2	84.6	84.8
Hispanic or Latino origin	14.9	15.7	15.7	15.8	15.4	15.2
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	2.6	2.7	2.7	2.6
Not a veteran	<i>n/c</i>	<i>n/c</i>	97.4	97.3	97.3	97.4
Total	<i>n/c</i>	<i>n/c</i>	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals; ‡ No data submitted; n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.46a. Vermont – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	24,580	31,244	30,924	31,263	31,333	27,451
Mental health diagnosis¹						
Trauma- and stressor-related disorders	6,552	7,888	8,604	9,585	9,826	9,847
Anxiety disorders	3,039	3,703	4,021	4,385	4,721	4,892
Attention deficit/hyperactivity disorder	1,879	2,322	2,384	2,464	2,483	2,593
Mood disorders	5,814	5,741	6,105	6,441	6,637	6,425
Bipolar disorders	1,639	1,265	1,353	1,382	1,411	1,334
Depressive disorders	4,242	4,518	4,804	5,127	5,289	5,154
Personality disorders	871	906	871	858	825	763
Schizophrenia and other psychotic disorders	1,218	1,786	1,823	1,863	1,901	1,847
Alcohol or substance-related disorders	41	2,950	3,210	6,924	6,612	2,919
Other mental disorders ²	4,053	3,934	3,920	6,905	6,804	6,573
Total	16,691	20,118	21,017	29,046	29,240	24,940
Gender						
Male	12,234	16,182	16,112	16,332	16,385	13,848
Female	12,324	14,988	14,772	14,865	14,862	13,281
Total	24,558	31,170	30,884	31,197	31,247	27,129
Age³						
0 to 11 years	5,223	5,323	5,279	5,607	5,503	5,532
12 to 14 years	2,231	2,207	2,232	2,284	2,400	2,503
15 to 17 years	2,278	2,507	2,434	2,658	2,506	2,481
18 to 20 years	1,408	1,781	1,766	1,766	1,798	1,677
21 to 24 years	1,434	2,115	2,006	1,995	1,876	1,615
25 to 29 years	1,589	2,820	2,711	2,621	2,629	1,830
30 to 34 years	1,521	2,655	2,627	2,466	2,540	1,797
35 to 39 years	1,297	1,989	2,092	2,092	2,203	1,575
40 to 44 years	1,199	1,666	1,605	1,574	1,561	1,229
45 to 49 years	1,319	1,715	1,691	1,683	1,611	1,261
50 to 54 years	1,385	1,823	1,764	1,690	1,665	1,294
55 to 59 years	1,237	1,616	1,653	1,624	1,725	1,427
60 to 64 years	940	1,098	1,153	1,212	1,321	1,196
65 years and older	1,494	1,929	1,911	1,991	1,993	2,034
Total	24,555	31,244	30,924	31,263	31,331	27,451

Continued. See notes at end of table.

Table 9.46a. Vermont – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	280	256	242	246	226	250
Asian ⁴	155	160	159	191	179	180
Black or African American	613	586	592	564	635	610
Native Hawaiian or Other Pacific Islander	--	--	--	--	--	--
White	21,671	20,519	20,097	20,106	19,488	18,941
Other	519	503	542	612	570	611
Total	23,238	22,024	21,632	21,719	21,098	20,592
Ethnicity						
Not of Hispanic or Latino origin	21,290	19,792	19,092	19,087	18,160	17,579
Hispanic or Latino origin	705	605	585	642	823	776
Total	21,995	20,397	19,677	19,729	18,983	18,355
Specific Hispanic or Latino origin						
Mexican	106	93	106	123	123	124
Puerto Rican	121	124	123	129	142	158
Cuban	22	25	20	24	22	18
Other specific	182	175	185	180	217	197
Total	431	417	434	456	504	497
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	‡	518	479	446
Not a veteran	<i>n/c</i>	<i>n/c</i>	‡	10,717	10,866	9,579
Total	<i>n/c</i>	<i>n/c</i>	‡	11,235	11,345	10,025

-- Quantity is zero; ‡ No data submitted; n/c Not collected.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.46b. Vermont – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	39.3	39.2	40.9	33.0	33.6	39.5
Anxiety disorders	18.2	18.4	19.1	15.1	16.1	19.6
Attention deficit/hyperactivity disorder	11.3	11.5	11.3	8.5	8.5	10.4
Mood disorders	34.8	28.5	29.0	22.2	22.7	25.8
Bipolar disorders	9.8	6.3	6.4	4.8	4.8	5.3
Depressive disorders	25.4	22.5	22.9	17.7	18.1	20.7
Personality disorders	5.2	4.5	4.1	3.0	2.8	3.1
Schizophrenia and other psychotic disorders	7.3	8.9	8.7	6.4	6.5	7.4
Alcohol or substance-related disorders	†	14.7	15.3	23.8	22.6	11.7
Other mental disorders ²	24.3	19.6	18.7	23.8	23.3	26.4
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	49.8	51.9	52.2	52.4	52.4	51.0
Female	50.2	48.1	47.8	47.6	47.6	49.0
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	21.3	17.0	17.1	17.9	17.6	20.2
12 to 14 years	9.1	7.1	7.2	7.3	7.7	9.1
15 to 17 years	9.3	8.0	7.9	8.5	8.0	9.0
18 to 20 years	5.7	5.7	5.7	5.6	5.7	6.1
21 to 24 years	5.8	6.8	6.5	6.4	6.0	5.9
25 to 29 years	6.5	9.0	8.8	8.4	8.4	6.7
30 to 34 years	6.2	8.5	8.5	7.9	8.1	6.5
35 to 39 years	5.3	6.4	6.8	6.7	7.0	5.7
40 to 44 years	4.9	5.3	5.2	5.0	5.0	4.5
45 to 49 years	5.4	5.5	5.5	5.4	5.1	4.6
50 to 54 years	5.6	5.8	5.7	5.4	5.3	4.7
55 to 59 years	5.0	5.2	5.3	5.2	5.5	5.2
60 to 64 years	3.8	3.5	3.7	3.9	4.2	4.4
65 years and older	6.1	6.2	6.2	6.4	6.4	7.4
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.46b. Vermont – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	1.2	1.2	1.1	1.1	1.1	1.2
Asian ⁴	†	†	†	†	†	†
Black or African American	2.6	2.7	2.7	2.6	3.0	3.0
Native Hawaiian or Other Pacific Islander	--	--	--	--	--	--
White	93.3	93.2	92.9	92.6	92.4	92.0
Other	2.2	2.3	2.5	2.8	2.7	3.0
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	96.8	97.0	97.0	96.7	95.7	95.8
Hispanic or Latino origin	3.2	3.0	3.0	3.3	4.3	4.2
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	24.6	22.3	24.4	27.0	24.4	24.9
Puerto Rican	28.1	29.7	28.3	28.3	28.2	31.8
Cuban	5.1	6.0	4.6	5.3	4.4	3.6
Other specific	42.2	42.0	42.6	39.5	43.1	39.6
Total	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	‡	4.6	4.2	4.4
Not a veteran	<i>n/c</i>	<i>n/c</i>	‡	95.4	95.8	95.6
Total	<i>n/c</i>	<i>n/c</i>	‡	100.0	100.0	100.0

† Less than 1 percent of individuals; -- Quantity is zero; ‡ No data submitted; *n/c* Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.47a. Virginia – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	20.2	19.7	21.2	23.7	25.1	25.9
Anxiety disorders	23.5	24.5	25.4	26.8	27.9	29.2
Attention deficit/hyperactivity disorder	18.7	18.6	18.3	18.5	18.7	18.1
Mood disorders	59.6	51.6	51.8	52.8	53.8	54.3
Bipolar disorders	28.4	20.3	19.6	20.0	20.1	20.2
Depressive disorders	36.0	34.3	35.3	36.4	37.7	38.2
Personality disorders	11.7	9.9	9.3	8.9	8.5	8.2
Schizophrenia and other psychotic disorders	21.5	21.4	21.2	21.1	21.4	21.3
Alcohol or substance-related disorders	--	†	†	†	†	†
Other mental disorders ²	24.6	34.2	31.3	29.6	29.0	28.0
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	50.4	50.3	50.1	50.3	50.3	50.2
Female	49.6	49.7	49.9	49.7	49.7	49.8
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	13.1	13.0	13.0	12.8	12.6	11.6
12 to 14 years	7.4	7.2	7.2	7.4	7.8	7.6
15 to 17 years	8.8	9.1	8.8	8.7	8.7	8.5
18 to 20 years	4.1	4.1	4.3	4.4	4.6	4.6
21 to 24 years	5.3	5.2	5.2	5.2	5.2	5.4
25 to 29 years	7.8	7.8	8.0	8.1	8.1	8.3
30 to 34 years	7.6	7.8	7.9	8.0	8.1	8.5
35 to 39 years	6.8	7.0	7.3	7.5	7.6	7.9
40 to 44 years	6.8	6.5	6.3	6.4	6.3	6.7
45 to 49 years	7.7	7.3	7.1	6.9	6.5	6.3
50 to 54 years	8.4	8.3	7.9	7.5	7.1	6.7
55 to 59 years	6.9	6.9	6.9	7.0	7.0	7.0
60 to 64 years	4.5	4.7	4.8	4.8	5.0	5.1
65 years and older	4.9	5.0	5.1	5.3	5.5	5.7
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.47a. Virginia – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	†	†	†	†	†	†
Asian ⁴	1.1	1.1	1.1	1.2	1.2	1.3
Black or African American	28.7	28.8	28.9	27.9	27.9	28.0
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	61.6	61.2	60.9	61.5	61.0	60.7
Other	8.2	8.5	8.8	9.0	9.6	9.7
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	93.5	93.4	93.1	92.6	92.6	92.5
Hispanic or Latino origin	6.5	6.6	6.9	7.4	7.4	7.5
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	15.0	15.4	15.3	15.3	16.6	15.9
Puerto Rican	15.7	15.1	15.2	15.2	15.8	15.4
Cuban	2.1	2.2	2.0	1.7	1.9	1.9
Other specific	67.1	67.3	67.6	67.8	65.8	66.8
Total	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡	‡
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡	‡
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡	‡

† Less than 1 percent of individuals; - - Quantity is zero; ‡ No data submitted; *n/c* Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.47b. Virginia – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	20.2	19.7	21.2	23.7	25.1	25.9
Anxiety disorders	23.5	24.5	25.4	26.8	27.9	29.2
Attention deficit/hyperactivity disorder	18.7	18.6	18.3	18.5	18.7	18.1
Mood disorders	59.6	51.6	51.8	52.8	53.8	54.3
Bipolar disorders	28.4	20.3	19.6	20.0	20.1	20.2
Depressive disorders	36.0	34.3	35.3	36.4	37.7	38.2
Personality disorders	11.7	9.9	9.3	8.9	8.5	8.2
Schizophrenia and other psychotic disorders	21.5	21.4	21.2	21.1	21.4	21.3
Alcohol or substance-related disorders	--	†	†	†	†	†
Other mental disorders ²	24.6	34.2	31.3	29.6	29.0	28.0
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	50.4	50.3	50.1	50.3	50.3	50.2
Female	49.6	49.7	49.9	49.7	49.7	49.8
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	13.1	13.0	13.0	12.8	12.6	11.6
12 to 14 years	7.4	7.2	7.2	7.4	7.8	7.6
15 to 17 years	8.8	9.1	8.8	8.7	8.7	8.5
18 to 20 years	4.1	4.1	4.3	4.4	4.6	4.6
21 to 24 years	5.3	5.2	5.2	5.2	5.2	5.4
25 to 29 years	7.8	7.8	8.0	8.1	8.1	8.3
30 to 34 years	7.6	7.8	7.9	8.0	8.1	8.5
35 to 39 years	6.8	7.0	7.3	7.5	7.6	7.9
40 to 44 years	6.8	6.5	6.3	6.4	6.3	6.7
45 to 49 years	7.7	7.3	7.1	6.9	6.5	6.3
50 to 54 years	8.4	8.3	7.9	7.5	7.1	6.7
55 to 59 years	6.9	6.9	6.9	7.0	7.0	7.0
60 to 64 years	4.5	4.7	4.8	4.8	5.0	5.1
65 years and older	4.9	5.0	5.1	5.3	5.5	5.7
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.47b. Virginia – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	†	†	†	†	†	†
Asian ⁴	1.1	1.1	1.1	1.2	1.2	1.3
Black or African American	28.7	28.8	28.9	27.9	27.9	28.0
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	61.6	61.2	60.9	61.5	61.0	60.7
Other	8.2	8.5	8.8	9.0	9.6	9.7
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	93.5	93.4	93.1	92.6	92.6	92.5
Hispanic or Latino origin	6.5	6.6	6.9	7.4	7.4	7.5
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	15.0	15.4	15.3	15.3	16.6	15.9
Puerto Rican	15.7	15.1	15.2	15.2	15.8	15.4
Cuban	2.1	2.2	2.0	1.7	1.9	1.9
Other specific	67.1	67.3	67.6	67.8	65.8	66.8
Total	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡
Not a veteran	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡
Total	<i>n/c</i>	<i>n/c</i>	‡	‡	‡	‡

† Less than 1 percent of individuals; - - Quantity is zero; ‡ No data submitted; *n/c* Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.48b. Washington – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	75,828	92,176	107,264	111,458	161,448	238,447
Mental health diagnosis¹						
Trauma- and stressor-related disorders	15,607	20,250	23,690	29,270	43,787	60,843
Anxiety disorders	10,144	16,081	20,739	27,446	49,313	78,654
Attention deficit/hyperactivity disorder	3,365	5,029	6,525	8,754	17,517	26,597
Mood disorders	38,423	44,364	44,927	47,929	68,086	92,468
Bipolar disorders	13,580	13,741	12,709	12,816	16,213	20,155
Depressive disorders	26,375	32,287	33,767	36,672	54,017	75,157
Personality disorders	1,179	1,771	3,131	4,025	3,996	4,118
Schizophrenia and other psychotic disorders	16,790	18,608	18,477	17,865	20,391	22,261
Alcohol or substance-related disorders	5,220	6,417	7,831	11,240	19,954	34,365
Other mental disorders ²	4,873	10,147	12,848	17,476	31,562	52,265
Total	70,821	87,283	95,765	101,767	150,904	214,013
Gender						
Male	30,600	36,862	45,492	49,042	69,802	100,009
Female	36,938	44,623	56,240	60,358	89,705	134,991
Total	67,538	81,485	101,732	109,400	159,507	235,000
Age³						
0 to 11 years	8,085	9,752	13,061	13,285	18,903	26,397
12 to 14 years	4,167	5,316	7,051	8,015	13,349	19,804
15 to 17 years	4,097	5,394	7,061	8,161	13,689	20,641
18 to 20 years	2,748	3,580	4,803	5,392	8,620	14,017
21 to 24 years	3,966	4,577	5,591	6,039	9,044	14,648
25 to 29 years	6,214	7,422	9,409	9,864	14,231	21,655
30 to 34 years	6,479	7,714	9,445	10,078	14,685	23,026
35 to 39 years	5,671	7,029	8,620	9,439	13,393	19,925
40 to 44 years	4,866	5,600	6,784	7,321	10,505	15,766
45 to 49 years	5,424	6,300	7,288	7,350	9,543	12,709
50 to 54 years	5,759	6,331	7,156	7,283	9,613	13,283
55 to 59 years	4,554	5,507	6,681	7,226	9,801	13,020
60 to 64 years	2,785	3,488	4,338	4,977	7,238	10,513
65 years and older	2,771	3,547	4,497	5,015	6,936	9,622
Total	67,586	81,557	101,785	109,445	159,550	235,026

Continued. See notes at end of table.

Table 9.48b. Washington – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	2,720	3,231	3,957	3,546	6,785	11,920
Asian ⁴	1,747	1,966	2,437	2,707	4,144	6,427
Black or African American	6,116	6,906	7,985	8,440	11,946	16,560
Native Hawaiian or Other Pacific Islander	904	1,073	1,394	1,553	2,368	3,764
White	48,946	59,616	74,697	80,384	113,120	163,308
Other	7,014	8,495	10,694	12,137	19,472	28,987
Total	67,447	81,287	101,164	108,767	157,835	230,966
Ethnicity						
Not of Hispanic or Latino origin	56,896	68,192	84,125	89,012	124,700	181,695
Hispanic or Latino origin	10,428	12,705	16,194	18,818	31,056	45,061
Total	67,324	80,897	100,319	107,830	155,756	226,756
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	1,621	1,822	1,380	1,462
Not a veteran	<i>n/c</i>	<i>n/c</i>	36,902	48,954	40,352	41,912
Total	<i>n/c</i>	<i>n/c</i>	38,523	50,776	41,732	43,374

‡ No data submitted; n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.48b. Washington – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	22.0	23.2	24.7	28.8	29.0	28.4
Anxiety disorders	14.3	18.4	21.7	27.0	32.7	36.8
Attention deficit/hyperactivity disorder	4.8	5.8	6.8	8.6	11.6	12.4
Mood disorders	54.3	50.8	46.9	47.1	45.1	43.2
Bipolar disorders	19.2	15.7	13.3	12.6	10.7	9.4
Depressive disorders	37.2	37.0	35.3	36.0	35.8	35.1
Personality disorders	1.7	2.0	3.3	4.0	2.6	1.9
Schizophrenia and other psychotic disorders	23.7	21.3	19.3	17.6	13.5	10.4
Alcohol or substance-related disorders	7.4	7.4	8.2	11.0	13.2	16.1
Other mental disorders ²	6.9	11.6	13.4	17.2	20.9	24.4
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	45.3	45.2	44.7	44.8	43.8	42.6
Female	54.7	54.8	55.3	55.2	56.2	57.4
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	12.0	12.0	12.8	12.1	11.8	11.2
12 to 14 years	6.2	6.5	6.9	7.3	8.4	8.4
15 to 17 years	6.1	6.6	6.9	7.5	8.6	8.8
18 to 20 years	4.1	4.4	4.7	4.9	5.4	6.0
21 to 24 years	5.9	5.6	5.5	5.5	5.7	6.2
25 to 29 years	9.2	9.1	9.2	9.0	8.9	9.2
30 to 34 years	9.6	9.5	9.3	9.2	9.2	9.8
35 to 39 years	8.4	8.6	8.5	8.6	8.4	8.5
40 to 44 years	7.2	6.9	6.7	6.7	6.6	6.7
45 to 49 years	8.0	7.7	7.2	6.7	6.0	5.4
50 to 54 years	8.5	7.8	7.0	6.7	6.0	5.7
55 to 59 years	6.7	6.8	6.6	6.6	6.1	5.5
60 to 64 years	4.1	4.3	4.3	4.5	4.5	4.5
65 years and older	4.1	4.3	4.4	4.6	4.3	4.1
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.48b. Washington – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	4.0	4.0	3.9	3.3	4.3	5.2
Asian ⁴	2.6	2.4	2.4	2.5	2.6	2.8
Black or African American	9.1	8.5	7.9	7.8	7.6	7.2
Native Hawaiian or Other Pacific Islander	1.3	1.3	1.4	1.4	1.5	1.6
White	72.6	73.3	73.8	73.9	71.7	70.7
Other	10.4	10.5	10.6	11.2	12.3	12.6
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	84.5	84.3	83.9	82.5	80.1	80.1
Hispanic or Latino origin	15.5	15.7	16.1	17.5	19.9	19.9
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	4.2	3.6	3.3	3.4
Not a veteran	<i>n/c</i>	<i>n/c</i>	95.8	96.4	96.7	96.6
Total	<i>n/c</i>	<i>n/c</i>	100.0	100.0	100.0	100.0

‡ No data submitted; n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

**Table 9.49a. West Virginia – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Number, 2015–2020**

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	‡	‡	‡	50,605	‡	‡
Mental health diagnosis¹						
Trauma- and stressor-related disorders	‡	‡	‡	6,019	‡	‡
Anxiety disorders	‡	‡	‡	17,562	‡	‡
Attention deficit/hyperactivity disorder	‡	‡	‡	4,899	‡	‡
Mood disorders	‡	‡	‡	23,605	‡	‡
Bipolar disorders	‡	‡	‡	6,938	‡	‡
Depressive disorders	‡	‡	‡	16,898	‡	‡
Personality disorders	‡	‡	‡	869	‡	‡
Schizophrenia and other psychotic disorders	‡	‡	‡	2,492	‡	‡
Alcohol or substance-related disorders	‡	‡	‡	7,877	‡	‡
Other mental disorders ²	‡	‡	‡	28,232	‡	‡
Total	‡	‡	‡	42,984	‡	‡
Gender						
Male	‡	‡	‡	22,982	‡	‡
Female	‡	‡	‡	27,606	‡	‡
Total	‡	‡	‡	50,588	‡	‡
Age³						
0 to 11 years	‡	‡	‡	3,475	‡	‡
12 to 14 years	‡	‡	‡	2,325	‡	‡
15 to 17 years	‡	‡	‡	2,949	‡	‡
18 to 20 years	‡	‡	‡	2,726	‡	‡
21 to 24 years	‡	‡	‡	3,264	‡	‡
25 to 29 years	‡	‡	‡	5,185	‡	‡
30 to 34 years	‡	‡	‡	4,796	‡	‡
35 to 39 years	‡	‡	‡	4,901	‡	‡
40 to 44 years	‡	‡	‡	4,296	‡	‡
45 to 49 years	‡	‡	‡	4,034	‡	‡
50 to 54 years	‡	‡	‡	3,822	‡	‡
55 to 59 years	‡	‡	‡	3,523	‡	‡
60 to 64 years	‡	‡	‡	2,513	‡	‡
65 years and older	‡	‡	‡	2,795	‡	‡
Total	‡	‡	‡	50,604	‡	‡

Continued. See notes at end of table.

**Table 9.49a. West Virginia – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Number, 2015–2020 (continued)**

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	‡	‡	‡	59	‡	‡
Asian ⁴	‡	‡	‡	58	‡	‡
Black or African American	‡	‡	‡	1,967	‡	‡
Native Hawaiian or Other Pacific Islander	‡	‡	‡	22	‡	‡
White	‡	‡	‡	42,520	‡	‡
Other	‡	‡	‡	5,227	‡	‡
Total	‡	‡	‡	49,853	‡	‡
Ethnicity						
Not of Hispanic or Latino origin	‡	‡	‡	‡	‡	‡
Hispanic or Latino origin	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	‡	968	‡	‡
Not a veteran	<i>n/c</i>	<i>n/c</i>	‡	34,678	‡	‡
Total	<i>n/c</i>	<i>n/c</i>	‡	35,646	‡	‡

‡ No data submitted; n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

**Table 9.49b. West Virginia – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Column percent distribution, 2015–2020**

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	‡	‡	‡	100.0	‡	‡
Mental health diagnosis¹						
Trauma- and stressor-related disorders	‡	‡	‡	14.0	‡	‡
Anxiety disorders	‡	‡	‡	40.9	‡	‡
Attention deficit/hyperactivity disorder	‡	‡	‡	11.4	‡	‡
Mood disorders	‡	‡	‡	54.9	‡	‡
Bipolar disorders	‡	‡	‡	16.1	‡	‡
Depressive disorders	‡	‡	‡	39.3	‡	‡
Personality disorders	‡	‡	‡	2.0	‡	‡
Schizophrenia and other psychotic disorders	‡	‡	‡	5.8	‡	‡
Alcohol or substance-related disorders	‡	‡	‡	18.3	‡	‡
Other mental disorders ²	‡	‡	‡	65.7	‡	‡
Total	‡	‡	‡	100.0	‡	‡
Gender						
Male	‡	‡	‡	45.4	‡	‡
Female	‡	‡	‡	54.6	‡	‡
Total	‡	‡	‡	100.0	‡	‡
Age³						
0 to 11 years	‡	‡	‡	6.9	‡	‡
12 to 14 years	‡	‡	‡	4.6	‡	‡
15 to 17 years	‡	‡	‡	5.8	‡	‡
18 to 20 years	‡	‡	‡	5.4	‡	‡
21 to 24 years	‡	‡	‡	6.5	‡	‡
25 to 29 years	‡	‡	‡	10.2	‡	‡
30 to 34 years	‡	‡	‡	9.5	‡	‡
35 to 39 years	‡	‡	‡	9.7	‡	‡
40 to 44 years	‡	‡	‡	8.5	‡	‡
45 to 49 years	‡	‡	‡	8.0	‡	‡
50 to 54 years	‡	‡	‡	7.6	‡	‡
55 to 59 years	‡	‡	‡	7.0	‡	‡
60 to 64 years	‡	‡	‡	5.0	‡	‡
65 years and older	‡	‡	‡	5.5	‡	‡
Total	‡	‡	‡	100.0	‡	‡

Continued. See notes at end of table.

**Table 9.49b. West Virginia – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Column percent distribution, 2015–2020 (continued)**

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	‡	‡	‡	†	‡	‡
Asian ⁴	‡	‡	‡	†	‡	‡
Black or African American	‡	‡	‡	3.9	‡	‡
Native Hawaiian or Other Pacific Islander	‡	‡	‡	†	‡	‡
White	‡	‡	‡	85.3	‡	‡
Other	‡	‡	‡	10.5	‡	‡
Total	‡	‡	‡	100.0	‡	‡
Ethnicity						
Not of Hispanic or Latino origin	‡	‡	‡	‡	‡	‡
Hispanic or Latino origin	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	‡	2.7	‡	‡
Not a veteran	<i>n/c</i>	<i>n/c</i>	‡	97.3	‡	‡
Total	<i>n/c</i>	<i>n/c</i>	‡	100.0	‡	‡

† Less than 1 percent of individuals; ‡ No data submitted; n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.50a. Wisconsin – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	72,929	77,345	65,079	71,193	74,279	75,905
Mental health diagnosis¹						
Trauma- and stressor-related disorders	9,859	9,750	7,444	7,817	7,811	7,857
Anxiety disorders	7,996	9,790	11,154	12,464	13,130	14,667
Attention deficit/hyperactivity disorder	3,706	3,763	3,184	3,621	3,925	3,908
Mood disorders	27,957	26,335	18,121	18,192	18,157	18,616
Bipolar disorders	11,948	10,816	6,322	6,214	6,210	5,978
Depressive disorders	16,616	16,010	12,000	12,198	12,149	12,829
Personality disorders	1,401	3,068	3,983	6,986	8,091	8,679
Schizophrenia and other psychotic disorders	10,921	10,869	8,865	9,430	9,744	9,736
Alcohol or substance-related disorders	1,193	931	370	242	173	162
Other mental disorders ²	14,274	17,151	11,478	12,685	14,100	14,077
Total	65,528	69,410	57,392	63,670	66,535	68,646
Gender						
Male	36,195	38,405	32,093	35,462	37,396	38,739
Female	36,727	38,927	32,974	35,723	36,869	37,143
Total	72,922	77,332	65,067	71,185	74,265	75,882
Age³						
0 to 11 years	3,089	3,427	3,076	3,437	3,861	4,141
12 to 14 years	3,399	3,475	3,154	3,633	3,938	4,380
15 to 17 years	4,769	5,265	4,808	5,405	5,392	5,414
18 to 20 years	3,487	3,863	3,415	3,969	4,248	4,183
21 to 24 years	5,244	5,481	4,572	4,914	5,063	5,078
25 to 29 years	7,031	7,413	6,121	6,747	7,034	7,268
30 to 34 years	7,018	7,313	6,055	6,557	6,842	7,086
35 to 39 years	6,046	6,522	5,343	6,009	6,397	6,634
40 to 44 years	5,892	6,018	4,722	4,954	5,153	5,267
45 to 49 years	6,297	6,360	5,104	5,381	5,324	5,157
50 to 54 years	7,138	7,193	5,664	5,663	5,540	5,451
55 to 59 years	5,549	6,012	5,104	5,551	5,721	5,849
60 to 64 years	3,636	3,979	3,456	3,907	4,151	4,283
65 years and older	4,334	5,024	4,484	5,066	5,615	5,714
Total	72,929	77,345	65,078	71,193	74,279	75,905

Continued. See notes at end of table.

Table 9.50a. Wisconsin – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	1,521	1,505	1,294	1,372	1,419	1,346
Asian ⁴	908	944	786	933	1,091	881
Black or African American	6,977	7,521	5,981	7,156	7,978	8,603
Native Hawaiian or Other Pacific Islander	85	71	67	79	69	88
White	60,581	63,262	51,983	55,006	56,994	57,521
Other	1,039	1,395	1,490	2,402	2,837	2,842
Total	71,111	74,698	61,601	66,948	70,388	71,281
Ethnicity						
Not of Hispanic or Latino origin	63,796	65,821	53,263	59,453	63,626	63,644
Hispanic or Latino origin	2,707	3,046	2,502	3,057	3,330	3,518
Total	66,503	68,867	55,765	62,510	66,956	67,162
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	‡	‡	582	593
Not a veteran	<i>n/c</i>	<i>n/c</i>	‡	‡	18,025	18,234
Total	<i>n/c</i>	<i>n/c</i>	‡	‡	18,607	18,827

‡ No data submitted; n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.50b. Wisconsin – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	15.0	14.0	13.0	12.3	11.7	11.4
Anxiety disorders	12.2	14.1	19.4	19.6	19.7	21.4
Attention deficit/hyperactivity disorder	5.7	5.4	5.5	5.7	5.9	5.7
Mood disorders	42.7	37.9	31.6	28.6	27.3	27.1
Bipolar disorders	18.2	15.6	11.0	9.8	9.3	8.7
Depressive disorders	25.4	23.1	20.9	19.2	18.3	18.7
Personality disorders	2.1	4.4	6.9	11.0	12.2	12.6
Schizophrenia and other psychotic disorders	16.7	15.7	15.4	14.8	14.6	14.2
Alcohol or substance-related disorders	1.8	1.3	†	†	†	†
Other mental disorders ²	21.8	24.7	20.0	19.9	21.2	20.5
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	49.6	49.7	49.3	49.8	50.4	51.1
Female	50.4	50.3	50.7	50.2	49.6	48.9
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	4.2	4.4	4.7	4.8	5.2	5.5
12 to 14 years	4.7	4.5	4.8	5.1	5.3	5.8
15 to 17 years	6.5	6.8	7.4	7.6	7.3	7.1
18 to 20 years	4.8	5.0	5.2	5.6	5.7	5.5
21 to 24 years	7.2	7.1	7.0	6.9	6.8	6.7
25 to 29 years	9.6	9.6	9.4	9.5	9.5	9.6
30 to 34 years	9.6	9.5	9.3	9.2	9.2	9.3
35 to 39 years	8.3	8.4	8.2	8.4	8.6	8.7
40 to 44 years	8.1	7.8	7.3	7.0	6.9	6.9
45 to 49 years	8.6	8.2	7.8	7.6	7.2	6.8
50 to 54 years	9.8	9.3	8.7	8.0	7.5	7.2
55 to 59 years	7.6	7.8	7.8	7.8	7.7	7.7
60 to 64 years	5.0	5.1	5.3	5.5	5.6	5.6
65 years and older	5.9	6.5	6.9	7.1	7.6	7.5
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.50b. Wisconsin – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	2.1	2.0	2.1	2.0	2.0	1.9
Asian ⁴	1.3	1.3	1.3	1.4	1.5	1.2
Black or African American	9.8	10.1	9.7	10.7	11.3	12.1
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	85.2	84.7	84.4	82.2	81.0	80.7
Other	1.5	1.9	2.4	3.6	4.0	4.0
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	95.9	95.6	95.5	95.1	95.0	94.8
Hispanic or Latino origin	4.1	4.4	4.5	4.9	5.0	5.2
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	‡	‡	3.1	3.1
Not a veteran	<i>n/c</i>	<i>n/c</i>	‡	‡	96.9	96.9
Total	<i>n/c</i>	<i>n/c</i>	‡	‡	100.0	100.0

† Less than 1 percent of individuals; ‡ No data submitted; *n/c* Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.51a. Wyoming – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	17,798	17,921	17,365	16,256	16,710	16,169
Mental health diagnosis¹						
Trauma- and stressor-related disorders	4,282	4,495	4,173	3,889	4,024	3,863
Anxiety disorders	3,803	4,055	4,106	4,149	4,764	4,632
Attention deficit/hyperactivity disorder	1,713	1,592	1,497	1,392	1,387	1,323
Mood disorders	8,451	8,027	7,182	7,237	7,551	7,496
Bipolar disorders	2,488	2,174	1,707	1,672	1,719	1,671
Depressive disorders	6,033	5,938	5,513	5,628	5,878	5,861
Personality disorders	1,042	869	748	696	641	628
Schizophrenia and other psychotic disorders	1,047	1,037	1,033	1,034	1,039	1,088
Alcohol or substance-related disorders	--	--	--	--	--	--
Other mental disorders ²	5,613	4,325	3,235	3,078	3,011	2,727
Total	17,280	17,238	16,665	15,771	16,259	15,765
Gender						
Male	7,938	7,940	7,738	7,316	7,479	7,120
Female	9,860	9,981	9,627	8,940	9,231	9,049
Total	17,798	17,921	17,365	16,256	16,710	16,169
Age³						
0 to 11 years	2,365	2,333	2,145	1,912	1,775	1,507
12 to 14 years	1,218	1,198	1,227	1,215	1,210	1,031
15 to 17 years	1,485	1,396	1,430	1,391	1,404	1,341
18 to 20 years	748	826	851	832	898	930
21 to 24 years	1,239	1,246	1,165	1,029	1,198	1,209
25 to 29 years	1,673	1,796	1,733	1,531	1,559	1,554
30 to 34 years	1,631	1,742	1,606	1,445	1,542	1,482
35 to 39 years	1,480	1,518	1,497	1,424	1,496	1,444
40 to 44 years	1,280	1,236	1,158	1,081	1,158	1,203
45 to 49 years	1,097	1,089	1,071	1,041	1,063	1,031
50 to 54 years	1,173	1,112	1,012	875	865	855
55 to 59 years	969	968	927	880	885	871
60 to 64 years	691	668	663	675	687	717
65 years and older	748	793	880	923	970	994
Total	17,797	17,921	17,365	16,254	16,710	16,169

Continued. See notes at end of table.

Table 9.51a. Wyoming – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	380	374	341	327	416	389
Asian ⁴	76	69	78	71	60	65
Black or African American	259	274	276	250	263	264
Native Hawaiian or Other Pacific Islander	37	40	42	28	51	33
White	15,539	15,561	15,039	14,098	14,321	13,735
Other	564	656	663	618	607	786
Total	16,855	16,974	16,439	15,392	15,718	15,272
Ethnicity						
Not of Hispanic or Latino origin	15,541	15,555	14,967	14,068	14,246	13,808
Hispanic or Latino origin	1,714	1,834	1,800	1,602	1,548	1,526
Total	17,255	17,389	16,767	15,670	15,794	15,334
Specific Hispanic or Latino origin						
Mexican	923	949	921	786	802	781
Puerto Rican	54	56	82	62	53	58
Cuban	24	24	25	12	11	11
Other specific	691	770	737	710	655	651
Total	1,692	1,799	1,765	1,570	1,521	1,501
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	250	217	293	286
Not a veteran	<i>n/c</i>	<i>n/c</i>	4,301	3,822	4,899	5,091
Total	<i>n/c</i>	<i>n/c</i>	4,551	4,039	5,192	5,377

- - Quantity is zero; n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 9.51b. Wyoming – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	24.8	26.1	25.0	24.7	24.7	24.5
Anxiety disorders	22.0	23.5	24.6	26.3	29.3	29.4
Attention deficit/hyperactivity disorder	9.9	9.2	9.0	8.8	8.5	8.4
Mood disorders	48.9	46.6	43.1	45.9	46.4	47.5
Bipolar disorders	14.4	12.6	10.2	10.6	10.6	10.6
Depressive disorders	34.9	34.4	33.1	35.7	36.2	37.2
Personality disorders	6.0	5.0	4.5	4.4	3.9	4.0
Schizophrenia and other psychotic disorders	6.1	6.0	6.2	6.6	6.4	6.9
Alcohol or substance-related disorders	--	--	--	--	--	--
Other mental disorders ²	32.5	25.1	19.4	19.5	18.5	17.3
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	44.6	44.3	44.6	45.0	44.8	44.0
Female	55.4	55.7	55.4	55.0	55.2	56.0
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	13.3	13.0	12.4	11.8	10.6	9.3
12 to 14 years	6.8	6.7	7.1	7.5	7.2	6.4
15 to 17 years	8.3	7.8	8.2	8.6	8.4	8.3
18 to 20 years	4.2	4.6	4.9	5.1	5.4	5.8
21 to 24 years	7.0	7.0	6.7	6.3	7.2	7.5
25 to 29 years	9.4	10.0	10.0	9.4	9.3	9.6
30 to 34 years	9.2	9.7	9.2	8.9	9.2	9.2
35 to 39 years	8.3	8.5	8.6	8.8	9.0	8.9
40 to 44 years	7.2	6.9	6.7	6.7	6.9	7.4
45 to 49 years	6.2	6.1	6.2	6.4	6.4	6.4
50 to 54 years	6.6	6.2	5.8	5.4	5.2	5.3
55 to 59 years	5.4	5.4	5.3	5.4	5.3	5.4
60 to 64 years	3.9	3.7	3.8	4.2	4.1	4.4
65 years and older	4.2	4.4	5.1	5.7	5.8	6.1
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.51b. Wyoming – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2015–2020 (continued)

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	2.3	2.2	2.1	2.1	2.6	2.5
Asian ⁴	†	†	†	†	†	†
Black or African American	1.5	1.6	1.7	1.6	1.7	1.7
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	92.2	91.7	91.5	91.6	91.1	89.9
Other	3.3	3.9	4.0	4.0	3.9	5.1
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	90.1	89.5	89.3	89.8	90.2	90.0
Hispanic or Latino origin	9.9	10.5	10.7	10.2	9.8	10.0
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	54.6	52.8	52.2	50.1	52.7	52.0
Puerto Rican	3.2	3.1	4.6	3.9	3.5	3.9
Cuban	1.4	1.3	1.4	†	†	†
Other specific	40.8	42.8	41.8	45.2	43.1	43.4
Total	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	5.5	5.4	5.6	5.3
Not a veteran	<i>n/c</i>	<i>n/c</i>	94.5	94.6	94.4	94.7
Total	<i>n/c</i>	<i>n/c</i>	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals; - - Quantity is zero; n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

**Table 9.52a. Other jurisdictions – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Number, 2015–2020**

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	175	376	389	392	526	594
Mental health diagnosis¹						
Trauma- and stressor-related disorders	5	11	21	8	35	48
Anxiety disorders	2	12	9	10	20	31
Attention deficit/hyperactivity disorder	3	14	7	12	42	28
Mood disorders	17	42	35	24	54	69
Bipolar disorders	7	20	15	5	18	32
Depressive disorders	10	22	20	19	36	37
Personality disorders	1	--	1	--	7	6
Schizophrenia and other psychotic disorders	14	99	87	116	95	141
Alcohol or substance-related disorders	24	48	63	103	123	94
Other mental disorders ²	10	41	23	30	61	45
Total	67	258	234	278	397	434
Gender						
Male	118	243	247	265	346	366
Female	57	133	142	127	180	228
Total	175	376	389	392	526	594
Age³						
0 to 11 years	--	4	13	--	29	28
12 to 14 years	2	8	9	3	8	13
15 to 17 years	6	14	5	7	16	17
18 to 20 years	12	14	19	14	17	32
21 to 24 years	18	31	32	31	42	42
25 to 29 years	26	45	55	50	54	75
30 to 34 years	26	40	47	56	67	68
35 to 39 years	23	31	44	60	63	70
40 to 44 years	13	43	36	37	60	59
45 to 49 years	16	35	26	36	40	44
50 to 54 years	10	42	39	45	40	40
55 to 59 years	13	36	27	22	40	42
60 to 64 years	6	18	22	17	22	28
65 years and older	--	10	13	6	28	36
Total	171	371	387	384	526	594

Continued. See notes at end of table.

**Table 9.52a. Other jurisdictions – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Number, 2015–2020 (continued)**

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	--	--	--	--	--	--
Asian ⁴	18	240	214	202	47	62
Black or African American	--	--	--	1	--	1
Native Hawaiian or Other Pacific Islander	24	73	115	127	397	437
White	8	14	9	2	13	11
Other	47	48	49	57	65	83
Total	97	375	387	389	522	594
Ethnicity						
Not of Hispanic or Latino origin	‡	234	178	182	258	286
Hispanic or Latino origin	‡	--	--	--	--	--
Total	‡	234	178	182	258	286
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	‡	4	4	4
Not a veteran	<i>n/c</i>	<i>n/c</i>	‡	46	41	102
Total	<i>n/c</i>	<i>n/c</i>	‡	50	45	106

-- Quantity is zero; ‡ No data submitted; n/c Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

**Table 9.52b. Other jurisdictions – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Column percent distribution, 2015–2020**

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	7.5	4.3	9.0	2.9	8.8	11.1
Anxiety disorders	3.0	4.7	3.8	3.6	5.0	7.1
Attention deficit/hyperactivity disorder	4.5	5.4	3.0	4.3	10.6	6.5
Mood disorders	25.4	16.3	15.0	8.6	13.6	15.9
Bipolar disorders	10.4	7.8	6.4	1.8	4.5	7.4
Depressive disorders	14.9	8.5	8.5	6.8	9.1	8.5
Personality disorders	1.5	--	†	--	1.8	1.4
Schizophrenia and other psychotic disorders	20.9	38.4	37.2	41.7	23.9	32.5
Alcohol or substance-related disorders	35.8	18.6	26.9	37.1	31.0	21.7
Other mental disorders ²	14.9	15.9	9.8	10.8	15.4	10.4
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	67.4	64.6	63.5	67.6	65.8	61.6
Female	32.6	35.4	36.5	32.4	34.2	38.4
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	--	1.1	3.4	--	5.5	4.7
12 to 14 years	1.2	2.2	2.3	†	1.5	2.2
15 to 17 years	3.5	3.8	1.3	1.8	3.0	2.9
18 to 20 years	7.0	3.8	4.9	3.6	3.2	5.4
21 to 24 years	10.5	8.4	8.3	8.1	8.0	7.1
25 to 29 years	15.2	12.1	14.2	13.0	10.3	12.6
30 to 34 years	15.2	10.8	12.1	14.6	12.7	11.4
35 to 39 years	13.5	8.4	11.4	15.6	12.0	11.8
40 to 44 years	7.6	11.6	9.3	9.6	11.4	9.9
45 to 49 years	9.4	9.4	6.7	9.4	7.6	7.4
50 to 54 years	5.8	11.3	10.1	11.7	7.6	6.7
55 to 59 years	7.6	9.7	7.0	5.7	7.6	7.1
60 to 64 years	3.5	4.9	5.7	4.4	4.2	4.7
65 years and older	--	2.7	3.4	1.6	5.3	6.1
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

**Table 9.52b. Other jurisdictions – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Column percent distribution, 2015–2020 (continued)**

<i>Selected characteristics</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Race						
American Indian or Alaska Native	--	--	--	--	--	--
Asian ⁴	18.6	64.0	55.3	51.9	9.0	10.4
Black or African American	--	--	--	†	--	†
Native Hawaiian or Other Pacific Islander	24.7	19.5	29.7	32.6	76.1	73.6
White	8.2	3.7	2.3	†	2.5	1.9
Other	48.5	12.8	12.7	14.7	12.5	14.0
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	‡	100.0	100.0	100.0	100.0	100.0
Hispanic or Latino origin	‡	--	--	--	--	--
Total	‡	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	‡	8.0	8.9	3.8
Not a veteran	<i>n/c</i>	<i>n/c</i>	‡	92.0	91.1	96.2
Total	<i>n/c</i>	<i>n/c</i>	‡	100.0	100.0	100.0

† Less than 1 percent of individuals; -- Quantity is zero; ‡ No data submitted; *n/c* Not collected.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to multiple diagnoses per client and/or rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Table 10.1a. Alabama – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	97,473	13,116	16,620	15,798	12,058	31,105	2,135	17,851	16,441
Gender									
Male	45,672	5,376	5,765	10,955	3,910	10,048	669	10,649	9,796
Female	51,798	7,740	10,855	4,843	8,148	21,056	1,466	7,201	6,644
Total	97,470	13,116	16,620	15,798	12,058	31,104	2,135	17,850	16,440
Age²									
0 to 11 years	14,461	3,478	1,659	7,856	29	618	3	27	4,560
12 to 14 years	8,765	1,836	1,360	3,731	73	1,775	4	37	2,711
15 to 17 years	8,692	1,643	1,816	2,426	185	2,712	11	83	2,612
18 to 20 years	4,411	595	935	796	415	1,830	76	305	899
21 to 24 years	4,558	485	940	273	749	1,988	147	796	646
25 to 29 years	6,768	698	1,326	200	1,269	2,735	233	1,569	812
30 to 34 years	7,042	750	1,386	149	1,382	2,684	248	1,785	754
35 to 39 years	6,838	678	1,317	120	1,351	2,457	256	2,018	620
40 to 44 years	6,461	677	1,133	65	1,332	2,447	254	1,896	533
45 to 49 years	6,025	636	1,133	62	1,320	2,435	241	1,637	434
50 to 54 years	6,157	561	1,083	39	1,197	2,594	221	1,775	382
55 to 59 years	6,261	459	971	32	1,145	2,591	196	1,993	449
60 to 64 years	5,077	301	746	32	817	1,957	135	1,905	338
65 years and older	5,955	319	815	17	794	2,282	110	2,024	690
Total	97,471	13,116	16,620	15,798	12,058	31,105	2,135	17,850	16,440
Race									
American Indian or Alaska Native	310	59	55	35	53	103	9	55	38
Asian ³	216	25	33	17	18	64	3	58	33
Black or African American	36,590	4,072	4,426	6,378	2,636	10,301	550	10,481	6,595
Native Hawaiian or Other Pacific Islander	51	7	7	5	11	17	1	4	8
White	53,729	7,998	11,079	8,111	8,798	18,886	1,494	6,597	8,030
Other	6,576	955	1,020	1,252	542	1,734	78	655	1,737
Total	97,472	13,116	16,620	15,798	12,058	31,105	2,135	17,850	16,441

Continued. See notes at end of table.

Table 10.1a. Alabama – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	93,108	12,246	15,717	14,984	11,601	29,550	2,066	17,511	15,651
Hispanic or Latino origin	4,365	870	903	814	457	1,555	69	340	790
Total	97,473	13,116	16,620	15,798	12,058	31,105	2,135	17,851	16,441
Specific Hispanic or Latino origin									
Mexican	1,045	229	224	232	48	301	12	57	247
Puerto Rican	296	51	58	52	43	100	6	39	55
Cuban	49	5	12	8	4	18	2	11	4
Other specific	2,956	582	606	521	359	1,134	48	223	479
Total	4,346	867	900	813	454	1,553	68	330	785
Veteran status⁴									
Veteran	883	175	147	20	141	338	18	199	99
Not a veteran	50,982	5,179	9,917	1,369	9,359	21,996	1,507	10,856	5,212
Total	51,865	5,354	10,064	1,389	9,500	22,334	1,525	11,055	5,311

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.1b. Alabama – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020

Selected characteristics		Mental health diagnosis ¹								
		Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender										
Male	46.9	41.0	34.7	69.3	32.4	32.3	31.3	59.7	59.6	
Female	53.1	59.0	65.3	30.7	67.6	67.7	68.7	40.3	40.4	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age ²										
0 to 11 years	14.8	26.5	10.0	49.7	†	2.0	†	†	27.7	
12 to 14 years	9.0	14.0	8.2	23.6	†	5.7	†	†	16.5	
15 to 17 years	8.9	12.5	10.9	15.4	1.5	8.7	†	†	15.9	
18 to 20 years	4.5	4.5	5.6	5.0	3.4	5.9	3.6	1.7	5.5	
21 to 24 years	4.7	3.7	5.7	1.7	6.2	6.4	6.9	4.5	3.9	
25 to 29 years	6.9	5.3	8.0	1.3	10.5	8.8	10.9	8.8	4.9	
30 to 34 years	7.2	5.7	8.3	†	11.5	8.6	11.6	10.0	4.6	
35 to 39 years	7.0	5.2	7.9	†	11.2	7.9	12.0	11.3	3.8	
40 to 44 years	6.6	5.2	6.8	†	11.0	7.9	11.9	10.6	3.2	
45 to 49 years	6.2	4.8	6.8	†	10.9	7.8	11.3	9.2	2.6	
50 to 54 years	6.3	4.3	6.5	†	9.9	8.3	10.4	9.9	2.3	
55 to 59 years	6.4	3.5	5.8	†	9.5	8.3	9.2	11.2	2.7	
60 to 64 years	5.2	2.3	4.5	†	6.8	6.3	6.3	10.7	2.1	
65 years and older	6.1	2.4	4.9	†	6.6	7.3	5.2	11.3	4.2	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race										
American Indian or Alaska Native	†	†	†	†	†	†	†	†	†	
Asian ³	†	†	†	†	†	†	†	†	†	
Black or African American	37.5	31.0	26.6	40.4	21.9	33.1	25.8	58.7	40.1	
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	†	†	†	
White	55.1	61.0	66.7	51.3	73.0	60.7	70.0	37.0	48.8	
Other	6.7	7.3	6.1	7.9	4.5	5.6	3.7	3.7	10.6	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.1b. Alabama – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	95.5	93.4	94.6	94.8	96.2	95.0	96.8	98.1	95.2
Hispanic or Latino origin	4.5	6.6	5.4	5.2	3.8	5.0	3.2	1.9	4.8
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	24.0	26.4	24.9	28.5	10.6	19.4	17.6	17.3	31.5
Puerto Rican	6.8	5.9	6.4	6.4	9.5	6.4	8.8	11.8	7.0
Cuban	1.1	†	1.3	†	†	1.2	2.9	3.3	†
Other specific	68.0	67.1	67.3	64.1	79.1	73.0	70.6	67.6	61.0
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁴									
Veteran	1.7	3.3	1.5	1.4	1.5	1.5	1.2	1.8	1.9
Not a veteran	98.3	96.7	98.5	98.6	98.5	98.5	98.8	98.2	98.1
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.2a. Alaska – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	10,202	3,329	1,521	512	527	2,613	97	648	2,203
Gender									
Male	4,408	1,265	525	343	202	882	35	440	1,228
Female	5,786	2,064	996	169	325	1,726	62	208	972
Total	10,194	3,329	1,521	512	527	2,608	97	648	2,200
Age²									
0 to 11 years	1,416	830	162	182	2	56	--	1	237
12 to 14 years	984	371	123	112	5	237	--	3	253
15 to 17 years	1,170	376	144	89	18	361	2	10	345
18 to 20 years	556	152	101	25	18	178	8	20	130
21 to 24 years	624	158	119	12	33	225	9	39	121
25 to 29 years	981	233	166	25	81	287	8	97	235
30 to 34 years	1,059	301	190	23	72	280	12	93	216
35 to 39 years	817	217	143	11	74	216	9	92	166
40 to 44 years	577	160	82	13	41	154	10	61	128
45 to 49 years	429	125	74	4	47	114	12	38	86
50 to 54 years	457	115	68	4	35	137	9	61	80
55 to 59 years	466	128	70	5	52	130	9	56	102
60 to 64 years	294	69	30	3	24	105	6	38	52
65 years and older	372	94	49	4	25	133	3	39	52
Total	10,202	3,329	1,521	512	527	2,613	97	648	2,203
Race									
American Indian or Alaska Native	2,866	1,040	379	132	53	769	22	168	729
Asian ³	124	35	25	5	11	28	1	15	17
Black or African American	326	82	27	7	28	71	--	64	65
Native Hawaiian or Other Pacific Islander	83	22	14	2	--	24	--	11	17
White	4,286	1,276	671	241	344	1,093	54	257	887
Other	2,324	813	368	114	90	579	17	122	445
Total	10,009	3,268	1,484	501	526	2,564	94	637	2,160

Continued. See notes at end of table.

Table 10.2a. Alaska – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	9,175	2,970	1,370	459	473	2,357	87	563	2,013
Hispanic or Latino origin	626	221	99	37	37	159	4	48	110
Total	9,801	3,191	1,469	496	510	2,516	91	611	2,123
Specific Hispanic or Latino origin									
Mexican	119	40	25	9	7	27	--	7	25
Puerto Rican	60	21	6	3	7	13	--	8	6
Cuban	11	3	1	1	2	3	--	--	1
Other specific	259	90	46	14	9	82	2	17	50
Total	449	154	78	27	25	125	2	32	82
Veteran status⁴									
Veteran	240	87	31	2	15	64	3	24	43
Not a veteran	5,896	1,556	982	110	450	1,735	84	555	1,226
Total	6,136	1,643	1,013	112	465	1,799	87	579	1,269

-- Quantity is zero.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.2b. Alaska – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020

Selected characteristics		Mental health diagnosis ¹								
		Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender										
Male	43.2	38.0	34.5	67.0	38.3	33.8	36.1	67.9	55.8	
Female	56.8	62.0	65.5	33.0	61.7	66.2	63.9	32.1	44.2	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age ²										
0 to 11 years	13.9	24.9	10.7	35.5	†	2.1	--	†	10.8	
12 to 14 years	9.6	11.1	8.1	21.9	†	9.1	--	†	11.5	
15 to 17 years	11.5	11.3	9.5	17.4	3.4	13.8	2.1	1.5	15.7	
18 to 20 years	5.4	4.6	6.6	4.9	3.4	6.8	8.2	3.1	5.9	
21 to 24 years	6.1	4.7	7.8	2.3	6.3	8.6	9.3	6.0	5.5	
25 to 29 years	9.6	7.0	10.9	4.9	15.4	11.0	8.2	15.0	10.7	
30 to 34 years	10.4	9.0	12.5	4.5	13.7	10.7	12.4	14.4	9.8	
35 to 39 years	8.0	6.5	9.4	2.1	14.0	8.3	9.3	14.2	7.5	
40 to 44 years	5.7	4.8	5.4	2.5	7.8	5.9	10.3	9.4	5.8	
45 to 49 years	4.2	3.8	4.9	†	8.9	4.4	12.4	5.9	3.9	
50 to 54 years	4.5	3.5	4.5	†	6.6	5.2	9.3	9.4	3.6	
55 to 59 years	4.6	3.8	4.6	†	9.9	5.0	9.3	8.6	4.6	
60 to 64 years	2.9	2.1	2.0	†	4.6	4.0	6.2	5.9	2.4	
65 years and older	3.6	2.8	3.2	†	4.7	5.1	3.1	6.0	2.4	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race										
American Indian or Alaska Native	28.6	31.8	25.5	26.3	10.1	30.0	23.4	26.4	33.8	
Asian ³	1.2	1.1	1.7	†	2.1	1.1	1.1	2.4	†	
Black or African American	3.3	2.5	1.8	1.4	5.3	2.8	--	10.0	3.0	
Native Hawaiian or Other Pacific Islander	†	†	†	†	--	†	--	1.7	†	
White	42.8	39.0	45.2	48.1	65.4	42.6	57.4	40.3	41.1	
Other	23.2	24.9	24.8	22.8	17.1	22.6	18.1	19.2	20.6	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.2b. Alaska – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020 (continued)

Selected characteristics		Mental health diagnosis ¹							
					Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD					
Ethnicity									
Not of Hispanic or Latino origin	93.6	93.1	93.3	92.5	92.7	93.7	95.6	92.1	94.8
Hispanic or Latino origin	6.4	6.9	6.7	7.5	7.3	6.3	4.4	7.9	5.2
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	26.5	26.0	32.1	33.3	28.0	21.6	--	21.9	30.5
Puerto Rican	13.4	13.6	7.7	11.1	28.0	10.4	--	25.0	7.3
Cuban	2.4	1.9	1.3	3.7	8.0	2.4	--	--	1.2
Other specific	57.7	58.4	59.0	51.9	36.0	65.6	100.0	53.1	61.0
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁴									
Veteran	3.9	5.3	3.1	1.8	3.2	3.6	3.4	4.1	3.4
Not a veteran	96.1	94.7	96.9	98.2	96.8	96.4	96.6	95.9	96.6
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

-- Quantity is zero; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.3a. Arizona – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	350,300	69,162	75,608	32,715	30,140	71,665	3,430	27,491	136,118
Gender									
Male	167,507	30,243	25,961	21,712	11,502	24,520	1,084	16,788	78,205
Female	182,793	38,919	49,647	11,003	18,638	47,145	2,346	10,703	57,913
Total	350,300	69,162	75,608	32,715	30,140	71,665	3,430	27,491	136,118
Age²									
0 to 11 years	54,197	18,908	4,173	13,822	209	1,003	48	206	21,769
12 to 14 years	24,948	7,727	3,482	6,264	416	3,328	31	185	8,037
15 to 17 years	23,803	6,079	4,680	3,813	877	5,106	52	228	8,554
18 to 20 years	15,480	2,562	4,259	1,292	1,097	4,122	181	667	6,073
21 to 24 years	21,050	3,127	5,718	1,023	1,806	5,197	298	1,514	9,191
25 to 29 years	32,662	4,766	8,492	1,557	2,957	7,382	407	2,612	15,094
30 to 34 years	31,424	4,388	8,304	1,369	3,234	6,773	438	2,953	14,215
35 to 39 years	27,814	4,263	7,207	1,061	3,144	5,949	388	3,018	11,963
40 to 44 years	22,063	3,337	5,716	777	2,881	5,129	317	2,680	8,695
45 to 49 years	20,585	3,073	5,192	578	3,058	5,057	298	2,838	7,342
50 to 54 years	20,413	3,019	5,025	413	3,034	5,519	301	2,908	6,927
55 to 59 years	20,633	2,866	5,104	370	3,025	5,987	286	2,969	6,597
60 to 64 years	15,678	2,036	3,944	244	2,219	4,954	191	2,187	4,649
65 years and older	19,550	3,011	4,312	132	2,183	6,159	194	2,526	7,012
Total	350,300	69,162	75,608	32,715	30,140	71,665	3,430	27,491	136,118
Race									
American Indian or Alaska Native	23,067	4,841	3,399	1,583	808	5,332	111	1,622	11,018
Asian ³	3,002	584	773	282	189	677	26	327	899
Black or African American	24,736	5,653	4,461	2,548	1,969	4,659	197	2,817	8,792
Native Hawaiian or Other Pacific Islander	616	131	143	73	37	129	5	43	205
White	180,429	32,644	44,245	17,802	17,405	39,173	2,026	11,845	69,095
Other	7	1	2	--	--	--	--	3	2
Total	231,857	43,854	53,023	22,288	20,408	49,970	2,365	16,657	90,011

Continued. See notes at end of table.

Table 10.3a. Arizona – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	--	--	--	--	--	--	--	--	--
Hispanic or Latino origin	236,798	44,622	54,004	22,569	20,927	51,090	2,413	17,648	91,610
Total	236,798	44,622	54,004	22,569	20,927	51,090	2,413	17,648	91,610
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴									
Veteran	1,903	369	336	41	324	397	20	340	628
Not a veteran	75,216	11,490	16,741	2,428	10,999	17,755	1,134	9,788	28,486
Total	77,119	11,859	17,077	2,469	11,323	18,152	1,154	10,128	29,114

-- Quantity is zero; ‡ No data submitted.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.3b. Arizona – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender									
Male	47.8	43.7	34.3	66.4	38.2	34.2	31.6	61.1	57.5
Female	52.2	56.3	65.7	33.6	61.8	65.8	68.4	38.9	42.5
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age²									
0 to 11 years	15.5	27.3	5.5	42.2	†	1.4	1.4	†	16.0
12 to 14 years	7.1	11.2	4.6	19.1	1.4	4.6	†	†	5.9
15 to 17 years	6.8	8.8	6.2	11.7	2.9	7.1	1.5	†	6.3
18 to 20 years	4.4	3.7	5.6	3.9	3.6	5.8	5.3	2.4	4.5
21 to 24 years	6.0	4.5	7.6	3.1	6.0	7.3	8.7	5.5	6.8
25 to 29 years	9.3	6.9	11.2	4.8	9.8	10.3	11.9	9.5	11.1
30 to 34 years	9.0	6.3	11.0	4.2	10.7	9.5	12.8	10.7	10.4
35 to 39 years	7.9	6.2	9.5	3.2	10.4	8.3	11.3	11.0	8.8
40 to 44 years	6.3	4.8	7.6	2.4	9.6	7.2	9.2	9.7	6.4
45 to 49 years	5.9	4.4	6.9	1.8	10.1	7.1	8.7	10.3	5.4
50 to 54 years	5.8	4.4	6.6	1.3	10.1	7.7	8.8	10.6	5.1
55 to 59 years	5.9	4.1	6.8	1.1	10.0	8.4	8.3	10.8	4.8
60 to 64 years	4.5	2.9	5.2	†	7.4	6.9	5.6	8.0	3.4
65 years and older	5.6	4.4	5.7	†	7.2	8.6	5.7	9.2	5.2
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race									
American Indian or Alaska Native	9.9	11.0	6.4	7.1	4.0	10.7	4.7	9.7	12.2
Asian ³	1.3	1.3	1.5	1.3	†	1.4	1.1	2.0	†
Black or African American	10.7	12.9	8.4	11.4	9.6	9.3	8.3	16.9	9.8
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	†	†	†
White	77.8	74.4	83.4	79.9	85.3	78.4	85.7	71.1	76.8
Other	†	†	†	--	--	--	--	†	†
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.3b. Arizona – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020 (continued)

		Mental health diagnosis¹								
Selected characteristics		Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders¹
						Bipolar disorders	Depressive disorders			
Ethnicity										
Not of Hispanic or Latino origin	--	--	--	--	--	--	--	--	--	--
Hispanic or Latino origin	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin										
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴										
Veteran	2.5	3.1	2.0	1.7	2.9	2.2	1.7	3.4	2.2	
Not a veteran	97.5	96.9	98.0	98.3	97.1	97.8	98.3	96.6	97.8	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	

-- Quantity is zero; ‡ No data submitted; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.4a. Arkansas – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	73,611	12,099	9,586	2,795	6,851	19,271	390	8,543	14,148
Gender									
Male	35,309	5,533	4,028	1,979	2,322	7,164	129	5,256	8,934
Female	38,257	6,560	5,550	813	4,525	12,092	261	3,285	5,205
Total	73,566	12,093	9,578	2,792	6,847	19,256	390	8,541	14,139
Age²									
0 to 11 years	10,257	3,728	1,329	1,410	11	405	--	8	3,367
12 to 14 years	5,844	1,534	978	602	20	961	1	11	1,742
15 to 17 years	6,021	1,301	1,081	317	62	1,660	4	35	1,566
18 to 20 years	4,428	683	790	167	218	1,465	11	117	985
21 to 24 years	4,043	449	582	64	421	1,440	36	406	649
25 to 29 years	5,622	639	767	66	612	1,757	59	746	983
30 to 34 years	6,124	707	798	41	779	1,665	65	963	1,113
35 to 39 years	5,800	672	656	33	790	1,610	53	1,077	920
40 to 44 years	5,368	574	593	23	855	1,487	50	1,030	765
45 to 49 years	4,875	508	541	19	793	1,528	44	864	582
50 to 54 years	4,473	439	443	21	698	1,475	28	883	489
55 to 59 years	4,413	371	416	13	719	1,545	22	892	437
60 to 64 years	3,130	232	290	10	439	1,087	12	764	299
65 years and older	3,211	260	322	9	434	1,186	5	747	251
Total	73,609	12,097	9,586	2,795	6,851	19,271	390	8,543	14,148
Race									
American Indian or Alaska Native	300	49	30	10	41	87	--	31	60
Asian ³	190	28	18	4	19	57	2	35	30
Black or African American	14,099	1,585	1,073	741	863	3,287	76	3,306	3,182
Native Hawaiian or Other Pacific Islander	86	11	17	3	6	21	--	9	20
White	43,851	7,144	6,057	1,536	5,042	11,648	243	4,260	7,941
Other	674	179	96	14	58	143	3	42	150
Total	59,200	8,996	7,291	2,308	6,029	15,243	324	7,683	11,383

Continued. See notes at end of table.

Table 10.4a. Arkansas – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	62,117	9,627	7,878	2,435	6,073	16,170	342	7,664	11,958
Hispanic or Latino origin	3,427	767	544	137	168	912	7	203	711
Total	65,544	10,394	8,422	2,572	6,241	17,082	349	7,867	12,669
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴									
Veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Not a veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡

- - Quantity is zero; ‡ No data submitted.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.4b. Arkansas – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender									
Male	48.0	45.8	42.1	70.9	33.9	37.2	33.1	61.5	63.2
Female	52.0	54.2	57.9	29.1	66.1	62.8	66.9	38.5	36.8
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age²									
0 to 11 years	13.9	30.8	13.9	50.4	†	2.1	--	†	23.8
12 to 14 years	7.9	12.7	10.2	21.5	†	5.0	†	†	12.3
15 to 17 years	8.2	10.8	11.3	11.3	†	8.6	1.0	†	11.1
18 to 20 years	6.0	5.6	8.2	6.0	3.2	7.6	2.8	1.4	7.0
21 to 24 years	5.5	3.7	6.1	2.3	6.1	7.5	9.2	4.8	4.6
25 to 29 years	7.6	5.3	8.0	2.4	8.9	9.1	15.1	8.7	6.9
30 to 34 years	8.3	5.8	8.3	1.5	11.4	8.6	16.7	11.3	7.9
35 to 39 years	7.9	5.6	6.8	1.2	11.5	8.4	13.6	12.6	6.5
40 to 44 years	7.3	4.7	6.2	†	12.5	7.7	12.8	12.1	5.4
45 to 49 years	6.6	4.2	5.6	†	11.6	7.9	11.3	10.1	4.1
50 to 54 years	6.1	3.6	4.6	†	10.2	7.7	7.2	10.3	3.5
55 to 59 years	6.0	3.1	4.3	†	10.5	8.0	5.6	10.4	3.1
60 to 64 years	4.3	1.9	3.0	†	6.4	5.6	3.1	8.9	2.1
65 years and older	4.4	2.1	3.4	†	6.3	6.2	1.3	8.7	1.8
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race									
American Indian or Alaska Native	†	†	†	†	†	†	--	†	†
Asian ³	†	†	†	†	†	†	†	†	†
Black or African American	23.8	17.6	14.7	32.1	14.3	21.6	23.5	43.0	28.0
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	--	†	†
White	74.1	79.4	83.1	66.6	83.6	76.4	75.0	55.4	69.8
Other	1.1	2.0	1.3	†	†	†	†	†	1.3
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.4b. Arkansas – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020 (continued)

Selected characteristics		Mental health diagnosis ¹							
		Trauma- and stressor-related disorders			Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total									
Ethnicity									
Not of Hispanic or Latino origin	94.8	92.6	93.5	94.7	97.3	94.7	98.0	97.4	94.4
Hispanic or Latino origin	5.2	7.4	6.5	5.3	2.7	5.3	2.0	2.6	5.6
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status ⁴									
Veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Not a veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡

- - Quantity is zero; ‡ No data submitted; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.5a. California – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	626,774	108,332	102,964	42,047	53,238	182,518	8,024	117,846	175,338
Gender									
Male	321,303	48,427	42,433	32,051	22,009	70,902	3,258	77,315	106,045
Female	304,724	59,780	60,410	9,977	31,166	111,341	4,755	40,418	69,076
Total	626,027	108,207	102,843	42,028	53,175	182,243	8,013	117,733	175,121
Age²									
0 to 11 years	104,034	32,725	17,715	22,504	342	10,727	5	263	33,549
12 to 14 years	58,100	11,739	10,374	9,339	758	18,879	26	468	18,509
15 to 17 years	63,557	11,719	13,331	5,826	1,726	25,435	93	1,169	21,060
18 to 20 years	30,451	5,175	6,889	1,583	2,193	11,726	285	2,930	9,083
21 to 24 years	32,827	4,873	5,890	564	3,488	10,257	556	7,597	9,598
25 to 29 years	50,516	6,950	8,084	619	6,142	14,175	897	14,306	14,865
30 to 34 years	46,303	6,492	7,287	432	6,076	12,488	999	13,933	13,387
35 to 39 years	42,100	5,975	6,503	308	5,525	11,278	951	13,416	11,647
40 to 44 years	34,354	4,761	5,242	213	4,852	9,854	739	10,913	8,932
45 to 49 years	34,305	4,419	4,842	181	4,992	10,621	760	10,944	8,477
50 to 54 years	37,451	4,555	5,212	175	5,347	12,694	797	11,886	8,674
55 to 59 years	37,986	4,035	4,912	156	5,161	13,717	811	12,345	7,980
60 to 64 years	28,295	2,605	3,488	86	3,608	10,768	528	9,222	5,111
65 years and older	26,066	2,110	3,161	59	3,018	9,863	573	8,419	4,308
Total	626,345	108,133	102,930	42,045	53,228	182,482	8,020	117,811	175,180
Race									
American Indian or Alaska Native	5,088	1,130	930	295	580	1,465	115	961	1,488
Asian ³	22,964	3,228	2,906	737	1,593	7,499	174	7,225	4,239
Black or African American	85,373	14,725	8,138	5,406	6,787	21,995	906	24,011	21,825
Native Hawaiian or Other Pacific Islander	1,967	428	327	98	151	551	15	420	502
White	177,157	28,812	32,813	10,967	24,353	49,908	4,308	37,037	55,727
Other	236,154	43,224	42,118	18,398	13,951	72,743	1,663	32,621	63,626
Total	528,703	91,547	87,232	35,901	47,415	154,161	7,181	102,275	147,407

Continued. See notes at end of table.

Table 10.5a. California – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	267,652	44,174	42,071	16,316	31,950	76,300	4,851	61,805	74,148
Hispanic or Latino origin	242,584	45,886	44,107	19,814	13,471	74,605	1,636	32,222	66,497
Total	510,236	90,060	86,178	36,130	45,421	150,905	6,487	94,027	140,645
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴									
Veteran	426	13	5	- -	32	25	28	214	300
Not a veteran	5,034	183	111	12	400	363	547	3,243	3,438
Total	5,460	196	116	12	432	388	575	3,457	3,738

- - Quantity is zero; ‡ No data submitted.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.5b. California – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020

		Mental health diagnosis ¹								
Selected characteristics						Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Bipolar disorders	Depressive disorders				
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender										
Male	51.3	44.8	41.3	76.3	41.4	38.9	40.7	65.7	60.6	
Female	48.7	55.2	58.7	23.7	58.6	61.1	59.3	34.3	39.4	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age ²										
0 to 11 years	16.6	30.3	17.2	53.5	†	5.9	†	†	19.2	
12 to 14 years	9.3	10.9	10.1	22.2	1.4	10.3	†	†	10.6	
15 to 17 years	10.1	10.8	13.0	13.9	3.2	13.9	1.2	†	12.0	
18 to 20 years	4.9	4.8	6.7	3.8	4.1	6.4	3.6	2.5	5.2	
21 to 24 years	5.2	4.5	5.7	1.3	6.6	5.6	6.9	6.4	5.5	
25 to 29 years	8.1	6.4	7.9	1.5	11.5	7.8	11.2	12.1	8.5	
30 to 34 years	7.4	6.0	7.1	1.0	11.4	6.8	12.5	11.8	7.6	
35 to 39 years	6.7	5.5	6.3	†	10.4	6.2	11.9	11.4	6.6	
40 to 44 years	5.5	4.4	5.1	†	9.1	5.4	9.2	9.3	5.1	
45 to 49 years	5.5	4.1	4.7	†	9.4	5.8	9.5	9.3	4.8	
50 to 54 years	6.0	4.2	5.1	†	10.0	7.0	9.9	10.1	5.0	
55 to 59 years	6.1	3.7	4.8	†	9.7	7.5	10.1	10.5	4.6	
60 to 64 years	4.5	2.4	3.4	†	6.8	5.9	6.6	7.8	2.9	
65 years and older	4.2	2.0	3.1	†	5.7	5.4	7.1	7.1	2.5	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race										
American Indian or Alaska Native	†	1.2	1.1	†	1.2	†	1.6	†	1.0	
Asian ³	4.3	3.5	3.3	2.1	3.4	4.9	2.4	7.1	2.9	
Black or African American	16.1	16.1	9.3	15.1	14.3	14.3	12.6	23.5	14.8	
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	†	†	†	
White	33.5	31.5	37.6	30.5	51.4	32.4	60.0	36.2	37.8	
Other	44.7	47.2	48.3	51.2	29.4	47.2	23.2	31.9	43.2	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.5b. California – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020 (continued)

Selected characteristics		Mental health diagnosis ¹								
		Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Ethnicity										
Not of Hispanic or Latino origin	52.5		49.0	48.8	45.2	70.3	50.6	74.8	65.7	52.7
Hispanic or Latino origin	47.5		51.0	51.2	54.8	29.7	49.4	25.2	34.3	47.3
Total	100.0		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin										
Mexican	‡		‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡		‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡		‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡		‡	‡	‡	‡	‡	‡	‡	‡
Total	‡		‡	‡	‡	‡	‡	‡	‡	‡
Veteran status ⁴										
Veteran	7.8		6.6	4.3	--	7.4	6.4	4.9	6.2	8.0
Not a veteran	92.2		93.4	95.7	100.0	92.6	93.6	95.1	93.8	92.0
Total	100.0		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

- - Quantity is zero; ‡ No data submitted; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.6a. Colorado – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	126,315	40,976	26,738	9,644	14,828	39,422	2,604	12,549	25,857
Gender									
Male	58,296	16,405	10,585	6,416	5,882	15,089	885	8,314	15,274
Female	68,019	24,571	16,153	3,228	8,946	24,333	1,719	4,235	10,583
Total	126,315	40,976	26,738	9,644	14,828	39,422	2,604	12,549	25,857
Age²									
0 to 11 years	13,729	7,671	1,834	2,862	25	936	2	14	2,458
12 to 14 years	10,135	3,995	1,950	1,755	88	3,022	10	41	1,837
15 to 17 years	11,748	3,799	2,990	1,365	334	4,748	31	177	2,355
18 to 20 years	6,674	1,968	1,788	565	560	2,687	119	377	1,402
21 to 24 years	7,407	2,085	1,870	353	1,024	2,670	252	822	1,663
25 to 29 years	12,047	3,436	2,821	584	1,766	3,866	349	1,500	2,868
30 to 34 years	12,057	3,593	2,728	614	1,915	3,650	332	1,504	2,873
35 to 39 years	10,962	3,324	2,453	443	1,725	3,286	315	1,506	2,652
40 to 44 years	8,734	2,679	1,874	332	1,531	2,680	241	1,274	1,894
45 to 49 years	7,563	2,222	1,544	240	1,383	2,478	251	1,102	1,573
50 to 54 years	7,285	2,111	1,409	193	1,383	2,438	253	1,156	1,436
55 to 59 years	7,012	1,793	1,402	144	1,291	2,594	188	1,186	1,291
60 to 64 years	5,178	1,162	965	108	911	2,004	139	926	825
65 years and older	5,784	1,138	1,110	86	892	2,363	122	964	730
Total	126,315	40,976	26,738	9,644	14,828	39,422	2,604	12,549	25,857
Race									
American Indian or Alaska Native	2,907	1,059	551	154	300	890	49	299	657
Asian ³	1,335	380	295	64	156	429	25	220	199
Black or African American	7,585	2,436	1,029	472	914	2,065	143	1,611	1,608
Native Hawaiian or Other Pacific Islander	434	147	89	25	42	151	9	52	97
White	91,134	28,478	19,949	7,120	11,206	28,743	1,980	8,569	19,002
Other	22,905	8,474	4,825	1,807	2,209	7,143	398	1,791	4,288
Total	126,300	40,974	26,738	9,642	14,827	39,421	2,604	12,542	25,851

Continued. See notes at end of table.

Table 10.6a. Colorado – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	81,433	24,777	17,213	6,139	11,203	25,536	1,954	9,115	17,088
Hispanic or Latino origin	32,565	11,604	6,745	2,430	2,284	10,064	410	2,422	6,208
Total	113,998	36,381	23,958	8,569	13,487	35,600	2,364	11,537	23,296
Specific Hispanic or Latino origin									
Mexican	20,680	7,404	4,281	1,611	1,274	6,555	236	1,384	3,841
Puerto Rican	816	312	176	90	114	228	21	69	161
Cuban	373	128	94	30	39	106	8	36	85
Other specific	10,574	3,750	2,182	692	845	3,161	127	860	2,082
Total	32,443	11,594	6,733	2,423	2,272	10,050	392	2,349	6,169
Veteran status⁴									
Veteran	2,666	993	512	98	347	858	67	322	550
Not a veteran	87,444	24,492	19,401	3,548	13,935	29,812	2,426	11,572	18,541
Total	90,110	25,485	19,913	3,646	14,282	30,670	2,493	11,894	19,091

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.6b. Colorado – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender									
Male	46.2	40.0	39.6	66.5	39.7	38.3	34.0	66.3	59.1
Female	53.8	60.0	60.4	33.5	60.3	61.7	66.0	33.7	40.9
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age²									
0 to 11 years	10.9	18.7	6.9	29.7	†	2.4	†	†	9.5
12 to 14 years	8.0	9.7	7.3	18.2	†	7.7	†	†	7.1
15 to 17 years	9.3	9.3	11.2	14.2	2.3	12.0	1.2	1.4	9.1
18 to 20 years	5.3	4.8	6.7	5.9	3.8	6.8	4.6	3.0	5.4
21 to 24 years	5.9	5.1	7.0	3.7	6.9	6.8	9.7	6.6	6.4
25 to 29 years	9.5	8.4	10.6	6.1	11.9	9.8	13.4	12.0	11.1
30 to 34 years	9.5	8.8	10.2	6.4	12.9	9.3	12.7	12.0	11.1
35 to 39 years	8.7	8.1	9.2	4.6	11.6	8.3	12.1	12.0	10.3
40 to 44 years	6.9	6.5	7.0	3.4	10.3	6.8	9.3	10.2	7.3
45 to 49 years	6.0	5.4	5.8	2.5	9.3	6.3	9.6	8.8	6.1
50 to 54 years	5.8	5.2	5.3	2.0	9.3	6.2	9.7	9.2	5.6
55 to 59 years	5.6	4.4	5.2	1.5	8.7	6.6	7.2	9.5	5.0
60 to 64 years	4.1	2.8	3.6	1.1	6.1	5.1	5.3	7.4	3.2
65 years and older	4.6	2.8	4.2	†	6.0	6.0	4.7	7.7	2.8
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race									
American Indian or Alaska Native	2.3	2.6	2.1	1.6	2.0	2.3	1.9	2.4	2.5
Asian ³	1.1	†	1.1	†	1.1	1.1	†	1.8	†
Black or African American	6.0	5.9	3.8	4.9	6.2	5.2	5.5	12.8	6.2
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	†	†	†
White	72.2	69.5	74.6	73.8	75.6	72.9	76.0	68.3	73.5
Other	18.1	20.7	18.0	18.7	14.9	18.1	15.3	14.3	16.6
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.6b. Colorado – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	71.4	68.1	71.8	71.6	83.1	71.7	82.7	79.0	73.4
Hispanic or Latino origin	28.6	31.9	28.2	28.4	16.9	28.3	17.3	21.0	26.6
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	63.7	63.9	63.6	66.5	56.1	65.2	60.2	58.9	62.3
Puerto Rican	2.5	2.7	2.6	3.7	5.0	2.3	5.4	2.9	2.6
Cuban	1.1	1.1	1.4	1.2	1.7	1.1	2.0	1.5	1.4
Other specific	32.6	32.3	32.4	28.6	37.2	31.5	32.4	36.6	33.7
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁴									
Veteran	3.0	3.9	2.6	2.7	2.4	2.8	2.7	2.7	2.9
Not a veteran	97.0	96.1	97.4	97.3	97.6	97.2	97.3	97.3	97.1
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.7a. Connecticut – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	41,625	9,682	10,148	2,130	7,356	14,395	2,057	9,827	6,468
Gender									
Male	20,267	3,805	4,157	1,219	2,999	5,810	696	6,380	3,523
Female	21,311	5,871	5,984	910	4,347	8,571	1,355	3,443	2,929
Total	41,578	9,676	10,141	2,129	7,346	14,381	2,051	9,823	6,452
Age²									
0 to 11 years	293	110	20	99	6	26	- -	14	149
12 to 14 years	315	88	32	99	13	49	1	12	193
15 to 17 years	536	147	70	121	24	126	5	28	312
18 to 20 years	1,760	444	460	231	213	645	55	214	473
21 to 24 years	2,819	726	803	257	447	986	166	473	604
25 to 29 years	3,740	1,046	960	305	676	1,164	223	809	677
30 to 34 years	3,880	1,100	1,078	253	671	1,222	195	892	567
35 to 39 years	3,757	1,093	977	222	738	1,139	167	880	448
40 to 44 years	3,356	950	835	133	658	1,116	173	797	398
45 to 49 years	3,468	902	908	98	747	1,161	202	823	432
50 to 54 years	4,505	1,056	1,076	91	878	1,704	222	1,119	558
55 to 59 years	4,897	964	1,117	108	911	1,824	257	1,317	596
60 to 64 years	4,023	583	898	66	708	1,539	204	1,193	504
65 years and older	4,276	473	914	47	666	1,694	187	1,256	557
Total	41,625	9,682	10,148	2,130	7,356	14,395	2,057	9,827	6,468
Race									
American Indian or Alaska Native	263	96	62	11	46	93	15	53	38
Asian ³	436	85	75	14	57	110	12	169	57
Black or African American	6,989	1,572	1,009	221	935	1,847	229	2,660	963
Native Hawaiian or Other Pacific Islander	139	35	22	5	19	62	3	32	19
White	26,765	6,019	7,367	1,580	5,259	9,658	1,528	5,429	4,188
Other	5,665	1,588	1,346	237	882	2,121	221	1,263	934
Total	40,257	9,395	9,881	2,068	7,198	13,891	2,008	9,606	6,199

Continued. See notes at end of table.

Table 10.7a. Connecticut – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	31,424	6,849	7,726	1,686	5,808	10,460	1,662	7,843	4,786
Hispanic or Latino origin	8,595	2,442	2,068	372	1,318	3,361	330	1,734	1,363
Total	40,019	9,291	9,794	2,058	7,126	13,821	1,992	9,577	6,149
Specific Hispanic or Latino origin									
Mexican	215	74	50	6	15	91	4	37	51
Puerto Rican	4,757	1,446	1,264	170	753	1,975	185	976	690
Cuban	60	15	18	1	8	18	7	16	8
Other specific	3,391	860	724	170	528	1,257	125	674	503
Total	8,423	2,395	2,056	347	1,304	3,341	321	1,703	1,252
Veteran status⁴									
Veteran	1,070	230	172	34	138	331	52	385	136
Not a veteran	37,078	8,698	9,508	1,706	6,841	13,087	1,901	8,891	5,213
Total	38,148	8,928	9,680	1,740	6,979	13,418	1,953	9,276	5,349

- - Quantity is zero.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.7b. Connecticut – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020

Selected characteristics		Mental health diagnosis ¹								
		Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender										
Male	48.7	39.3	41.0	57.3	40.8	40.4	33.9	64.9	54.6	
Female	51.3	60.7	59.0	42.7	59.2	59.6	66.1	35.1	45.4	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age ²										
0 to 11 years	†	1.1	†	4.6	†	†	--	†	2.3	
12 to 14 years	†	†	†	4.6	†	†	†	†	3.0	
15 to 17 years	1.3	1.5	†	5.7	†	†	†	†	4.8	
18 to 20 years	4.2	4.6	4.5	10.8	2.9	4.5	2.7	2.2	7.3	
21 to 24 years	6.8	7.5	7.9	12.1	6.1	6.8	8.1	4.8	9.3	
25 to 29 years	9.0	10.8	9.5	14.3	9.2	8.1	10.8	8.2	10.5	
30 to 34 years	9.3	11.4	10.6	11.9	9.1	8.5	9.5	9.1	8.8	
35 to 39 years	9.0	11.3	9.6	10.4	10.0	7.9	8.1	9.0	6.9	
40 to 44 years	8.1	9.8	8.2	6.2	8.9	7.8	8.4	8.1	6.2	
45 to 49 years	8.3	9.3	8.9	4.6	10.2	8.1	9.8	8.4	6.7	
50 to 54 years	10.8	10.9	10.6	4.3	11.9	11.8	10.8	11.4	8.6	
55 to 59 years	11.8	10.0	11.0	5.1	12.4	12.7	12.5	13.4	9.2	
60 to 64 years	9.7	6.0	8.8	3.1	9.6	10.7	9.9	12.1	7.8	
65 years and older	10.3	4.9	9.0	2.2	9.1	11.8	9.1	12.8	8.6	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race										
American Indian or Alaska Native	†	1.0	†	†	†	†	†	†	†	
Asian ³	1.1	†	†	†	†	†	†	1.8	†	
Black or African American	17.4	16.7	10.2	10.7	13.0	13.3	11.4	27.7	15.5	
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	†	†	†	
White	66.5	64.1	74.6	76.4	73.1	69.5	76.1	56.5	67.6	
Other	14.1	16.9	13.6	11.5	12.3	15.3	11.0	13.1	15.1	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.7b. Connecticut – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	78.5	73.7	78.9	81.9	81.5	75.7	83.4	81.9	77.8
Hispanic or Latino origin	21.5	26.3	21.1	18.1	18.5	24.3	16.6	18.1	22.2
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	2.6	3.1	2.4	1.7	1.2	2.7	1.2	2.2	4.1
Puerto Rican	56.5	60.4	61.5	49.0	57.7	59.1	57.6	57.3	55.1
Cuban	†	†	†	†	†	†	2.2	†	†
Other specific	40.3	35.9	35.2	49.0	40.5	37.6	38.9	39.6	40.2
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁴									
Veteran	2.8	2.6	1.8	2.0	2.0	2.5	2.7	4.2	2.5
Not a veteran	97.2	97.4	98.2	98.0	98.0	97.5	97.3	95.8	97.5
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

- - Quantity is zero; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.8a. Delaware – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	10,559	1,388	1,356	636	2,007	3,383	368	2,083	5,517
Gender									
Male	5,713	610	576	416	943	1,663	154	1,283	3,269
Female	4,734	776	777	220	1,061	1,714	214	795	2,147
Total	10,447	1,386	1,353	636	2,004	3,377	368	2,078	5,416
Age²									
0 to 11 years	208	51	21	101	10	19	--	1	120
12 to 14 years	230	57	28	96	26	45	--	3	140
15 to 17 years	290	64	32	118	50	88	1	7	153
18 to 20 years	611	116	102	89	99	270	12	51	242
21 to 24 years	872	129	134	70	169	353	26	123	383
25 to 29 years	1,310	157	153	49	231	400	36	235	751
30 to 34 years	1,207	144	140	36	223	343	24	219	721
35 to 39 years	1,093	136	136	18	218	305	24	212	614
40 to 44 years	922	125	127	21	201	268	31	197	542
45 to 49 years	836	115	96	11	183	276	34	163	455
50 to 54 years	835	102	131	12	188	276	33	199	430
55 to 59 years	860	101	108	11	189	300	42	236	421
60 to 64 years	637	50	73	2	117	216	48	205	286
65 years and older	647	41	75	2	102	224	57	232	259
Total	10,558	1,388	1,356	636	2,006	3,383	368	2,083	5,517
Race									
American Indian or Alaska Native	69	11	8	2	15	15	1	25	40
Asian ³	80	7	17	5	14	29	2	24	27
Black or African American	3,212	473	305	250	586	917	107	999	1,620
Native Hawaiian or Other Pacific Islander	53	6	6	1	12	20	3	15	24
White	6,111	782	871	334	1,248	1,927	237	901	3,443
Other	177	20	27	15	24	56	5	40	71
Total	9,702	1,299	1,234	607	1,899	2,964	355	2,004	5,225

Continued. See notes at end of table.

Table 10.8a. Delaware – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020 (continued)

Mental health diagnosis ¹									
Selected characteristics	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	8,373	1,113	1,016	538	1,666	2,435	320	1,723	4,674
Hispanic or Latino origin	535	97	79	46	89	187	21	106	220
Total	8,908	1,210	1,095	584	1,755	2,622	341	1,829	4,894
Specific Hispanic or Latino origin									
Mexican	56	5	12	1	4	20	--	11	24
Puerto Rican	145	19	21	--	28	53	7	39	56
Cuban	10	2	--	--	--	7	--	2	2
Other specific	179	32	28	5	31	71	10	42	75
Total	390	58	61	6	63	151	17	94	157
Veteran status⁴									
Veteran	347	66	39	7	45	164	8	70	148
Not a veteran	7,277	944	1,032	162	1,505	2,512	330	1,583	3,664
Total	7,624	1,010	1,071	169	1,550	2,676	338	1,653	3,812

-- Quantity is zero.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.8b. Delaware – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020

Selected characteristics		Mental health diagnosis ¹								
		Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender										
Male	54.7	44.0	42.6	65.4	47.1	49.2	41.8	61.7	60.4	
Female	45.3	56.0	57.4	34.6	52.9	50.8	58.2	38.3	39.6	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age ²										
0 to 11 years	2.0	3.7	1.5	15.9	†	†	--	†	2.2	
12 to 14 years	2.2	4.1	2.1	15.1	1.3	1.3	--	†	2.5	
15 to 17 years	2.7	4.6	2.4	18.6	2.5	2.6	†	†	2.8	
18 to 20 years	5.8	8.4	7.5	14.0	4.9	8.0	3.3	2.4	4.4	
21 to 24 years	8.3	9.3	9.9	11.0	8.4	10.4	7.1	5.9	6.9	
25 to 29 years	12.4	11.3	11.3	7.7	11.5	11.8	9.8	11.3	13.6	
30 to 34 years	11.4	10.4	10.3	5.7	11.1	10.1	6.5	10.5	13.1	
35 to 39 years	10.4	9.8	10.0	2.8	10.9	9.0	6.5	10.2	11.1	
40 to 44 years	8.7	9.0	9.4	3.3	10.0	7.9	8.4	9.5	9.8	
45 to 49 years	7.9	8.3	7.1	1.7	9.1	8.2	9.2	7.8	8.2	
50 to 54 years	7.9	7.3	9.7	1.9	9.4	8.2	9.0	9.6	7.8	
55 to 59 years	8.1	7.3	8.0	1.7	9.4	8.9	11.4	11.3	7.6	
60 to 64 years	6.0	3.6	5.4	†	5.8	6.4	13.0	9.8	5.2	
65 years and older	6.1	3.0	5.5	†	5.1	6.6	15.5	11.1	4.7	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race										
American Indian or Alaska Native	†	†	†	†	†	†	†	1.2	†	
Asian ³	†	†	1.4	†	†	†	†	1.2	†	
Black or African American	33.1	36.4	24.7	41.2	30.9	30.9	30.1	49.9	31.0	
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	†	†	†	
White	63.0	60.2	70.6	55.0	65.7	65.0	66.8	45.0	65.9	
Other	1.8	1.5	2.2	2.5	1.3	1.9	1.4	2.0	1.4	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.8b. Delaware – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020 (continued)

Mental health diagnosis ¹													
Selected characteristics		Trauma- and stressor-related disorders				Mood disorders		Personality disorders		Schizophrenia, other psychotic disorders		Other mental disorders ¹	
						Bipolar disorders	Depressive disorders						
Total		Total		Total		Total		Total		Total		Total	
Ethnicity		Total		Total		Total		Total		Total		Total	
Not of Hispanic or Latino origin		94.0	92.0	92.8	92.1	94.9	92.9	93.8	94.2	95.5			
Hispanic or Latino origin		6.0	8.0	7.2	7.9	5.1	7.1	6.2	5.8	4.5			
Total		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0			
Specific Hispanic or Latino origin		Total		Total		Total		Total		Total		Total	
Mexican		14.4	8.6	19.7	16.7	6.3	13.2	--	11.7	15.3			
Puerto Rican		37.2	32.8	34.4	--	44.4	35.1	41.2	41.5	35.7			
Cuban		2.6	3.4	--	--	--	4.6	--	2.1	1.3			
Other specific		45.9	55.2	45.9	83.3	49.2	47.0	58.8	44.7	47.8			
Total		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0			
Veteran status ⁴		Total		Total		Total		Total		Total		Total	
Veteran		4.6	6.5	3.6	4.1	2.9	6.1	2.4	4.2	3.9			
Not a veteran		95.4	93.5	96.4	95.9	97.1	93.9	97.6	95.8	96.1			
Total		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0			

-- Quantity is zero; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.9a. District of Columbia – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	35,161	3,871	1,525	2,157	5,839	13,701	73	6,372	3,893
Gender									
Male	18,399	1,721	618	1,438	2,621	6,745	30	3,802	2,365
Female	16,709	2,148	904	718	3,212	6,938	43	2,557	1,513
Total	35,108	3,869	1,522	2,156	5,833	13,683	73	6,359	3,878
Age²									
0 to 11 years	635	173	42	230	4	61	--	1	134
12 to 14 years	1,133	246	48	396	20	258	--	7	196
15 to 17 years	1,221	261	62	348	50	350	--	9	190
18 to 20 years	1,574	245	64	350	128	548	1	25	274
21 to 24 years	1,917	239	65	329	301	717	--	90	286
25 to 29 years	2,811	298	123	161	562	1,177	3	333	299
30 to 34 years	3,141	330	168	78	696	1,246	8	508	307
35 to 39 years	2,961	312	147	54	624	1,148	13	534	332
40 to 44 years	2,657	305	125	39	522	1,106	6	525	235
45 to 49 years	2,369	248	119	33	446	1,005	2	475	218
50 to 54 years	3,472	334	142	41	737	1,473	11	688	330
55 to 59 years	4,322	401	162	43	803	1,867	9	978	396
60 to 64 years	3,557	260	127	31	546	1,534	8	921	360
65 years and older	3,391	219	131	24	400	1,211	12	1,278	336
Total	35,161	3,871	1,525	2,157	5,839	13,701	73	6,372	3,893
Race									
American Indian or Alaska Native	55	4	3	1	8	19	--	18	6
Asian ³	105	8	9	1	16	30	1	37	11
Black or African American	31,943	3,564	1,227	2,037	5,318	12,713	62	5,476	3,520
Native Hawaiian or Other Pacific Islander	1	--	--	--	--	--	--	1	--
White	1,148	89	87	48	252	331	7	365	106
Other	61	8	11	2	12	18	--	10	11
Total	33,313	3,673	1,337	2,089	5,606	13,111	70	5,907	3,654

Continued. See notes at end of table.

Table 10.9a. District of Columbia – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	13,078	1,398	611	572	2,353	4,738	32	2,501	1,694
Hispanic or Latino origin	679	71	58	23	110	239	1	156	84
Total	13,757	1,469	669	595	2,463	4,977	33	2,657	1,778
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴									
Veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Not a veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡

- - Quantity is zero; ‡ No data submitted.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.9b. District of Columbia – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020

Selected characteristics		Mental health diagnosis ¹								
		Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender										
Male	52.4	44.5	40.6	66.7	44.9	49.3	41.1	59.8	61.0	
Female	47.6	55.5	59.4	33.3	55.1	50.7	58.9	40.2	39.0	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age ²										
0 to 11 years	1.8	4.5	2.8	10.7	†	†	--	†	3.4	
12 to 14 years	3.2	6.4	3.1	18.4	†	1.9	--	†	5.0	
15 to 17 years	3.5	6.7	4.1	16.1	†	2.6	--	†	4.9	
18 to 20 years	4.5	6.3	4.2	16.2	2.2	4.0	1.4	†	7.0	
21 to 24 years	5.5	6.2	4.3	15.3	5.2	5.2	--	1.4	7.3	
25 to 29 years	8.0	7.7	8.1	7.5	9.6	8.6	4.1	5.2	7.7	
30 to 34 years	8.9	8.5	11.0	3.6	11.9	9.1	11.0	8.0	7.9	
35 to 39 years	8.4	8.1	9.6	2.5	10.7	8.4	17.8	8.4	8.5	
40 to 44 years	7.6	7.9	8.2	1.8	8.9	8.1	8.2	8.2	6.0	
45 to 49 years	6.7	6.4	7.8	1.5	7.6	7.3	2.7	7.5	5.6	
50 to 54 years	9.9	8.6	9.3	1.9	12.6	10.8	15.1	10.8	8.5	
55 to 59 years	12.3	10.4	10.6	2.0	13.8	13.6	12.3	15.3	10.2	
60 to 64 years	10.1	6.7	8.3	1.4	9.4	11.2	11.0	14.5	9.2	
65 years and older	9.6	5.7	8.6	1.1	6.9	8.8	16.4	20.1	8.6	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race										
American Indian or Alaska Native	†	†	†	†	†	†	--	†	†	
Asian ³	†	†	†	†	†	†	1.4	†	†	
Black or African American	95.9	97.0	91.8	97.5	94.9	97.0	88.6	92.7	96.3	
Native Hawaiian or Other Pacific Islander	†	--	--	--	--	--	--	†	--	
White	3.4	2.4	6.5	2.3	4.5	2.5	10.0	6.2	2.9	
Other	†	†	†	†	†	†	--	†	†	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.9b. District of Columbia – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020
(continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	95.1	95.2	91.3	96.1	95.5	95.2	97.0	94.1	95.3
Hispanic or Latino origin	4.9	4.8	8.7	3.9	4.5	4.8	3.0	5.9	4.7
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴									
Veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Not a veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡

- - Quantity is zero; ‡ No data submitted; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.10a. Florida – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	244,240	44,036	21,148	13,991	32,579	76,895	952	35,609	27,475
Gender									
Male	119,983	20,536	8,271	9,660	13,710	32,545	484	23,050	15,718
Female	124,257	23,500	12,877	4,331	18,869	44,350	468	12,559	11,757
Total	244,240	44,036	21,148	13,991	32,579	76,895	952	35,609	27,475
Age²									
0 to 11 years	17,470	5,854	749	5,674	61	1,206	2	47	4,136
12 to 14 years	13,717	3,520	761	3,014	194	3,289	12	74	3,110
15 to 17 years	16,051	3,675	1,161	2,113	535	5,239	15	231	3,416
18 to 20 years	12,231	2,344	1,225	865	1,091	4,296	53	869	1,695
21 to 24 years	16,540	2,831	1,622	489	2,405	5,414	100	2,520	1,521
25 to 29 years	24,973	4,304	2,429	475	4,010	7,713	136	4,474	2,091
30 to 34 years	25,154	4,453	2,480	395	4,153	7,535	146	4,672	2,110
35 to 39 years	22,846	3,978	2,138	304	3,999	6,872	126	4,367	1,762
40 to 44 years	18,432	2,945	1,728	213	3,423	5,764	92	3,629	1,345
45 to 49 years	17,500	2,492	1,523	149	3,305	6,001	73	3,365	1,311
50 to 54 years	17,875	2,491	1,475	98	3,201	6,757	63	3,377	1,305
55 to 59 years	17,856	2,249	1,494	80	3,039	7,151	67	3,458	1,347
60 to 64 years	12,152	1,541	1,159	72	1,838	4,936	36	2,428	988
65 years and older	11,443	1,359	1,204	50	1,325	4,722	31	2,098	1,338
Total	244,240	44,036	21,148	13,991	32,579	76,895	952	35,609	27,475
Race									
American Indian or Alaska Native	812	134	56	30	135	267	3	145	90
Asian ³	1,466	299	128	43	160	465	4	300	125
Black or African American	51,692	9,068	2,355	3,215	5,438	13,725	247	13,918	5,941
Native Hawaiian or Other Pacific Islander	459	72	37	17	55	166	4	72	58
White	159,662	28,703	15,998	8,614	23,356	52,135	622	17,441	17,755
Other	30,142	5,758	2,573	2,072	3,432	10,137	72	3,730	3,503
Total	244,233	44,034	21,147	13,991	32,576	76,895	952	35,606	27,472

Continued. See notes at end of table.

Table 10.10a. Florida – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	196,161	37,142	16,509	11,121	27,313	60,904	857	28,930	20,175
Hispanic or Latino origin	46,037	6,626	4,546	2,796	5,097	15,437	89	5,829	7,098
Total	242,198	43,768	21,055	13,917	32,410	76,341	946	34,759	27,273
Specific Hispanic or Latino origin									
Mexican	3,627	854	343	303	277	1,191	8	335	406
Puerto Rican	9,237	1,258	754	630	1,372	3,106	16	1,262	1,022
Cuban	7,981	534	867	255	810	2,795	18	1,142	1,862
Other specific	7,084	1,018	870	495	859	2,581	7	841	612
Total	27,929	3,664	2,834	1,683	3,318	9,673	49	3,580	3,902
Veteran status⁴									
Veteran	44,143	5,826	3,627	594	6,592	17,811	266	7,119	2,667
Not a veteran	146,137	24,560	14,417	2,553	24,356	47,672	408	25,528	13,111
Total	190,280	30,386	18,044	3,147	30,948	65,483	674	32,647	15,778

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.10b. Florida – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020

Selected characteristics		Mental health diagnosis¹							
		Trauma- and stressor-related disorders			Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders¹
					Bipolar disorders	Depressive disorders			
Total									
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender									
Male	49.1	46.6	39.1	69.0	42.1	42.3	50.8	64.7	57.2
Female	50.9	53.4	60.9	31.0	57.9	57.7	49.2	35.3	42.8
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age²									
0 to 11 years	7.2	13.3	3.5	40.6	†	1.6	†	†	15.1
12 to 14 years	5.6	8.0	3.6	21.5	†	4.3	1.3	†	11.3
15 to 17 years	6.6	8.3	5.5	15.1	1.6	6.8	1.6	†	12.4
18 to 20 years	5.0	5.3	5.8	6.2	3.3	5.6	5.6	2.4	6.2
21 to 24 years	6.8	6.4	7.7	3.5	7.4	7.0	10.5	7.1	5.5
25 to 29 years	10.2	9.8	11.5	3.4	12.3	10.0	14.3	12.6	7.6
30 to 34 years	10.3	10.1	11.7	2.8	12.7	9.8	15.3	13.1	7.7
35 to 39 years	9.4	9.0	10.1	2.2	12.3	8.9	13.2	12.3	6.4
40 to 44 years	7.5	6.7	8.2	1.5	10.5	7.5	9.7	10.2	4.9
45 to 49 years	7.2	5.7	7.2	1.1	10.1	7.8	7.7	9.4	4.8
50 to 54 years	7.3	5.7	7.0	†	9.8	8.8	6.6	9.5	4.7
55 to 59 years	7.3	5.1	7.1	†	9.3	9.3	7.0	9.7	4.9
60 to 64 years	5.0	3.5	5.5	†	5.6	6.4	3.8	6.8	3.6
65 years and older	4.7	3.1	5.7	†	4.1	6.1	3.3	5.9	4.9
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race									
American Indian or Alaska Native	†	†	†	†	†	†	†	†	†
Asian³	†	†	†	†	†	†	†	†	†
Black or African American	21.2	20.6	11.1	23.0	16.7	17.8	25.9	39.1	21.6
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	†	†	†
White	65.4	65.2	75.7	61.6	71.7	67.8	65.3	49.0	64.6
Other	12.3	13.1	12.2	14.8	10.5	13.2	7.6	10.5	12.8
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.10b. Florida – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	81.0	84.9	78.4	79.9	84.3	79.8	90.6	83.2	74.0
Hispanic or Latino origin	19.0	15.1	21.6	20.1	15.7	20.2	9.4	16.8	26.0
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	13.0	23.3	12.1	18.0	8.3	12.3	16.3	9.4	10.4
Puerto Rican	33.1	34.3	26.6	37.4	41.4	32.1	32.7	35.3	26.2
Cuban	28.6	14.6	30.6	15.2	24.4	28.9	36.7	31.9	47.7
Other specific	25.4	27.8	30.7	29.4	25.9	26.7	14.3	23.5	15.7
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁴									
Veteran	23.2	19.2	20.1	18.9	21.3	27.2	39.5	21.8	16.9
Not a veteran	76.8	80.8	79.9	81.1	78.7	72.8	60.5	78.2	83.1
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.11a. Georgia – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	140,657	25,596	31,929	12,127	25,989	45,549	3,432	29,419	55,686
Gender									
Male	69,312	10,158	12,063	7,981	9,975	18,105	1,317	17,772	32,228
Female	71,071	15,406	19,824	4,121	15,989	27,397	2,111	11,591	23,296
Total	140,383	25,564	31,887	12,102	25,964	45,502	3,428	29,363	55,524
Age²									
0 to 11 years	4,892	1,238	408	3,240	112	294	4	65	2,176
12 to 14 years	4,024	838	500	2,174	216	723	6	95	2,050
15 to 17 years	4,364	861	725	1,652	396	1,141	31	199	2,220
18 to 20 years	5,982	1,196	1,536	1,193	947	2,074	131	605	2,417
21 to 24 years	9,597	1,748	2,488	826	1,785	3,178	237	1,671	3,979
25 to 29 years	15,575	2,760	3,758	872	3,074	4,838	439	3,137	6,927
30 to 34 years	15,798	2,977	3,717	630	3,290	4,663	433	3,333	7,327
35 to 39 years	14,378	2,842	3,405	448	3,025	4,320	426	3,212	6,378
40 to 44 years	12,432	2,395	2,993	312	2,757	3,848	409	2,950	5,155
45 to 49 years	12,340	2,510	3,020	256	2,785	4,256	337	2,828	4,606
50 to 54 years	12,940	2,332	3,091	203	2,772	4,917	343	3,122	4,438
55 to 59 years	13,104	2,150	3,037	168	2,444	5,279	327	3,410	4,244
60 to 64 years	8,722	1,121	1,891	94	1,443	3,540	192	2,570	2,407
65 years and older	6,507	628	1,360	59	943	2,478	117	2,222	1,360
Total	140,655	25,596	31,929	12,127	25,989	45,549	3,432	29,419	55,684
Race									
American Indian or Alaska Native	263	55	63	22	51	76	7	59	95
Asian ³	1,686	206	278	233	242	451	19	439	714
Black or African American	59,355	9,463	8,569	5,718	8,524	17,472	1,129	18,524	22,210
Native Hawaiian or Other Pacific Islander	162	20	40	23	26	41	6	22	68
White	68,844	13,750	20,806	4,725	15,788	24,559	2,079	8,650	28,209
Other	5,737	1,305	1,393	661	835	1,997	124	888	1,979
Total	136,047	24,799	31,149	11,382	25,466	44,596	3,364	28,582	53,275

Continued. See notes at end of table.

Table 10.11a. Georgia – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	125,791	23,023	29,086	10,552	23,954	41,511	3,152	26,449	48,926
Hispanic or Latino origin	5,804	1,268	1,444	481	789	2,027	110	878	2,076
Total	131,595	24,291	30,530	11,033	24,743	43,538	3,262	27,327	51,002
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴									
Veteran	2,458	686	555	68	460	900	70	515	925
Not a veteran	105,283	19,548	26,369	4,380	22,012	37,624	2,918	24,555	36,532
Total	107,741	20,234	26,924	4,448	22,472	38,524	2,988	25,070	37,457

‡ No data submitted.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.11b. Georgia – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020

		Mental health diagnosis ¹								
Selected characteristics		Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Total individuals		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender										
Male		49.4	39.7	37.8	65.9	38.4	39.8	38.4	60.5	58.0
Female		50.6	60.3	62.2	34.1	61.6	60.2	61.6	39.5	42.0
Total		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age ²										
0 to 11 years		3.5	4.8	1.3	26.7	†	†	†	†	3.9
12 to 14 years		2.9	3.3	1.6	17.9	†	1.6	†	†	3.7
15 to 17 years		3.1	3.4	2.3	13.6	1.5	2.5	†	†	4.0
18 to 20 years		4.3	4.7	4.8	9.8	3.6	4.6	3.8	2.1	4.3
21 to 24 years		6.8	6.8	7.8	6.8	6.9	7.0	6.9	5.7	7.1
25 to 29 years		11.1	10.8	11.8	7.2	11.8	10.6	12.8	10.7	12.4
30 to 34 years		11.2	11.6	11.6	5.2	12.7	10.2	12.6	11.3	13.2
35 to 39 years		10.2	11.1	10.7	3.7	11.6	9.5	12.4	10.9	11.5
40 to 44 years		8.8	9.4	9.4	2.6	10.6	8.4	11.9	10.0	9.3
45 to 49 years		8.8	9.8	9.5	2.1	10.7	9.3	9.8	9.6	8.3
50 to 54 years		9.2	9.1	9.7	1.7	10.7	10.8	10.0	10.6	8.0
55 to 59 years		9.3	8.4	9.5	1.4	9.4	11.6	9.5	11.6	7.6
60 to 64 years		6.2	4.4	5.9	†	5.6	7.8	5.6	8.7	4.3
65 years and older		4.6	2.5	4.3	†	3.6	5.4	3.4	7.6	2.4
Total		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race										
American Indian or Alaska Native		†	†	†	†	†	†	†	†	†
Asian ³		1.2	†	†	2.0	†	1.0	†	1.5	1.3
Black or African American		43.6	38.2	27.5	50.2	33.5	39.2	33.6	64.8	41.7
Native Hawaiian or Other Pacific Islander		†	†	†	†	†	†	†	†	†
White		50.6	55.4	66.8	41.5	62.0	55.1	61.8	30.3	52.9
Other		4.2	5.3	4.5	5.8	3.3	4.5	3.7	3.1	3.7
Total		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.11b. Georgia – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020 (continued)

Selected characteristics		Mental health diagnosis¹								
		Trauma- and stressor-related disorders			Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders¹	
					Bipolar disorders	Depressive disorders				
Total										
Ethnicity										
Not of Hispanic or Latino origin	95.6		94.8	95.3	95.6	96.8	95.3	96.6	96.8	95.9
Hispanic or Latino origin	4.4		5.2	4.7	4.4	3.2	4.7	3.4	3.2	4.1
Total	100.0		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin										
Mexican	‡		‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡		‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡		‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡		‡	‡	‡	‡	‡	‡	‡	‡
Total	‡		‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴										
Veteran	2.3		3.4	2.1	1.5	2.0	2.3	2.3	2.1	2.5
Not a veteran	97.7		96.6	97.9	98.5	98.0	97.7	97.7	97.9	97.5
Total	100.0		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

‡ No data submitted; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.12a. Hawaii – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	6,083	1,120	501	512	784	1,352	148	2,354	1,221
Gender									
Male	3,634	563	250	404	393	645	71	1,570	833
Female	2,434	554	250	108	391	703	77	780	383
Total	6,068	1,117	500	512	784	1,348	148	2,350	1,216
Age²									
0 to 11 years	545	204	75	250	1	36	--	1	261
12 to 14 years	460	153	75	132	9	112	1	7	240
15 to 17 years	665	209	115	109	24	199	4	26	347
18 to 20 years	161	38	19	17	7	48	2	21	72
21 to 24 years	123	30	5	2	19	17	--	54	7
25 to 29 years	244	46	6	--	34	50	7	110	23
30 to 34 years	366	47	20	1	47	74	8	200	35
35 to 39 years	380	50	10	--	86	72	8	194	31
40 to 44 years	366	39	20	--	68	73	20	205	29
45 to 49 years	415	52	30	--	78	89	9	225	26
50 to 54 years	480	48	23	--	81	119	18	279	30
55 to 59 years	565	71	32	--	87	136	24	326	33
60 to 64 years	549	67	36	--	86	126	15	314	36
65 years and older	762	66	35	1	157	201	32	390	51
Total	6,081	1,120	501	512	784	1,352	148	2,352	1,221
Race									
American Indian or Alaska Native	40	11	6	--	5	11	2	15	4
Asian ³	1,038	73	72	35	101	177	21	691	129
Black or African American	119	19	5	13	15	15	3	62	17
Native Hawaiian or Other Pacific Islander	728	154	44	53	69	154	26	286	184
White	1,486	208	140	75	328	376	53	576	218
Other	1,781	415	189	298	155	338	36	525	555
Total	5,192	880	456	474	673	1,071	141	2,155	1,107

Continued. See notes at end of table.

Table 10.12a. Hawaii – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020 (continued)

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	4,634	749	411	391	623	938	128	2,016	940
Hispanic or Latino origin	557	136	58	94	54	131	13	138	178
Total	5,191	885	469	485	677	1,069	141	2,154	1,118
Specific Hispanic or Latino origin									
Mexican	104	30	10	23	12	30	2	10	35
Puerto Rican	255	63	25	42	21	61	7	60	94
Cuban	4	1	1	- -	1	1	- -	1	- -
Other specific	86	12	8	- -	14	15	3	56	5
Total	449	106	44	65	48	107	12	127	134
Veteran status⁴									
Veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Not a veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡

- - Quantity is zero; ‡ No data submitted.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.12b. Hawaii – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020

		Mental health diagnosis¹								
Selected characteristics		Total	Trauma- and stressor-related disorders			Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders¹
						Bipolar disorders	Depressive disorders			
Total individuals		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender										
Male		59.9	50.4	50.0	78.9	50.1	47.8	48.0	66.8	68.5
Female		40.1	49.6	50.0	21.1	49.9	52.2	52.0	33.2	31.5
Total		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age²										
0 to 11 years		9.0	18.2	15.0	48.8	†	2.7	--	†	21.4
12 to 14 years		7.6	13.7	15.0	25.8	1.1	8.3	†	†	19.7
15 to 17 years		10.9	18.7	23.0	21.3	3.1	14.7	2.7	1.1	28.4
18 to 20 years		2.6	3.4	3.8	3.3	†	3.6	1.4	†	5.9
21 to 24 years		2.0	2.7	†	†	2.4	1.3	--	2.3	†
25 to 29 years		4.0	4.1	1.2	--	4.3	3.7	4.7	4.7	1.9
30 to 34 years		6.0	4.2	4.0	†	6.0	5.5	5.4	8.5	2.9
35 to 39 years		6.2	4.5	2.0	--	11.0	5.3	5.4	8.2	2.5
40 to 44 years		6.0	3.5	4.0	--	8.7	5.4	13.5	8.7	2.4
45 to 49 years		6.8	4.6	6.0	--	9.9	6.6	6.1	9.6	2.1
50 to 54 years		7.9	4.3	4.6	--	10.3	8.8	12.2	11.9	2.5
55 to 59 years		9.3	6.3	6.4	--	11.1	10.1	16.2	13.9	2.7
60 to 64 years		9.0	6.0	7.2	--	11.0	9.3	10.1	13.4	2.9
65 years and older		12.5	5.9	7.0	†	20.0	14.9	21.6	16.6	4.2
Total		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race										
American Indian or Alaska Native		†	1.3	1.3	--	†	1.0	1.4	†	†
Asian³		20.0	8.3	15.8	7.4	15.0	16.5	14.9	32.1	11.7
Black or African American		2.3	2.2	1.1	2.7	2.2	1.4	2.1	2.9	1.5
Native Hawaiian or Other Pacific Islander		14.0	17.5	9.6	11.2	10.3	14.4	18.4	13.3	16.6
White		28.6	23.6	30.7	15.8	48.7	35.1	37.6	26.7	19.7
Other		34.3	47.2	41.4	62.9	23.0	31.6	25.5	24.4	50.1
Total		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.12b. Hawaii – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020 (continued)

Mental health diagnosis ¹									
Selected characteristics	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	89.3	84.6	87.6	80.6	92.0	87.7	90.8	93.6	84.1
Hispanic or Latino origin	10.7	15.4	12.4	19.4	8.0	12.3	9.2	6.4	15.9
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	23.2	28.3	22.7	35.4	25.0	28.0	16.7	7.9	26.1
Puerto Rican	56.8	59.4	56.8	64.6	43.8	57.0	58.3	47.2	70.1
Cuban	†	†	2.3	--	2.1	†	--	†	--
Other specific	19.2	11.3	18.2	--	29.2	14.0	25.0	44.1	3.7
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁴									
Veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Not a veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡

-- Quantity is zero; ‡ No data submitted; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.13a. Idaho – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	7,082	805	1,237	691	1,440	2,389	514	1,993	1,306
Gender									
Male	3,857	319	552	520	618	1,127	225	1,274	873
Female	3,218	486	682	169	821	1,262	289	719	430
Total	7,075	805	1,234	689	1,439	2,389	514	1,993	1,303
Age²									
0 to 11 years	169	23	40	86	--	11	--	2	105
12 to 14 years	426	78	95	178	10	128	1	5	252
15 to 17 years	707	138	224	228	38	305	9	17	371
18 to 20 years	475	81	116	71	60	205	28	75	161
21 to 24 years	520	64	80	28	106	176	51	172	86
25 to 29 years	692	73	97	24	161	229	81	253	67
30 to 34 years	748	83	107	16	193	252	83	259	49
35 to 39 years	725	64	119	21	182	229	67	252	61
40 to 44 years	578	49	82	10	145	201	48	201	41
45 to 49 years	517	38	83	10	150	162	45	181	34
50 to 54 years	495	50	74	8	136	175	31	163	19
55 to 59 years	458	37	61	6	117	144	33	182	18
60 to 64 years	281	18	32	5	72	95	19	105	12
65 years and older	290	9	27	--	69	77	18	126	30
Total	7,081	805	1,237	691	1,439	2,389	514	1,993	1,306
Race									
American Indian or Alaska Native	141	19	18	15	22	45	11	40	30
Asian ³	37	1	8	2	7	9	1	20	4
Black or African American	139	13	15	14	18	30	5	62	33
Native Hawaiian or Other Pacific Islander	28	5	6	2	4	10	2	5	9
White	5,471	606	972	540	1,162	1,857	424	1,511	950
Other	523	63	100	49	94	182	33	152	108
Total	6,339	707	1,119	622	1,307	2,133	476	1,790	1,134

Continued. See notes at end of table.

Table 10.13a. Idaho – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	5,436	601	955	532	1,163	1,820	420	1,511	950
Hispanic or Latino origin	701	86	130	76	111	231	34	220	154
Total	6,137	687	1,085	608	1,274	2,051	454	1,731	1,104
Specific Hispanic or Latino origin									
Mexican	104	10	16	10	22	28	6	42	23
Puerto Rican	14	2	1	--	2	3	--	10	--
Cuban	1	--	--	--	--	1	--	--	--
Other specific	76	4	15	5	14	17	5	28	11
Total	195	16	32	15	38	49	11	80	34
Veteran status⁴									
Veteran	191	22	25	4	29	65	10	78	14
Not a veteran	4,488	443	728	172	1,083	1,538	399	1,472	470
Total	4,679	465	753	176	1,112	1,603	409	1,550	484

-- Quantity is zero.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.13b. Idaho – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020

Selected characteristics		Mental health diagnosis¹							
		Trauma- and stressor-related disorders			Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders¹
					Bipolar disorders	Depressive disorders			
Total									
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender									
Male	54.5	39.6	44.7	75.5	42.9	47.2	43.8	63.9	67.0
Female	45.5	60.4	55.3	24.5	57.1	52.8	56.2	36.1	33.0
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age²									
0 to 11 years	2.4	2.9	3.2	12.4	--	†	--	†	8.0
12 to 14 years	6.0	9.7	7.7	25.8	†	5.4	†	†	19.3
15 to 17 years	10.0	17.1	18.1	33.0	2.6	12.8	1.8	†	28.4
18 to 20 years	6.7	10.1	9.4	10.3	4.2	8.6	5.4	3.8	12.3
21 to 24 years	7.3	8.0	6.5	4.1	7.4	7.4	9.9	8.6	6.6
25 to 29 years	9.8	9.1	7.8	3.5	11.2	9.6	15.8	12.7	5.1
30 to 34 years	10.6	10.3	8.6	2.3	13.4	10.5	16.1	13.0	3.8
35 to 39 years	10.2	8.0	9.6	3.0	12.6	9.6	13.0	12.6	4.7
40 to 44 years	8.2	6.1	6.6	1.4	10.1	8.4	9.3	10.1	3.1
45 to 49 years	7.3	4.7	6.7	1.4	10.4	6.8	8.8	9.1	2.6
50 to 54 years	7.0	6.2	6.0	1.2	9.5	7.3	6.0	8.2	1.5
55 to 59 years	6.5	4.6	4.9	†	8.1	6.0	6.4	9.1	1.4
60 to 64 years	4.0	2.2	2.6	†	5.0	4.0	3.7	5.3	†
65 years and older	4.1	1.1	2.2	--	4.8	3.2	3.5	6.3	2.3
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race									
American Indian or Alaska Native	2.2	2.7	1.6	2.4	1.7	2.1	2.3	2.2	2.6
Asian³	†	†	†	†	†	†	†	1.1	†
Black or African American	2.2	1.8	1.3	2.3	1.4	1.4	1.1	3.5	2.9
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	†	†	†
White	86.3	85.7	86.9	86.8	88.9	87.1	89.1	84.4	83.8
Other	8.3	8.9	8.9	7.9	7.2	8.5	6.9	8.5	9.5
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.13b. Idaho – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	88.6	87.5	88.0	87.5	91.3	88.7	92.5	87.3	86.1
Hispanic or Latino origin	11.4	12.5	12.0	12.5	8.7	11.3	7.5	12.7	13.9
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	53.3	62.5	50.0	66.7	57.9	57.1	54.5	52.5	67.6
Puerto Rican	7.2	12.5	3.1	--	5.3	6.1	--	12.5	--
Cuban	†	--	--	--	--	2.0	--	--	--
Other specific	39.0	25.0	46.9	33.3	36.8	34.7	45.5	35.0	32.4
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁴									
Veteran	4.1	4.7	3.3	2.3	2.6	4.1	2.4	5.0	2.9
Not a veteran	95.9	95.3	96.7	97.7	97.4	95.9	97.6	95.0	97.1
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

-- Quantity is zero; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.14a. Illinois – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	39,267	8,298	8,079	4,198	6,111	11,329	1,239	7,682	8,625
Gender									
Male	19,743	3,387	3,048	2,849	2,685	4,295	439	5,108	5,367
Female	19,524	4,911	5,031	1,349	3,426	7,034	800	2,574	3,258
Total	39,267	8,298	8,079	4,198	6,111	11,329	1,239	7,682	8,625
Age²									
0 to 11 years	4,472	1,941	483	1,535	22	234	1	8	1,343
12 to 14 years	3,127	933	578	899	60	800	3	13	929
15 to 17 years	3,256	760	779	690	140	1,267	7	41	864
18 to 20 years	2,233	487	593	281	282	910	41	180	588
21 to 24 years	2,102	392	474	140	406	674	64	443	551
25 to 29 years	3,025	538	640	144	651	876	122	702	768
30 to 34 years	2,844	498	606	132	609	782	121	717	666
35 to 39 years	2,769	501	576	104	597	780	116	758	564
40 to 44 years	2,656	445	600	67	622	739	115	796	443
45 to 49 years	2,542	434	535	65	615	758	144	737	438
50 to 54 years	2,693	417	622	50	613	880	157	844	404
55 to 59 years	3,028	419	654	40	679	988	151	976	465
60 to 64 years	2,217	251	456	27	437	770	114	751	306
65 years and older	2,303	282	483	24	378	871	83	716	296
Total	39,267	8,298	8,079	4,198	6,111	11,329	1,239	7,682	8,625
Race									
American Indian or Alaska Native	148	37	35	12	30	46	7	25	26
Asian ³	541	114	110	32	80	161	7	147	99
Black or African American	8,517	1,604	1,003	801	1,259	2,040	163	2,756	2,080
Native Hawaiian or Other Pacific Islander	33	4	4	2	4	10	2	9	5
White	25,636	5,515	6,104	2,823	4,287	7,780	1,008	4,161	5,287
Other	386	121	50	53	50	84	8	68	96
Total	35,261	7,395	7,306	3,723	5,710	10,121	1,195	7,166	7,593

Continued. See notes at end of table.

Table 10.14a. Illinois – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	31,128	6,314	6,171	3,258	5,208	8,713	1,108	6,560	6,991
Hispanic or Latino origin	4,204	1,044	1,107	454	443	1,438	71	583	847
Total	35,332	7,358	7,278	3,712	5,651	10,151	1,179	7,143	7,838
Specific Hispanic or Latino origin									
Mexican	2,500	660	703	275	227	849	35	294	500
Puerto Rican	414	94	96	56	64	138	7	75	86
Cuban	32	6	10	3	6	12	1	7	4
Other specific	1,166	280	291	120	130	427	24	155	216
Total	4,112	1,040	1,100	454	427	1,426	67	531	806
Veteran status⁴									
Veteran	588	107	109	11	129	165	28	219	88
Not a veteran	24,743	4,026	5,526	985	5,162	7,841	982	6,633	4,902
Total	25,331	4,133	5,635	996	5,291	8,006	1,010	6,852	4,990

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.14b. Illinois – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020

		Mental health diagnosis ¹								
Selected characteristics		Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Total individuals		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender										
Male		50.3	40.8	37.7	67.9	43.9	37.9	35.4	66.5	62.2
Female		49.7	59.2	62.3	32.1	56.1	62.1	64.6	33.5	37.8
Total		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age ²										
0 to 11 years		11.4	23.4	6.0	36.6	†	2.1	†	†	15.6
12 to 14 years		8.0	11.2	7.2	21.4	†	7.1	†	†	10.8
15 to 17 years		8.3	9.2	9.6	16.4	2.3	11.2	†	†	10.0
18 to 20 years		5.7	5.9	7.3	6.7	4.6	8.0	3.3	2.3	6.8
21 to 24 years		5.4	4.7	5.9	3.3	6.6	5.9	5.2	5.8	6.4
25 to 29 years		7.7	6.5	7.9	3.4	10.7	7.7	9.8	9.1	8.9
30 to 34 years		7.2	6.0	7.5	3.1	10.0	6.9	9.8	9.3	7.7
35 to 39 years		7.1	6.0	7.1	2.5	9.8	6.9	9.4	9.9	6.5
40 to 44 years		6.8	5.4	7.4	1.6	10.2	6.5	9.3	10.4	5.1
45 to 49 years		6.5	5.2	6.6	1.5	10.1	6.7	11.6	9.6	5.1
50 to 54 years		6.9	5.0	7.7	1.2	10.0	7.8	12.7	11.0	4.7
55 to 59 years		7.7	5.0	8.1	†	11.1	8.7	12.2	12.7	5.4
60 to 64 years		5.6	3.0	5.6	†	7.2	6.8	9.2	9.8	3.5
65 years and older		5.9	3.4	6.0	†	6.2	7.7	6.7	9.3	3.4
Total		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race										
American Indian or Alaska Native		†	†	†	†	†	†	†	†	†
Asian ³		1.5	1.5	1.5	†	1.4	1.6	†	2.1	1.3
Black or African American		24.2	21.7	13.7	21.5	22.0	20.2	13.6	38.5	27.4
Native Hawaiian or Other Pacific Islander		†	†	†	†	†	†	†	†	†
White		72.7	74.6	83.5	75.8	75.1	76.9	84.4	58.1	69.6
Other		1.1	1.6	†	1.4	†	†	†	†	1.3
Total		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.14b. Illinois – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	88.1	85.8	84.8	87.8	92.2	85.8	94.0	91.8	89.2
Hispanic or Latino origin	11.9	14.2	15.2	12.2	7.8	14.2	6.0	8.2	10.8
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	60.8	63.5	63.9	60.6	53.2	59.5	52.2	55.4	62.0
Puerto Rican	10.1	9.0	8.7	12.3	15.0	9.7	10.4	14.1	10.7
Cuban	†	†	†	†	1.4	†	1.5	1.3	†
Other specific	28.4	26.9	26.5	26.4	30.4	29.9	35.8	29.2	26.8
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁴									
Veteran	2.3	2.6	1.9	1.1	2.4	2.1	2.8	3.2	1.8
Not a veteran	97.7	97.4	98.1	98.9	97.6	97.9	97.2	96.8	98.2
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.15a. Indiana – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	137,905	31,334	50,737	27,231	13,981	51,612	6,161	14,278	36,630
Gender									
Male	64,806	11,450	19,615	18,104	4,958	18,702	1,775	8,609	22,830
Female	73,099	19,884	31,122	9,127	9,023	32,910	4,386	5,669	13,800
Total	137,905	31,334	50,737	27,231	13,981	51,612	6,161	14,278	36,630
Age²									
0 to 11 years	29,292	7,109	7,439	12,889	65	3,307	4	66	15,585
12 to 14 years	14,479	3,122	4,608	5,818	154	5,042	21	90	6,595
15 to 17 years	12,354	2,700	4,666	3,533	360	6,070	86	146	4,640
18 to 20 years	6,318	1,416	2,809	1,123	581	3,363	275	276	1,567
21 to 24 years	7,073	1,669	3,164	667	971	3,463	501	700	1,110
25 to 29 years	10,006	2,428	4,376	717	1,517	4,533	748	1,294	1,278
30 to 34 years	9,469	2,418	4,124	595	1,476	4,130	704	1,334	1,066
35 to 39 years	9,119	2,322	3,923	514	1,566	3,869	723	1,521	926
40 to 44 years	7,861	1,886	3,283	399	1,486	3,342	663	1,454	793
45 to 49 years	7,500	1,853	3,004	313	1,528	3,233	612	1,483	715
50 to 54 years	7,260	1,622	2,855	232	1,416	3,306	583	1,545	725
55 to 59 years	7,322	1,434	2,803	224	1,342	3,372	575	1,806	662
60 to 64 years	5,143	781	1,969	132	855	2,355	363	1,357	508
65 years and older	4,709	574	1,714	75	664	2,227	303	1,206	460
Total	137,905	31,334	50,737	27,231	13,981	51,612	6,161	14,278	36,630
Race									
American Indian or Alaska Native	420	104	144	61	55	163	22	56	90
Asian ³	553	106	184	64	41	242	29	108	95
Black or African American	18,988	3,658	4,985	4,393	1,430	5,755	600	3,680	5,777
Native Hawaiian or Other Pacific Islander	108	21	41	18	17	29	2	13	28
White	105,962	24,475	41,331	19,879	11,482	41,212	5,208	9,639	27,031
Other	11,871	2,970	4,051	2,816	956	4,210	300	779	3,607
Total	137,902	31,334	50,736	27,231	13,981	51,611	6,161	14,275	36,628

Continued. See notes at end of table.

Table 10.15a. Indiana – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	126,210	28,649	46,554	24,595	13,092	47,282	5,830	13,409	33,079
Hispanic or Latino origin	9,829	2,200	3,552	2,179	645	3,730	234	624	2,985
Total	136,039	30,849	50,106	26,774	13,737	51,012	6,064	14,033	36,064
Specific Hispanic or Latino origin									
Mexican	3,227	690	1,133	704	165	1,325	39	169	912
Puerto Rican	602	114	225	136	74	211	20	69	154
Cuban	86	22	28	17	7	37	1	9	18
Other specific	5,886	1,365	2,160	1,317	396	2,152	174	366	1,889
Total	9,801	2,191	3,546	2,174	642	3,725	234	613	2,973
Veteran status⁴									
Veteran	2,144	600	753	86	323	919	137	434	193
Not a veteran	78,888	17,654	32,992	4,695	13,037	35,898	5,905	13,507	9,302
Total	81,032	18,254	33,745	4,781	13,360	36,817	6,042	13,941	9,495

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.15b. Indiana – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020

Selected characteristics		Mental health diagnosis ¹							
		Trauma- and stressor-related disorders			Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total									
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender									
Male	47.0	36.5	38.7	66.5	35.5	36.2	28.8	60.3	62.3
Female	53.0	63.5	61.3	33.5	64.5	63.8	71.2	39.7	37.7
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age ²									
0 to 11 years	21.2	22.7	14.7	47.3	†	6.4	†	†	42.5
12 to 14 years	10.5	10.0	9.1	21.4	1.1	9.8	†	†	18.0
15 to 17 years	9.0	8.6	9.2	13.0	2.6	11.8	1.4	1.0	12.7
18 to 20 years	4.6	4.5	5.5	4.1	4.2	6.5	4.5	1.9	4.3
21 to 24 years	5.1	5.3	6.2	2.4	6.9	6.7	8.1	4.9	3.0
25 to 29 years	7.3	7.7	8.6	2.6	10.9	8.8	12.1	9.1	3.5
30 to 34 years	6.9	7.7	8.1	2.2	10.6	8.0	11.4	9.3	2.9
35 to 39 years	6.6	7.4	7.7	1.9	11.2	7.5	11.7	10.7	2.5
40 to 44 years	5.7	6.0	6.5	1.5	10.6	6.5	10.8	10.2	2.2
45 to 49 years	5.4	5.9	5.9	1.1	10.9	6.3	9.9	10.4	2.0
50 to 54 years	5.3	5.2	5.6	†	10.1	6.4	9.5	10.8	2.0
55 to 59 years	5.3	4.6	5.5	†	9.6	6.5	9.3	12.6	1.8
60 to 64 years	3.7	2.5	3.9	†	6.1	4.6	5.9	9.5	1.4
65 years and older	3.4	1.8	3.4	†	4.7	4.3	4.9	8.4	1.3
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race									
American Indian or Alaska Native	†	†	†	†	†	†	†	†	†
Asian ³	†	†	†	†	†	†	†	†	†
Black or African American	13.8	11.7	9.8	16.1	10.2	11.2	9.7	25.8	15.8
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	†	†	†
White	76.8	78.1	81.5	73.0	82.1	79.9	84.5	67.5	73.8
Other	8.6	9.5	8.0	10.3	6.8	8.2	4.9	5.5	9.8
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.15b. Indiana – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	92.8	92.9	92.9	91.9	95.3	92.7	96.1	95.6	91.7
Hispanic or Latino origin	7.2	7.1	7.1	8.1	4.7	7.3	3.9	4.4	8.3
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	32.9	31.5	32.0	32.4	25.7	35.6	16.7	27.6	30.7
Puerto Rican	6.1	5.2	6.3	6.3	11.5	5.7	8.5	11.3	5.2
Cuban	†	1.0	†	†	1.1	†	†	1.5	†
Other specific	60.1	62.3	60.9	60.6	61.7	57.8	74.4	59.7	63.5
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁴									
Veteran	2.6	3.3	2.2	1.8	2.4	2.5	2.3	3.1	2.0
Not a veteran	97.4	96.7	97.8	98.2	97.6	97.5	97.7	96.9	98.0
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.16a. Iowa – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	56,738	4,714	22,784	7,147	6,501	20,852	1,157	7,155	9,473
Gender									
Male	24,403	1,843	8,009	4,747	2,568	7,572	345	4,146	5,047
Female	32,335	2,871	14,775	2,400	3,933	13,280	812	3,009	4,426
Total	56,738	4,714	22,784	7,147	6,501	20,852	1,157	7,155	9,473
Age²									
0 to 11 years	5,689	519	1,626	2,993	20	358	5	23	2,251
12 to 14 years	3,525	359	1,052	1,325	80	1,311	11	39	1,108
15 to 17 years	4,077	425	1,482	921	195	1,954	23	66	1,053
18 to 20 years	3,127	285	1,251	388	326	1,368	82	206	458
21 to 24 years	3,557	289	1,584	298	408	1,432	110	352	456
25 to 29 years	5,299	438	2,396	371	803	2,028	161	725	624
30 to 34 years	4,974	415	2,241	266	817	1,835	160	734	559
35 to 39 years	4,651	421	2,138	191	717	1,733	109	786	508
40 to 44 years	3,845	321	1,786	144	684	1,442	134	603	406
45 to 49 years	3,427	275	1,511	86	590	1,356	78	632	378
50 to 54 years	3,496	292	1,488	69	518	1,430	95	629	429
55 to 59 years	3,705	280	1,500	46	551	1,551	81	750	433
60 to 64 years	2,778	171	1,111	30	354	1,137	54	594	327
65 years and older	4,588	224	1,618	19	438	1,917	54	1,016	483
Total	56,738	4,714	22,784	7,147	6,501	20,852	1,157	7,155	9,473
Race									
American Indian or Alaska Native	932	101	366	94	145	363	25	108	117
Asian ³	545	52	207	42	47	213	6	91	98
Black or African American	3,490	382	1,211	373	375	1,179	50	751	550
Native Hawaiian or Other Pacific Islander	5	2	--	2	--	1	--	--	--
White	38,053	3,015	15,763	4,241	4,902	14,336	898	5,135	5,966
Other	1,671	190	591	360	133	602	34	117	378
Total	44,696	3,742	18,138	5,112	5,602	16,694	1,013	6,202	7,109

Continued. See notes at end of table.

Table 10.16a. Iowa – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	43,421	3,637	17,662	4,919	5,449	16,252	980	6,039	6,871
Hispanic or Latino origin	2,567	252	1,069	360	195	906	40	207	442
Total	45,988	3,889	18,731	5,279	5,644	17,158	1,020	6,246	7,313
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴									
Veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Not a veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡

- - Quantity is zero; ‡ No data submitted.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.16b. Iowa – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020

Selected characteristics		Mental health diagnosis ¹							
		Trauma- and stressor-related disorders			Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total									
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender									
Male	43.0	39.1	35.2	66.4	39.5	36.3	29.8	57.9	53.3
Female	57.0	60.9	64.8	33.6	60.5	63.7	70.2	42.1	46.7
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age ²									
0 to 11 years	10.0	11.0	7.1	41.9	†	1.7	†	†	23.8
12 to 14 years	6.2	7.6	4.6	18.5	1.2	6.3	†	†	11.7
15 to 17 years	7.2	9.0	6.5	12.9	3.0	9.4	2.0	†	11.1
18 to 20 years	5.5	6.0	5.5	5.4	5.0	6.6	7.1	2.9	4.8
21 to 24 years	6.3	6.1	7.0	4.2	6.3	6.9	9.5	4.9	4.8
25 to 29 years	9.3	9.3	10.5	5.2	12.4	9.7	13.9	10.1	6.6
30 to 34 years	8.8	8.8	9.8	3.7	12.6	8.8	13.8	10.3	5.9
35 to 39 years	8.2	8.9	9.4	2.7	11.0	8.3	9.4	11.0	5.4
40 to 44 years	6.8	6.8	7.8	2.0	10.5	6.9	11.6	8.4	4.3
45 to 49 years	6.0	5.8	6.6	1.2	9.1	6.5	6.7	8.8	4.0
50 to 54 years	6.2	6.2	6.5	†	8.0	6.9	8.2	8.8	4.5
55 to 59 years	6.5	5.9	6.6	†	8.5	7.4	7.0	10.5	4.6
60 to 64 years	4.9	3.6	4.9	†	5.4	5.5	4.7	8.3	3.5
65 years and older	8.1	4.8	7.1	†	6.7	9.2	4.7	14.2	5.1
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race									
American Indian or Alaska Native	2.1	2.7	2.0	1.8	2.6	2.2	2.5	1.7	1.6
Asian ³	1.2	1.4	1.1	†	†	1.3	†	1.5	1.4
Black or African American	7.8	10.2	6.7	7.3	6.7	7.1	4.9	12.1	7.7
Native Hawaiian or Other Pacific Islander	†	†	--	†	--	†	--	--	--
White	85.1	80.6	86.9	83.0	87.5	85.9	88.6	82.8	83.9
Other	3.7	5.1	3.3	7.0	2.4	3.6	3.4	1.9	5.3
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.16b. Iowa – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020 (continued)

Selected characteristics		Mental health diagnosis¹								
		Trauma- and stressor-related disordersAnxiety disordersADHD				Mood disorders		Personality disordersSchizophrenia, other psychotic disordersOther mental disorders¹		
						Bipolar disorders	Depressive disorders			
Total		94.4	93.5	94.3	93.2	96.5	94.7	96.1	96.7	94.0
Ethnicity										
Not of Hispanic or Latino origin		94.4	93.5	94.3	93.2	96.5	94.7	96.1	96.7	94.0
Hispanic or Latino origin		5.6	6.5	5.7	6.8	3.5	5.3	3.9	3.3	6.0
Total		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin										
Mexican		‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican		‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban		‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific		‡	‡	‡	‡	‡	‡	‡	‡	‡
Total		‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴										
Veteran		‡	‡	‡	‡	‡	‡	‡	‡	‡
Not a veteran		‡	‡	‡	‡	‡	‡	‡	‡	‡
Total		‡	‡	‡	‡	‡	‡	‡	‡	‡

- - Quantity is zero; ‡ No data submitted; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.17a. Kansas – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	139,114	30,712	33,232	15,678	15,838	47,536	2,328	8,746	34,069
Gender									
Male	65,200	12,686	12,913	10,971	6,213	17,542	905	5,380	20,875
Female	73,351	17,932	20,217	4,664	9,598	29,782	1,422	3,350	13,126
Total	138,551	30,618	33,130	15,635	15,811	47,324	2,327	8,730	34,001
Age²									
0 to 11 years	16,754	5,588	2,471	4,953	56	882	- -	17	6,414
12 to 14 years	10,351	2,685	1,967	2,931	165	2,128	6	34	3,544
15 to 17 years	11,115	2,472	2,606	2,270	338	3,747	16	74	3,347
18 to 20 years	8,727	1,842	2,276	1,415	710	3,247	73	205	2,282
21 to 24 years	10,125	2,072	2,753	1,153	1,194	3,873	164	475	2,454
25 to 29 years	12,488	2,627	3,498	1,018	1,783	4,787	251	798	2,875
30 to 34 years	11,853	2,543	3,312	597	1,957	4,442	287	905	2,536
35 to 39 years	11,105	2,369	3,017	408	1,906	4,120	303	1,043	2,249
40 to 44 years	9,311	1,946	2,560	274	1,699	3,611	278	934	1,727
45 to 49 years	7,846	1,678	2,013	210	1,452	3,168	215	849	1,378
50 to 54 years	7,514	1,464	1,805	150	1,345	3,246	208	878	1,316
55 to 59 years	7,510	1,294	1,768	132	1,274	3,489	221	916	1,308
60 to 64 years	5,801	875	1,340	76	931	2,756	134	730	1,048
65 years and older	8,614	1,257	1,846	91	1,028	4,040	172	888	1,591
Total	139,114	30,712	33,232	15,678	15,838	47,536	2,328	8,746	34,069
Race									
American Indian or Alaska Native	3,543	813	619	376	479	1,173	42	223	841
Asian ³	1,438	291	247	110	131	437	14	155	330
Black or African American	10,328	2,460	1,926	1,323	1,178	3,148	141	1,619	2,832
Native Hawaiian or Other Pacific Islander	248	64	49	32	29	69	4	22	53
White	105,532	22,689	25,987	11,778	12,354	36,822	1,875	5,394	25,357
Other	17,285	4,245	4,277	2,011	1,601	5,560	244	1,250	4,466
Total	138,374	30,562	33,105	15,630	15,772	47,209	2,320	8,663	33,879

Continued. See notes at end of table.

Table 10.17a. Kansas – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	126,982	27,691	30,578	14,399	15,044	43,216	2,183	8,089	30,990
Hispanic or Latino origin	10,258	2,674	2,280	1,127	625	3,571	124	533	2,695
Total	137,240	30,365	32,858	15,526	15,669	46,787	2,307	8,622	33,685
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴									
Veteran	1,761	603	513	59	180	709	16	115	280
Not a veteran	53,995	11,669	16,150	3,837	7,037	21,541	1,150	4,212	12,488
Total	55,756	12,272	16,663	3,896	7,217	22,250	1,166	4,327	12,768

- - Quantity is zero; ‡ No data submitted.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.17b. Kansas – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020

Selected characteristics		Mental health diagnosis ¹								
		Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender										
Male	47.1	41.4	39.0	70.2	39.3	37.1	38.9	61.6	61.4	
Female	52.9	58.6	61.0	29.8	60.7	62.9	61.1	38.4	38.6	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age ²										
0 to 11 years	12.0	18.2	7.4	31.6	†	1.9	--	†	18.8	
12 to 14 years	7.4	8.7	5.9	18.7	1.0	4.5	†	†	10.4	
15 to 17 years	8.0	8.0	7.8	14.5	2.1	7.9	†	†	9.8	
18 to 20 years	6.3	6.0	6.8	9.0	4.5	6.8	3.1	2.3	6.7	
21 to 24 years	7.3	6.7	8.3	7.4	7.5	8.1	7.0	5.4	7.2	
25 to 29 years	9.0	8.6	10.5	6.5	11.3	10.1	10.8	9.1	8.4	
30 to 34 years	8.5	8.3	10.0	3.8	12.4	9.3	12.3	10.3	7.4	
35 to 39 years	8.0	7.7	9.1	2.6	12.0	8.7	13.0	11.9	6.6	
40 to 44 years	6.7	6.3	7.7	1.7	10.7	7.6	11.9	10.7	5.1	
45 to 49 years	5.6	5.5	6.1	1.3	9.2	6.7	9.2	9.7	4.0	
50 to 54 years	5.4	4.8	5.4	†	8.5	6.8	8.9	10.0	3.9	
55 to 59 years	5.4	4.2	5.3	†	8.0	7.3	9.5	10.5	3.8	
60 to 64 years	4.2	2.8	4.0	†	5.9	5.8	5.8	8.3	3.1	
65 years and older	6.2	4.1	5.6	†	6.5	8.5	7.4	10.2	4.7	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race										
American Indian or Alaska Native	2.6	2.7	1.9	2.4	3.0	2.5	1.8	2.6	2.5	
Asian ³	1.0	†	†	†	†	†	†	1.8	†	
Black or African American	7.5	8.0	5.8	8.5	7.5	6.7	6.1	18.7	8.4	
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	†	†	†	
White	76.3	74.2	78.5	75.4	78.3	78.0	80.8	62.3	74.8	
Other	12.5	13.9	12.9	12.9	10.2	11.8	10.5	14.4	13.2	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.17b. Kansas – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	92.5	91.2	93.1	92.7	96.0	92.4	94.6	93.8	92.0
Hispanic or Latino origin	7.5	8.8	6.9	7.3	4.0	7.6	5.4	6.2	8.0
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴									
Veteran	3.2	4.9	3.1	1.5	2.5	3.2	1.4	2.7	2.2
Not a veteran	96.8	95.1	96.9	98.5	97.5	96.8	98.6	97.3	97.8
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

- - Quantity is zero; ‡ No data submitted; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.18a. Kentucky – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	153,854	35,466	37,282	19,408	12,056	43,158	3,599	11,274	73,239
Gender									
Male	76,559	14,284	14,652	13,363	4,266	15,997	1,224	6,644	43,262
Female	76,535	20,977	22,485	6,000	7,750	26,932	2,364	4,600	29,598
Total	153,094	35,261	37,137	19,363	12,016	42,929	3,588	11,244	72,860
Age²									
0 to 11 years	24,933	8,411	3,867	8,864	49	1,284	4	43	10,753
12 to 14 years	13,053	3,472	3,032	4,029	109	2,811	13	37	5,454
15 to 17 years	12,194	2,867	3,549	2,652	247	3,960	42	119	4,892
18 to 20 years	7,483	1,494	2,126	1,048	439	2,689	170	265	3,585
21 to 24 years	8,918	1,734	2,262	674	789	2,813	286	572	4,810
25 to 29 years	13,264	2,615	3,076	655	1,239	3,806	427	1,039	7,743
30 to 34 years	12,846	2,601	2,989	442	1,358	3,507	408	1,095	7,726
35 to 39 years	12,203	2,577	2,829	313	1,305	3,454	382	1,237	7,242
40 to 44 years	10,553	2,337	2,624	227	1,315	3,338	363	1,167	5,764
45 to 49 years	9,561	2,219	2,549	183	1,300	3,424	336	1,197	4,498
50 to 54 years	9,144	1,941	2,630	132	1,296	3,757	374	1,243	3,791
55 to 59 years	8,337	1,613	2,448	97	1,211	3,493	324	1,241	3,184
60 to 64 years	5,686	898	1,702	62	731	2,486	254	1,014	1,956
65 years and older	5,673	686	1,599	30	668	2,334	216	1,003	1,838
Total	153,848	35,465	37,282	19,408	12,056	43,156	3,599	11,272	73,236
Race									
American Indian or Alaska Native	282	80	56	25	26	68	11	29	124
Asian ³	397	74	71	30	17	109	10	69	173
Black or African American	14,674	3,427	2,016	2,020	965	3,199	302	1,933	7,048
Native Hawaiian or Other Pacific Islander	128	49	22	7	7	30	2	9	55
White	121,450	27,369	32,020	15,401	10,188	35,796	3,000	8,313	57,730
Other	14,695	3,956	2,864	1,823	724	3,477	227	728	6,943
Total	151,626	34,955	37,049	19,306	11,927	42,679	3,552	11,081	72,073

Continued. See notes at end of table.

Table 10.18a. Kentucky – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	146,056	33,682	35,898	18,653	11,496	41,237	3,120	10,357	69,468
Hispanic or Latino origin	3,853	1,096	850	597	182	1,009	52	175	1,505
Total	149,909	34,778	36,748	19,250	11,678	42,246	3,172	10,532	70,973
Specific Hispanic or Latino origin									
Mexican	1,471	422	287	271	43	355	8	46	572
Puerto Rican	475	139	109	78	42	122	13	30	182
Cuban	298	68	72	27	13	102	2	27	99
Other specific	1,578	460	377	221	80	424	28	65	640
Total	3,822	1,089	845	597	178	1,003	51	168	1,493
Veteran status⁴									
Veteran	2,375	656	567	57	231	787	53	207	1,080
Not a veteran	97,770	19,549	25,967	3,788	11,108	33,494	3,029	10,151	48,941
Total	100,145	20,205	26,534	3,845	11,339	34,281	3,082	10,358	50,021

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.18b. Kentucky – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020

Selected characteristics		Mental health diagnosis¹										
		Trauma- and stressor-related disorders				Mood disorders		Personality disorders			Schizophrenia, other psychotic disorders	Other mental disorders¹
						Bipolar disorders	Depressive disorders					
Total												
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	
Gender												
Male	50.0	40.5	39.5	69.0	35.5	37.3	34.1	59.1	59.4			
Female	50.0	59.5	60.5	31.0	64.5	62.7	65.9	40.9	40.6			
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0		
Age²												
0 to 11 years	16.2	23.7	10.4	45.7	†	3.0	†	†	14.7			
12 to 14 years	8.5	9.8	8.1	20.8	†	6.5	†	†	7.4			
15 to 17 years	7.9	8.1	9.5	13.7	2.0	9.2	1.2	1.1	6.7			
18 to 20 years	4.9	4.2	5.7	5.4	3.6	6.2	4.7	2.4	4.9			
21 to 24 years	5.8	4.9	6.1	3.5	6.5	6.5	7.9	5.1	6.6			
25 to 29 years	8.6	7.4	8.3	3.4	10.3	8.8	11.9	9.2	10.6			
30 to 34 years	8.3	7.3	8.0	2.3	11.3	8.1	11.3	9.7	10.5			
35 to 39 years	7.9	7.3	7.6	1.6	10.8	8.0	10.6	11.0	9.9			
40 to 44 years	6.9	6.6	7.0	1.2	10.9	7.7	10.1	10.4	7.9			
45 to 49 years	6.2	6.3	6.8	†	10.8	7.9	9.3	10.6	6.1			
50 to 54 years	5.9	5.5	7.1	†	10.7	8.7	10.4	11.0	5.2			
55 to 59 years	5.4	4.5	6.6	†	10.0	8.1	9.0	11.0	4.3			
60 to 64 years	3.7	2.5	4.6	†	6.1	5.8	7.1	9.0	2.7			
65 years and older	3.7	1.9	4.3	†	5.5	5.4	6.0	8.9	2.5			
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0		
Race												
American Indian or Alaska Native	†	†	†	†	†	†	†	†	†			
Asian³	†	†	†	†	†	†	†	†	†			
Black or African American	9.7	9.8	5.4	10.5	8.1	7.5	8.5	17.4	9.8			
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	†	†	†			
White	80.1	78.3	86.4	79.8	85.4	83.9	84.5	75.0	80.1			
Other	9.7	11.3	7.7	9.4	6.1	8.1	6.4	6.6	9.6			
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0		

Continued. See notes at end of table.

Table 10.18b. Kentucky – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020 (continued)

Selected characteristics		Mental health diagnosis¹							
		Trauma- and stressor-related disorders			Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders¹
					Bipolar disorders	Depressive disorders			
Total									
Ethnicity									
Not of Hispanic or Latino origin	97.4	96.8	97.7	96.9	98.4	97.6	98.4	98.3	97.9
Hispanic or Latino origin	2.6	3.2	2.3	3.1	1.6	2.4	1.6	1.7	2.1
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	38.5	38.8	34.0	45.4	24.2	35.4	15.7	27.4	38.3
Puerto Rican	12.4	12.8	12.9	13.1	23.6	12.2	25.5	17.9	12.2
Cuban	7.8	6.2	8.5	4.5	7.3	10.2	3.9	16.1	6.6
Other specific	41.3	42.2	44.6	37.0	44.9	42.3	54.9	38.7	42.9
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁴									
Veteran	2.4	3.2	2.1	1.5	2.0	2.3	1.7	2.0	2.2
Not a veteran	97.6	96.8	97.9	98.5	98.0	97.7	98.3	98.0	97.8
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.19a. Louisiana – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	17,526	1,907	3,904	1,879	3,228	6,656	288	3,934	2,174
Gender									
Male	7,776	701	1,230	1,202	1,056	2,171	87	2,441	1,305
Female	9,695	1,202	2,666	667	2,165	4,467	199	1,475	859
Total	17,471	1,903	3,896	1,869	3,221	6,638	286	3,916	2,164
Age²									
0 to 11 years	1,224	239	165	788	2	70	- -	6	487
12 to 14 years	852	152	130	435	20	223	3	10	278
15 to 17 years	885	127	211	309	46	356	2	24	246
18 to 20 years	660	104	216	97	97	295	16	68	110
21 to 24 years	919	95	281	54	196	403	21	178	108
25 to 29 years	1,510	159	441	35	366	590	41	358	145
30 to 34 years	1,640	186	418	48	370	608	35	464	141
35 to 39 years	1,615	167	389	32	352	612	38	462	121
40 to 44 years	1,437	143	352	21	375	558	24	398	108
45 to 49 years	1,342	132	302	22	342	543	30	355	101
50 to 54 years	1,566	162	318	14	366	683	29	399	80
55 to 59 years	1,700	135	323	10	334	765	21	491	112
60 to 64 years	1,261	66	221	8	214	564	21	403	75
65 years and older	915	40	137	6	148	386	7	318	62
Total	17,526	1,907	3,904	1,879	3,228	6,656	288	3,934	2,174
Race									
American Indian or Alaska Native	150	13	26	11	31	65	1	20	14
Asian ³	69	6	16	3	6	32	1	19	9
Black or African American	7,509	781	1,243	969	930	2,414	67	2,518	1,149
Native Hawaiian or Other Pacific Islander	36	4	10	3	9	14	1	1	2
White	8,207	885	2,254	707	1,920	3,535	172	1,088	806
Other	316	45	74	44	63	121	14	52	31
Total	16,287	1,734	3,623	1,737	2,959	6,181	256	3,698	2,011

Continued. See notes at end of table.

Table 10.19a. Louisiana – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	15,990	1,704	3,556	1,694	2,921	6,116	251	3,610	1,970
Hispanic or Latino origin	491	84	116	53	83	190	10	67	66
Total	16,481	1,788	3,672	1,747	3,004	6,306	261	3,677	2,036
Specific Hispanic or Latino origin									
Mexican	17	2	5	2	4	5	--	2	2
Puerto Rican	10	3	4	--	1	5	--	--	3
Cuban	6	1	--	--	1	3	--	1	1
Other specific	14	5	3	2	2	3	--	4	4
Total	47	11	12	4	8	16	--	7	10
Veteran status⁴									
Veteran	1,134	190	364	36	301	484	35	206	67
Not a veteran	11,532	1,031	2,595	244	2,421	4,754	208	3,200	929
Total	12,666	1,221	2,959	280	2,722	5,238	243	3,406	996

-- Quantity is zero.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.19b. Louisiana – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020

Selected characteristics		Mental health diagnosis¹							
		Trauma- and stressor-related disorders			Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders¹
					Bipolar disorders	Depressive disorders			
Total									
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender									
Male	44.5	36.8	31.6	64.3	32.8	32.7	30.4	62.3	60.3
Female	55.5	63.2	68.4	35.7	67.2	67.3	69.6	37.7	39.7
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age²									
0 to 11 years	7.0	12.5	4.2	41.9	†	1.1	--	†	22.4
12 to 14 years	4.9	8.0	3.3	23.2	†	3.4	1.0	†	12.8
15 to 17 years	5.0	6.7	5.4	16.4	1.4	5.3	†	†	11.3
18 to 20 years	3.8	5.5	5.5	5.2	3.0	4.4	5.6	1.7	5.1
21 to 24 years	5.2	5.0	7.2	2.9	6.1	6.1	7.3	4.5	5.0
25 to 29 years	8.6	8.3	11.3	1.9	11.3	8.9	14.2	9.1	6.7
30 to 34 years	9.4	9.8	10.7	2.6	11.5	9.1	12.2	11.8	6.5
35 to 39 years	9.2	8.8	10.0	1.7	10.9	9.2	13.2	11.7	5.6
40 to 44 years	8.2	7.5	9.0	1.1	11.6	8.4	8.3	10.1	5.0
45 to 49 years	7.7	6.9	7.7	1.2	10.6	8.2	10.4	9.0	4.6
50 to 54 years	8.9	8.5	8.1	†	11.3	10.3	10.1	10.1	3.7
55 to 59 years	9.7	7.1	8.3	†	10.3	11.5	7.3	12.5	5.2
60 to 64 years	7.2	3.5	5.7	†	6.6	8.5	7.3	10.2	3.4
65 years and older	5.2	2.1	3.5	†	4.6	5.8	2.4	8.1	2.9
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race									
American Indian or Alaska Native	†	†	†	†	1.0	1.1	†	†	†
Asian³	†	†	†	†	†	†	†	†	†
Black or African American	46.1	45.0	34.3	55.8	31.4	39.1	26.2	68.1	57.1
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	†	†	†
White	50.4	51.0	62.2	40.7	64.9	57.2	67.2	29.4	40.1
Other	1.9	2.6	2.0	2.5	2.1	2.0	5.5	1.4	1.5
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.19b. Louisiana – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020 (continued)

Selected characteristics		Mental health diagnosis ¹								
		Trauma- and stressor-related disorders			Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹	
					Bipolar disorders	Depressive disorders				
Total			Anxiety disorders	ADHD						
Ethnicity										
Not of Hispanic or Latino origin	97.0		95.3	96.8	97.0	97.2	97.0	96.2	98.2	96.8
Hispanic or Latino origin	3.0		4.7	3.2	3.0	2.8	3.0	3.8	1.8	3.2
Total	100.0		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin										
Mexican	36.2		18.2	41.7	50.0	50.0	31.3	--	28.6	20.0
Puerto Rican	21.3		27.3	33.3	--	12.5	31.3	--	--	30.0
Cuban	12.8		9.1	--	--	12.5	18.8	--	14.3	10.0
Other specific	29.8		45.5	25.0	50.0	25.0	18.8	--	57.1	40.0
Total	100.0		100.0	100.0	100.0	100.0	100.0	--	100.0	100.0
Veteran status ⁴										
Veteran	9.0		15.6	12.3	12.9	11.1	9.2	14.4	6.0	6.7
Not a veteran	91.0		84.4	87.7	87.1	88.9	90.8	85.6	94.0	93.3
Total	100.0		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

-- Quantity is zero; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.20a. Massachusetts – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	6,898	1,138	1,413	481	1,501	1,644	206	3,156	939
Gender									
Male	3,756	391	604	286	699	631	56	2,120	486
Female	3,142	747	809	195	802	1,013	150	1,036	453
Total	6,898	1,138	1,413	481	1,501	1,644	206	3,156	939
Age²									
0 to 11 years	148	39	61	68	5	7	--	--	59
12 to 14 years	181	44	62	92	7	37	--	6	71
15 to 17 years	290	66	108	83	40	122	3	24	103
18 to 20 years	394	91	123	62	75	168	13	83	115
21 to 24 years	620	116	170	64	145	167	21	250	140
25 to 29 years	753	93	115	30	203	130	30	420	95
30 to 34 years	631	87	119	23	153	124	19	337	47
35 to 39 years	568	84	92	19	139	106	23	311	57
40 to 44 years	519	105	98	10	118	99	12	270	41
45 to 49 years	495	84	79	9	124	121	14	245	36
50 to 54 years	575	105	123	3	141	173	23	250	41
55 to 59 years	703	105	127	9	145	185	27	361	59
60 to 64 years	558	78	80	6	120	123	12	311	36
65 years and older	463	41	56	3	86	82	9	288	39
Total	6,898	1,138	1,413	481	1,501	1,644	206	3,156	939
Race									
American Indian or Alaska Native	13	2	2	2	6	1	--	7	1
Asian ³	162	21	24	10	21	46	3	96	21
Black or African American	1,054	141	105	49	148	118	17	728	93
Native Hawaiian or Other Pacific Islander	3	--	--	--	--	1	1	1	2
White	4,200	734	914	289	1,015	1,015	157	1,863	594
Other	134	30	33	19	22	34	5	44	35
Total	5,566	928	1,078	369	1,212	1,215	183	2,739	746

Continued. See notes at end of table.

Table 10.20a. Massachusetts – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	4,755	764	962	313	1,087	1,061	165	2,281	641
Hispanic or Latino origin	629	138	100	38	118	154	16	314	88
Total	5,384	902	1,062	351	1,205	1,215	181	2,595	729
Specific Hispanic or Latino origin									
Mexican	7	2	2	1	2	2	--	1	2
Puerto Rican	241	64	35	13	40	51	7	122	39
Cuban	--	--	--	--	--	--	--	--	--
Other specific	110	18	18	6	20	29	2	64	16
Total	358	84	55	20	62	82	9	187	57
Veteran status⁴									
Veteran	130	16	22	4	29	24	--	78	8
Not a veteran	5,593	898	1,060	217	1,301	1,268	192	2,859	657
Total	5,723	914	1,082	221	1,330	1,292	192	2,937	665

-- Quantity is zero.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.20b. Massachusetts – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020

Selected characteristics		Mental health diagnosis¹										
		Trauma- and stressor-related disorders				Mood disorders		Personality disorders			Schizophrenia, other psychotic disorders	Other mental disorders¹
						Bipolar disorders	Depressive disorders					
Total												
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	
Gender												
Male	54.5	34.4	42.7	59.5	46.6	38.4	27.2	67.2	51.8			
Female	45.5	65.6	57.3	40.5	53.4	61.6	72.8	32.8	48.2			
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	
Age²												
0 to 11 years	2.1	3.4	4.3	14.1	†	†	--	--	6.3			
12 to 14 years	2.6	3.9	4.4	19.1	†	2.3	--	†	7.6			
15 to 17 years	4.2	5.8	7.6	17.3	2.7	7.4	1.5	†	11.0			
18 to 20 years	5.7	8.0	8.7	12.9	5.0	10.2	6.3	2.6	12.2			
21 to 24 years	9.0	10.2	12.0	13.3	9.7	10.2	10.2	7.9	14.9			
25 to 29 years	10.9	8.2	8.1	6.2	13.5	7.9	14.6	13.3	10.1			
30 to 34 years	9.1	7.6	8.4	4.8	10.2	7.5	9.2	10.7	5.0			
35 to 39 years	8.2	7.4	6.5	4.0	9.3	6.4	11.2	9.9	6.1			
40 to 44 years	7.5	9.2	6.9	2.1	7.9	6.0	5.8	8.6	4.4			
45 to 49 years	7.2	7.4	5.6	1.9	8.3	7.4	6.8	7.8	3.8			
50 to 54 years	8.3	9.2	8.7	†	9.4	10.5	11.2	7.9	4.4			
55 to 59 years	10.2	9.2	9.0	1.9	9.7	11.3	13.1	11.4	6.3			
60 to 64 years	8.1	6.9	5.7	1.2	8.0	7.5	5.8	9.9	3.8			
65 years and older	6.7	3.6	4.0	†	5.7	5.0	4.4	9.1	4.2			
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	
Race												
American Indian or Alaska Native	†	†	†	†	†	†	--	†	†			
Asian³	2.9	2.3	2.2	2.7	1.7	3.8	1.6	3.5	2.8			
Black or African American	18.9	15.2	9.7	13.3	12.2	9.7	9.3	26.6	12.5			
Native Hawaiian or Other Pacific Islander	†	--	--	--	--	†	†	†	†			
White	75.5	79.1	84.8	78.3	83.7	83.5	85.8	68.0	79.6			
Other	2.4	3.2	3.1	5.1	1.8	2.8	2.7	1.6	4.7			
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	

Continued. See notes at end of table.

Table 10.20b. Massachusetts – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020 (continued)

Mental health diagnosis ¹										
Selected characteristics		Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Ethnicity										
Not of Hispanic or Latino origin	88.3	84.7	90.6	89.2	90.2	87.3	91.2	87.9	87.9	
Hispanic or Latino origin	11.7	15.3	9.4	10.8	9.8	12.7	8.8	12.1	12.1	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	
Specific Hispanic or Latino origin										
Mexican	2.0	2.4	3.6	5.0	3.2	2.4	--	†	3.5	
Puerto Rican	67.3	76.2	63.6	65.0	64.5	62.2	77.8	65.2	68.4	
Cuban	--	--	--	--	--	--	--	--	--	
Other specific	30.7	21.4	32.7	30.0	32.3	35.4	22.2	34.2	28.1	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	
Veteran status ⁴										
Veteran	2.3	1.8	2.0	1.8	2.2	1.9	--	2.7	1.2	
Not a veteran	97.7	98.2	98.0	98.2	97.8	98.1	100.0	97.3	98.8	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	

-- Quantity is zero; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.21a. Michigan – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	207,770	40,535	45,722	28,347	32,606	61,284	15,136	34,612	90,889
Gender									
Male	107,851	16,854	18,923	19,226	13,793	24,507	5,632	21,043	54,886
Female	99,919	23,681	26,799	9,121	18,813	36,777	9,504	13,569	36,003
Total	207,770	40,535	45,722	28,347	32,606	61,284	15,136	34,612	90,889
Age²									
0 to 11 years	27,755	8,192	2,669	9,883	137	1,343	8	102	19,114
12 to 14 years	12,065	3,156	2,051	4,755	306	3,027	58	146	7,164
15 to 17 years	12,574	3,024	2,802	3,625	699	4,504	178	256	6,851
18 to 20 years	9,323	1,806	2,215	1,803	1,151	3,294	551	734	4,551
21 to 24 years	12,480	2,132	3,031	1,656	2,121	4,181	1,109	1,596	5,765
25 to 29 years	19,206	3,480	4,638	1,811	3,777	6,302	1,772	3,076	7,975
30 to 34 years	17,513	3,374	4,544	1,353	3,889	5,636	1,719	3,116	6,700
35 to 39 years	15,931	3,161	4,290	1,039	3,771	5,288	1,771	3,215	5,430
40 to 44 years	14,026	2,721	3,753	677	3,370	4,870	1,593	3,073	4,528
45 to 49 years	14,545	2,794	3,900	578	3,478	5,290	1,666	3,362	4,523
50 to 54 years	15,234	2,517	3,782	455	3,386	5,469	1,606	3,961	4,872
55 to 59 years	15,453	2,149	3,647	348	3,085	5,451	1,528	4,554	5,007
60 to 64 years	11,214	1,220	2,432	233	1,938	3,767	967	3,751	3,792
65 years and older	10,451	809	1,968	131	1,498	2,862	610	3,670	4,617
Total	207,770	40,535	45,722	28,347	32,606	61,284	15,136	34,612	90,889
Race									
American Indian or Alaska Native	1,224	337	277	140	189	403	109	205	422
Asian ³	1,045	119	201	85	110	223	40	213	543
Black or African American	48,321	8,286	7,459	6,232	6,507	13,786	2,421	12,322	19,815
Native Hawaiian or Other Pacific Islander	130	24	30	11	18	44	10	24	44
White	124,839	25,828	31,248	16,447	21,105	38,033	10,767	16,198	55,043
Other	31,044	5,691	6,311	5,366	4,563	8,418	1,730	5,488	14,637
Total	206,603	40,285	45,526	28,281	32,492	60,907	15,077	34,450	90,504

Continued. See notes at end of table.

Table 10.21a. Michigan – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	178,864	34,880	40,027	24,676	28,989	53,394	13,494	30,896	77,218
Hispanic or Latino origin	12,631	2,860	2,782	2,046	1,613	3,909	698	1,380	5,383
Total	191,495	37,740	42,809	26,722	30,602	57,303	14,192	32,276	82,601
Specific Hispanic or Latino origin									
Mexican	3,355	852	763	546	302	1,020	131	231	1,460
Puerto Rican	1,042	245	238	206	118	336	54	111	417
Cuban	139	27	27	16	15	48	5	24	60
Other specific	2,239	529	488	317	289	738	119	222	851
Total	6,775	1,653	1,516	1,085	724	2,142	309	588	2,788
Veteran status⁴									
Veteran	3,556	872	814	118	701	1,320	345	910	762
Not a veteran	142,662	24,333	35,870	9,549	29,566	48,831	14,007	31,483	52,090
Total	146,218	25,205	36,684	9,667	30,267	50,151	14,352	32,393	52,852

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.21b. Michigan – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020

Selected characteristics		Mental health diagnosis¹								
		Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders¹
						Bipolar disorders	Depressive disorders			
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender										
Male	51.9	41.6	41.4	67.8	42.3	40.0	37.2	60.8	60.4	
Female	48.1	58.4	58.6	32.2	57.7	60.0	62.8	39.2	39.6	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age²										
0 to 11 years	13.4	20.2	5.8	34.9	†	2.2	†	†	21.0	
12 to 14 years	5.8	7.8	4.5	16.8	†	4.9	†	†	7.9	
15 to 17 years	6.1	7.5	6.1	12.8	2.1	7.3	1.2	†	7.5	
18 to 20 years	4.5	4.5	4.8	6.4	3.5	5.4	3.6	2.1	5.0	
21 to 24 years	6.0	5.3	6.6	5.8	6.5	6.8	7.3	4.6	6.3	
25 to 29 years	9.2	8.6	10.1	6.4	11.6	10.3	11.7	8.9	8.8	
30 to 34 years	8.4	8.3	9.9	4.8	11.9	9.2	11.4	9.0	7.4	
35 to 39 years	7.7	7.8	9.4	3.7	11.6	8.6	11.7	9.3	6.0	
40 to 44 years	6.8	6.7	8.2	2.4	10.3	7.9	10.5	8.9	5.0	
45 to 49 years	7.0	6.9	8.5	2.0	10.7	8.6	11.0	9.7	5.0	
50 to 54 years	7.3	6.2	8.3	1.6	10.4	8.9	10.6	11.4	5.4	
55 to 59 years	7.4	5.3	8.0	1.2	9.5	8.9	10.1	13.2	5.5	
60 to 64 years	5.4	3.0	5.3	†	5.9	6.1	6.4	10.8	4.2	
65 years and older	5.0	2.0	4.3	†	4.6	4.7	4.0	10.6	5.1	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race										
American Indian or Alaska Native	†	†	†	†	†	†	†	†	†	
Asian³	†	†	†	†	†	†	†	†	†	
Black or African American	23.4	20.6	16.4	22.0	20.0	22.6	16.1	35.8	21.9	
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	†	†	†	
White	60.4	64.1	68.6	58.2	65.0	62.4	71.4	47.0	60.8	
Other	15.0	14.1	13.9	19.0	14.0	13.8	11.5	15.9	16.2	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.21b. Michigan – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020 (continued)

Selected characteristics		Mental health diagnosis ¹							
					Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD					
Ethnicity									
Not of Hispanic or Latino origin	93.4	92.4	93.5	92.3	94.7	93.2	95.1	95.7	93.5
Hispanic or Latino origin	6.6	7.6	6.5	7.7	5.3	6.8	4.9	4.3	6.5
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	49.5	51.5	50.3	50.3	41.7	47.6	42.4	39.3	52.4
Puerto Rican	15.4	14.8	15.7	19.0	16.3	15.7	17.5	18.9	15.0
Cuban	2.1	1.6	1.8	1.5	2.1	2.2	1.6	4.1	2.2
Other specific	33.0	32.0	32.2	29.2	39.9	34.5	38.5	37.8	30.5
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁴									
Veteran	2.4	3.5	2.2	1.2	2.3	2.6	2.4	2.8	1.4
Not a veteran	97.6	96.5	97.8	98.8	97.7	97.4	97.6	97.2	98.6
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.22a. Minnesota – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	284,601	107,500	147,707	48,532	24,968	126,039	17,267	25,217	58,153
Gender									
Male	124,489	42,306	56,288	29,389	10,177	46,429	5,612	15,513	30,216
Female	160,112	65,194	91,419	19,143	14,791	79,610	11,655	9,704	27,937
Total	284,601	107,500	147,707	48,532	24,968	126,039	17,267	25,217	58,153
Age²									
0 to 11 years	39,478	21,388	15,276	15,280	69	3,589	15	30	11,960
12 to 14 years	19,371	8,879	9,224	7,119	119	6,311	27	62	5,942
15 to 17 years	18,771	7,479	10,017	5,507	342	8,607	112	201	5,828
18 to 20 years	13,024	4,504	7,757	2,917	651	6,583	587	489	3,149
21 to 24 years	15,212	5,171	9,200	2,433	1,357	7,617	1,323	1,221	2,834
25 to 29 years	26,463	9,214	15,939	3,743	2,933	12,800	2,376	2,393	4,587
30 to 34 years	25,849	9,502	15,089	3,334	3,207	12,338	2,464	2,719	4,154
35 to 39 years	23,729	8,874	13,676	2,595	3,040	11,397	2,335	2,772	3,628
40 to 44 years	18,960	7,051	10,822	1,685	2,590	9,590	1,823	2,260	2,938
45 to 49 years	17,352	6,211	9,691	1,337	2,526	9,200	1,589	2,201	2,700
50 to 54 years	17,480	5,855	9,052	939	2,477	9,879	1,517	2,439	2,821
55 to 59 years	18,298	5,525	9,169	884	2,462	10,539	1,385	2,948	2,945
60 to 64 years	13,869	3,703	6,501	510	1,634	8,065	900	2,465	2,255
65 years and older	16,744	4,143	6,294	249	1,561	9,524	814	3,017	2,412
Total	284,600	107,499	147,707	48,532	24,968	126,039	17,267	25,217	58,153
Race									
American Indian or Alaska Native	11,181	5,243	5,394	1,531	941	5,275	577	823	2,332
Asian ³	7,709	2,714	3,113	461	252	4,830	214	855	1,143
Black or African American	34,299	14,997	14,931	4,017	2,929	15,551	1,735	5,320	6,056
Native Hawaiian or Other Pacific Islander	309	131	150	47	24	132	24	35	49
White	169,279	57,127	92,230	29,480	17,491	76,032	12,504	16,068	35,357
Other	13,317	6,070	6,834	3,194	1,014	5,146	657	665	3,109
Total	236,094	86,282	122,652	38,730	22,651	106,966	15,711	23,766	48,046

Continued. See notes at end of table.

Table 10.22a. Minnesota – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	237,177	88,664	124,630	41,282	21,001	105,204	13,772	21,645	49,757
Hispanic or Latino origin	26,497	9,959	11,853	3,151	2,836	12,164	2,702	2,974	4,058
Total	263,674	98,623	136,483	44,433	23,837	117,368	16,474	24,619	53,815
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴									
Veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Not a veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡

‡ No data submitted.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.22b. Minnesota – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender									
Male	43.7	39.4	38.1	60.6	40.8	36.8	32.5	61.5	52.0
Female	56.3	60.6	61.9	39.4	59.2	63.2	67.5	38.5	48.0
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age²									
0 to 11 years	13.9	19.9	10.3	31.5	†	2.8	†	†	20.6
12 to 14 years	6.8	8.3	6.2	14.7	†	5.0	†	†	10.2
15 to 17 years	6.6	7.0	6.8	11.3	1.4	6.8	†	†	10.0
18 to 20 years	4.6	4.2	5.3	6.0	2.6	5.2	3.4	1.9	5.4
21 to 24 years	5.3	4.8	6.2	5.0	5.4	6.0	7.7	4.8	4.9
25 to 29 years	9.3	8.6	10.8	7.7	11.7	10.2	13.8	9.5	7.9
30 to 34 years	9.1	8.8	10.2	6.9	12.8	9.8	14.3	10.8	7.1
35 to 39 years	8.3	8.3	9.3	5.3	12.2	9.0	13.5	11.0	6.2
40 to 44 years	6.7	6.6	7.3	3.5	10.4	7.6	10.6	9.0	5.1
45 to 49 years	6.1	5.8	6.6	2.8	10.1	7.3	9.2	8.7	4.6
50 to 54 years	6.1	5.4	6.1	1.9	9.9	7.8	8.8	9.7	4.9
55 to 59 years	6.4	5.1	6.2	1.8	9.9	8.4	8.0	11.7	5.1
60 to 64 years	4.9	3.4	4.4	1.1	6.5	6.4	5.2	9.8	3.9
65 years and older	5.9	3.9	4.3	†	6.3	7.6	4.7	12.0	4.1
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race									
American Indian or Alaska Native	4.7	6.1	4.4	4.0	4.2	4.9	3.7	3.5	4.9
Asian ³	3.3	3.1	2.5	1.2	1.1	4.5	1.4	3.6	2.4
Black or African American	14.5	17.4	12.2	10.4	12.9	14.5	11.0	22.4	12.6
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	†	†	†
White	71.7	66.2	75.2	76.1	77.2	71.1	79.6	67.6	73.6
Other	5.6	7.0	5.6	8.2	4.5	4.8	4.2	2.8	6.5
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.22b. Minnesota – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	90.0	89.9	91.3	92.9	88.1	89.6	83.6	87.9	92.5
Hispanic or Latino origin	10.0	10.1	8.7	7.1	11.9	10.4	16.4	12.1	7.5
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴									
Veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Not a veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡

‡ No data submitted; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.23a. Mississippi – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	101,461	14,273	17,736	20,989	12,179	29,870	2,231	14,574	38,370
Gender									
Male	48,844	5,674	5,587	13,923	4,556	9,428	725	8,633	23,014
Female	52,573	8,587	12,141	7,054	7,621	20,436	1,505	5,938	15,342
Total	101,417	14,261	17,728	20,977	12,177	29,864	2,230	14,571	38,356
Age²									
0 to 11 years	15,510	3,571	746	9,237	438	458	2	41	6,867
12 to 14 years	9,551	1,869	701	5,108	477	1,115	5	54	4,436
15 to 17 years	8,172	1,528	845	3,206	578	1,808	19	94	3,660
18 to 20 years	5,208	778	814	1,423	643	1,544	69	243	2,192
21 to 24 years	4,907	567	1,016	552	758	1,685	145	668	2,024
25 to 29 years	7,435	811	1,681	413	1,220	2,596	223	1,337	3,205
30 to 34 years	7,416	827	1,716	281	1,248	2,558	263	1,462	3,215
35 to 39 years	7,441	828	1,728	238	1,235	2,590	233	1,684	3,030
40 to 44 years	6,465	773	1,523	152	1,071	2,475	254	1,516	2,342
45 to 49 years	5,957	745	1,449	117	1,112	2,471	227	1,279	1,855
50 to 54 years	6,188	676	1,466	108	1,034	2,691	234	1,531	1,778
55 to 59 years	6,560	582	1,589	87	995	2,949	231	1,730	1,717
60 to 64 years	5,114	402	1,168	44	682	2,381	176	1,417	1,054
65 years and older	5,536	316	1,294	23	688	2,548	150	1,518	995
Total	101,460	14,273	17,736	20,989	12,179	29,869	2,231	14,574	38,370
Race									
American Indian or Alaska Native	225	46	34	38	33	68	6	27	77
Asian ³	168	13	36	19	18	51	5	43	44
Black or African American	50,687	5,629	5,285	11,968	4,258	13,600	801	10,187	20,643
Native Hawaiian or Other Pacific Islander	21	4	4	2	3	10	1	3	5
White	47,347	7,879	11,928	8,084	7,626	15,501	1,380	4,148	16,456
Other	1,734	404	262	597	140	370	9	87	609
Total	100,182	13,975	17,549	20,708	12,078	29,600	2,202	14,495	37,834

Continued. See notes at end of table.

Table 10.23a. Mississippi – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	94,465	12,895	16,560	19,341	11,675	28,043	2,133	14,140	35,600
Hispanic or Latino origin	1,811	389	300	437	135	479	21	119	689
Total	96,276	13,284	16,860	19,778	11,810	28,522	2,154	14,259	36,289
Specific Hispanic or Latino origin									
Mexican	707	175	132	165	42	197	6	40	236
Puerto Rican	115	23	29	38	14	31	-	10	32
Cuban	44	4	6	9	8	10	3	12	11
Other specific	209	25	27	65	22	52	4	19	102
Total	1,075	227	194	277	86	290	13	81	381
Veteran status⁴									
Veteran	1,456	293	282	38	190	541	40	320	476
Not a veteran	64,457	6,852	14,794	3,329	10,192	25,322	2,097	13,333	22,232
Total	65,913	7,145	15,076	3,367	10,382	25,863	2,137	13,653	22,708

- - Quantity is zero.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.23b. Mississippi – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender									
Male	48.2	39.8	31.5	66.4	37.4	31.6	32.5	59.2	60.0
Female	51.8	60.2	68.5	33.6	62.6	68.4	67.5	40.8	40.0
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age²									
0 to 11 years	15.3	25.0	4.2	44.0	3.6	1.5	†	†	17.9
12 to 14 years	9.4	13.1	4.0	24.3	3.9	3.7	†	†	11.6
15 to 17 years	8.1	10.7	4.8	15.3	4.7	6.1	†	†	9.5
18 to 20 years	5.1	5.5	4.6	6.8	5.3	5.2	3.1	1.7	5.7
21 to 24 years	4.8	4.0	5.7	2.6	6.2	5.6	6.5	4.6	5.3
25 to 29 years	7.3	5.7	9.5	2.0	10.0	8.7	10.0	9.2	8.4
30 to 34 years	7.3	5.8	9.7	1.3	10.2	8.6	11.8	10.0	8.4
35 to 39 years	7.3	5.8	9.7	1.1	10.1	8.7	10.4	11.6	7.9
40 to 44 years	6.4	5.4	8.6	†	8.8	8.3	11.4	10.4	6.1
45 to 49 years	5.9	5.2	8.2	†	9.1	8.3	10.2	8.8	4.8
50 to 54 years	6.1	4.7	8.3	†	8.5	9.0	10.5	10.5	4.6
55 to 59 years	6.5	4.1	9.0	†	8.2	9.9	10.4	11.9	4.5
60 to 64 years	5.0	2.8	6.6	†	5.6	8.0	7.9	9.7	2.7
65 years and older	5.5	2.2	7.3	†	5.6	8.5	6.7	10.4	2.6
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race									
American Indian or Alaska Native	†	†	†	†	†	†	†	†	†
Asian ³	†	†	†	†	†	†	†	†	†
Black or African American	50.6	40.3	30.1	57.8	35.3	45.9	36.4	70.3	54.6
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	†	†	†
White	47.3	56.4	68.0	39.0	63.1	52.4	62.7	28.6	43.5
Other	1.7	2.9	1.5	2.9	1.2	1.3	†	†	1.6
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.23b. Mississippi – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020 (continued)

Mental health diagnosis ¹										
Selected characteristics		Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Ethnicity										
Not of Hispanic or Latino origin	98.1		97.1	98.2	97.8	98.9	98.3	99.0	99.2	98.1
Hispanic or Latino origin	1.9		2.9	1.8	2.2	1.1	1.7	†	†	1.9
Total	100.0		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin										
Mexican	65.8		77.1	68.0	59.6	48.8	67.9	46.2	49.4	61.9
Puerto Rican	10.7		10.1	14.9	13.7	16.3	10.7	--	12.3	8.4
Cuban	4.1		1.8	3.1	3.2	9.3	3.4	23.1	14.8	2.9
Other specific	19.4		11.0	13.9	23.5	25.6	17.9	30.8	23.5	26.8
Total	100.0		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁴										
Veteran	2.2		4.1	1.9	1.1	1.8	2.1	1.9	2.3	2.1
Not a veteran	97.8		95.9	98.1	98.9	98.2	97.9	98.1	97.7	97.9
Total	100.0		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

- - Quantity is zero; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.24a. Missouri – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	75,768	6,638	10,943	7,338	13,219	25,147	1,111	13,252	5,827
Gender									
Male	36,002	2,347	4,475	5,352	4,934	9,465	339	8,510	3,737
Female	39,764	4,291	6,468	1,986	8,285	15,680	772	4,742	2,090
Total	75,766	6,638	10,943	7,338	13,219	25,145	1,111	13,252	5,827
Age²									
0 to 11 years	8,510	883	1,331	3,790	79	655	- -	16	2,428
12 to 14 years	5,462	570	923	1,800	152	1,342	12	43	1,218
15 to 17 years	4,574	494	718	1,017	302	1,664	18	77	825
18 to 20 years	2,744	271	489	319	463	1,102	57	223	218
21 to 24 years	3,482	319	572	135	707	1,308	101	616	128
25 to 29 years	5,585	546	920	85	1,285	1,818	146	1,231	168
30 to 34 years	6,053	572	917	59	1,447	2,018	149	1,406	161
35 to 39 years	6,285	606	943	47	1,525	2,077	157	1,478	127
40 to 44 years	5,896	567	822	28	1,402	2,083	121	1,435	112
45 to 49 years	6,013	514	842	20	1,450	2,248	113	1,367	99
50 to 54 years	6,349	484	861	17	1,396	2,587	85	1,457	111
55 to 59 years	6,853	444	791	10	1,440	2,910	77	1,688	105
60 to 64 years	4,757	235	533	5	962	2,012	44	1,279	73
65 years and older	3,205	133	281	6	609	1,323	31	936	54
Total	75,768	6,638	10,943	7,338	13,219	25,147	1,111	13,252	5,827
Race									
American Indian or Alaska Native	150	13	28	10	27	49	3	23	11
Asian ³	202	18	24	5	28	59	1	76	5
Black or African American	16,292	1,074	1,278	1,944	2,204	4,744	179	5,104	1,215
Native Hawaiian or Other Pacific Islander	44	6	10	2	5	15	2	6	4
White	54,114	4,994	8,845	4,656	10,293	18,865	858	7,242	3,972
Other	3,003	324	444	419	465	816	46	558	378
Total	73,805	6,429	10,629	7,036	13,022	24,548	1,089	13,009	5,585

Continued. See notes at end of table.

Table 10.24a. Missouri – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	73,807	6,431	10,606	7,095	12,986	24,509	1,082	12,998	5,619
Hispanic or Latino origin	1,959	207	337	243	233	638	28	253	208
Total	75,766	6,638	10,943	7,338	13,219	25,147	1,110	13,251	5,827
Specific Hispanic or Latino origin									
Mexican	959	109	156	128	122	320	16	127	93
Puerto Rican	170	16	25	12	27	59	6	31	13
Cuban	80	7	14	4	16	22	- -	12	5
Other specific	741	74	141	98	66	233	6	80	95
Total	1,950	206	336	242	231	634	28	250	206
Veteran status⁴									
Veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Not a veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡

- - Quantity is zero; ‡ No data submitted.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.24b. Missouri – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020

Selected characteristics		Mental health diagnosis¹							
		Trauma- and stressor-related disorders			Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders¹
					Bipolar disorders	Depressive disorders			
Total									
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender									
Male	47.5	35.4	40.9	72.9	37.3	37.6	30.5	64.2	64.1
Female	52.5	64.6	59.1	27.1	62.7	62.4	69.5	35.8	35.9
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age²									
0 to 11 years	11.2	13.3	12.2	51.6	†	2.6	--	†	41.7
12 to 14 years	7.2	8.6	8.4	24.5	1.1	5.3	1.1	†	20.9
15 to 17 years	6.0	7.4	6.6	13.9	2.3	6.6	1.6	†	14.2
18 to 20 years	3.6	4.1	4.5	4.3	3.5	4.4	5.1	1.7	3.7
21 to 24 years	4.6	4.8	5.2	1.8	5.3	5.2	9.1	4.6	2.2
25 to 29 years	7.4	8.2	8.4	1.2	9.7	7.2	13.1	9.3	2.9
30 to 34 years	8.0	8.6	8.4	†	10.9	8.0	13.4	10.6	2.8
35 to 39 years	8.3	9.1	8.6	†	11.5	8.3	14.1	11.2	2.2
40 to 44 years	7.8	8.5	7.5	†	10.6	8.3	10.9	10.8	1.9
45 to 49 years	7.9	7.7	7.7	†	11.0	8.9	10.2	10.3	1.7
50 to 54 years	8.4	7.3	7.9	†	10.6	10.3	7.7	11.0	1.9
55 to 59 years	9.0	6.7	7.2	†	10.9	11.6	6.9	12.7	1.8
60 to 64 years	6.3	3.5	4.9	†	7.3	8.0	4.0	9.7	1.3
65 years and older	4.2	2.0	2.6	†	4.6	5.3	2.8	7.1	†
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race									
American Indian or Alaska Native	†	†	†	†	†	†	†	†	†
Asian³	†	†	†	†	†	†	†	†	†
Black or African American	22.1	16.7	12.0	27.6	16.9	19.3	16.4	39.2	21.8
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	†	†	†
White	73.3	77.7	83.2	66.2	79.0	76.8	78.8	55.7	71.1
Other	4.1	5.0	4.2	6.0	3.6	3.3	4.2	4.3	6.8
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.24b. Missouri – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	97.4	96.9	96.9	96.7	98.2	97.5	97.5	98.1	96.4
Hispanic or Latino origin	2.6	3.1	3.1	3.3	1.8	2.5	2.5	1.9	3.6
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	49.2	52.9	46.4	52.9	52.8	50.5	57.1	50.8	45.1
Puerto Rican	8.7	7.8	7.4	5.0	11.7	9.3	21.4	12.4	6.3
Cuban	4.1	3.4	4.2	1.7	6.9	3.5	- -	4.8	2.4
Other specific	38.0	35.9	42.0	40.5	28.6	36.8	21.4	32.0	46.1
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁴									
Veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Not a veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡

- - Quantity is zero; ‡ No data submitted; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.25a. Montana – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	73,240	14,170	29,733	7,241	6,423	23,201	860	4,003	17,276
Gender									
Male	30,438	5,237	10,295	4,619	2,378	7,968	244	2,398	9,336
Female	42,802	8,933	19,438	2,622	4,045	15,233	616	1,605	7,940
Total	73,240	14,170	29,733	7,241	6,423	23,201	860	4,003	17,276
Age²									
0 to 11 years	12,047	3,251	4,659	3,083	61	749	- -	14	5,096
12 to 14 years	5,990	1,431	2,394	1,203	155	1,624	5	29	2,073
15 to 17 years	5,790	1,195	2,365	685	288	2,368	9	52	1,728
18 to 20 years	3,718	565	1,550	262	304	1,490	44	123	869
21 to 24 years	4,451	768	1,936	216	461	1,687	76	236	870
25 to 29 years	6,848	1,219	3,084	418	743	2,461	111	391	1,100
30 to 34 years	6,806	1,315	3,118	390	782	2,395	140	397	1,022
35 to 39 years	6,156	1,165	2,680	360	744	2,199	93	465	931
40 to 44 years	4,811	911	2,105	215	636	1,698	91	385	690
45 to 49 years	3,846	664	1,520	160	585	1,436	92	355	557
50 to 54 years	3,411	535	1,332	101	475	1,349	70	322	496
55 to 59 years	3,759	540	1,324	75	541	1,560	55	477	540
60 to 64 years	2,868	390	983	44	354	1,191	41	341	445
65 years and older	2,739	221	683	29	294	994	33	416	859
Total	73,240	14,170	29,733	7,241	6,423	23,201	860	4,003	17,276
Race									
American Indian or Alaska Native	8,666	2,139	3,609	612	531	2,931	70	510	2,021
Asian ³	228	54	82	13	18	64	2	16	61
Black or African American	764	188	281	83	56	205	9	64	199
Native Hawaiian or Other Pacific Islander	67	11	24	10	6	17	3	3	19
White	55,603	10,175	22,545	5,639	5,258	17,576	699	3,132	13,053
Other	1,793	394	723	229	137	505	22	88	492
Total	67,121	12,961	27,264	6,586	6,006	21,298	805	3,813	15,845

Continued. See notes at end of table.

Table 10.25a. Montana – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	70,380	13,556	28,524	6,962	6,203	22,363	836	3,842	16,569
Hispanic or Latino origin	2,855	614	1,208	279	219	838	24	159	704
Total	73,235	14,170	29,732	7,241	6,422	23,201	860	4,001	17,273
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴									
Veteran	159	29	41	5	34	42	4	52	32
Not a veteran	14,887	2,173	5,557	651	2,619	5,622	402	2,261	2,667
Total	15,046	2,202	5,598	656	2,653	5,664	406	2,313	2,699

- - Quantity is zero; ‡ No data submitted.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.25b. Montana – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020

		Mental health diagnosis ¹								
Selected characteristics		Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Total individuals		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender										
Male		41.6	37.0	34.6	63.8	37.0	34.3	28.4	59.9	54.0
Female		58.4	63.0	65.4	36.2	63.0	65.7	71.6	40.1	46.0
Total		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age ²										
0 to 11 years		16.4	22.9	15.7	42.6	†	3.2	--	†	29.5
12 to 14 years		8.2	10.1	8.1	16.6	2.4	7.0	†	†	12.0
15 to 17 years		7.9	8.4	8.0	9.5	4.5	10.2	1.0	1.3	10.0
18 to 20 years		5.1	4.0	5.2	3.6	4.7	6.4	5.1	3.1	5.0
21 to 24 years		6.1	5.4	6.5	3.0	7.2	7.3	8.8	5.9	5.0
25 to 29 years		9.4	8.6	10.4	5.8	11.6	10.6	12.9	9.8	6.4
30 to 34 years		9.3	9.3	10.5	5.4	12.2	10.3	16.3	9.9	5.9
35 to 39 years		8.4	8.2	9.0	5.0	11.6	9.5	10.8	11.6	5.4
40 to 44 years		6.6	6.4	7.1	3.0	9.9	7.3	10.6	9.6	4.0
45 to 49 years		5.3	4.7	5.1	2.2	9.1	6.2	10.7	8.9	3.2
50 to 54 years		4.7	3.8	4.5	1.4	7.4	5.8	8.1	8.0	2.9
55 to 59 years		5.1	3.8	4.5	1.0	8.4	6.7	6.4	11.9	3.1
60 to 64 years		3.9	2.8	3.3	†	5.5	5.1	4.8	8.5	2.6
65 years and older		3.7	1.6	2.3	†	4.6	4.3	3.8	10.4	5.0
Total		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race										
American Indian or Alaska Native		12.9	16.5	13.2	9.3	8.8	13.8	8.7	13.4	12.8
Asian ³		†	†	†	†	†	†	†	†	†
Black or African American		1.1	1.5	1.0	1.3	†	†	1.1	1.7	1.3
Native Hawaiian or Other Pacific Islander		†	†	†	†	†	†	†	†	†
White		82.8	78.5	82.7	85.6	87.5	82.5	86.8	82.1	82.4
Other		2.7	3.0	2.7	3.5	2.3	2.4	2.7	2.3	3.1
Total		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.25b. Montana – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	96.1	95.7	95.9	96.1	96.6	96.4	97.2	96.0	95.9
Hispanic or Latino origin	3.9	4.3	4.1	3.9	3.4	3.6	2.8	4.0	4.1
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴									
Veteran	1.1	1.3	†	†	1.3	†	†	2.2	1.2
Not a veteran	98.9	98.7	99.3	99.2	98.7	99.3	99.0	97.8	98.8
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

- - Quantity is zero; ‡ No data submitted; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.26a. Nebraska – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	21,981	4,265	4,709	1,203	3,828	7,126	626	2,671	7,011
Gender									
Male	11,335	1,801	1,960	758	1,706	3,102	272	1,803	4,217
Female	10,559	2,453	2,724	435	2,104	3,985	353	864	2,773
Total	21,894	4,254	4,684	1,193	3,810	7,087	625	2,667	6,990
Age²									
0 to 11 years	698	218	65	220	13	40	1	--	315
12 to 14 years	726	176	88	158	12	119	1	2	373
15 to 17 years	922	196	138	139	20	189	5	5	484
18 to 20 years	1,162	281	253	97	137	402	40	82	400
21 to 24 years	1,869	376	457	106	331	652	69	191	561
25 to 29 years	2,899	565	677	128	506	951	65	397	868
30 to 34 years	2,725	526	625	113	553	842	83	368	831
35 to 39 years	2,473	444	580	76	513	811	78	350	763
40 to 44 years	2,036	400	491	66	430	669	69	291	623
45 to 49 years	1,759	350	384	39	380	592	66	260	507
50 to 54 years	1,649	284	353	24	380	637	60	223	462
55 to 59 years	1,506	234	319	17	283	620	43	248	423
60 to 64 years	903	131	191	15	154	379	25	146	212
65 years and older	654	84	88	5	116	223	21	108	189
Total	21,981	4,265	4,709	1,203	3,828	7,126	626	2,671	7,011
Race									
American Indian or Alaska Native	574	159	106	24	76	176	14	51	203
Asian ³	172	30	20	13	20	38	7	40	56
Black or African American	2,053	295	275	79	263	483	39	471	779
Native Hawaiian or Other Pacific Islander	88	21	20	--	15	27	--	13	33
White	17,135	3,344	3,924	1,003	3,129	5,817	503	1,802	5,359
Other	1,679	361	287	69	273	458	51	268	539
Total	21,701	4,210	4,632	1,188	3,776	6,999	614	2,645	6,969

Continued. See notes at end of table.

Table 10.26a. Nebraska – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	16,981	3,472	3,991	1,010	3,100	5,854	489	1,970	5,237
Hispanic or Latino origin	2,148	530	410	106	267	667	42	235	718
Total	19,129	4,002	4,401	1,116	3,367	6,521	531	2,205	5,955
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴									
Veteran	308	57	47	6	65	77	10	68	78
Not a veteran	17,510	3,128	4,030	621	3,403	6,207	471	2,284	4,936
Total	17,818	3,185	4,077	627	3,468	6,284	481	2,352	5,014

- - Quantity is zero; ‡ No data submitted.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.26b. Nebraska – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020

		Mental health diagnosis ¹								
Selected characteristics		Total	Trauma- and stressor-related disorders			Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Total individuals		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender										
Male		51.8	42.3	41.8	63.5	44.8	43.8	43.5	67.6	60.3
Female		48.2	57.7	58.2	36.5	55.2	56.2	56.5	32.4	39.7
Total		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age ²										
0 to 11 years		3.2	5.1	1.4	18.3	†	†	†	--	4.5
12 to 14 years		3.3	4.1	1.9	13.1	†	1.7	†	†	5.3
15 to 17 years		4.2	4.6	2.9	11.6	†	2.7	†	†	6.9
18 to 20 years		5.3	6.6	5.4	8.1	3.6	5.6	6.4	3.1	5.7
21 to 24 years		8.5	8.8	9.7	8.8	8.6	9.1	11.0	7.2	8.0
25 to 29 years		13.2	13.2	14.4	10.6	13.2	13.3	10.4	14.9	12.4
30 to 34 years		12.4	12.3	13.3	9.4	14.4	11.8	13.3	13.8	11.9
35 to 39 years		11.3	10.4	12.3	6.3	13.4	11.4	12.5	13.1	10.9
40 to 44 years		9.3	9.4	10.4	5.5	11.2	9.4	11.0	10.9	8.9
45 to 49 years		8.0	8.2	8.2	3.2	9.9	8.3	10.5	9.7	7.2
50 to 54 years		7.5	6.7	7.5	2.0	9.9	8.9	9.6	8.3	6.6
55 to 59 years		6.9	5.5	6.8	1.4	7.4	8.7	6.9	9.3	6.0
60 to 64 years		4.1	3.1	4.1	1.2	4.0	5.3	4.0	5.5	3.0
65 years and older		3.0	2.0	1.9	†	3.0	3.1	3.4	4.0	2.7
Total		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race										
American Indian or Alaska Native		2.6	3.8	2.3	2.0	2.0	2.5	2.3	1.9	2.9
Asian ³		†	†	†	1.1	†	†	1.1	1.5	†
Black or African American		9.5	7.0	5.9	6.6	7.0	6.9	6.4	17.8	11.2
Native Hawaiian or Other Pacific Islander		†	†	†	--	†	†	--	†	†
White		79.0	79.4	84.7	84.4	82.9	83.1	81.9	68.1	76.9
Other		7.7	8.6	6.2	5.8	7.2	6.5	8.3	10.1	7.7
Total		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.26b. Nebraska – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020 (continued)

Mental health diagnosis ¹										
Selected characteristics		Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Ethnicity										
Not of Hispanic or Latino origin		88.8	86.8	90.7	90.5	92.1	89.8	92.1	89.3	87.9
Hispanic or Latino origin		11.2	13.2	9.3	9.5	7.9	10.2	7.9	10.7	12.1
Total		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin										
Mexican		‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican		‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban		‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific		‡	‡	‡	‡	‡	‡	‡	‡	‡
Total		‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status ⁴										
Veteran		1.7	1.8	1.2	†	1.9	1.2	2.1	2.9	1.6
Not a veteran		98.3	98.2	98.8	99.0	98.1	98.8	97.9	97.1	98.4
Total		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

- - Quantity is zero; ‡ No data submitted; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.27a. Nevada – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	13,036	1,930	2,244	669	1,123	2,395	409	2,321	4,183
Gender									
Male	6,576	792	903	438	474	969	164	1,512	2,381
Female	6,445	1,138	1,336	231	646	1,424	245	806	1,795
Total	13,021	1,930	2,239	669	1,120	2,393	409	2,318	4,176
Age²									
0 to 11 years	1,161	433	186	226	23	147	1	6	397
12 to 14 years	1,026	247	200	156	46	347	2	9	326
15 to 17 years	1,180	274	228	123	78	401	8	37	415
18 to 20 years	524	78	80	35	35	113	24	55	223
21 to 24 years	779	63	108	22	71	125	44	168	285
25 to 29 years	1,159	117	175	23	104	142	47	280	414
30 to 34 years	1,128	100	190	12	78	155	35	296	400
35 to 39 years	996	103	156	9	116	108	49	265	324
40 to 44 years	934	97	160	17	95	117	40	248	281
45 to 49 years	945	104	163	12	107	132	42	235	272
50 to 54 years	971	101	184	14	110	129	51	244	280
55 to 59 years	994	108	188	3	130	169	29	222	277
60 to 64 years	701	53	137	10	80	161	25	152	168
65 years and older	534	52	89	5	50	149	11	104	120
Total	13,032	1,930	2,244	667	1,123	2,395	408	2,321	4,182
Race									
American Indian or Alaska Native	247	41	37	20	18	46	12	44	84
Asian ³	330	36	34	8	22	53	4	131	66
Black or African American	1,707	239	176	54	92	227	47	543	584
Native Hawaiian or Other Pacific Islander	95	17	16	9	6	21	2	23	31
White	6,948	1,090	1,242	409	716	1,403	230	1,005	2,173
Other	1,107	73	192	12	81	146	49	304	363
Total	10,434	1,496	1,697	512	935	1,896	344	2,050	3,301

Continued. See notes at end of table.

Table 10.27a. Nevada – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	7,999	1,104	1,236	397	802	1,376	279	1,685	2,496
Hispanic or Latino origin	2,284	365	419	86	133	544	65	387	697
Total	10,283	1,469	1,655	483	935	1,920	344	2,072	3,193
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴									
Veteran	31	2	2	- -	4	8	2	14	3
Not a veteran	1,039	123	141	9	146	287	62	382	157
Total	1,070	125	143	9	150	295	64	396	160

- - Quantity is zero; ‡ No data submitted.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.27b. Nevada – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020

Selected characteristics		Mental health diagnosis ¹								
		Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender										
Male	50.5	41.0	40.3	65.5	42.3	40.5	40.1	65.2	57.0	
Female	49.5	59.0	59.7	34.5	57.7	59.5	59.9	34.8	43.0	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age ²										
0 to 11 years	8.9	22.4	8.3	33.9	2.0	6.1	†	†	9.5	
12 to 14 years	7.9	12.8	8.9	23.4	4.1	14.5	†	†	7.8	
15 to 17 years	9.1	14.2	10.2	18.4	6.9	16.7	2.0	1.6	9.9	
18 to 20 years	4.0	4.0	3.6	5.2	3.1	4.7	5.9	2.4	5.3	
21 to 24 years	6.0	3.3	4.8	3.3	6.3	5.2	10.8	7.2	6.8	
25 to 29 years	8.9	6.1	7.8	3.4	9.3	5.9	11.5	12.1	9.9	
30 to 34 years	8.7	5.2	8.5	1.8	6.9	6.5	8.6	12.8	9.6	
35 to 39 years	7.6	5.3	7.0	1.3	10.3	4.5	12.0	11.4	7.7	
40 to 44 years	7.2	5.0	7.1	2.5	8.5	4.9	9.8	10.7	6.7	
45 to 49 years	7.3	5.4	7.3	1.8	9.5	5.5	10.3	10.1	6.5	
50 to 54 years	7.5	5.2	8.2	2.1	9.8	5.4	12.5	10.5	6.7	
55 to 59 years	7.6	5.6	8.4	†	11.6	7.1	7.1	9.6	6.6	
60 to 64 years	5.4	2.7	6.1	1.5	7.1	6.7	6.1	6.5	4.0	
65 years and older	4.1	2.7	4.0	†	4.5	6.2	2.7	4.5	2.9	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race										
American Indian or Alaska Native	2.4	2.7	2.2	3.9	1.9	2.4	3.5	2.1	2.5	
Asian ³	3.2	2.4	2.0	1.6	2.4	2.8	1.2	6.4	2.0	
Black or African American	16.4	16.0	10.4	10.5	9.8	12.0	13.7	26.5	17.7	
Native Hawaiian or Other Pacific Islander	†	1.1	†	1.8	†	1.1	†	1.1	†	
White	66.6	72.9	73.2	79.9	76.6	74.0	66.9	49.0	65.8	
Other	10.6	4.9	11.3	2.3	8.7	7.7	14.2	14.8	11.0	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.27b. Nevada – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020 (continued)

Mental health diagnosis ¹														
Selected characteristics			Trauma- and stressor-related disorders			Mood disorders		Personality disorders			Schizophrenia, other psychotic disorders		Other mental disorders ¹	
						Bipolar disorders	Depressive disorders							
Total														
Ethnicity														
Not of Hispanic or Latino origin			77.8	75.2	74.7	82.2	85.8	71.7	81.1	81.3	78.2			
Hispanic or Latino origin			22.2	24.8	25.3	17.8	14.2	28.3	18.9	18.7	21.8			
Total			100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0			
Specific Hispanic or Latino origin														
Mexican			‡	‡	‡	‡	‡	‡	‡	‡	‡			
Puerto Rican			‡	‡	‡	‡	‡	‡	‡	‡	‡			
Cuban			‡	‡	‡	‡	‡	‡	‡	‡	‡			
Other specific			‡	‡	‡	‡	‡	‡	‡	‡	‡			
Total			‡	‡	‡	‡	‡	‡	‡	‡	‡			
Veteran status ⁴														
Veteran			2.9	1.6	1.4	--	2.7	2.7	3.1	3.5	1.9			
Not a veteran			97.1	98.4	98.6	100.0	97.3	97.3	96.9	96.5	98.1			
Total			100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0			

- - Quantity is zero; ‡ No data submitted; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.28a. New Jersey – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	315,988	61,405	35,098	25,562	31,660	79,778	9,266	37,502	176,753
Gender									
Male	149,885	25,298	13,111	19,150	13,124	27,680	3,672	20,785	92,710
Female	165,552	36,009	21,943	6,395	18,492	51,985	5,570	16,599	83,711
Total	315,437	61,307	35,054	25,545	31,616	79,665	9,242	37,384	176,421
Age²									
0 to 11 years	13,081	5,564	1,367	4,866	106	303	5	39	5,768
12 to 14 years	10,207	4,008	1,495	3,160	179	1,230	17	50	4,282
15 to 17 years	13,175	4,231	2,251	3,445	494	2,547	61	113	5,957
18 to 20 years	10,555	2,662	1,659	2,711	693	1,839	85	179	5,650
21 to 24 years	10,398	2,020	1,242	2,391	992	1,652	57	258	6,047
25 to 29 years	18,021	3,550	1,741	3,294	1,945	3,035	144	729	11,132
30 to 34 years	21,979	4,664	2,096	2,727	2,580	4,728	323	1,306	13,222
35 to 39 years	23,119	4,528	2,449	1,398	2,826	5,759	510	1,900	13,713
40 to 44 years	21,680	4,190	2,482	518	2,814	5,950	692	2,275	12,392
45 to 49 years	22,553	4,113	2,841	319	2,922	6,424	909	2,640	12,558
50 to 54 years	26,824	4,569	3,039	225	3,378	7,815	1,105	3,774	15,455
55 to 59 years	31,212	4,760	3,352	210	3,752	9,160	1,366	5,067	18,156
60 to 64 years	28,845	4,136	2,963	144	3,083	8,707	1,276	5,667	16,545
65 years and older	64,192	8,390	6,117	154	5,894	20,615	2,710	13,495	35,759
Total	315,841	61,385	35,094	25,562	31,658	79,764	9,260	37,492	176,636
Race									
American Indian or Alaska Native	2,857	404	237	109	328	639	135	444	1,738
Asian ³	4,953	850	480	316	397	1,068	90	734	2,920
Black or African American	61,957	11,198	3,458	5,827	5,045	12,189	1,328	11,621	39,809
Native Hawaiian or Other Pacific Islander	27	10	8	5	4	9	--	6	7
White	165,562	31,811	21,565	10,688	19,344	44,638	6,384	17,788	89,599
Other	17,958	3,630	2,170	2,447	2,326	4,096	228	1,477	8,532
Total	253,314	47,903	27,918	19,392	27,444	62,639	8,165	32,070	142,605

Continued. See notes at end of table.

Table 10.28a. New Jersey – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	239,413	42,204	25,928	15,340	27,064	60,774	8,112	31,921	136,623
Hispanic or Latino origin	56,305	12,535	6,265	6,223	3,688	15,169	1,016	4,903	30,875
Total	295,718	54,739	32,193	21,563	30,752	75,943	9,128	36,824	167,498
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴									
Veteran	6,108	974	517	52	769	1,769	232	1,123	3,498
Not a veteran	238,970	39,560	26,298	12,147	27,807	67,556	7,037	30,962	137,914
Total	245,078	40,534	26,815	12,199	28,576	69,325	7,269	32,085	141,412

- - Quantity is zero; ‡ No data submitted.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.28b. New Jersey – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020

Selected characteristics		Mental health diagnosis ¹								
		Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender										
Male	47.5	41.3	37.4	75.0	41.5	34.7	39.7	55.6	52.6	
Female	52.5	58.7	62.6	25.0	58.5	65.3	60.3	44.4	47.4	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age ²										
0 to 11 years	4.1	9.1	3.9	19.0	†	†	†	†	3.3	
12 to 14 years	3.2	6.5	4.3	12.4	†	1.5	†	†	2.4	
15 to 17 years	4.2	6.9	6.4	13.5	1.6	3.2	†	†	3.4	
18 to 20 years	3.3	4.3	4.7	10.6	2.2	2.3	†	†	3.2	
21 to 24 years	3.3	3.3	3.5	9.4	3.1	2.1	†	†	3.4	
25 to 29 years	5.7	5.8	5.0	12.9	6.1	3.8	1.6	1.9	6.3	
30 to 34 years	7.0	7.6	6.0	10.7	8.1	5.9	3.5	3.5	7.5	
35 to 39 years	7.3	7.4	7.0	5.5	8.9	7.2	5.5	5.1	7.8	
40 to 44 years	6.9	6.8	7.1	2.0	8.9	7.5	7.5	6.1	7.0	
45 to 49 years	7.1	6.7	8.1	1.2	9.2	8.1	9.8	7.0	7.1	
50 to 54 years	8.5	7.4	8.7	†	10.7	9.8	11.9	10.1	8.7	
55 to 59 years	9.9	7.8	9.6	†	11.9	11.5	14.8	13.5	10.3	
60 to 64 years	9.1	6.7	8.4	†	9.7	10.9	13.8	15.1	9.4	
65 years and older	20.3	13.7	17.4	†	18.6	25.8	29.3	36.0	20.2	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race										
American Indian or Alaska Native	1.1	†	†	†	1.2	1.0	1.7	1.4	1.2	
Asian ³	2.0	1.8	1.7	1.6	1.4	1.7	1.1	2.3	2.0	
Black or African American	24.5	23.4	12.4	30.0	18.4	19.5	16.3	36.2	27.9	
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	--	†	†	
White	65.4	66.4	77.2	55.1	70.5	71.3	78.2	55.5	62.8	
Other	7.1	7.6	7.8	12.6	8.5	6.5	2.8	4.6	6.0	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.28b. New Jersey – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020 (continued)

Selected characteristics		Mental health diagnosis¹								
		Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders¹
						Bipolar disorders	Depressive disorders			
Ethnicity										
Not of Hispanic or Latino origin	81.0	77.1	80.5	71.1	88.0	80.0	88.9	86.7	81.6	
Hispanic or Latino origin	19.0	22.9	19.5	28.9	12.0	20.0	11.1	13.3	18.4	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	
Specific Hispanic or Latino origin										
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡	
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡	
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡	
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡	
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡	
Veteran status⁴										
Veteran	2.5	2.4	1.9	†	2.7	2.6	3.2	3.5	2.5	
Not a veteran	97.5	97.6	98.1	99.6	97.3	97.4	96.8	96.5	97.5	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	

- - Quantity is zero; ‡ No data submitted; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.29a. New Mexico – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	179,605	43,583	41,569	13,116	9,932	35,556	565	7,082	81,353
Gender									
Male	85,964	17,393	13,950	8,886	3,724	11,963	230	4,357	48,682
Female	93,530	26,175	27,606	4,228	6,201	23,578	335	2,716	32,603
Total	179,494	43,568	41,556	13,114	9,925	35,541	565	7,073	81,285
Age²									
0 to 11 years	40,938	9,767	2,683	5,985	131	1,095	22	58	28,566
12 to 14 years	13,275	4,794	2,274	2,414	202	2,569	7	109	5,670
15 to 17 years	12,587	4,181	3,075	1,396	425	3,433	8	186	4,945
18 to 20 years	8,032	2,022	2,441	534	447	2,237	24	262	2,675
21 to 24 years	9,649	2,247	3,092	386	649	2,462	49	452	3,431
25 to 29 years	15,409	3,540	4,765	557	1,091	3,413	81	779	6,205
30 to 34 years	15,425	3,528	4,655	545	1,149	3,139	66	762	6,660
35 to 39 years	13,825	3,247	4,197	389	1,160	2,836	57	780	5,740
40 to 44 years	10,861	2,493	3,314	283	1,019	2,477	62	665	4,270
45 to 49 years	9,007	2,114	2,611	180	939	2,215	44	588	3,401
50 to 54 years	8,415	1,792	2,445	164	861	2,337	55	610	3,004
55 to 59 years	8,706	1,726	2,505	140	810	2,495	41	677	2,997
60 to 64 years	6,487	1,180	1,802	104	526	2,018	17	491	2,105
65 years and older	6,987	952	1,710	39	523	2,830	32	662	1,683
Total	179,603	43,583	41,569	13,116	9,932	35,556	565	7,081	81,352
Race									
American Indian or Alaska Native	17,312	3,487	2,609	652	305	2,974	29	447	10,421
Asian ³	1,757	402	342	98	53	309	6	61	897
Black or African American	3,804	1,081	869	287	266	723	11	285	1,504
Native Hawaiian or Other Pacific Islander	229	61	48	7	13	57	- -	24	90
White	129,950	32,520	32,398	9,406	7,628	26,662	420	4,719	55,115
Other	4,259	1,111	931	323	283	812	26	276	1,892
Total	157,311	38,662	37,197	10,773	8,548	31,537	492	5,812	69,919

Continued. See notes at end of table.

Table 10.29a. New Mexico – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020 (continued)

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	73,727	17,883	17,425	5,031	5,054	15,414	303	3,148	32,131
Hispanic or Latino origin	84,081	20,788	20,033	5,773	3,578	16,325	192	2,822	37,759
Total	157,808	38,671	37,458	10,804	8,632	31,739	495	5,970	69,890
Specific Hispanic or Latino origin									
Mexican	683	172	155	81	59	161	3	69	254
Puerto Rican	66	20	16	10	10	7	-	5	28
Cuban	28	8	5	4	4	5	1	2	9
Other specific	371	100	80	37	45	74	3	43	144
Total	1,148	300	256	132	118	247	7	119	435
Veteran status⁴									
Veteran	871	226	170	23	80	228	4	55	311
Not a veteran	108,032	24,046	32,965	3,266	8,743	27,502	474	6,204	39,985
Total	108,903	24,272	33,135	3,289	8,823	27,730	478	6,259	40,296

- - Quantity is zero.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.29b. New Mexico – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020

Selected characteristics		Mental health diagnosis¹										
		Trauma- and stressor-related disorders				Mood disorders		Personality disorders			Schizophrenia, other psychotic disorders	Other mental disorders¹
						Bipolar disorders	Depressive disorders					
Total		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	
Total individuals		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	
Gender												
Male		47.9	39.9	33.6	67.8	37.5	33.7	40.7	61.6	59.9		
Female		52.1	60.1	66.4	32.2	62.5	66.3	59.3	38.4	40.1		
	Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0		
Age²												
0 to 11 years		22.8	22.4	6.5	45.6	1.3	3.1	3.9	†	35.1		
12 to 14 years		7.4	11.0	5.5	18.4	2.0	7.2	1.2	1.5	7.0		
15 to 17 years		7.0	9.6	7.4	10.6	4.3	9.7	1.4	2.6	6.1		
18 to 20 years		4.5	4.6	5.9	4.1	4.5	6.3	4.2	3.7	3.3		
21 to 24 years		5.4	5.2	7.4	2.9	6.5	6.9	8.7	6.4	4.2		
25 to 29 years		8.6	8.1	11.5	4.2	11.0	9.6	14.3	11.0	7.6		
30 to 34 years		8.6	8.1	11.2	4.2	11.6	8.8	11.7	10.8	8.2		
35 to 39 years		7.7	7.5	10.1	3.0	11.7	8.0	10.1	11.0	7.1		
40 to 44 years		6.0	5.7	8.0	2.2	10.3	7.0	11.0	9.4	5.2		
45 to 49 years		5.0	4.9	6.3	1.4	9.5	6.2	7.8	8.3	4.2		
50 to 54 years		4.7	4.1	5.9	1.3	8.7	6.6	9.7	8.6	3.7		
55 to 59 years		4.8	4.0	6.0	1.1	8.2	7.0	7.3	9.6	3.7		
60 to 64 years		3.6	2.7	4.3	†	5.3	5.7	3.0	6.9	2.6		
65 years and older		3.9	2.2	4.1	†	5.3	8.0	5.7	9.3	2.1		
	Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0		
Race												
American Indian or Alaska Native		11.0	9.0	7.0	6.1	3.6	9.4	5.9	7.7	14.9		
Asian³		1.1	1.0	†	†	†	†	1.2	1.0	1.3		
Black or African American		2.4	2.8	2.3	2.7	3.1	2.3	2.2	4.9	2.2		
Native Hawaiian or Other Pacific Islander		†	†	†	†	†	†	--	†	†		
White		82.6	84.1	87.1	87.3	89.2	84.5	85.4	81.2	78.8		
Other		2.7	2.9	2.5	3.0	3.3	2.6	5.3	4.7	2.7		
	Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0		

Continued. See notes at end of table.

Table 10.29b. New Mexico – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020 (continued)

Selected characteristics		Mental health diagnosis ¹							
					Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD					
Ethnicity									
Not of Hispanic or Latino origin	46.7	46.2	46.5	46.6	58.5	48.6	61.2	52.7	46.0
Hispanic or Latino origin	53.3	53.8	53.5	53.4	41.5	51.4	38.8	47.3	54.0
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	59.5	57.3	60.5	61.4	50.0	65.2	42.9	58.0	58.4
Puerto Rican	5.7	6.7	6.3	7.6	8.5	2.8	- -	4.2	6.4
Cuban	2.4	2.7	2.0	3.0	3.4	2.0	14.3	1.7	2.1
Other specific	32.3	33.3	31.3	28.0	38.1	30.0	42.9	36.1	33.1
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁴									
Veteran	†	†	†	†	†	†	†	†	†
Not a veteran	99.2	99.1	99.5	99.3	99.1	99.2	99.2	99.1	99.2
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

- - Quantity is zero; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.30a. New York – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals²	54,840	14,035	7,606	3,779	6,359	11,289	9,177	19,827	12,875
Gender									
Male	36,037	9,768	3,706	2,642	3,091	5,039	7,686	13,602	9,422
Female	18,803	4,267	3,900	1,137	3,268	6,250	1,491	6,225	3,453
Total	54,840	14,035	7,606	3,779	6,359	11,289	9,177	19,827	12,875
Age³									
0 to 11 years	1,982	545	424	1,230	44	225	1	25	1,181
12 to 14 years	2,285	622	647	934	185	828	22	75	1,112
15 to 17 years	2,737	689	929	666	324	1,323	58	202	1,091
18 to 20 years	1,309	459	263	177	168	360	230	251	382
21 to 24 years	2,609	1,020	329	122	268	444	663	718	647
25 to 29 years	5,084	2,000	587	196	454	690	1,309	1,488	1,182
30 to 34 years	5,516	2,009	657	142	590	790	1,385	1,689	1,307
35 to 39 years	5,272	1,737	641	122	569	799	1,298	1,750	1,210
40 to 44 years	4,512	1,265	565	68	529	770	969	1,746	939
45 to 49 years	4,126	1,016	518	44	581	769	782	1,690	839
50 to 54 years	4,955	1,063	585	30	669	1,051	908	2,171	927
55 to 59 years	5,208	824	534	28	759	1,089	706	2,660	836
60 to 64 years	4,193	440	401	11	548	917	483	2,413	578
65 years and older	5,051	346	526	9	671	1,234	363	2,948	644
Total	54,839	14,035	7,606	3,779	6,359	11,289	9,177	19,826	12,875
Race									
American Indian or Alaska Native	347	144	41	14	39	72	92	61	101
Asian ⁴	1,012	101	116	36	99	211	45	660	112
Black or African American	15,822	4,375	871	666	1,022	1,868	3,678	7,252	3,563
Native Hawaiian or Other Pacific Islander	47	8	4	-	7	3	5	28	8
White	27,292	6,687	5,181	2,289	4,019	6,562	3,842	8,366	6,712
Other	6,559	1,852	841	567	657	1,449	1,135	2,230	1,702
Total	51,079	13,167	7,054	3,572	5,843	10,165	8,797	18,597	12,198

Continued. See notes at end of table.

Table 10.30a. New York – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	40,858	10,640	5,779	2,958	4,715	7,993	7,113	14,738	9,809
Hispanic or Latino origin	8,101	2,192	968	459	800	1,828	1,493	2,877	1,875
Total	48,959	12,832	6,747	3,417	5,515	9,821	8,606	17,615	11,684
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁵									
Veteran	286	35	16	1	26	39	51	179	48
Not a veteran	10,834	1,452	740	68	1,153	1,490	2,215	6,400	1,793
Total	11,120	1,487	756	69	1,179	1,529	2,266	6,579	1,841

- - Quantity is zero; ‡ No data submitted.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² New York reports only a subset of the total individuals served through their SMHAs.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.30b. New York – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020

Selected characteristics		Mental health diagnosis¹								
		Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders¹
						Bipolar disorders	Depressive disorders			
Total individuals²	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender										
Male	65.7	69.6	48.7	69.9	48.6	44.6	83.8	68.6	73.2	
Female	34.3	30.4	51.3	30.1	51.4	55.4	16.2	31.4	26.8	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age³										
0 to 11 years	3.6	3.9	5.6	32.5	†	2.0	†	†	9.2	
12 to 14 years	4.2	4.4	8.5	24.7	2.9	7.3	†	†	8.6	
15 to 17 years	5.0	4.9	12.2	17.6	5.1	11.7	†	1.0	8.5	
18 to 20 years	2.4	3.3	3.5	4.7	2.6	3.2	2.5	1.3	3.0	
21 to 24 years	4.8	7.3	4.3	3.2	4.2	3.9	7.2	3.6	5.0	
25 to 29 years	9.3	14.3	7.7	5.2	7.1	6.1	14.3	7.5	9.2	
30 to 34 years	10.1	14.3	8.6	3.8	9.3	7.0	15.1	8.5	10.2	
35 to 39 years	9.6	12.4	8.4	3.2	8.9	7.1	14.1	8.8	9.4	
40 to 44 years	8.2	9.0	7.4	1.8	8.3	6.8	10.6	8.8	7.3	
45 to 49 years	7.5	7.2	6.8	1.2	9.1	6.8	8.5	8.5	6.5	
50 to 54 years	9.0	7.6	7.7	†	10.5	9.3	9.9	11.0	7.2	
55 to 59 years	9.5	5.9	7.0	†	11.9	9.6	7.7	13.4	6.5	
60 to 64 years	7.6	3.1	5.3	†	8.6	8.1	5.3	12.2	4.5	
65 years and older	9.2	2.5	6.9	†	10.6	10.9	4.0	14.9	5.0	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race										
American Indian or Alaska Native	†	1.1	†	†	†	†	1.0	†	†	
Asian⁴	2.0	†	1.6	1.0	1.7	2.1	†	3.5	†	
Black or African American	31.0	33.2	12.3	18.6	17.5	18.4	41.8	39.0	29.2	
Native Hawaiian or Other Pacific Islander	†	†	†	--	†	†	†	†	†	
White	53.4	50.8	73.4	64.1	68.8	64.6	43.7	45.0	55.0	
Other	12.8	14.1	11.9	15.9	11.2	14.3	12.9	12.0	14.0	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.30b. New York – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	83.5	82.9	85.7	86.6	85.5	81.4	82.7	83.7	84.0
Hispanic or Latino origin	16.5	17.1	14.3	13.4	14.5	18.6	17.3	16.3	16.0
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status ⁵									
Veteran	2.6	2.4	2.1	1.4	2.2	2.6	2.3	2.7	2.6
Not a veteran	97.4	97.6	97.9	98.6	97.8	97.4	97.7	97.3	97.4
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

- - Quantity is zero; ‡ No data submitted; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² New York reports only a subset of the total individuals served through their SMHAs.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.31a. North Carolina – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	99,827	24,649	22,486	5,430	19,466	41,645	3,499	16,122	14,627
Gender									
Male	49,063	10,095	9,282	3,490	8,612	18,423	1,332	10,284	8,469
Female	50,753	14,553	13,204	1,936	10,852	23,220	2,166	5,835	6,156
Total	99,816	24,648	22,486	5,426	19,464	41,643	3,498	16,119	14,625
Age²									
0 to 11 years	2,925	1,046	252	1,059	38	238	- -	26	1,316
12 to 14 years	3,549	1,078	403	1,049	111	773	7	40	1,672
15 to 17 years	3,850	1,037	542	815	219	1,171	29	65	1,798
18 to 20 years	4,269	1,072	869	465	586	1,772	130	388	1,044
21 to 24 years	8,523	1,996	1,953	409	1,646	3,673	372	1,257	1,187
25 to 29 years	13,329	3,231	3,286	413	2,760	5,707	597	2,311	1,431
30 to 34 years	12,531	3,129	3,198	354	2,831	5,247	549	2,245	1,211
35 to 39 years	10,742	2,728	2,671	289	2,432	4,481	455	1,976	1,032
40 to 44 years	9,190	2,450	2,270	172	2,162	4,064	373	1,648	769
45 to 49 years	9,047	2,374	2,251	150	2,155	4,260	357	1,538	793
50 to 54 years	8,336	1,992	1,959	126	1,965	4,058	279	1,487	788
55 to 59 years	7,005	1,416	1,575	86	1,493	3,433	196	1,420	637
60 to 64 years	4,040	735	913	32	723	1,947	106	959	428
65 years and older	2,491	365	344	11	345	821	49	762	521
Total	99,827	24,649	22,486	5,430	19,466	41,645	3,499	16,122	14,627
Race									
American Indian or Alaska Native	1,653	438	324	80	472	627	64	257	200
Asian ³	509	109	92	15	71	184	19	134	67
Black or African American	29,734	6,564	4,462	1,796	4,913	11,268	718	8,121	4,534
Native Hawaiian or Other Pacific Islander	73	22	16	2	13	26	1	10	14
White	63,202	16,082	16,656	3,257	13,391	27,603	2,560	6,995	9,226
Other	4,333	1,380	875	255	550	1,834	120	513	509
Total	99,504	24,595	22,425	5,405	19,410	41,542	3,482	16,030	14,550

Continued. See notes at end of table.

Table 10.31a. North Carolina – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020 (continued)

Selected characteristics		Mental health diagnosis¹								
		Trauma- and stressor-related disorders			Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders¹	
					Bipolar disorders	Depressive disorders				
Total										
Ethnicity										
Not of Hispanic or Latino origin		94,632	22,912	21,301	5,112	18,909	39,396	3,389	15,534	13,936
Hispanic or Latino origin		4,969	1,693	1,137	286	516	2,173	104	535	635
Total		99,601	24,605	22,438	5,398	19,425	41,569	3,493	16,069	14,571
Specific Hispanic or Latino origin										
Mexican		1,327	437	319	91	113	626	24	137	183
Puerto Rican		400	122	114	24	89	171	18	62	46
Cuban		86	24	23	7	14	41	4	15	15
Other specific		3,090	1,101	670	160	291	1,315	55	305	379
Total		4,903	1,684	1,126	282	507	2,153	101	519	623
Veteran status⁴										
Veteran		1,270	340	233	22	200	531	37	236	181
Not a veteran		87,505	21,040	20,940	2,469	18,754	38,742	3,376	15,407	9,550
Total		88,775	21,380	21,173	2,491	18,954	39,273	3,413	15,643	9,731

- - Quantity is zero.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.31b. North Carolina – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020

Selected characteristics		Mental health diagnosis¹							
		Trauma- and stressor-related disorders			Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders¹
					Bipolar disorders	Depressive disorders			
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender									
Male	49.2	41.0	41.3	64.3	44.2	44.2	38.1	63.8	57.9
Female	50.8	59.0	58.7	35.7	55.8	55.8	61.9	36.2	42.1
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age²									
0 to 11 years	2.9	4.2	1.1	19.5	†	†	--	†	9.0
12 to 14 years	3.6	4.4	1.8	19.3	†	1.9	†	†	11.4
15 to 17 years	3.9	4.2	2.4	15.0	1.1	2.8	†	†	12.3
18 to 20 years	4.3	4.3	3.9	8.6	3.0	4.3	3.7	2.4	7.1
21 to 24 years	8.5	8.1	8.7	7.5	8.5	8.8	10.6	7.8	8.1
25 to 29 years	13.4	13.1	14.6	7.6	14.2	13.7	17.1	14.3	9.8
30 to 34 years	12.6	12.7	14.2	6.5	14.5	12.6	15.7	13.9	8.3
35 to 39 years	10.8	11.1	11.9	5.3	12.5	10.8	13.0	12.3	7.1
40 to 44 years	9.2	9.9	10.1	3.2	11.1	9.8	10.7	10.2	5.3
45 to 49 years	9.1	9.6	10.0	2.8	11.1	10.2	10.2	9.5	5.4
50 to 54 years	8.4	8.1	8.7	2.3	10.1	9.7	8.0	9.2	5.4
55 to 59 years	7.0	5.7	7.0	1.6	7.7	8.2	5.6	8.8	4.4
60 to 64 years	4.0	3.0	4.1	†	3.7	4.7	3.0	5.9	2.9
65 years and older	2.5	1.5	1.5	†	1.8	2.0	1.4	4.7	3.6
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race									
American Indian or Alaska Native	1.7	1.8	1.4	1.5	2.4	1.5	1.8	1.6	1.4
Asian³	†	†	†	†	†	†	†	†	†
Black or African American	29.9	26.7	19.9	33.2	25.3	27.1	20.6	50.7	31.2
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	†	†	†
White	63.5	65.4	74.3	60.3	69.0	66.4	73.5	43.6	63.4
Other	4.4	5.6	3.9	4.7	2.8	4.4	3.4	3.2	3.5
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

**Table 10.31b. North Carolina – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020
(continued)**

Selected characteristics		Mental health diagnosis ¹							
					Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD					
Ethnicity									
Not of Hispanic or Latino origin	95.0	93.1	94.9	94.7	97.3	94.8	97.0	96.7	95.6
Hispanic or Latino origin	5.0	6.9	5.1	5.3	2.7	5.2	3.0	3.3	4.4
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	27.1	26.0	28.3	32.3	22.3	29.1	23.8	26.4	29.4
Puerto Rican	8.2	7.2	10.1	8.5	17.6	7.9	17.8	11.9	7.4
Cuban	1.8	1.4	2.0	2.5	2.8	1.9	4.0	2.9	2.4
Other specific	63.0	65.4	59.5	56.7	57.4	61.1	54.5	58.8	60.8
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁴									
Veteran	1.4	1.6	1.1	†	1.1	1.4	1.1	1.5	1.9
Not a veteran	98.6	98.4	98.9	99.1	98.9	98.6	98.9	98.5	98.1
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

- - Quantity is zero; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.32a. North Dakota – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	9,341	2,547	3,880	1,308	1,073	3,624	937	1,490	2,616
Gender									
Male	4,792	922	1,728	841	515	1,556	378	929	1,633
Female	4,532	1,616	2,146	466	557	2,060	558	561	980
Total	9,324	2,538	3,874	1,307	1,072	3,616	936	1,490	2,613
Age²									
0 to 11 years	660	255	126	290	9	58	--	1	374
12 to 14 years	524	165	160	239	22	152	2	3	300
15 to 17 years	581	167	248	164	39	280	16	7	231
18 to 20 years	428	123	195	84	53	174	36	27	129
21 to 24 years	610	189	250	68	63	281	75	63	134
25 to 29 years	1,013	318	477	102	119	406	113	149	202
30 to 34 years	981	301	443	90	107	383	126	174	179
35 to 39 years	865	269	427	69	116	354	115	143	143
40 to 44 years	751	199	355	55	103	319	110	145	137
45 to 49 years	608	159	253	39	103	249	95	129	149
50 to 54 years	606	147	257	44	106	261	72	130	145
55 to 59 years	623	134	244	27	84	266	75	181	141
60 to 64 years	493	81	198	20	77	198	46	153	142
65 years and older	598	40	247	17	72	243	56	185	210
Total	9,341	2,547	3,880	1,308	1,073	3,624	937	1,490	2,616
Race									
American Indian or Alaska Native	1,236	496	432	135	77	504	93	148	263
Asian ³	76	19	23	6	8	26	7	22	27
Black or African American	355	124	120	34	29	119	27	97	83
Native Hawaiian or Other Pacific Islander	--	--	--	--	--	--	--	--	--
White	7,059	1,697	3,089	1,041	905	2,754	760	1,134	2,055
Other	562	206	208	90	51	212	48	87	148
Total	9,288	2,542	3,872	1,306	1,070	3,615	935	1,488	2,576

Continued. See notes at end of table.

Table 10.32a. North Dakota – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	9,024	2,426	3,749	1,273	1,047	3,489	915	1,451	2,544
Hispanic or Latino origin	310	119	130	35	26	131	22	39	71
Total	9,334	2,545	3,879	1,308	1,073	3,620	937	1,490	2,615
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴									
Veteran	185	65	78	9	24	79	20	37	26
Not a veteran	6,594	1,803	2,995	549	894	2,784	809	1,214	1,314
Total	6,779	1,868	3,073	558	918	2,863	829	1,251	1,340

- - Quantity is zero; ‡ No data submitted.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.32b. North Dakota – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020

Selected characteristics		Mental health diagnosis ¹								
		Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender										
Male	51.4	36.3	44.6	64.3	48.0	43.0	40.4	62.3	62.5	
Female	48.6	63.7	55.4	35.7	52.0	57.0	59.6	37.7	37.5	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age ²										
0 to 11 years	7.1	10.0	3.2	22.2	†	1.6	--	†	14.3	
12 to 14 years	5.6	6.5	4.1	18.3	2.1	4.2	†	†	11.5	
15 to 17 years	6.2	6.6	6.4	12.5	3.6	7.7	1.7	†	8.8	
18 to 20 years	4.6	4.8	5.0	6.4	4.9	4.8	3.8	1.8	4.9	
21 to 24 years	6.5	7.4	6.4	5.2	5.9	7.8	8.0	4.2	5.1	
25 to 29 years	10.8	12.5	12.3	7.8	11.1	11.2	12.1	10.0	7.7	
30 to 34 years	10.5	11.8	11.4	6.9	10.0	10.6	13.4	11.7	6.8	
35 to 39 years	9.3	10.6	11.0	5.3	10.8	9.8	12.3	9.6	5.5	
40 to 44 years	8.0	7.8	9.1	4.2	9.6	8.8	11.7	9.7	5.2	
45 to 49 years	6.5	6.2	6.5	3.0	9.6	6.9	10.1	8.7	5.7	
50 to 54 years	6.5	5.8	6.6	3.4	9.9	7.2	7.7	8.7	5.5	
55 to 59 years	6.7	5.3	6.3	2.1	7.8	7.3	8.0	12.1	5.4	
60 to 64 years	5.3	3.2	5.1	1.5	7.2	5.5	4.9	10.3	5.4	
65 years and older	6.4	1.6	6.4	1.3	6.7	6.7	6.0	12.4	8.0	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race										
American Indian or Alaska Native	13.3	19.5	11.2	10.3	7.2	13.9	9.9	9.9	10.2	
Asian ³	†	†	†	†	†	†	†	1.5	1.0	
Black or African American	3.8	4.9	3.1	2.6	2.7	3.3	2.9	6.5	3.2	
Native Hawaiian or Other Pacific Islander	--	--	--	--	--	--	--	--	--	
White	76.0	66.8	79.8	79.7	84.6	76.2	81.3	76.2	79.8	
Other	6.1	8.1	5.4	6.9	4.8	5.9	5.1	5.8	5.7	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.32b. North Dakota – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020 (continued)

Mental health diagnosis ¹									
Selected characteristics	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	96.7	95.3	96.6	97.3	97.6	96.4	97.7	97.4	97.3
Hispanic or Latino origin	3.3	4.7	3.4	2.7	2.4	3.6	2.3	2.6	2.7
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status ⁴									
Veteran	2.7	3.5	2.5	1.6	2.6	2.8	2.4	3.0	1.9
Not a veteran	97.3	96.5	97.5	98.4	97.4	97.2	97.6	97.0	98.1
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

- - Quantity is zero; ‡ No data submitted; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.33a. Ohio – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	429,022	107,550	110,594	56,771	54,712	126,003	7,226	39,576	73,300
Gender									
Male	193,101	43,908	40,085	36,793	19,910	44,798	2,776	23,589	41,784
Female	226,724	60,907	68,509	18,155	33,724	78,961	4,284	15,431	29,520
Total	419,825	104,815	108,594	54,948	53,634	123,759	7,060	39,020	71,304
Age²									
0 to 11 years	83,694	32,332	12,681	31,083	420	4,683	22	101	24,064
12 to 14 years	35,546	10,573	7,601	9,723	752	8,741	43	131	9,545
15 to 17 years	28,211	7,501	7,330	5,017	1,386	9,251	171	399	6,709
18 to 20 years	16,965	3,829	5,075	1,738	2,189	6,251	375	1,148	3,024
21 to 24 years	23,902	5,421	7,259	1,584	3,999	8,534	660	2,322	3,400
25 to 29 years	36,820	8,569	11,632	2,034	6,723	12,811	988	3,820	4,569
30 to 34 years	35,900	8,468	11,614	1,737	6,903	12,007	966	3,886	4,072
35 to 39 years	32,967	7,437	10,483	1,361	6,860	10,993	927	4,257	3,424
40 to 44 years	26,579	5,879	8,032	878	5,754	9,109	713	3,773	2,781
45 to 49 years	26,224	5,359	7,603	640	5,604	9,673	686	4,124	2,768
50 to 54 years	25,956	4,575	7,268	431	5,307	10,271	608	4,393	2,684
55 to 59 years	24,518	3,628	6,278	326	4,476	9,915	546	4,935	2,487
60 to 64 years	16,268	2,083	4,012	160	2,551	6,757	332	3,447	1,761
65 years and older	15,467	1,896	3,724	59	1,787	7,006	189	2,838	2,009
Total	429,017	107,550	110,592	56,771	54,711	126,002	7,226	39,574	73,297
Race									
American Indian or Alaska Native	1,592	465	383	156	202	472	24	139	270
Asian ³	2,323	558	530	168	185	772	20	370	382
Black or African American	100,427	25,017	16,641	15,261	9,229	25,087	1,385	14,313	20,169
Native Hawaiian or Other Pacific Islander	425	104	106	60	38	144	5	27	60
White	282,833	71,663	82,790	36,861	39,155	87,099	4,928	20,321	46,204
Other	1,605	396	384	184	175	468	17	207	238
Total	389,205	98,203	100,834	52,690	48,984	114,042	6,379	35,377	67,323

Continued. See notes at end of table.

Table 10.33a. Ohio – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	389,704	99,131	103,118	53,529	49,219	114,822	6,179	32,725	67,939
Hispanic or Latino origin	12,325	3,251	2,935	1,905	980	3,665	118	756	2,282
Total	402,029	102,382	106,053	55,434	50,199	118,487	6,297	33,481	70,221
Specific Hispanic or Latino origin									
Mexican	321	85	67	29	46	123	3	28	40
Puerto Rican	228	46	40	19	24	73	4	28	42
Cuban	21	5	9	1	3	6	- -	3	4
Other specific	1,193	323	214	154	104	345	13	132	249
Total	1,763	459	330	203	177	547	20	191	335
Veteran status⁴									
Veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Not a veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡

- - Quantity is zero; ‡ No data submitted.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.33b. Ohio – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020

Selected characteristics		Mental health diagnosis ¹										
		Trauma- and stressor-related disorders				Mood disorders		Personality disorders			Schizophrenia, other psychotic disorders	Other mental disorders ¹
						Bipolar disorders	Depressive disorders					
Total		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	
Total individuals		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	
Gender												
Male		46.0	41.9	36.9	67.0	37.1	36.2	39.3	60.5	58.6		
Female		54.0	58.1	63.1	33.0	62.9	63.8	60.7	39.5	41.4		
Total		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	
Age ²												
0 to 11 years		19.5	30.1	11.5	54.8	†	3.7	†	†	32.8		
12 to 14 years		8.3	9.8	6.9	17.1	1.4	6.9	†	†	13.0		
15 to 17 years		6.6	7.0	6.6	8.8	2.5	7.3	2.4	1.0	9.2		
18 to 20 years		4.0	3.6	4.6	3.1	4.0	5.0	5.2	2.9	4.1		
21 to 24 years		5.6	5.0	6.6	2.8	7.3	6.8	9.1	5.9	4.6		
25 to 29 years		8.6	8.0	10.5	3.6	12.3	10.2	13.7	9.7	6.2		
30 to 34 years		8.4	7.9	10.5	3.1	12.6	9.5	13.4	9.8	5.6		
35 to 39 years		7.7	6.9	9.5	2.4	12.5	8.7	12.8	10.8	4.7		
40 to 44 years		6.2	5.5	7.3	1.5	10.5	7.2	9.9	9.5	3.8		
45 to 49 years		6.1	5.0	6.9	1.1	10.2	7.7	9.5	10.4	3.8		
50 to 54 years		6.1	4.3	6.6	†	9.7	8.2	8.4	11.1	3.7		
55 to 59 years		5.7	3.4	5.7	†	8.2	7.9	7.6	12.5	3.4		
60 to 64 years		3.8	1.9	3.6	†	4.7	5.4	4.6	8.7	2.4		
65 years and older		3.6	1.8	3.4	†	3.3	5.6	2.6	7.2	2.7		
Total		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	
Race												
American Indian or Alaska Native		†	†	†	†	†	†	†	†	†		
Asian ³		†	†	†	†	†	†	†	1.0	†		
Black or African American		25.8	25.5	16.5	29.0	18.8	22.0	21.7	40.5	30.0		
Native Hawaiian or Other Pacific Islander		†	†	†	†	†	†	†	†	†		
White		72.7	73.0	82.1	70.0	79.9	76.4	77.3	57.4	68.6		
Other		†	†	†	†	†	†	†	†	†		
Total		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	

Continued. See notes at end of table.

Table 10.33b. Ohio – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	96.9	96.8	97.2	96.6	98.0	96.9	98.1	97.7	96.8
Hispanic or Latino origin	3.1	3.2	2.8	3.4	2.0	3.1	1.9	2.3	3.2
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	18.2	18.5	20.3	14.3	26.0	22.5	15.0	14.7	11.9
Puerto Rican	12.9	10.0	12.1	9.4	13.6	13.3	20.0	14.7	12.5
Cuban	1.2	1.1	2.7	†	1.7	1.1	- -	1.6	1.2
Other specific	67.7	70.4	64.8	75.9	58.8	63.1	65.0	69.1	74.3
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁴									
Veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Not a veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡

- - Quantity is zero; ‡ No data submitted; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.34a. Oklahoma – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	94,809	20,896	16,707	4,459	10,594	37,554	397	10,429	9,080
Gender									
Male	44,352	8,887	7,034	3,250	3,887	15,310	151	6,598	5,641
Female	50,457	12,009	9,673	1,209	6,707	22,244	246	3,831	3,439
Total	94,809	20,896	16,707	4,459	10,594	37,554	397	10,429	9,080
Age²									
0 to 11 years	12,807	6,017	1,900	2,409	8	1,052	3	4	2,673
12 to 14 years	6,588	2,289	1,197	1,021	27	1,706	3	13	1,264
15 to 17 years	6,133	1,858	1,287	541	87	2,211	2	52	976
18 to 20 years	4,638	998	931	186	313	2,196	20	222	543
21 to 24 years	6,004	1,039	1,095	79	688	3,037	36	633	465
25 to 29 years	9,010	1,458	1,732	57	1,219	4,301	48	1,194	618
30 to 34 years	9,017	1,540	1,639	37	1,362	4,142	57	1,332	568
35 to 39 years	8,522	1,447	1,588	48	1,388	3,756	51	1,370	454
40 to 44 years	7,302	1,160	1,269	28	1,276	3,227	38	1,238	411
45 to 49 years	6,571	943	1,078	19	1,229	3,016	46	1,077	370
50 to 54 years	6,128	858	1,040	14	1,056	2,918	35	1,088	261
55 to 59 years	5,870	694	983	10	960	2,927	33	1,047	233
60 to 64 years	3,749	367	562	5	624	1,859	15	715	121
65 years and older	2,470	228	406	5	357	1,206	10	444	123
Total	94,809	20,896	16,707	4,459	10,594	37,554	397	10,429	9,080
Race									
American Indian or Alaska Native	8,403	1,991	1,262	423	718	3,394	35	923	842
Asian ³	593	86	91	10	57	231	7	139	44
Black or African American	12,044	2,275	1,750	489	1,020	4,576	37	2,428	1,073
Native Hawaiian or Other Pacific Islander	420	82	57	20	44	179	- -	49	33
White	67,805	14,790	12,335	3,063	8,263	27,456	302	6,508	6,302
Other	5,541	1,672	1,212	454	492	1,717	16	380	784
Total	94,806	20,896	16,707	4,459	10,594	37,553	397	10,427	9,078

Continued. See notes at end of table.

Table 10.34a. Oklahoma – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	86,755	18,813	15,068	4,085	10,084	34,524	380	9,746	8,257
Hispanic or Latino origin	8,046	2,083	1,638	374	509	3,028	16	679	822
Total	94,801	20,896	16,706	4,459	10,593	37,552	396	10,425	9,079
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴									
Veteran	1,735	289	248	3	257	753	7	310	96
Not a veteran	65,996	10,172	11,866	472	10,067	31,165	367	9,839	3,858
Total	67,731	10,461	12,114	475	10,324	31,918	374	10,149	3,954

- - Quantity is zero; ‡ No data submitted.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.34b. Oklahoma – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender									
Male	46.8	42.5	42.1	72.9	36.7	40.8	38.0	63.3	62.1
Female	53.2	57.5	57.9	27.1	63.3	59.2	62.0	36.7	37.9
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age²									
0 to 11 years	13.5	28.8	11.4	54.0	†	2.8	†	†	29.4
12 to 14 years	6.9	11.0	7.2	22.9	†	4.5	†	†	13.9
15 to 17 years	6.5	8.9	7.7	12.1	†	5.9	†	†	10.7
18 to 20 years	4.9	4.8	5.6	4.2	3.0	5.8	5.0	2.1	6.0
21 to 24 years	6.3	5.0	6.6	1.8	6.5	8.1	9.1	6.1	5.1
25 to 29 years	9.5	7.0	10.4	1.3	11.5	11.5	12.1	11.4	6.8
30 to 34 years	9.5	7.4	9.8	†	12.9	11.0	14.4	12.8	6.3
35 to 39 years	9.0	6.9	9.5	1.1	13.1	10.0	12.8	13.1	5.0
40 to 44 years	7.7	5.6	7.6	†	12.0	8.6	9.6	11.9	4.5
45 to 49 years	6.9	4.5	6.5	†	11.6	8.0	11.6	10.3	4.1
50 to 54 years	6.5	4.1	6.2	†	10.0	7.8	8.8	10.4	2.9
55 to 59 years	6.2	3.3	5.9	†	9.1	7.8	8.3	10.0	2.6
60 to 64 years	4.0	1.8	3.4	†	5.9	5.0	3.8	6.9	1.3
65 years and older	2.6	1.1	2.4	†	3.4	3.2	2.5	4.3	1.4
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race									
American Indian or Alaska Native	8.9	9.5	7.6	9.5	6.8	9.0	8.8	8.9	9.3
Asian ³	†	†	†	†	†	†	1.8	1.3	†
Black or African American	12.7	10.9	10.5	11.0	9.6	12.2	9.3	23.3	11.8
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	--	†	†
White	71.5	70.8	73.8	68.7	78.0	73.1	76.1	62.4	69.4
Other	5.8	8.0	7.3	10.2	4.6	4.6	4.0	3.6	8.6
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.34b. Oklahoma – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	91.5	90.0	90.2	91.6	95.2	91.9	96.0	93.5	90.9
Hispanic or Latino origin	8.5	10.0	9.8	8.4	4.8	8.1	4.0	6.5	9.1
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴									
Veteran	2.6	2.8	2.0	†	2.5	2.4	1.9	3.1	2.4
Not a veteran	97.4	97.2	98.0	99.4	97.5	97.6	98.1	96.9	97.6
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

- - Quantity is zero; ‡ No data submitted; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.35a. Oregon – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	145,349	54,561	57,606	16,169	12,514	45,310	3,299	13,150	30,378
Gender									
Male	59,798	19,956	19,181	9,779	4,451	15,410	959	8,264	14,282
Female	85,547	34,602	38,425	6,390	8,062	29,899	2,340	4,886	16,096
Total	145,345	54,558	57,606	16,169	12,513	45,309	3,299	13,150	30,378
Age²									
0 to 11 years	22,372	11,690	5,203	5,902	41	1,142	17	9	6,353
12 to 14 years	11,594	4,740	3,942	2,397	96	2,988	21	53	3,012
15 to 17 years	12,140	4,312	5,105	1,717	261	4,693	59	183	3,126
18 to 20 years	6,904	2,231	3,241	664	413	2,807	160	386	1,709
21 to 24 years	8,453	2,777	4,025	565	816	3,265	295	784	1,776
25 to 29 years	13,857	4,605	6,536	1,036	1,527	5,059	456	1,398	2,685
30 to 34 years	13,407	4,641	6,284	1,026	1,559	4,637	443	1,491	2,303
35 to 39 years	11,896	4,401	5,409	868	1,480	4,026	403	1,488	1,881
40 to 44 years	9,398	3,512	4,079	610	1,269	3,196	336	1,255	1,455
45 to 49 years	8,564	3,108	3,664	473	1,245	3,088	319	1,297	1,336
50 to 54 years	7,980	2,846	3,267	337	1,151	3,066	249	1,233	1,268
55 to 59 years	8,092	2,770	3,067	319	1,173	3,190	257	1,378	1,303
60 to 64 years	5,855	1,820	2,236	175	803	2,260	171	1,077	1,011
65 years and older	4,837	1,108	1,548	80	680	1,893	113	1,118	1,160
Total	145,349	54,561	57,606	16,169	12,514	45,310	3,299	13,150	30,378
Race									
American Indian or Alaska Native	5,779	2,442	2,254	561	472	1,767	161	548	1,127
Asian ³	1,900	507	709	120	86	681	27	331	389
Black or African American	5,719	2,190	1,951	740	381	1,666	86	718	1,108
Native Hawaiian or Other Pacific Islander	646	279	209	62	52	184	11	72	120
White	93,105	34,756	37,591	10,611	8,743	29,186	2,196	7,969	19,849
Other	25,524	9,615	9,717	2,965	2,330	8,057	700	3,075	5,488
Total	132,673	49,789	52,431	15,059	12,064	41,541	3,181	12,713	28,081

Continued. See notes at end of table.

Table 10.35a. Oregon – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	111,138	41,120	44,466	12,483	10,717	34,985	2,791	11,140	23,440
Hispanic or Latino origin	20,882	8,413	7,429	2,357	1,194	6,226	352	1,533	4,445
Total	132,020	49,533	51,895	14,840	11,911	41,211	3,143	12,673	27,885
Specific Hispanic or Latino origin									
Mexican	7,139	2,879	2,525	751	402	2,237	130	528	1,565
Puerto Rican	511	187	184	64	55	158	21	55	120
Cuban	471	227	128	102	16	93	5	48	108
Other specific	2,679	1,075	1,014	315	224	812	65	251	540
Total	10,800	4,368	3,851	1,232	697	3,300	221	882	2,333
Veteran status⁴									
Veteran	1,775	680	577	102	271	640	59	317	260
Not a veteran	10,538	3,511	3,536	521	1,610	3,232	488	2,336	1,732
Total	12,313	4,191	4,113	623	1,881	3,872	547	2,653	1,992

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.35b. Oregon – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020

Selected characteristics		Mental health diagnosis ¹								
		Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender										
Male	41.1	36.6	33.3	60.5	35.6	34.0	29.1	62.8	47.0	
Female	58.9	63.4	66.7	39.5	64.4	66.0	70.9	37.2	53.0	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age ²										
0 to 11 years	15.4	21.4	9.0	36.5	†	2.5	†	†	20.9	
12 to 14 years	8.0	8.7	6.8	14.8	†	6.6	†	†	9.9	
15 to 17 years	8.4	7.9	8.9	10.6	2.1	10.4	1.8	1.4	10.3	
18 to 20 years	4.7	4.1	5.6	4.1	3.3	6.2	4.8	2.9	5.6	
21 to 24 years	5.8	5.1	7.0	3.5	6.5	7.2	8.9	6.0	5.8	
25 to 29 years	9.5	8.4	11.3	6.4	12.2	11.2	13.8	10.6	8.8	
30 to 34 years	9.2	8.5	10.9	6.3	12.5	10.2	13.4	11.3	7.6	
35 to 39 years	8.2	8.1	9.4	5.4	11.8	8.9	12.2	11.3	6.2	
40 to 44 years	6.5	6.4	7.1	3.8	10.1	7.1	10.2	9.5	4.8	
45 to 49 years	5.9	5.7	6.4	2.9	9.9	6.8	9.7	9.9	4.4	
50 to 54 years	5.5	5.2	5.7	2.1	9.2	6.8	7.5	9.4	4.2	
55 to 59 years	5.6	5.1	5.3	2.0	9.4	7.0	7.8	10.5	4.3	
60 to 64 years	4.0	3.3	3.9	1.1	6.4	5.0	5.2	8.2	3.3	
65 years and older	3.3	2.0	2.7	†	5.4	4.2	3.4	8.5	3.8	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race										
American Indian or Alaska Native	4.4	4.9	4.3	3.7	3.9	4.3	5.1	4.3	4.0	
Asian ³	1.4	1.0	1.4	†	†	1.6	†	2.6	1.4	
Black or African American	4.3	4.4	3.7	4.9	3.2	4.0	2.7	5.6	3.9	
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	†	†	†	
White	70.2	69.8	71.7	70.5	72.5	70.3	69.0	62.7	70.7	
Other	19.2	19.3	18.5	19.7	19.3	19.4	22.0	24.2	19.5	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.35b. Oregon – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020 (continued)

Selected characteristics		Mental health diagnosis¹							
		Trauma- and stressor-related disorders			Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders¹
					Bipolar disorders	Depressive disorders			
Total									
Ethnicity									
Not of Hispanic or Latino origin	84.2	83.0	85.7	84.1	90.0	84.9	88.8	87.9	84.1
Hispanic or Latino origin	15.8	17.0	14.3	15.9	10.0	15.1	11.2	12.1	15.9
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	66.1	65.9	65.6	61.0	57.7	67.8	58.8	59.9	67.1
Puerto Rican	4.7	4.3	4.8	5.2	7.9	4.8	9.5	6.2	5.1
Cuban	4.4	5.2	3.3	8.3	2.3	2.8	2.3	5.4	4.6
Other specific	24.8	24.6	26.3	25.6	32.1	24.6	29.4	28.5	23.1
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁴									
Veteran	14.4	16.2	14.0	16.4	14.4	16.5	10.8	11.9	13.1
Not a veteran	85.6	83.8	86.0	83.6	85.6	83.5	89.2	88.1	86.9
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.36a. Pennsylvania – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020

Selected characteristics		Mental health diagnosis ¹								
		Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Total individuals	514,459	96,097	99,060	71,529	57,474	145,626	4,892	40,906	88,199	
Gender										
Male	236,428	39,562	34,567	48,313	21,176	50,726	1,593	24,207	51,590	
Female	278,031	56,535	64,493	23,216	36,298	94,900	3,299	16,699	36,609	
Total	514,459	96,097	99,060	71,529	57,474	145,626	4,892	40,906	88,199	
Age ²										
0 to 11 years	86,955	26,516	8,221	34,204	358	2,725	14	91	26,752	
12 to 14 years	42,110	11,078	6,217	13,457	685	6,403	45	145	11,222	
15 to 17 years	39,697	9,117	7,537	8,714	1,548	9,972	129	268	9,699	
18 to 20 years	24,033	4,212	5,441	3,611	2,161	7,682	280	792	4,327	
21 to 24 years	26,169	4,396	6,226	2,127	3,618	8,948	498	1,823	3,504	
25 to 29 years	42,304	6,954	10,622	2,548	6,841	14,212	725	3,513	4,947	
30 to 34 years	41,963	6,976	10,701	2,091	7,288	14,082	683	3,665	4,368	
35 to 39 years	38,555	6,183	9,554	1,634	7,113	13,203	603	3,850	3,751	
40 to 44 years	31,052	4,580	7,229	1,073	5,969	11,297	424	3,278	2,898	
45 to 49 years	31,667	4,287	6,864	784	5,883	12,231	394	3,896	2,944	
50 to 54 years	34,112	4,248	6,687	563	5,996	13,915	355	4,741	3,151	
55 to 59 years	32,277	3,504	5,916	441	5,007	13,471	352	5,509	2,990	
60 to 64 years	22,784	2,199	3,926	218	3,046	9,477	208	4,519	2,446	
65 years and older	20,778	1,846	3,919	64	1,960	8,008	182	4,815	5,200	
Total	514,456	96,096	99,060	71,529	57,473	145,626	4,892	40,905	88,199	
Race										
American Indian or Alaska Native	1,142	258	236	117	145	334	17	62	183	
Asian ³	5,662	1,164	877	408	320	1,812	23	724	1,036	
Black or African American	107,611	23,218	12,586	14,879	9,736	26,880	660	14,295	19,921	
Native Hawaiian or Other Pacific Islander	431	102	94	41	44	134	2	33	68	
White	337,370	60,169	74,086	45,595	42,700	96,659	3,786	22,525	56,441	
Other	54,390	9,465	9,451	9,472	3,931	17,409	345	2,843	9,176	
Total	506,606	94,376	97,330	70,512	56,876	143,228	4,833	40,482	86,825	

Continued. See notes at end of table.

Table 10.36a. Pennsylvania – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	447,516	84,211	88,196	60,072	52,543	124,196	4,365	36,881	77,395
Hispanic or Latino origin	65,434	11,801	10,718	11,402	4,802	21,277	342	3,046	10,446
Total	512,950	96,012	98,914	71,474	57,345	145,473	4,707	39,927	87,841
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴									
Veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Not a veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡

‡ No data submitted.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.36b. Pennsylvania – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020

Selected characteristics		Mental health diagnosis ¹								
		Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender										
Male	46.0	41.2	34.9	67.5	36.8	34.8	32.6	59.2	58.5	
Female	54.0	58.8	65.1	32.5	63.2	65.2	67.4	40.8	41.5	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age ²										
0 to 11 years	16.9	27.6	8.3	47.8	†	1.9	†	†	30.3	
12 to 14 years	8.2	11.5	6.3	18.8	1.2	4.4	†	†	12.7	
15 to 17 years	7.7	9.5	7.6	12.2	2.7	6.8	2.6	†	11.0	
18 to 20 years	4.7	4.4	5.5	5.0	3.8	5.3	5.7	1.9	4.9	
21 to 24 years	5.1	4.6	6.3	3.0	6.3	6.1	10.2	4.5	4.0	
25 to 29 years	8.2	7.2	10.7	3.6	11.9	9.8	14.8	8.6	5.6	
30 to 34 years	8.2	7.3	10.8	2.9	12.7	9.7	14.0	9.0	5.0	
35 to 39 years	7.5	6.4	9.6	2.3	12.4	9.1	12.3	9.4	4.3	
40 to 44 years	6.0	4.8	7.3	1.5	10.4	7.8	8.7	8.0	3.3	
45 to 49 years	6.2	4.5	6.9	1.1	10.2	8.4	8.1	9.5	3.3	
50 to 54 years	6.6	4.4	6.8	†	10.4	9.6	7.3	11.6	3.6	
55 to 59 years	6.3	3.6	6.0	†	8.7	9.3	7.2	13.5	3.4	
60 to 64 years	4.4	2.3	4.0	†	5.3	6.5	4.3	11.0	2.8	
65 years and older	4.0	1.9	4.0	†	3.4	5.5	3.7	11.8	5.9	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race										
American Indian or Alaska Native	†	†	†	†	†	†	†	†	†	
Asian ³	1.1	1.2	†	†	†	1.3	†	1.8	1.2	
Black or African American	21.2	24.6	12.9	21.1	17.1	18.8	13.7	35.3	22.9	
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	†	†	†	
White	66.6	63.8	76.1	64.7	75.1	67.5	78.3	55.6	65.0	
Other	10.7	10.0	9.7	13.4	6.9	12.2	7.1	7.0	10.6	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.36b. Pennsylvania – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020 (continued)

Selected characteristics		Mental health diagnosis¹							
		Trauma- and stressor-related disorders			Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders¹
					Bipolar disorders	Depressive disorders			
Total									
Ethnicity									
Not of Hispanic or Latino origin	87.2	87.7	89.2	84.0	91.6	85.4	92.7	92.4	88.1
Hispanic or Latino origin	12.8	12.3	10.8	16.0	8.4	14.6	7.3	7.6	11.9
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴									
Veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Not a veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡

‡ No data submitted; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.37a. Puerto Rico – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	2,950	88	291	192	226	601	29	1,379	792
Gender									
Male	1,823	53	128	148	92	260	23	888	602
Female	1,127	35	163	44	134	341	6	491	190
Total	2,950	88	291	192	226	601	29	1,379	792
Age²									
0 to 11 years	154	27	11	78	--	6	--	9	86
12 to 14 years	62	8	5	39	3	6	--	--	31
15 to 17 years	112	8	4	44	3	15	--	2	68
18 to 20 years	70	5	5	19	4	16	--	4	44
21 to 24 years	102	5	14	4	16	21	2	36	41
25 to 29 years	190	4	14	2	16	32	1	72	83
30 to 34 years	207	4	17	4	23	32	5	100	59
35 to 39 years	209	7	20	1	19	43	6	105	64
40 to 44 years	212	6	18	1	23	45	6	106	60
45 to 49 years	249	3	30	--	32	53	3	135	44
50 to 54 years	293	2	34	--	21	70	3	172	47
55 to 59 years	371	3	46	--	32	93	2	203	68
60 to 64 years	309	1	29	--	19	71	--	192	38
65 years and older	410	5	44	--	15	98	1	243	59
Total	2,950	88	291	192	226	601	29	1,379	792
Race									
American Indian or Alaska Native	40	3	--	9	1	5	1	14	15
Asian ³	4	1	--	--	--	1	--	2	1
Black or African American	32	1	3	1	4	3	--	14	14
Native Hawaiian or Other Pacific Islander	--	--	--	--	--	--	--	--	--
White	408	35	28	87	19	75	7	102	181
Other	2,323	45	242	82	186	483	19	1,198	531
Total	2,807	85	273	179	210	567	27	1,330	742

Continued. See notes at end of table.

Table 10.37a. Puerto Rico – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020 (continued)

		Mental health diagnosis ¹							
Selected characteristics	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	40	3	3	--	3	10	3	18	15
Hispanic or Latino origin	2,894	85	288	190	223	589	26	1,354	770
Total	2,934	88	291	190	226	599	29	1,372	785
Specific Hispanic or Latino origin									
Mexican	8	--	--	--	1	1	--	7	1
Puerto Rican	2,782	78	279	188	212	571	25	1,290	737
Cuban	10	--	--	--	2	2	--	6	--
Other specific	37	2	3	1	3	6	--	16	12
Total	2,837	80	282	189	218	580	25	1,319	750
Veteran status ⁴									
Veteran	84	8	8	2	4	20	--	32	31
Not a veteran	2,228	23	245	11	192	516	23	1,161	493
Total	2,312	31	253	13	196	536	23	1,193	524

-- Quantity is zero.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.37b. Puerto Rico – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020

Selected characteristics		Mental health diagnosis ¹								
		Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender										
Male	61.8	60.2	44.0	77.1	40.7	43.3	79.3	64.4	76.0	
Female	38.2	39.8	56.0	22.9	59.3	56.7	20.7	35.6	24.0	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age ²										
0 to 11 years	5.2	30.7	3.8	40.6	--	†	--	†	10.9	
12 to 14 years	2.1	9.1	1.7	20.3	1.3	†	--	--	3.9	
15 to 17 years	3.8	9.1	1.4	22.9	1.3	2.5	--	†	8.6	
18 to 20 years	2.4	5.7	1.7	9.9	1.8	2.7	--	†	5.6	
21 to 24 years	3.5	5.7	4.8	2.1	7.1	3.5	6.9	2.6	5.2	
25 to 29 years	6.4	4.5	4.8	1.0	7.1	5.3	3.4	5.2	10.5	
30 to 34 years	7.0	4.5	5.8	2.1	10.2	5.3	17.2	7.3	7.4	
35 to 39 years	7.1	8.0	6.9	†	8.4	7.2	20.7	7.6	8.1	
40 to 44 years	7.2	6.8	6.2	†	10.2	7.5	20.7	7.7	7.6	
45 to 49 years	8.4	3.4	10.3	--	14.2	8.8	10.3	9.8	5.6	
50 to 54 years	9.9	2.3	11.7	--	9.3	11.6	10.3	12.5	5.9	
55 to 59 years	12.6	3.4	15.8	--	14.2	15.5	6.9	14.7	8.6	
60 to 64 years	10.5	1.1	10.0	--	8.4	11.8	--	13.9	4.8	
65 years and older	13.9	5.7	15.1	--	6.6	16.3	3.4	17.6	7.4	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race										
American Indian or Alaska Native	1.4	3.5	--	5.0	†	†	3.7	1.1	2.0	
Asian ³	†	1.2	--	--	--	†	--	†	†	
Black or African American	1.1	1.2	1.1	†	1.9	†	--	1.1	1.9	
Native Hawaiian or Other Pacific Islander	--	--	--	--	--	--	--	--	--	
White	14.5	41.2	10.3	48.6	9.0	13.2	25.9	7.7	24.4	
Other	82.8	52.9	88.6	45.8	88.6	85.2	70.4	90.1	71.6	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.37b. Puerto Rico – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020 (continued)

Selected characteristics		Mental health diagnosis¹								
		Trauma- and stressor-related disorders			Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders¹	
					Bipolar disorders	Depressive disorders				
Total										
Ethnicity										
Not of Hispanic or Latino origin		1.4	3.4	1.0	--	1.3	1.7	10.3	1.3	1.9
Hispanic or Latino origin		98.6	96.6	99.0	100.0	98.7	98.3	89.7	98.7	98.1
Total		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin										
Mexican		†	--	--	--	†	†	--	†	†
Puerto Rican		98.1	97.5	98.9	99.5	97.2	98.4	100.0	97.8	98.3
Cuban		†	--	--	--	†	†	--	†	--
Other specific		1.3	2.5	1.1	†	1.4	1.0	--	1.2	1.6
Total		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁴										
Veteran		3.6	25.8	3.2	15.4	2.0	3.7	--	2.7	5.9
Not a veteran		96.4	74.2	96.8	84.6	98.0	96.3	100.0	97.3	94.1
Total		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

-- Quantity is zero; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.38a. Rhode Island – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	29,366	5,568	10,635	3,724	4,176	13,251	1,487	4,466	5,434
Gender									
Male	14,338	2,492	4,255	2,392	1,715	5,418	531	2,850	3,093
Female	15,027	3,076	6,380	1,332	2,460	7,833	956	1,616	2,341
Total	29,365	5,568	10,635	3,724	4,175	13,251	1,487	4,466	5,434
Age²									
0 to 11 years	2,679	1,051	625	1,036	4	170	2	1	1,074
12 to 14 years	1,759	517	599	689	10	449	1	5	609
15 to 17 years	1,835	481	722	551	28	759	10	21	513
18 to 20 years	1,572	330	695	377	123	779	38	87	400
21 to 24 years	1,583	296	614	235	253	741	96	222	315
25 to 29 years	2,124	386	753	202	386	967	154	407	326
30 to 34 years	2,261	395	845	146	461	1,047	152	425	278
35 to 39 years	2,208	367	830	159	403	1,063	153	423	253
40 to 44 years	1,940	309	738	84	412	1,026	134	356	234
45 to 49 years	2,128	328	801	82	453	1,127	139	397	246
50 to 54 years	2,534	334	927	65	500	1,456	169	485	336
55 to 59 years	2,655	357	977	45	462	1,500	184	578	310
60 to 64 years	1,997	213	730	35	353	1,087	139	471	248
65 years and older	2,091	204	779	18	328	1,080	116	588	292
Total	29,366	5,568	10,635	3,724	4,176	13,251	1,487	4,466	5,434
Race									
American Indian or Alaska Native	225	44	78	27	48	114	17	39	31
Asian ³	304	63	92	15	20	149	5	73	65
Black or African American	3,451	575	1,109	409	451	1,555	136	726	668
Native Hawaiian or Other Pacific Islander	66	17	23	9	6	30	2	8	11
White	20,164	3,683	7,539	2,503	3,161	9,012	1,147	3,039	3,539
Other	1,051	141	381	145	141	572	55	163	195
Total	25,261	4,523	9,222	3,108	3,827	11,432	1,362	4,048	4,509

Continued. See notes at end of table.

Table 10.38a. Rhode Island – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	20,036	3,764	7,238	2,324	3,376	8,703	1,162	3,416	3,254
Hispanic or Latino origin	6,516	1,126	2,525	938	579	3,500	266	767	1,554
Total	26,552	4,890	9,763	3,262	3,955	12,203	1,428	4,183	4,808
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴									
Veteran	474	98	139	21	106	217	33	110	30
Not a veteran	19,319	2,919	7,395	1,241	3,478	10,004	1,343	3,720	2,733
Total	19,793	3,017	7,534	1,262	3,584	10,221	1,376	3,830	2,763

‡ No data submitted.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.38b. Rhode Island – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020

Selected characteristics		Mental health diagnosis ¹								
		Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender										
Male	48.8	44.8	40.0	64.2	41.1	40.9	35.7	63.8	56.9	
Female	51.2	55.2	60.0	35.8	58.9	59.1	64.3	36.2	43.1	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age ²										
0 to 11 years	9.1	18.9	5.9	27.8	†	1.3	†	†	19.8	
12 to 14 years	6.0	9.3	5.6	18.5	†	3.4	†	†	11.2	
15 to 17 years	6.2	8.6	6.8	14.8	†	5.7	†	†	9.4	
18 to 20 years	5.4	5.9	6.5	10.1	2.9	5.9	2.6	1.9	7.4	
21 to 24 years	5.4	5.3	5.8	6.3	6.1	5.6	6.5	5.0	5.8	
25 to 29 years	7.2	6.9	7.1	5.4	9.2	7.3	10.4	9.1	6.0	
30 to 34 years	7.7	7.1	7.9	3.9	11.0	7.9	10.2	9.5	5.1	
35 to 39 years	7.5	6.6	7.8	4.3	9.7	8.0	10.3	9.5	4.7	
40 to 44 years	6.6	5.5	6.9	2.3	9.9	7.7	9.0	8.0	4.3	
45 to 49 years	7.2	5.9	7.5	2.2	10.8	8.5	9.3	8.9	4.5	
50 to 54 years	8.6	6.0	8.7	1.7	12.0	11.0	11.4	10.9	6.2	
55 to 59 years	9.0	6.4	9.2	1.2	11.1	11.3	12.4	12.9	5.7	
60 to 64 years	6.8	3.8	6.9	†	8.5	8.2	9.3	10.5	4.6	
65 years and older	7.1	3.7	7.3	†	7.9	8.2	7.8	13.2	5.4	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race										
American Indian or Alaska Native	†	†	†	†	1.3	†	1.2	†	†	
Asian ³	1.2	1.4	†	†	†	1.3	†	1.8	1.4	
Black or African American	13.7	12.7	12.0	13.2	11.8	13.6	10.0	17.9	14.8	
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	†	†	†	
White	79.8	81.4	81.8	80.5	82.6	78.8	84.2	75.1	78.5	
Other	4.2	3.1	4.1	4.7	3.7	5.0	4.0	4.0	4.3	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.38b. Rhode Island – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020 (continued)

Selected characteristics		Mental health diagnosis¹								
		Trauma- and stressor-related disorders			Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders¹	
					Bipolar disorders	Depressive disorders				
Total										
Ethnicity										
Not of Hispanic or Latino origin	75.5		77.0	74.1	71.2	85.4	71.3	81.4	81.7	67.7
Hispanic or Latino origin	24.5		23.0	25.9	28.8	14.6	28.7	18.6	18.3	32.3
Total	100.0		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin										
Mexican	‡		‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡		‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡		‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡		‡	‡	‡	‡	‡	‡	‡	‡
Total	‡		‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴										
Veteran	2.4		3.2	1.8	1.7	3.0	2.1	2.4	2.9	1.1
Not a veteran	97.6		96.8	98.2	98.3	97.0	97.9	97.6	97.1	98.9
Total	100.0		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

‡ No data submitted; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.39a. South Carolina – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	95,263	25,244	22,886	16,762	12,157	29,930	4,420	17,323	34,166
Gender									
Male	44,536	9,535	8,022	11,305	4,093	10,003	1,148	10,179	20,122
Female	50,725	15,709	14,863	5,457	8,064	19,927	3,272	7,144	14,043
Total	95,261	25,244	22,885	16,762	12,157	29,930	4,420	17,323	34,165
Age²									
0 to 11 years	13,898	4,923	2,087	7,786	28	830	1	39	5,832
12 to 14 years	9,568	3,157	2,070	3,907	93	2,375	9	65	3,835
15 to 17 years	9,610	2,951	2,580	2,740	247	3,530	35	158	3,933
18 to 20 years	4,453	1,180	1,391	773	405	1,970	177	365	1,711
21 to 24 years	4,661	1,175	1,393	315	773	1,908	398	894	1,789
25 to 29 years	6,488	1,623	1,791	271	1,178	2,432	551	1,560	2,557
30 to 34 years	6,628	1,778	1,807	230	1,292	2,322	548	1,667	2,563
35 to 39 years	6,271	1,633	1,690	202	1,262	2,142	460	1,799	2,297
40 to 44 years	5,582	1,433	1,377	129	1,236	1,908	463	1,623	1,912
45 to 49 years	5,634	1,422	1,444	127	1,300	2,107	467	1,561	1,809
50 to 54 years	5,981	1,438	1,463	103	1,254	2,328	463	1,732	1,857
55 to 59 years	6,353	1,249	1,573	94	1,294	2,443	398	2,005	1,795
60 to 64 years	4,749	738	1,072	43	940	1,698	235	1,726	1,201
65 years and older	5,387	544	1,148	42	855	1,937	215	2,129	1,075
Total	95,263	25,244	22,886	16,762	12,157	29,930	4,420	17,323	34,166
Race									
American Indian or Alaska Native	412	112	97	54	51	149	19	61	163
Asian ³	317	65	84	25	49	89	5	86	97
Black or African American	36,304	7,920	5,313	7,173	2,975	9,380	997	10,678	13,831
Native Hawaiian or Other Pacific Islander	13	4	1	1	3	2	2	2	1
White	48,826	13,996	15,157	7,473	8,517	17,327	3,191	5,883	16,896
Other	7,517	2,630	1,812	1,699	435	2,315	171	440	2,465
Total	93,389	24,727	22,464	16,425	12,030	29,262	4,385	17,150	33,453

Continued. See notes at end of table.

Table 10.39a. South Carolina – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	69,871	18,331	16,922	12,108	9,458	22,022	3,425	13,168	25,399
Hispanic or Latino origin	3,772	1,295	897	639	308	1,191	125	412	1,154
Total	73,643	19,626	17,819	12,747	9,766	23,213	3,550	13,580	26,553
Specific Hispanic or Latino origin									
Mexican	1,140	389	262	192	65	357	27	103	318
Puerto Rican	329	104	88	62	41	118	16	34	124
Cuban	348	85	80	26	65	96	17	106	108
Other specific	1,938	714	465	355	134	617	65	164	599
Total	3,755	1,292	895	635	305	1,188	125	407	1,149
Veteran status⁴									
Veteran	898	237	182	15	180	289	57	276	311
Not a veteran	61,263	13,973	15,966	2,314	11,602	22,902	4,318	16,771	20,251
Total	62,161	14,210	16,148	2,329	11,782	23,191	4,375	17,047	20,562

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.39b. South Carolina – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020

Selected characteristics		Mental health diagnosis¹								
		Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders¹
						Bipolar disorders	Depressive disorders			
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender										
Male	46.8	37.8	35.1	67.4	33.7	33.4	26.0	58.8	58.9	
Female	53.2	62.2	64.9	32.6	66.3	66.6	74.0	41.2	41.1	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age²										
0 to 11 years	14.6	19.5	9.1	46.5	†	2.8	†	†	17.1	
12 to 14 years	10.0	12.5	9.0	23.3	†	7.9	†	†	11.2	
15 to 17 years	10.1	11.7	11.3	16.3	2.0	11.8	†	†	11.5	
18 to 20 years	4.7	4.7	6.1	4.6	3.3	6.6	4.0	2.1	5.0	
21 to 24 years	4.9	4.7	6.1	1.9	6.4	6.4	9.0	5.2	5.2	
25 to 29 years	6.8	6.4	7.8	1.6	9.7	8.1	12.5	9.0	7.5	
30 to 34 years	7.0	7.0	7.9	1.4	10.6	7.8	12.4	9.6	7.5	
35 to 39 years	6.6	6.5	7.4	1.2	10.4	7.2	10.4	10.4	6.7	
40 to 44 years	5.9	5.7	6.0	†	10.2	6.4	10.5	9.4	5.6	
45 to 49 years	5.9	5.6	6.3	†	10.7	7.0	10.6	9.0	5.3	
50 to 54 years	6.3	5.7	6.4	†	10.3	7.8	10.5	10.0	5.4	
55 to 59 years	6.7	4.9	6.9	†	10.6	8.2	9.0	11.6	5.3	
60 to 64 years	5.0	2.9	4.7	†	7.7	5.7	5.3	10.0	3.5	
65 years and older	5.7	2.2	5.0	†	7.0	6.5	4.9	12.3	3.1	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race										
American Indian or Alaska Native	†	†	†	†	†	†	†	†	†	
Asian³	†	†	†	†	†	†	†	†	†	
Black or African American	38.9	32.0	23.7	43.7	24.7	32.1	22.7	62.3	41.3	
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	†	†	†	
White	52.3	56.6	67.5	45.5	70.8	59.2	72.8	34.3	50.5	
Other	8.0	10.6	8.1	10.3	3.6	7.9	3.9	2.6	7.4	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

**Table 10.39b. South Carolina – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020
(continued)**

Mental health diagnosis ¹										
Selected characteristics		Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Ethnicity										
Not of Hispanic or Latino origin	94.9	93.4	95.0	95.0	96.8	94.9	96.5	97.0	95.7	
Hispanic or Latino origin	5.1	6.6	5.0	5.0	3.2	5.1	3.5	3.0	4.3	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	
Specific Hispanic or Latino origin										
Mexican	30.4	30.1	29.3	30.2	21.3	30.1	21.6	25.3	27.7	
Puerto Rican	8.8	8.0	9.8	9.8	13.4	9.9	12.8	8.4	10.8	
Cuban	9.3	6.6	8.9	4.1	21.3	8.1	13.6	26.0	9.4	
Other specific	51.6	55.3	52.0	55.9	43.9	51.9	52.0	40.3	52.1	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	
Veteran status ⁴										
Veteran	1.4	1.7	1.1	†	1.5	1.2	1.3	1.6	1.5	
Not a veteran	98.6	98.3	98.9	99.4	98.5	98.8	98.7	98.4	98.5	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.40a. South Dakota – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	14,789	4,121	1,696	1,126	1,350	3,621	51	1,345	1,821
Gender									
Male	6,644	1,830	645	804	500	1,201	18	810	995
Female	8,145	2,291	1,051	322	850	2,420	33	535	826
Total	14,789	4,121	1,696	1,126	1,350	3,621	51	1,345	1,821
Age²									
0 to 11 years	2,316	1,469	131	390	10	58	--	--	286
12 to 14 years	1,483	717	122	256	17	189	--	1	205
15 to 17 years	1,377	540	150	163	29	303	--	2	211
18 to 20 years	674	162	100	69	33	192	2	10	122
21 to 24 years	824	154	127	54	75	222	1	37	171
25 to 29 years	1,074	187	175	71	117	323	2	81	148
30 to 34 years	1,040	194	170	50	129	294	3	110	116
35 to 39 years	1,064	151	175	17	149	334	1	120	136
40 to 44 years	850	123	121	18	143	254	1	121	83
45 to 49 years	805	104	117	12	133	253	11	122	78
50 to 54 years	752	104	71	11	138	245	8	140	60
55 to 59 years	889	83	66	7	159	337	8	197	59
60 to 64 years	696	61	64	2	102	240	11	195	51
65 years and older	945	72	107	6	116	377	3	209	95
Total	14,789	4,121	1,696	1,126	1,350	3,621	51	1,345	1,821
Race									
American Indian or Alaska Native	3	--	--	--	--	--	--	1	2
Asian ³	103	36	13	6	7	19	--	10	14
Black or African American	459	211	26	38	23	50	1	61	54
Native Hawaiian or Other Pacific Islander	23	3	7	2	2	8	--	1	2
White	10,684	2,555	1,334	823	1,117	2,785	49	1,028	1,250
Other	3,517	1,316	316	257	201	759	1	244	499
Total	14,789	4,121	1,696	1,126	1,350	3,621	51	1,345	1,821

Continued. See notes at end of table.

Table 10.40a. South Dakota – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	14,082	3,846	1,628	1,074	1,297	3,486	50	1,312	1,718
Hispanic or Latino origin	707	275	68	52	53	135	1	33	103
Total	14,789	4,121	1,696	1,126	1,350	3,621	51	1,345	1,821
Specific Hispanic or Latino origin									
Mexican	227	85	22	20	20	43	--	7	36
Puerto Rican	60	20	4	6	3	14	--	4	9
Cuban	11	3	1	2	1	1	--	3	--
Other specific	118	54	10	7	9	17	--	4	18
Total	416	162	37	35	33	75	--	18	63
Veteran status⁴									
Veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Not a veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡

-- Quantity is zero; ‡ No data submitted.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.40b. South Dakota – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020

Selected characteristics		Mental health diagnosis¹										
		Trauma- and stressor-related disorders				Mood disorders		Personality disorders			Schizophrenia, other psychotic disorders	Other mental disorders¹
						Bipolar disorders	Depressive disorders					
Total		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	
Total individuals		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	
Gender												
Male		44.9	44.4	38.0	71.4	37.0	33.2	35.3	60.2	54.6		
Female		55.1	55.6	62.0	28.6	63.0	66.8	64.7	39.8	45.4		
	Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0		
Age²												
0 to 11 years		15.7	35.6	7.7	34.6	†	1.6	--	--	15.7		
12 to 14 years		10.0	17.4	7.2	22.7	1.3	5.2	--	†	11.3		
15 to 17 years		9.3	13.1	8.8	14.5	2.1	8.4	--	†	11.6		
18 to 20 years		4.6	3.9	5.9	6.1	2.4	5.3	3.9	†	6.7		
21 to 24 years		5.6	3.7	7.5	4.8	5.6	6.1	2.0	2.8	9.4		
25 to 29 years		7.3	4.5	10.3	6.3	8.7	8.9	3.9	6.0	8.1		
30 to 34 years		7.0	4.7	10.0	4.4	9.6	8.1	5.9	8.2	6.4		
35 to 39 years		7.2	3.7	10.3	1.5	11.0	9.2	2.0	8.9	7.5		
40 to 44 years		5.7	3.0	7.1	1.6	10.6	7.0	2.0	9.0	4.6		
45 to 49 years		5.4	2.5	6.9	1.1	9.9	7.0	21.6	9.1	4.3		
50 to 54 years		5.1	2.5	4.2	†	10.2	6.8	15.7	10.4	3.3		
55 to 59 years		6.0	2.0	3.9	†	11.8	9.3	15.7	14.6	3.2		
60 to 64 years		4.7	1.5	3.8	†	7.6	6.6	21.6	14.5	2.8		
65 years and older		6.4	1.7	6.3	†	8.6	10.4	5.9	15.5	5.2		
	Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0		
Race												
American Indian or Alaska Native		†	--	--	--	--	--	--	†	†		
Asian³		†	†	†	†	†	†	--	†	†		
Black or African American		3.1	5.1	1.5	3.4	1.7	1.4	2.0	4.5	3.0		
Native Hawaiian or Other Pacific Islander		†	†	†	†	†	†	--	†	†		
White		72.2	62.0	78.7	73.1	82.7	76.9	96.1	76.4	68.6		
Other		23.8	31.9	18.6	22.8	14.9	21.0	2.0	18.1	27.4		
	Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0		

Continued. See notes at end of table.

Table 10.40b. South Dakota – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020 (continued)

Selected characteristics		Mental health diagnosis ¹								
		Trauma- and stressor-related disorders			Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹	
					Bipolar disorders	Depressive disorders				
Total										
Ethnicity										
Not of Hispanic or Latino origin		95.2	93.3	96.0	95.4	96.1	96.3	98.0	97.5	94.3
Hispanic or Latino origin		4.8	6.7	4.0	4.6	3.9	3.7	2.0	2.5	5.7
Total		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin										
Mexican		54.6	52.5	59.5	57.1	60.6	57.3	--	38.9	57.1
Puerto Rican		14.4	12.3	10.8	17.1	9.1	18.7	--	22.2	14.3
Cuban		2.6	1.9	2.7	5.7	3.0	1.3	--	16.7	--
Other specific		28.4	33.3	27.0	20.0	27.3	22.7	--	22.2	28.6
Total		100.0	100.0	100.0	100.0	100.0	100.0	--	100.0	100.0
Veteran status ⁴										
Veteran		‡	‡	‡	‡	‡	‡	‡	‡	‡
Not a veteran		‡	‡	‡	‡	‡	‡	‡	‡	‡
Total		‡	‡	‡	‡	‡	‡	‡	‡	‡

-- Quantity is zero; ‡ No data submitted; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.41a. Tennessee – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals²	62,819	13,862	16,477	3,959	12,487	27,003	1,175	8,154	7,358
Gender									
Male	30,748	5,820	6,794	2,610	5,483	12,173	549	5,391	4,292
Female	31,925	7,984	9,655	1,326	6,981	14,783	625	2,747	3,047
Total	62,673	13,804	16,449	3,936	12,464	26,956	1,174	8,138	7,339
Age³									
0 to 11 years	4,076	1,740	624	1,862	11	240	1	27	1,544
12 to 14 years	2,246	761	502	700	38	639	3	31	791
15 to 17 years	2,093	606	645	368	111	899	17	25	628
18 to 20 years	2,226	480	647	169	392	1,057	71	238	386
21 to 24 years	4,587	872	1,307	140	980	2,083	111	641	491
25 to 29 years	7,627	1,532	2,081	215	1,744	3,371	153	1,217	664
30 to 34 years	7,421	1,541	2,037	158	1,816	3,141	164	1,200	585
35 to 39 years	6,835	1,456	1,765	106	1,757	2,877	171	1,050	510
40 to 44 years	6,225	1,363	1,566	80	1,592	2,780	145	960	425
45 to 49 years	6,041	1,291	1,614	66	1,427	2,837	106	824	418
50 to 54 years	5,657	1,094	1,557	46	1,238	2,901	96	729	353
55 to 59 years	4,550	728	1,275	24	836	2,515	74	590	304
60 to 64 years	2,424	304	654	21	408	1,297	43	406	174
65 years and older	811	94	203	4	137	366	20	216	85
Total	62,819	13,862	16,477	3,959	12,487	27,003	1,175	8,154	7,358
Race									
American Indian or Alaska Native	149	39	25	9	29	52	3	25	28
Asian ⁴	98	19	25	1	13	44	2	24	15
Black or African American	16,795	4,168	3,057	1,989	2,376	5,718	219	3,528	2,911
Native Hawaiian or Other Pacific Islander	96	11	24	7	11	38	2	24	5
White	44,091	9,330	13,020	1,872	9,756	20,420	936	4,300	4,235
Other	450	91	80	32	99	179	6	107	54
Total	61,679	13,658	16,231	3,910	12,284	26,451	1,168	8,008	7,248

Continued. See notes at end of table.

Table 10.41a. Tennessee – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	27,754	6,018	6,292	1,973	6,384	11,522	815	5,416	4,230
Hispanic or Latino origin	584	125	123	40	112	246	10	135	96
Total	28,338	6,143	6,415	2,013	6,496	11,768	825	5,551	4,326
Specific Hispanic or Latino origin									
Mexican	24	7	1	--	4	10	--	10	2
Puerto Rican	--	--	--	--	--	--	--	--	--
Cuban	1	--	--	--	--	1	--	--	1
Other specific	54	16	4	--	8	26	--	25	7
Total	79	23	5	--	12	37	--	35	10
Veteran status⁵									
Veteran	372	86	82	7	88	178	14	71	32
Not a veteran	17,336	3,535	3,859	333	4,598	7,573	632	3,952	2,102
Total	17,708	3,621	3,941	340	4,686	7,751	646	4,023	2,134

-- Quantity is zero.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Tennessee reports only a subset of the total individuals served through their SMHAs.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.41b. Tennessee – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020

Selected characteristics		Mental health diagnosis ¹							
		Trauma- and stressor-related disorders			Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals ²	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender									
Male	49.1	42.2	41.3	66.3	44.0	45.2	46.8	66.2	58.5
Female	50.9	57.8	58.7	33.7	56.0	54.8	53.2	33.8	41.5
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age ³									
0 to 11 years	6.5	12.6	3.8	47.0	†	†	†	†	21.0
12 to 14 years	3.6	5.5	3.0	17.7	†	2.4	†	†	10.8
15 to 17 years	3.3	4.4	3.9	9.3	†	3.3	1.4	†	8.5
18 to 20 years	3.5	3.5	3.9	4.3	3.1	3.9	6.0	2.9	5.2
21 to 24 years	7.3	6.3	7.9	3.5	7.8	7.7	9.4	7.9	6.7
25 to 29 years	12.1	11.1	12.6	5.4	14.0	12.5	13.0	14.9	9.0
30 to 34 years	11.8	11.1	12.4	4.0	14.5	11.6	14.0	14.7	8.0
35 to 39 years	10.9	10.5	10.7	2.7	14.1	10.7	14.6	12.9	6.9
40 to 44 years	9.9	9.8	9.5	2.0	12.7	10.3	12.3	11.8	5.8
45 to 49 years	9.6	9.3	9.8	1.7	11.4	10.5	9.0	10.1	5.7
50 to 54 years	9.0	7.9	9.4	1.2	9.9	10.7	8.2	8.9	4.8
55 to 59 years	7.2	5.3	7.7	†	6.7	9.3	6.3	7.2	4.1
60 to 64 years	3.9	2.2	4.0	†	3.3	4.8	3.7	5.0	2.4
65 years and older	1.3	†	1.2	†	1.1	1.4	1.7	2.6	1.2
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race									
American Indian or Alaska Native	†	†	†	†	†	†	†	†	†
Asian ⁴	†	†	†	†	†	†	†	†	†
Black or African American	27.2	30.5	18.8	50.9	19.3	21.6	18.8	44.1	40.2
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	†	†	†
White	71.5	68.3	80.2	47.9	79.4	77.2	80.1	53.7	58.4
Other	†	†	†	†	†	†	†	1.3	†
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.41b. Tennessee – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020 (continued)

Selected characteristics		Mental health diagnosis ¹							
					Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD					
Ethnicity									
Not of Hispanic or Latino origin	97.9	98.0	98.1	98.0	98.3	97.9	98.8	97.6	97.8
Hispanic or Latino origin	2.1	2.0	1.9	2.0	1.7	2.1	1.2	2.4	2.2
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	30.4	30.4	20.0	--	33.3	27.0	--	28.6	20.0
Puerto Rican	--	--	--	--	--	--	--	--	--
Cuban	1.3	--	--	--	--	2.7	--	--	10.0
Other specific	68.4	69.6	80.0	--	66.7	70.3	--	71.4	70.0
Total	100.0	100.0	100.0	--	100.0	100.0	--	100.0	100.0
Veteran status⁵									
Veteran	2.1	2.4	2.1	2.1	1.9	2.3	2.2	1.8	1.5
Not a veteran	97.9	97.6	97.9	97.9	98.1	97.7	97.8	98.2	98.5
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

-- Quantity is zero; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Tennessee reports only a subset of the total individuals served through their SMHAs.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.42a. Texas – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	283,773	59,384	80,964	42,610	73,794	108,379	11,474	54,059	42,920
Gender									
Male	138,706	21,339	30,845	28,948	30,227	42,681	4,862	34,226	27,388
Female	145,067	38,045	50,119	13,662	43,567	65,698	6,612	19,833	15,532
Total	283,773	59,384	80,964	42,610	73,794	108,379	11,474	54,059	42,920
Age²									
0 to 11 years	27,882	6,581	3,285	19,891	321	2,199	6	218	13,493
12 to 14 years	18,608	3,906	3,921	9,821	829	6,183	33	250	7,875
15 to 17 years	19,348	3,904	5,805	6,868	1,836	9,158	97	501	6,656
18 to 20 years	12,621	2,395	4,247	1,783	3,408	6,689	409	1,381	1,788
21 to 24 years	17,413	3,665	5,758	911	5,565	8,150	809	3,128	1,696
25 to 29 years	26,223	5,639	8,458	892	8,756	10,904	1,365	5,935	1,907
30 to 34 years	26,751	6,077	8,590	740	9,438	10,293	1,525	6,442	1,660
35 to 39 years	25,617	6,106	8,107	538	9,264	9,527	1,542	6,454	1,457
40 to 44 years	21,777	5,164	6,754	319	7,851	8,087	1,302	5,634	1,165
45 to 49 years	20,445	4,822	6,615	271	7,065	8,134	1,151	5,225	1,027
50 to 54 years	21,135	4,523	6,387	237	6,908	8,736	1,161	5,579	1,126
55 to 59 years	20,763	3,646	5,981	190	6,264	8,850	1,069	5,707	1,119
60 to 64 years	14,516	2,027	4,104	104	3,822	6,496	661	4,287	867
65 years and older	10,674	929	2,952	45	2,467	4,973	344	3,318	1,084
Total	283,773	59,384	80,964	42,610	73,794	108,379	11,474	54,059	42,920
Race									
American Indian or Alaska Native	969	266	250	92	272	394	37	186	98
Asian ³	2,513	316	546	141	460	886	60	938	264
Black or African American	63,532	11,053	12,364	10,723	14,492	19,056	2,354	18,323	10,885
Native Hawaiian or Other Pacific Islander	291	57	91	32	71	123	6	56	27
White	203,123	44,676	64,256	29,007	55,727	82,797	8,623	32,457	29,242
Other	13,334	3,016	3,456	2,614	2,772	5,118	394	2,094	2,404
Total	283,762	59,384	80,963	42,609	73,794	108,374	11,474	54,054	42,920

Continued. See notes at end of table.

Table 10.42a. Texas – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	202,629	43,363	56,656	27,538	57,982	73,495	9,214	41,661	28,567
Hispanic or Latino origin	81,135	16,020	24,308	15,071	15,811	34,881	2,259	12,393	14,353
Total	283,764	59,383	80,964	42,609	73,793	108,376	11,473	54,054	42,920
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴									
Veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Not a veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡

‡ No data submitted.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.42b. Texas – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020

Selected characteristics		Mental health diagnosis¹							
		Trauma- and stressor-related disorders			Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders¹
					Bipolar disorders	Depressive disorders			
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender									
Male	48.9	35.9	38.1	67.9	41.0	39.4	42.4	63.3	63.8
Female	51.1	64.1	61.9	32.1	59.0	60.6	57.6	36.7	36.2
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age²									
0 to 11 years	9.8	11.1	4.1	46.7	†	2.0	†	†	31.4
12 to 14 years	6.6	6.6	4.8	23.0	1.1	5.7	†	†	18.3
15 to 17 years	6.8	6.6	7.2	16.1	2.5	8.4	†	†	15.5
18 to 20 years	4.4	4.0	5.2	4.2	4.6	6.2	3.6	2.6	4.2
21 to 24 years	6.1	6.2	7.1	2.1	7.5	7.5	7.1	5.8	4.0
25 to 29 years	9.2	9.5	10.4	2.1	11.9	10.1	11.9	11.0	4.4
30 to 34 years	9.4	10.2	10.6	1.7	12.8	9.5	13.3	11.9	3.9
35 to 39 years	9.0	10.3	10.0	1.3	12.6	8.8	13.4	11.9	3.4
40 to 44 years	7.7	8.7	8.3	†	10.6	7.5	11.3	10.4	2.7
45 to 49 years	7.2	8.1	8.2	†	9.6	7.5	10.0	9.7	2.4
50 to 54 years	7.4	7.6	7.9	†	9.4	8.1	10.1	10.3	2.6
55 to 59 years	7.3	6.1	7.4	†	8.5	8.2	9.3	10.6	2.6
60 to 64 years	5.1	3.4	5.1	†	5.2	6.0	5.8	7.9	2.0
65 years and older	3.8	1.6	3.6	†	3.3	4.6	3.0	6.1	2.5
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race									
American Indian or Alaska Native	†	†	†	†	†	†	†	†	†
Asian³	†	†	†	†	†	†	†	1.7	†
Black or African American	22.4	18.6	15.3	25.2	19.6	17.6	20.5	33.9	25.4
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	†	†	†
White	71.6	75.2	79.4	68.1	75.5	76.4	75.2	60.0	68.1
Other	4.7	5.1	4.3	6.1	3.8	4.7	3.4	3.9	5.6
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.42b. Texas – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020 (continued)

Selected characteristics		Mental health diagnosis ¹							
		Trauma- and stressor-related disorders			Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total									
Ethnicity									
Not of Hispanic or Latino origin	71.4	73.0	70.0	64.6	78.6	67.8	80.3	77.1	66.6
Hispanic or Latino origin	28.6	27.0	30.0	35.4	21.4	32.2	19.7	22.9	33.4
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status ⁴									
Veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Not a veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡

‡ No data submitted; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.43a. Utah – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	50,333	10,875	21,525	5,139	4,652	16,790	2,356	6,068	11,239
Gender									
Male	23,074	4,532	8,314	3,460	1,700	6,158	754	3,752	6,621
Female	27,255	6,342	13,210	1,679	2,951	10,632	1,602	2,316	4,617
Total	50,329	10,874	21,524	5,139	4,651	16,790	2,356	6,068	11,238
Age²									
0 to 11 years	10,211	3,767	3,507	2,286	31	1,135	2	22	3,096
12 to 14 years	5,180	1,309	2,172	860	72	1,930	16	53	1,395
15 to 17 years	4,162	829	1,931	467	168	1,939	46	115	1,013
18 to 20 years	1,985	320	985	171	209	876	104	168	509
21 to 24 years	2,639	467	1,237	150	324	1,053	154	338	616
25 to 29 years	4,047	699	1,920	260	509	1,512	227	622	799
30 to 34 years	3,967	674	1,871	220	567	1,444	278	636	695
35 to 39 years	3,988	719	1,862	209	562	1,483	276	722	649
40 to 44 years	3,326	580	1,457	184	566	1,174	268	723	540
45 to 49 years	2,693	444	1,185	120	444	994	255	626	457
50 to 54 years	2,352	351	1,012	81	381	869	186	585	415
55 to 59 years	2,323	317	995	66	360	911	222	603	396
60 to 64 years	1,613	201	651	36	238	630	184	441	291
65 years and older	1,847	198	740	29	221	840	138	414	368
Total	50,333	10,875	21,525	5,139	4,652	16,790	2,356	6,068	11,239
Race									
American Indian or Alaska Native	1,020	306	365	61	55	350	30	133	203
Asian ³	331	50	114	16	20	118	11	83	53
Black or African American	1,085	267	338	115	95	305	35	218	253
Native Hawaiian or Other Pacific Islander	366	88	121	42	30	106	14	60	78
White	40,242	8,745	17,961	4,277	3,976	13,825	2,027	4,626	8,488
Other	5,326	1,218	2,229	569	351	1,691	178	624	1,160
Total	48,370	10,674	21,128	5,080	4,527	16,395	2,295	5,744	10,235

Continued. See notes at end of table.

Table 10.43a. Utah – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	40,939	8,707	18,049	4,312	4,134	14,004	2,118	5,195	8,634
Hispanic or Latino origin	7,298	1,914	2,994	748	380	2,344	191	628	1,573
Total	48,237	10,621	21,043	5,060	4,514	16,348	2,309	5,823	10,207
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴									
Veteran	709	149	290	37	98	279	52	141	97
Not a veteran	28,056	4,621	13,390	1,481	4,175	11,174	2,177	5,398	4,454
Total	28,765	4,770	13,680	1,518	4,273	11,453	2,229	5,539	4,551

‡ No data submitted.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.43b. Utah – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender									
Male	45.8	41.7	38.6	67.3	36.6	36.7	32.0	61.8	58.9
Female	54.2	58.3	61.4	32.7	63.4	63.3	68.0	38.2	41.1
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age²									
0 to 11 years	20.3	34.6	16.3	44.5	†	6.8	†	†	27.5
12 to 14 years	10.3	12.0	10.1	16.7	1.5	11.5	†	†	12.4
15 to 17 years	8.3	7.6	9.0	9.1	3.6	11.5	2.0	1.9	9.0
18 to 20 years	3.9	2.9	4.6	3.3	4.5	5.2	4.4	2.8	4.5
21 to 24 years	5.2	4.3	5.7	2.9	7.0	6.3	6.5	5.6	5.5
25 to 29 years	8.0	6.4	8.9	5.1	10.9	9.0	9.6	10.3	7.1
30 to 34 years	7.9	6.2	8.7	4.3	12.2	8.6	11.8	10.5	6.2
35 to 39 years	7.9	6.6	8.7	4.1	12.1	8.8	11.7	11.9	5.8
40 to 44 years	6.6	5.3	6.8	3.6	12.2	7.0	11.4	11.9	4.8
45 to 49 years	5.4	4.1	5.5	2.3	9.5	5.9	10.8	10.3	4.1
50 to 54 years	4.7	3.2	4.7	1.6	8.2	5.2	7.9	9.6	3.7
55 to 59 years	4.6	2.9	4.6	1.3	7.7	5.4	9.4	9.9	3.5
60 to 64 years	3.2	1.8	3.0	†	5.1	3.8	7.8	7.3	2.6
65 years and older	3.7	1.8	3.4	†	4.8	5.0	5.9	6.8	3.3
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race									
American Indian or Alaska Native	2.1	2.9	1.7	1.2	1.2	2.1	1.3	2.3	2.0
Asian ³	†	†	†	†	†	†	†	1.4	†
Black or African American	2.2	2.5	1.6	2.3	2.1	1.9	1.5	3.8	2.5
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	†	1.0	†
White	83.2	81.9	85.0	84.2	87.8	84.3	88.3	80.5	82.9
Other	11.0	11.4	10.5	11.2	7.8	10.3	7.8	10.9	11.3
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.43b. Utah – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020 (continued)

Mental health diagnosis ¹												
Selected characteristics			Trauma- and stressor-related disorders			Mood disorders		Personality disorders				
						Schizophrenia, other psychotic disorders						
Total			Anxiety disorders			ADHD		Bipolar disorders		Depressive disorders		
Ethnicity												
Not of Hispanic or Latino origin			84.9			82.0			85.8		85.2	
Hispanic or Latino origin			15.1			18.0			14.2		14.8	
Total			100.0			100.0			100.0		100.0	
Specific Hispanic or Latino origin												
Mexican			‡			‡			‡		‡	
Puerto Rican			‡			‡			‡		‡	
Cuban			‡			‡			‡		‡	
Other specific			‡			‡			‡		‡	
Total			‡			‡			‡		‡	
Veteran status ⁴												
Veteran			2.5			3.1			2.1		2.4	
Not a veteran			97.5			96.9			97.9		97.6	
Total			100.0			100.0			100.0		100.0	

‡ No data submitted; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.44a. Vermont – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	24,940	9,847	4,892	2,593	1,334	5,154	763	1,847	9,288
Gender									
Male	12,645	4,411	2,129	1,778	605	2,033	202	1,157	5,454
Female	12,210	5,393	2,757	812	724	3,098	559	684	3,815
Total	24,855	9,804	4,886	2,590	1,329	5,131	761	1,841	9,269
Age²									
0 to 11 years	4,542	2,660	790	978	5	142	--	2	1,234
12 to 14 years	2,381	1,125	593	581	7	379	1	6	717
15 to 17 years	2,343	1,006	662	455	27	604	7	16	742
18 to 20 years	1,542	547	411	125	41	417	26	31	548
21 to 24 years	1,485	484	295	62	74	372	55	73	611
25 to 29 years	1,716	535	296	55	111	383	63	152	853
30 to 34 years	1,674	570	297	78	137	390	62	157	783
35 to 39 years	1,469	549	250	67	126	347	72	173	668
40 to 44 years	1,139	375	219	42	118	278	67	155	496
45 to 49 years	1,177	404	214	44	129	299	71	153	493
50 to 54 years	1,216	413	208	30	122	337	99	182	519
55 to 59 years	1,344	418	202	30	151	378	94	232	564
60 to 64 years	1,145	321	178	26	130	326	81	224	463
65 years and older	1,767	440	277	20	156	502	65	291	597
Total	24,940	9,847	4,892	2,593	1,334	5,154	763	1,847	9,288
Race									
American Indian or Alaska Native	228	105	57	25	9	51	7	17	68
Asian ³	170	70	26	17	8	34	2	18	57
Black or African American	557	273	89	101	19	92	9	55	149
Native Hawaiian or Other Pacific Islander	--	--	--	--	--	--	--	--	--
White	17,538	7,812	3,994	2,140	1,124	4,141	667	1,493	5,438
Other	545	286	111	81	19	109	15	29	153
Total	19,038	8,546	4,277	2,364	1,179	4,427	700	1,612	5,865

Continued. See notes at end of table.

Table 10.44a. Vermont – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	16,232	7,124	3,708	2,059	1,037	3,764	633	1,469	5,044
Hispanic or Latino origin	715	357	158	122	29	150	17	34	204
Total	16,947	7,481	3,866	2,181	1,066	3,914	650	1,503	5,248
Specific Hispanic or Latino origin									
Mexican	113	55	33	11	3	22	2	3	30
Puerto Rican	147	78	25	33	5	34	- -	4	46
Cuban	18	4	7	3	- -	2	- -	3	7
Other specific	174	91	40	30	6	41	6	5	41
Total	452	228	105	77	14	99	8	15	124
Veteran status⁴									
Veteran	432	160	67	15	48	117	20	83	152
Not a veteran	9,134	3,585	2,090	467	939	2,811	613	1,333	3,080
Total	9,566	3,745	2,157	482	987	2,928	633	1,416	3,232

- - Quantity is zero.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.44b. Vermont – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020

Selected characteristics		Mental health diagnosis¹										
		Trauma- and stressor-related disorders				Mood disorders		Personality disorders			Schizophrenia, other psychotic disorders	Other mental disorders¹
						Bipolar disorders	Depressive disorders					
Total		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	
Total individuals		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	
Gender												
Male	50.9	45.0	43.6	68.6	45.5	39.6	26.5	62.8	58.8			
Female	49.1	55.0	56.4	31.4	54.5	60.4	73.5	37.2	41.2			
Total		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0		
Age²												
0 to 11 years	18.2	27.0	16.1	37.7	†	2.8	--	†	13.3			
12 to 14 years	9.5	11.4	12.1	22.4	†	7.4	†	†	7.7			
15 to 17 years	9.4	10.2	13.5	17.5	2.0	11.7	†	†	8.0			
18 to 20 years	6.2	5.6	8.4	4.8	3.1	8.1	3.4	1.7	5.9			
21 to 24 years	6.0	4.9	6.0	2.4	5.5	7.2	7.2	4.0	6.6			
25 to 29 years	6.9	5.4	6.1	2.1	8.3	7.4	8.3	8.2	9.2			
30 to 34 years	6.7	5.8	6.1	3.0	10.3	7.6	8.1	8.5	8.4			
35 to 39 years	5.9	5.6	5.1	2.6	9.4	6.7	9.4	9.4	7.2			
40 to 44 years	4.6	3.8	4.5	1.6	8.8	5.4	8.8	8.4	5.3			
45 to 49 years	4.7	4.1	4.4	1.7	9.7	5.8	9.3	8.3	5.3			
50 to 54 years	4.9	4.2	4.3	1.2	9.1	6.5	13.0	9.9	5.6			
55 to 59 years	5.4	4.2	4.1	1.2	11.3	7.3	12.3	12.6	6.1			
60 to 64 years	4.6	3.3	3.6	1.0	9.7	6.3	10.6	12.1	5.0			
65 years and older	7.1	4.5	5.7	†	11.7	9.7	8.5	15.8	6.4			
Total		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0		
Race												
American Indian or Alaska Native	1.2	1.2	1.3	1.1	†	1.2	1.0	1.1	1.2			
Asian³	†	†	†	†	†	†	†	1.1	†			
Black or African American	2.9	3.2	2.1	4.3	1.6	2.1	1.3	3.4	2.5			
Native Hawaiian or Other Pacific Islander	--	--	--	--	--	--	--	--	--			
White	92.1	91.4	93.4	90.5	95.3	93.5	95.3	92.6	92.7			
Other	2.9	3.3	2.6	3.4	1.6	2.5	2.1	1.8	2.6			
Total		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0		

Continued. See notes at end of table.

Table 10.44b. Vermont – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020 (continued)

Mental health diagnosis ¹										
Selected characteristics		Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Ethnicity										
Not of Hispanic or Latino origin	95.8		95.2	95.9	94.4	97.3	96.2	97.4	97.7	96.1
Hispanic or Latino origin	4.2		4.8	4.1	5.6	2.7	3.8	2.6	2.3	3.9
Total	100.0		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin										
Mexican	25.0		24.1	31.4	14.3	21.4	22.2	25.0	20.0	24.2
Puerto Rican	32.5		34.2	23.8	42.9	35.7	34.3	--	26.7	37.1
Cuban	4.0		1.8	6.7	3.9	--	2.0	--	20.0	5.6
Other specific	38.5		39.9	38.1	39.0	42.9	41.4	75.0	33.3	33.1
Total	100.0		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁴										
Veteran	4.5		4.3	3.1	3.1	4.9	4.0	3.2	5.9	4.7
Not a veteran	95.5		95.7	96.9	96.9	95.1	96.0	96.8	94.1	95.3
Total	100.0		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

- - Quantity is zero; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.45a. Virginia – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	122,981	31,906	35,935	22,315	24,781	46,992	10,092	26,164	34,456
Gender									
Male	60,910	13,980	14,413	14,809	10,223	18,913	4,493	15,937	20,030
Female	62,044	17,917	21,514	7,504	14,552	28,072	5,598	10,225	14,423
Total	122,954	31,897	35,927	22,313	24,775	46,985	10,091	26,162	34,453
Age²									
0 to 11 years	13,315	5,567	2,500	7,787	151	1,021	5	75	5,287
12 to 14 years	9,410	3,461	2,603	4,435	328	2,758	20	131	3,730
15 to 17 years	10,517	3,399	3,542	3,769	699	4,466	89	268	4,072
18 to 20 years	5,695	1,492	2,089	1,455	867	2,788	248	590	1,864
21 to 24 years	6,700	1,643	2,294	1,004	1,596	3,061	578	1,400	1,611
25 to 29 years	10,166	2,366	3,441	976	2,807	4,463	990	2,517	2,010
30 to 34 years	10,393	2,578	3,338	876	3,074	4,423	1,097	2,712	2,131
35 to 39 years	9,773	2,415	3,046	647	2,839	3,997	1,158	2,787	1,991
40 to 44 years	8,271	2,048	2,617	449	2,503	3,404	1,029	2,387	1,675
45 to 49 years	7,832	1,818	2,521	299	2,349	3,362	1,063	2,284	1,621
50 to 54 years	8,380	1,784	2,409	232	2,283	3,797	1,099	2,587	1,957
55 to 59 years	8,802	1,605	2,365	160	2,208	3,961	1,165	3,042	2,155
60 to 64 years	6,476	950	1,588	139	1,508	2,691	810	2,520	1,779
65 years and older	7,247	778	1,582	87	1,569	2,800	741	2,864	2,571
Total	122,977	31,904	35,935	22,315	24,781	46,992	10,092	26,164	34,454
Race									
American Indian or Alaska Native	247	72	74	36	59	105	18	39	66
Asian ³	1,527	244	352	99	245	507	102	625	366
Black or African American	34,007	7,699	6,106	6,060	6,113	11,475	2,819	11,692	11,104
Native Hawaiian or Other Pacific Islander	102	23	28	7	21	41	6	27	26
White	74,000	19,571	25,757	13,090	16,785	30,111	6,595	11,997	19,380
Other	11,688	3,761	3,197	2,704	1,460	4,298	529	1,715	3,234
Total	121,571	31,370	35,514	21,996	24,683	46,537	10,069	26,095	34,176

Continued. See notes at end of table.

Table 10.45a. Virginia – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	110,524	28,019	32,279	19,872	23,330	42,257	9,611	24,597	31,288
Hispanic or Latino origin	8,818	2,745	2,502	1,647	1,105	3,357	386	1,298	2,410
Total	119,342	30,764	34,781	21,519	24,435	45,614	9,997	25,895	33,698
Specific Hispanic or Latino origin									
Mexican	1,081	388	324	202	108	430	48	121	301
Puerto Rican	1,056	315	312	276	189	379	81	168	276
Cuban	133	34	38	23	28	50	8	29	31
Other specific	4,411	1,264	1,197	717	582	1,652	192	803	1,277
Total	6,681	2,001	1,871	1,218	907	2,511	329	1,121	1,885
Veteran status⁴									
Veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Not a veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡

‡ No data submitted.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.45b. Virginia – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020

Selected characteristics		Mental health diagnosis ¹								
		Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender										
Male	49.5	43.8	40.1	66.4	41.3	40.3	44.5	60.9	58.1	
Female	50.5	56.2	59.9	33.6	58.7	59.7	55.5	39.1	41.9	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age ²										
0 to 11 years	10.8	17.4	7.0	34.9	†	2.2	†	†	15.3	
12 to 14 years	7.7	10.8	7.2	19.9	1.3	5.9	†	†	10.8	
15 to 17 years	8.6	10.7	9.9	16.9	2.8	9.5	†	1.0	11.8	
18 to 20 years	4.6	4.7	5.8	6.5	3.5	5.9	2.5	2.3	5.4	
21 to 24 years	5.4	5.1	6.4	4.5	6.4	6.5	5.7	5.4	4.7	
25 to 29 years	8.3	7.4	9.6	4.4	11.3	9.5	9.8	9.6	5.8	
30 to 34 years	8.5	8.1	9.3	3.9	12.4	9.4	10.9	10.4	6.2	
35 to 39 years	7.9	7.6	8.5	2.9	11.5	8.5	11.5	10.7	5.8	
40 to 44 years	6.7	6.4	7.3	2.0	10.1	7.2	10.2	9.1	4.9	
45 to 49 years	6.4	5.7	7.0	1.3	9.5	7.2	10.5	8.7	4.7	
50 to 54 years	6.8	5.6	6.7	1.0	9.2	8.1	10.9	9.9	5.7	
55 to 59 years	7.2	5.0	6.6	†	8.9	8.4	11.5	11.6	6.3	
60 to 64 years	5.3	3.0	4.4	†	6.1	5.7	8.0	9.6	5.2	
65 years and older	5.9	2.4	4.4	†	6.3	6.0	7.3	10.9	7.5	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race										
American Indian or Alaska Native	†	†	†	†	†	†	†	†	†	
Asian ³	1.3	†	†	†	†	1.1	1.0	2.4	1.1	
Black or African American	28.0	24.5	17.2	27.6	24.8	24.7	28.0	44.8	32.5	
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	†	†	†	
White	60.9	62.4	72.5	59.5	68.0	64.7	65.5	46.0	56.7	
Other	9.6	12.0	9.0	12.3	5.9	9.2	5.3	6.6	9.5	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.45b. Virginia – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020 (continued)

Mental health diagnosis¹										
Selected characteristics		Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders¹
						Bipolar disorders	Depressive disorders			
Ethnicity										
Not of Hispanic or Latino origin	92.6		91.1	92.8	92.3	95.5	92.6	96.1	95.0	92.8
Hispanic or Latino origin	7.4		8.9	7.2	7.7	4.5	7.4	3.9	5.0	7.2
Total	100.0		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin										
Mexican	16.2		19.4	17.3	16.6	11.9	17.1	14.6	10.8	16.0
Puerto Rican	15.8		15.7	16.7	22.7	20.8	15.1	24.6	15.0	14.6
Cuban	2.0		1.7	2.0	1.9	3.1	2.0	2.4	2.6	1.6
Other specific	66.0		63.2	64.0	58.9	64.2	65.8	58.4	71.6	67.7
Total	100.0		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁴										
Veteran	‡		‡	‡	‡	‡	‡	‡	‡	‡
Not a veteran	‡		‡	‡	‡	‡	‡	‡	‡	‡
Total	‡		‡	‡	‡	‡	‡	‡	‡	‡

‡ No data submitted; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.46a. Washington – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	214,013	60,843	78,654	26,597	20,155	75,157	4,118	22,261	83,185
Gender									
Male	90,483	20,751	26,846	16,314	7,253	25,407	1,114	13,641	42,848
Female	120,096	39,700	51,377	10,161	12,730	49,185	2,977	8,294	37,848
Total	210,579	60,451	78,223	26,475	19,983	74,592	4,091	21,935	80,696
Age²									
0 to 11 years	23,824	9,595	5,458	8,240	47	1,256	11	18	10,497
12 to 14 years	15,758	5,534	5,558	5,187	110	4,041	15	49	5,192
15 to 17 years	16,702	5,002	7,255	3,459	311	7,331	46	145	5,319
18 to 20 years	11,978	3,326	5,846	1,549	701	5,658	161	426	3,739
21 to 24 years	13,146	3,171	6,209	1,009	1,369	5,682	393	1,255	4,752
25 to 29 years	19,799	4,922	8,592	1,526	2,355	7,670	578	2,361	8,114
30 to 34 years	21,247	5,445	8,750	1,610	2,675	7,786	591	2,707	9,347
35 to 39 years	18,483	4,956	7,338	1,244	2,414	6,594	534	2,704	8,043
40 to 44 years	14,791	4,193	5,596	866	2,240	5,399	418	2,288	6,077
45 to 49 years	12,027	3,677	4,431	599	1,793	4,759	370	1,863	4,508
50 to 54 years	12,476	3,728	4,192	507	1,889	5,218	337	2,021	4,689
55 to 59 years	12,098	3,223	4,015	359	1,797	5,060	282	2,184	4,423
60 to 64 years	9,505	2,203	2,977	230	1,283	4,189	194	1,795	3,102
65 years and older	8,764	1,476	2,009	91	1,002	3,951	163	2,133	2,898
Total	210,598	60,451	78,226	26,476	19,986	74,594	4,093	21,949	80,700
Race									
American Indian or Alaska Native	11,193	4,177	3,701	1,128	746	3,439	132	826	4,064
Asian ³	5,173	1,163	1,699	330	362	2,414	52	837	1,563
Black or African American	14,626	4,566	4,081	1,406	1,371	4,822	227	2,806	5,798
Native Hawaiian or Other Pacific Islander	2,974	829	953	264	225	1,157	37	439	1,007
White	150,419	41,989	58,149	19,726	15,716	53,670	3,329	14,914	59,017
Other	23,486	7,065	8,677	3,171	1,465	8,381	298	2,014	8,183
Total	207,871	59,789	77,260	26,025	19,885	73,883	4,075	21,836	79,632

Continued. See notes at end of table.

Table 10.46a. Washington – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020 (continued)

Selected characteristics		Mental health diagnosis ¹							
					Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD					
Ethnicity									
Not of Hispanic or Latino origin	167,607	47,874	61,906	20,403	17,371	59,804	3,585	18,779	65,369
Hispanic or Latino origin	36,520	11,061	13,527	5,016	2,304	12,663	453	2,957	13,196
Total	204,127	58,935	75,433	25,419	19,675	72,467	4,038	21,736	78,565
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴									
Veteran	1,453	432	324	47	235	572	58	355	651
Not a veteran	41,609	12,432	12,776	1,987	6,465	16,985	1,591	8,427	19,095
Total	43,062	12,864	13,100	2,034	6,700	17,557	1,649	8,782	19,746

‡ No data submitted.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.46b. Washington – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020

Selected characteristics		Mental health diagnosis¹								
		Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders¹
						Bipolar disorders	Depressive disorders			
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender										
Male	43.0	34.3	34.3	61.6	36.3	34.1	27.2	62.2	53.1	
Female	57.0	65.7	65.7	38.4	63.7	65.9	72.8	37.8	46.9	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age²										
0 to 11 years	11.3	15.9	7.0	31.1	†	1.7	†	†	13.0	
12 to 14 years	7.5	9.2	7.1	19.6	†	5.4	†	†	6.4	
15 to 17 years	7.9	8.3	9.3	13.1	1.6	9.8	1.1	†	6.6	
18 to 20 years	5.7	5.5	7.5	5.9	3.5	7.6	3.9	1.9	4.6	
21 to 24 years	6.2	5.2	7.9	3.8	6.8	7.6	9.6	5.7	5.9	
25 to 29 years	9.4	8.1	11.0	5.8	11.8	10.3	14.1	10.8	10.1	
30 to 34 years	10.1	9.0	11.2	6.1	13.4	10.4	14.4	12.3	11.6	
35 to 39 years	8.8	8.2	9.4	4.7	12.1	8.8	13.0	12.3	10.0	
40 to 44 years	7.0	6.9	7.2	3.3	11.2	7.2	10.2	10.4	7.5	
45 to 49 years	5.7	6.1	5.7	2.3	9.0	6.4	9.0	8.5	5.6	
50 to 54 years	5.9	6.2	5.4	1.9	9.5	7.0	8.2	9.2	5.8	
55 to 59 years	5.7	5.3	5.1	1.4	9.0	6.8	6.9	10.0	5.5	
60 to 64 years	4.5	3.6	3.8	†	6.4	5.6	4.7	8.2	3.8	
65 years and older	4.2	2.4	2.6	†	5.0	5.3	4.0	9.7	3.6	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race										
American Indian or Alaska Native	5.4	7.0	4.8	4.3	3.8	4.7	3.2	3.8	5.1	
Asian³	2.5	1.9	2.2	1.3	1.8	3.3	1.3	3.8	2.0	
Black or African American	7.0	7.6	5.3	5.4	6.9	6.5	5.6	12.9	7.3	
Native Hawaiian or Other Pacific Islander	1.4	1.4	1.2	1.0	1.1	1.6	†	2.0	1.3	
White	72.4	70.2	75.3	75.8	79.0	72.6	81.7	68.3	74.1	
Other	11.3	11.8	11.2	12.2	7.4	11.3	7.3	9.2	10.3	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.46b. Washington – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020 (continued)

Mental health diagnosis¹										
Selected characteristics		Trauma- and stressor-related disorders			Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders¹	
					Bipolar disorders	Depressive disorders				
Total										
Ethnicity										
Not of Hispanic or Latino origin		82.1	81.2	82.1	80.3	88.3	82.5	88.8	86.4	83.2
Hispanic or Latino origin		17.9	18.8	17.9	19.7	11.7	17.5	11.2	13.6	16.8
Total		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin										
Mexican		‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican		‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban		‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific		‡	‡	‡	‡	‡	‡	‡	‡	‡
Total		‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴										
Veteran		3.4	3.4	2.5	2.3	3.5	3.3	3.5	4.0	3.3
Not a veteran		96.6	96.6	97.5	97.7	96.5	96.7	96.5	96.0	96.7
Total		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

‡ No data submitted; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.47a. Wisconsin – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	68,646	7,857	14,667	3,908	5,978	12,829	8,679	9,736	14,219
Gender									
Male	34,735	3,426	6,892	2,512	2,581	5,363	4,506	6,183	7,652
Female	33,890	4,429	7,773	1,395	3,397	7,457	4,172	3,552	6,557
Total	68,625	7,855	14,665	3,907	5,978	12,820	8,678	9,735	14,209
Age²									
0 to 11 years	3,809	1,264	671	912	22	137	33	7	1,332
12 to 14 years	4,001	769	1,000	717	49	443	57	20	1,543
15 to 17 years	4,891	791	1,284	525	138	813	135	78	1,885
18 to 20 years	3,748	387	905	245	190	758	535	180	1,089
21 to 24 years	4,538	407	1,060	183	349	887	780	405	1,002
25 to 29 years	6,457	620	1,446	262	587	1,240	1,070	806	1,172
30 to 34 years	6,249	631	1,391	241	702	1,189	999	869	1,008
35 to 39 years	5,963	602	1,347	230	639	1,150	890	969	931
40 to 44 years	4,789	514	1,039	135	591	962	702	774	749
45 to 49 years	4,720	497	943	136	539	1,022	727	839	692
50 to 54 years	5,004	452	954	109	615	1,084	797	1,028	672
55 to 59 years	5,381	403	991	96	629	1,168	787	1,348	746
60 to 64 years	3,958	248	675	52	447	919	530	1,115	525
65 years and older	5,138	272	961	65	481	1,057	637	1,298	873
Total	68,646	7,857	14,667	3,908	5,978	12,829	8,679	9,736	14,219
Race									
American Indian or Alaska Native	1,230	162	248	69	65	170	120	122	427
Asian ³	831	95	174	29	76	204	90	142	174
Black or African American	8,174	649	796	394	496	1,014	2,819	1,753	946
Native Hawaiian or Other Pacific Islander	76	10	13	5	3	18	5	9	27
White	51,769	6,369	12,190	3,116	4,902	9,870	4,614	7,144	10,937
Other	2,697	350	543	205	321	424	419	450	644
Total	64,777	7,635	13,964	3,818	5,863	11,700	8,067	9,620	13,155

Continued. See notes at end of table.

Table 10.47a. Wisconsin – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	57,799	6,833	12,694	3,511	5,450	10,113	6,932	8,813	11,799
Hispanic or Latino origin	3,215	461	606	213	201	632	526	389	640
Total	61,014	7,294	13,300	3,724	5,651	10,745	7,458	9,202	12,439
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴									
Veteran	553	38	105	8	49	73	157	111	74
Not a veteran	17,031	1,252	2,826	387	1,883	2,507	4,747	4,043	1,789
Total	17,584	1,290	2,931	395	1,932	2,580	4,904	4,154	1,863

‡ No data submitted.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.47b. Wisconsin – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020

Selected characteristics		Mental health diagnosis¹								
		Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders¹
						Bipolar disorders	Depressive disorders			
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender										
Male	50.6	43.6	47.0	64.3	43.2	41.8	51.9	63.5	53.9	
Female	49.4	56.4	53.0	35.7	56.8	58.2	48.1	36.5	46.1	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age²										
0 to 11 years	5.5	16.1	4.6	23.3	†	1.1	†	†	9.4	
12 to 14 years	5.8	9.8	6.8	18.3	†	3.5	†	†	10.9	
15 to 17 years	7.1	10.1	8.8	13.4	2.3	6.3	1.6	†	13.3	
18 to 20 years	5.5	4.9	6.2	6.3	3.2	5.9	6.2	1.8	7.7	
21 to 24 years	6.6	5.2	7.2	4.7	5.8	6.9	9.0	4.2	7.0	
25 to 29 years	9.4	7.9	9.9	6.7	9.8	9.7	12.3	8.3	8.2	
30 to 34 years	9.1	8.0	9.5	6.2	11.7	9.3	11.5	8.9	7.1	
35 to 39 years	8.7	7.7	9.2	5.9	10.7	9.0	10.3	10.0	6.5	
40 to 44 years	7.0	6.5	7.1	3.5	9.9	7.5	8.1	7.9	5.3	
45 to 49 years	6.9	6.3	6.4	3.5	9.0	8.0	8.4	8.6	4.9	
50 to 54 years	7.3	5.8	6.5	2.8	10.3	8.4	9.2	10.6	4.7	
55 to 59 years	7.8	5.1	6.8	2.5	10.5	9.1	9.1	13.8	5.2	
60 to 64 years	5.8	3.2	4.6	1.3	7.5	7.2	6.1	11.5	3.7	
65 years and older	7.5	3.5	6.6	1.7	8.0	8.2	7.3	13.3	6.1	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race										
American Indian or Alaska Native	1.9	2.1	1.8	1.8	1.1	1.5	1.5	1.3	3.2	
Asian³	1.3	1.2	1.2	†	1.3	1.7	1.1	1.5	1.3	
Black or African American	12.6	8.5	5.7	10.3	8.5	8.7	34.9	18.2	7.2	
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	†	†	†	
White	79.9	83.4	87.3	81.6	83.6	84.4	57.2	74.3	83.1	
Other	4.2	4.6	3.9	5.4	5.5	3.6	5.2	4.7	4.9	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.47b. Wisconsin – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020 (continued)

Mental health diagnosis ¹									
Selected characteristics	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	94.7	93.7	95.4	94.3	96.4	94.1	92.9	95.8	94.9
Hispanic or Latino origin	5.3	6.3	4.6	5.7	3.6	5.9	7.1	4.2	5.1
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status ⁴									
Veteran	3.1	2.9	3.6	2.0	2.5	2.8	3.2	2.7	4.0
Not a veteran	96.9	97.1	96.4	98.0	97.5	97.2	96.8	97.3	96.0
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

‡ No data submitted; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.48a. Wyoming – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	15,765	3,863	4,632	1,323	1,671	5,861	628	1,088	2,727
Gender									
Male	6,878	1,428	1,617	865	619	2,147	238	671	1,536
Female	8,887	2,435	3,015	458	1,052	3,714	390	417	1,191
Total	15,765	3,863	4,632	1,323	1,671	5,861	628	1,088	2,727
Age²									
0 to 11 years	1,453	581	236	347	6	116	--	2	419
12 to 14 years	1,010	306	208	203	10	323	2	1	262
15 to 17 years	1,314	250	322	192	27	583	10	3	368
18 to 20 years	917	184	261	76	71	467	42	27	169
21 to 24 years	1,184	286	434	55	124	501	57	63	186
25 to 29 years	1,527	364	541	82	162	608	74	90	193
30 to 34 years	1,438	355	503	82	196	534	82	113	196
35 to 39 years	1,402	340	492	87	210	481	61	160	176
40 to 44 years	1,165	278	377	62	194	435	71	103	145
45 to 49 years	1,006	241	339	42	151	388	45	113	144
50 to 54 years	840	211	250	33	128	351	54	101	106
55 to 59 years	845	178	236	30	150	346	59	111	104
60 to 64 years	700	128	202	19	101	299	36	104	106
65 years and older	964	161	231	13	141	429	35	97	153
Total	15,765	3,863	4,632	1,323	1,671	5,861	628	1,088	2,727
Race									
American Indian or Alaska Native	384	100	61	19	31	157	16	39	60
Asian ³	65	14	19	3	5	33	4	7	8
Black or African American	257	69	58	17	18	93	12	49	51
Native Hawaiian or Other Pacific Islander	30	8	7	2	1	13	1	2	5
White	13,404	3,250	3,980	1,169	1,462	5,001	528	861	2,325
Other	771	199	243	36	58	257	25	42	128
Total	14,911	3,640	4,368	1,246	1,575	5,554	586	1,000	2,577

Continued. See notes at end of table.

Table 10.48a. Wyoming – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	13,486	3,233	3,932	1,123	1,454	5,033	552	905	2,321
Hispanic or Latino origin	1,480	414	460	130	128	525	49	116	273
Total	14,966	3,647	4,392	1,253	1,582	5,558	601	1,021	2,594
Specific Hispanic or Latino origin									
Mexican	752	200	214	59	49	260	14	54	139
Puerto Rican	56	17	19	5	11	24	2	4	8
Cuban	10	4	5	- -	6	3	1	1	- -
Other specific	637	190	220	66	58	233	28	50	117
Total	1,455	411	458	130	124	520	45	109	264
Veteran status⁴									
Veteran	268	87	59	6	38	110	9	24	26
Not a veteran	4,918	1,059	1,469	267	722	2,059	234	439	653
Total	5,186	1,146	1,528	273	760	2,169	243	463	679

- - Quantity is zero.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.48b. Wyoming – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020

Selected characteristics		Mental health diagnosis¹										
		Trauma- and stressor-related disorders				Mood disorders		Personality disorders			Schizophrenia, other psychotic disorders	Other mental disorders¹
						Bipolar disorders	Depressive disorders					
Total		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	
Total individuals		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	
Gender												
Male		43.6	37.0	34.9	65.4	37.0	36.6	37.9	61.7	56.3		
Female		56.4	63.0	65.1	34.6	63.0	63.4	62.1	38.3	43.7		
Total		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	
Age²												
0 to 11 years		9.2	15.0	5.1	26.2	†	2.0	--	†	15.4		
12 to 14 years		6.4	7.9	4.5	15.3	†	5.5	†	†	9.6		
15 to 17 years		8.3	6.5	7.0	14.5	1.6	9.9	1.6	†	13.5		
18 to 20 years		5.8	4.8	5.6	5.7	4.2	8.0	6.7	2.5	6.2		
21 to 24 years		7.5	7.4	9.4	4.2	7.4	8.5	9.1	5.8	6.8		
25 to 29 years		9.7	9.4	11.7	6.2	9.7	10.4	11.8	8.3	7.1		
30 to 34 years		9.1	9.2	10.9	6.2	11.7	9.1	13.1	10.4	7.2		
35 to 39 years		8.9	8.8	10.6	6.6	12.6	8.2	9.7	14.7	6.5		
40 to 44 years		7.4	7.2	8.1	4.7	11.6	7.4	11.3	9.5	5.3		
45 to 49 years		6.4	6.2	7.3	3.2	9.0	6.6	7.2	10.4	5.3		
50 to 54 years		5.3	5.5	5.4	2.5	7.7	6.0	8.6	9.3	3.9		
55 to 59 years		5.4	4.6	5.1	2.3	9.0	5.9	9.4	10.2	3.8		
60 to 64 years		4.4	3.3	4.4	1.4	6.0	5.1	5.7	9.6	3.9		
65 years and older		6.1	4.2	5.0	†	8.4	7.3	5.6	8.9	5.6		
Total		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	
Race												
American Indian or Alaska Native		2.6	2.7	1.4	1.5	2.0	2.8	2.7	3.9	2.3		
Asian³		†	†	†	†	†	†	†	†	†		
Black or African American		1.7	1.9	1.3	1.4	1.1	1.7	2.0	4.9	2.0		
Native Hawaiian or Other Pacific Islander		†	†	†	†	†	†	†	†	†		
White		89.9	89.3	91.1	93.8	92.8	90.0	90.1	86.1	90.2		
Other		5.2	5.5	5.6	2.9	3.7	4.6	4.3	4.2	5.0		
Total		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	

Continued. See notes at end of table.

Table 10.48b. Wyoming – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	90.1	88.6	89.5	89.6	91.9	90.6	91.8	88.6	89.5
Hispanic or Latino origin	9.9	11.4	10.5	10.4	8.1	9.4	8.2	11.4	10.5
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	51.7	48.7	46.7	45.4	39.5	50.0	31.1	49.5	52.7
Puerto Rican	3.8	4.1	4.1	3.8	8.9	4.6	4.4	3.7	3.0
Cuban	†	†	1.1	--	4.8	†	2.2	†	--
Other specific	43.8	46.2	48.0	50.8	46.8	44.8	62.2	45.9	44.3
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁴									
Veteran	5.2	7.6	3.9	2.2	5.0	5.1	3.7	5.2	3.8
Not a veteran	94.8	92.4	96.1	97.8	95.0	94.9	96.3	94.8	96.2
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

-- Quantity is zero; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.49a. Other jurisdictions – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	434	48	31	28	32	37	6	141	137
Gender									
Male	280	23	9	22	16	16	4	92	115
Female	154	25	22	6	16	21	2	49	22
Total	434	48	31	28	32	37	6	141	137
Age²									
0 to 11 years	21	--	1	12	--	--	--	1	11
12 to 14 years	7	2	1	1	--	--	1	--	3
15 to 17 years	11	4	--	--	1	3	--	1	2
18 to 20 years	17	3	3	4	2	2	1	--	4
21 to 24 years	25	3	4	2	3	2	1	2	8
25 to 29 years	48	10	2	4	4	1	2	12	13
30 to 34 years	48	4	1	--	5	4	--	18	16
35 to 39 years	53	8	1	1	4	3	1	17	22
40 to 44 years	48	2	6	2	2	6	--	15	23
45 to 49 years	35	5	2	1	2	2	--	11	12
50 to 54 years	30	3	1	--	2	4	--	15	7
55 to 59 years	37	1	4	--	2	2	--	24	6
60 to 64 years	26	1	2	--	4	3	--	13	5
65 years and older	28	2	3	1	1	5	--	12	5
Total	434	48	31	28	32	37	6	141	137
Race									
American Indian or Alaska Native	--	--	--	--	--	--	--	--	--
Asian ³	32	7	8	3	2	3	1	3	7
Black or African American	--	--	--	--	--	--	--	--	--
Native Hawaiian or Other Pacific Islander	353	33	21	23	28	29	5	134	99
White	8	2	1	1	1	2	--	--	1
Other	41	6	1	1	1	3	--	4	30
Total	434	48	31	28	32	37	6	141	137

Continued. See notes at end of table.

Table 10.49a. Other jurisdictions – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2020 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	276	16	21	25	24	29	5	126	47
Hispanic or Latino origin	--	--	--	--	--	--	--	--	--
Total	276	16	21	25	24	29	5	126	47
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴									
Veteran	4	--	--	2	--	1	--	--	1
Not a veteran	101	4	7	3	13	13	3	53	10
Total	105	4	7	5	13	14	3	53	11

-- Quantity is zero; ‡ No data submitted.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Table 10.49b. Other jurisdictions – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020

Selected characteristics		Mental health diagnosis ¹								
		Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender										
Male	64.5	47.9	29.0	78.6	50.0	43.2	66.7	65.2	83.9	
Female	35.5	52.1	71.0	21.4	50.0	56.8	33.3	34.8	16.1	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age ²										
0 to 11 years	4.8	--	3.2	42.9	--	--	--	†	8.0	
12 to 14 years	1.6	4.2	3.2	3.6	--	--	16.7	--	2.2	
15 to 17 years	2.5	8.3	--	--	3.1	8.1	--	†	1.5	
18 to 20 years	3.9	6.3	9.7	14.3	6.3	5.4	16.7	--	2.9	
21 to 24 years	5.8	6.3	12.9	7.1	9.4	5.4	16.7	1.4	5.8	
25 to 29 years	11.1	20.8	6.5	14.3	12.5	2.7	33.3	8.5	9.5	
30 to 34 years	11.1	8.3	3.2	--	15.6	10.8	--	12.8	11.7	
35 to 39 years	12.2	16.7	3.2	3.6	12.5	8.1	16.7	12.1	16.1	
40 to 44 years	11.1	4.2	19.4	7.1	6.3	16.2	--	10.6	16.8	
45 to 49 years	8.1	10.4	6.5	3.6	6.3	5.4	--	7.8	8.8	
50 to 54 years	6.9	6.3	3.2	--	6.3	10.8	--	10.6	5.1	
55 to 59 years	8.5	2.1	12.9	--	6.3	5.4	--	17.0	4.4	
60 to 64 years	6.0	2.1	6.5	--	12.5	8.1	--	9.2	3.6	
65 years and older	6.5	4.2	9.7	3.6	3.1	13.5	--	8.5	3.6	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race										
American Indian or Alaska Native	--	--	--	--	--	--	--	--	--	
Asian ³	7.4	14.6	25.8	10.7	6.3	8.1	16.7	2.1	5.1	
Black or African American	--	--	--	--	--	--	--	--	--	
Native Hawaiian or Other Pacific Islander	81.3	68.8	67.7	82.1	87.5	78.4	83.3	95.0	72.3	
White	1.8	4.2	3.2	3.6	3.1	5.4	--	--	†	
Other	9.4	12.5	3.2	3.6	3.1	8.1	--	2.8	21.9	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

**Table 10.49b. Other jurisdictions – Individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2020
(continued)**

Mental health diagnosis ¹										
Selected characteristics		Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Ethnicity										
Not of Hispanic or Latino origin		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Hispanic or Latino origin		--	--	--	--	--	--	--	--	--
Total		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin										
Mexican		‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican		‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban		‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific		‡	‡	‡	‡	‡	‡	‡	‡	‡
Total		‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status ⁴										
Veteran		3.8	--	--	40.0	--	7.1	--	--	9.1
Not a veteran		96.2	100.0	100.0	60.0	100.0	92.9	100.0	100.0	90.9
Total		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

-- Quantity is zero; ‡ No data submitted; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" includes all mental health diagnoses, which is up to three diagnoses reported for each individual. For this reason, the numbers may not sum to total. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent due to rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2020 reporting period. Data as of 03.03.2022.

Appendix A

MH-TEDS/MH-CLD Crosswalk Tables

MH-TEDS					MH-CLD			
Variable Name	Data Field Reference No.			Categories	Variable Name	Data Field Reference No.		Categories
	Admission Record	Discharge/Update Record				Data Files		
		Value taken from admission record	Value taken at discharge or data update			BCI	SHR	
System Transaction Type	SDS 1		DIS 1	A=Add C=Change D=Delete	Transaction Type	C-00	R-00	A=Add C=Change D=Delete
State Code — Key field	SDS 2		DIS 2	2 alphanumeric	Reporting State Code	H-02	H-01	2 alphanumeric
Reporting Date	SDS 3		DIS 3	MMYYYY	contained in the file name algorithm			MMDDYYYY
CLIENT RECORD					CLIENT RECORD			
State Provider Identifier – Key field	MDS 1	DIS 11	DIS 4	1-15 alphanumeric	Not reported			
Client Identifier – Key field	MDS 2	DIS 12	DIS 5	1-15 alphanumeric	Client Identifier	C-01	R-01	1-15 alphanumeric
Codependent or Collateral – Key field	MDS 3	DIS 13	DIS 6	1=codependent/collateral 2=client	Not reported			
Client Transaction Type – Key field	MDS 4			A=Initial Admission (SA)	Not reported			
				T=Transfer or change in service (SA)				
				M=Initial Admission (MH)				
				X=Transfer or change in service (MH)				
Date of Admission – Key field	MDS 5	DIS 15		MMDDYYYY	Client treatment status at the start of the reporting period	C-02		1=New client; 2=Continuing client;
Date of Birth	MDS 8	DIS 17		MMDDYYYY	Age	C-05		
				01010007=unknown				
				01010008=not collected				
Gender	MDS 9	DIS 18		1=male	Sex	C-04		1=male
				2=female				2=female
				7=unknown				7=unknown
				8=not collected				8=not collected
Race	MDS 10	DIS 19		01=Alaska Native	Race	C-06		
				02=American Indian and Alaska Native				02=American Indian and Alaska Native
				13=Asian				13=Asian
				23=Native Hawaiian or other Pacific Islander				23=Native Hawaiian or other Pacific Islander

Appendix A

MH-TEDS/MH-CLD Crosswalk Tables

MH-TEDS				MH-CLD				
Variable Name	Data Field Reference No.			Categories	Variable Name	Data Field Reference No.		Categories
	Admission Record	Discharge/Update Record				Data Files		
		Value taken from admission record	Value taken at discharge or data update			BCI	SHR	
				03=Asian or Pacific Islander				03=Asian or Pacific Islander (temporary code)
				04= Black or African American				04= Black or African American
				05=White				05=White
				20=Other single race				20=Some other race alone
				21=Two or more races				21=Two or more races
								22=Hispanic (temporary code)
				97=unknown				97=unknown
				98=not collected				98=not collected
Hispanic or Latino Origin (Ethnicity)	MDS 11	DIS 20		01=Puerto Rico	Ethnicity	C-07		11=Puerto Rican (optional)
				02=Mexican				12=Mexican (optional)
				03=Cuban				13=Cuban (optional)
				04=Other Specific Hispanic or Latino				14=Other Specific Hispanic
				05 Not of Hispanic Origin				02=Not of Hispanic Origin
				06=Hispanic, specific origin not specified				01=Hispanic origin regardless of race
				97=unknown				97=unknown
				98=not collected				98=Not collected
Education	MDS 12		MHD 5 (reported only for ages 3-17 and up to 21 if in special education).	00= less than one grade completed or no schooling	School grade level	C-25		00=no years of schooling
				01-11=grades 1 thru 11th				01-11=grades 1 thru 11th
				12=12th grade or GED				12 = 12th grade
				13=Freshman or first year in a 2-year (Associate) Degree				17=freshman (1st year in College)
				14=Sophomore or second year in a 2-year (Associate) Degree				18=sophomore (2nd year in College)
				15=Junior in college				19=junior (3rd year in College)
				16=Senior in college or Bachelor's Degree				20=senior (4th year in College)
				17=Some Post-Graduate Study-Degree not completed				
				18= Master's Degree completed				
				19-25= Post graduate study				
				70= graduate or professional school (can be used instead of codes 17-25)				21=Graduate or professional school
				71=vocational school				16=vocational school

Appendix A

MH-TEDS/MH-CLD Crosswalk Tables

MH-TEDS				MH-CLD				
Variable Name	Data Field Reference No.			Categories	Variable Name	Data Field Reference No.		Categories
	Admission Record	Discharge/Update Record				Data Files		
		Value taken from admission record	Value taken at discharge or data update			BCI	SHR	
				72=nursery school, pre-school				13=nursery school, pre-school (head start)
				73=kindergarten				14=kindergarten
				74=self-contained special education class				15=self-contained special education class
				97=unknown				97 = unknown
				98=not collected				98=not collected
Employment Status	MDS 13		DIS 24	01=full time	Competitive Employment Status at admission/start of the reporting period and discharge/end of the reporting period (>= 16 yrs old)	C-09 and C-10		01=full time
				02=part-time				02=part time
				03=unemployed				03=unemployed
				04=not in the labor force				84=not in the labor force
				05=employed, full/part-time not specified (temporary code)				05=employed, full/part-time not specified (temporary code)
				97=unknown				97=unknown
				98=not collected				98=not collected
Type of Treatment Service/ Treatment Setting – Key field	MDS 18	DIS 16	DIS 7		Service Setting Status Throughout the Reporting Period	C-15		
				01=detox, 24-hr, hospital				
				02= detox, 24-hr, residential				
				03=rehabilitation/residential-hospital (not detox)				
				04=rehabilitation/residential-short term (<30 days)				
				05=rehabilitation/ residential-long term (>30 days)				
				06=ambulatory-intensive outpatient				
				07=ambulatory-non-intensive outpatient				
				08=ambulatory-detoxification				
				72=state psychiatric hospital				1=state psychiatric hospital
				73=SMHA funded/operated community- based program				2=SMHA funded/operated community-based program
				74=residential treatment center				3=residential treatment center
				75=other psychiatric inpatient				4=other psychiatric inpatient

Appendix A

MH-TEDS/MH-CLD Crosswalk Tables

MH-TEDS					MH-CLD			
Variable Name	Data Field Reference No.			Categories	Variable Name	Data Field Reference No.		Categories
	Admission Record	Discharge/Update Record				Data Files		
		Value taken from admission record	Value taken at discharge or data update			BCI	SHR	
				76=institutions under the justice system 96=Not applicable (use only for codependents or collateral clients (SA) and for MH clients receiving MH assessments, evaluation, or screening only)			5=institutions under the justice system	
Date of Last Contact or Data Update - Key field			DIS 8	MMDDYYYY 01010007 = Unknown 01010008 = Not Collected	Information used in determining client status			
Reason for Discharge, Transfer, or Discontinuance of Treatment			DIS 10		Discharge reasons	BCI (C-03)		SHR (R-03)
				01=treatment completed		12=treatment completed	01=completed SH inpatient treatment	
				02= dropped out of treatment (lost contact, administrative discharge, left against medical advise, eloped, failed to return from leave, and client choice)		22= lost contact or administrative discharge	03=left against medical advise, eloped, failed to return from leave, non-compliance with treatment and/or policy; 04=client choice	
				03=terminated by facility				
				04=transferred to another treatment program or facility provider			07=discharged to other inpatient provider	
				14=transferred to another treatment program/facility but did not report				
				05=incarcerated or released by or to courts		32=discharged to corrections, jail	02=released by or to courts	
				06=death		43=death of client	06=death	
				07=Other (includes aging out of the children's MH system, extended placement (conditional release), and all other reasons)		52=aged out; 62= other specified reasons	05=extended placement; 09=reason not classified elsewhere	
				08= Unknown - This code will continue to be accepted. However, states are encouraged to use the code 97 Unknown instead				
				24=transferred to another treatment program or facility that is not in the state reporting system				

Appendix A

MH-TEDS/MH-CLD Crosswalk Tables

MH-TEDS					MH-CLD				
Variable Name	Data Field Reference No.			Categories	Variable Name	Data Field Reference No.		Categories	
	Admission Record	Discharge/Update Record				Data Files			
		Value taken from admission record	Value taken at discharge or data update			BCI	SHR		
				34 =discharged temporarily to an acute medical facility for medical services 96 = Not applicable 97=unknown 98=not collected				08=discharged to an acute medical facility for medical services 72=reason unknown 82=reason not collected 97=unknown 98=not collected	
Client Discharge Transaction Type - Key field		DIS 14	DIS 28	D=Discharge (SA client) E=Discharge (MH client) ata update (MH client) U=D	Not reported				
Co-occurring Mental and Substance Use Disorders	SuDS 5			1=yes 2=no 7=unknown 8=not collected	Substance use problem	C-21		1=yes 2=no 7=unknown 8=not collected	
Date of Discharge - Key field			DIS 9	MMDDYYYY	Information used in determining Client status - at end of the reporting period	C-03			
Living Arrangements	SuDS 8		DIS 23	01=homeless 02=dependent living (under age 18 living with parents, relatives, or guardians) 22=dependent living: residential care and other residential status 32=dependent living: foster home/foster care 42=dependent living: crisis residence 52=dependent living:institutional setting 62=dependent living: jail and other institutions under the justice system 72=dependent living: adults in private residence who needs assistance in daily living 	Residential status at start and end of the reporting period	C-12 and C-13		01=homeless 37=private residence (children only) 03=residential care 08=other residential status 02=fosterhome/foster care 04=crisis residence 05=institutional setting 06=jail/correctional facility 27=private residence, dependent living (adult)	

Appendix A

MH-TEDS/MH-CLD Crosswalk Tables

MH-TEDS				MH-CLD				
Variable Name	Data Field Reference No.			Categories	Variable Name	Data Field Reference No.		Categories
	Admission Record	Discharge/Update Record				Data Files		
		Value taken from admission record	Value taken at discharge or data update			BCI	SHR	
				03=independent living 04=private residence, living arrangement not specified, adults (temporary code) 97=unknown 98=not collected				17=private residence, independent living (adult) 07=private residence, living arrangement not specified, adults (temporary code) 97=unknown 98=not collected
Detailed Not-in-Labor Force	SuDS 12		DIS 25	01=homemaker 02=student 03=retired 04=disabled 05= resident of institution or persons receiving services from institutional facilities such as hospitals, jails, prisons, etc. 06=other 07=sheltered/non-competitive employment 96=not applicable (if employment status is not 04, NILF) 97=unknown 98=not collected	Competitive Employment Status - at start and end of the reporting period	C-09 and C-10		14=homemaker 24=student 34=retired 44=disabled 64=other reported classification 74=sheltered/non-competitive employment 96=not applicable (if in institutional setting or if age <16) 97=unknown 84=NILF, classification not specified 98=not collected
Arrests in Past 30 Days	SuDS 16		DIS 26	00-96=number of arrests 97=unknown 98=not collected	Number of Arrests in Prior 30 days - at start and end of the reporting period	C-22 and C-23		00-96=number of arrests 97=unknown 98=not collected
Diagnostic Code Set Identifier	SuDS 18		MHD 1	1=DSM-IV 2=ICD-9 3=ICD-10 4=DSM-V 5=DSM-IIIR 7=unknown 8=not collected	Diagnostic Code Identifier	H-08		1=DSM IV 2=ICD-9 3=ICD 10

Appendix A

MH-TEDS/MH-CLD Crosswalk Tables

		MH-TEDS				MH-CLD		
Variable Name	Data Field Reference No.			Categories	Variable Name	Data Field Reference No.		Categories
	Admission Record	Discharge/Update Record				Data Files		
		Value taken from admission record	Value taken at discharge or data update			BCI	SHR	
SH-Legal Status at Admission	MHA 4			01=voluntary-self 02=voluntary-others (parents, guardians, etc) 03=involuntary-civil 04=involuntary-criminal 05=involuntary-juvenile justice 06=involuntary-civil, sexual 96=not applicable 97=unknown 98=not collected	State Hospital-Admission Legal Status		R-05	01=voluntary-self 02=voluntary-others (parents, guardians, etc) 03=involuntary-civil 04=involuntary-criminal 05=involuntary-juvenile justice 06=involuntary-civil, sexual 97=unknown 98=not collected
SMI/SED Status	MHA 2		MHD 3	1=SMI 2=SED 3=at risk for SED (optional) 4=Not SMI/SED 7=unknown 8=not collected	SMI/SED Status	C-08		1=SMI 2=SED 3=at risk for SED (optional) 4=Not SMI/SED 7=unknown 8=not collected
School Attendance Status	MHA 3		MHD 4	1=Yes, client has attended school at any time in the past 3 months 2=No, client has not attended school at any time in the past three months 6=not applicable (non-school age) 7=unknown 8=not collected	School Attendance Status (reported only for school age children, 3-17, and up to 21 if in special education).	C-24		1=Yes, client has attended school at any time in the past 3 months 2=No, client has not attended school at any time in the past three months 6=not applicable 7=unknown 8=not collected
MH Diagnostic Code - one (ICD-10 Compatible)	MHA 1a		MHD 2a	xxx.xxxx; xxx----; xxx----; xxx.x---; xxx.xx--; xxx.xxx-; 999.9997 unknown 999.9998=not collected	Mental Health Diagnosis - one	C-17		xxx.xxxx; xxx.xxx-; xxx.xx--; xxx.x---; xxx-- -- 999.9997=unknown 999.9998=not collected

Appendix A

MH-TEDS/MH-CLD Crosswalk Tables

		MH-TEDS				MH-CLD		
Variable Name	Data Field Reference No.			Categories	Variable Name	Data Field Reference No.		Categories
	Admission Record	Discharge/Update Record				Data Files		
		Value taken from admission record	Value taken at discharge or data update			BCI	SHR	
MH Diagnostic Code - two (ICD-10 Compatible)	MHA 1b		MHD 2b	xxx.xxxx; xxx-----; xxx-----; xxx.x- --; xxx.xx--; xxx.xxx-;	Mental Health Diagnosis - two	C-18		xxx.xxxx; xxx-----; xxx.----; xxx.x- --; xxx.xx--; xxx.xxx-
				999.9996=no second MH diagnosis				999.9996=no second diagnosis
				999.9997=unknown				999.9997=unknown
				999.998=not collected				999.9998=not collected
MH Diagnostic Code - three (ICD-10 Compatible)	MHA 1c		MHD 2c	xxx.xxxx; xxx-----; xxx-----; xxx.x- --; xxx.xx--; xxx.xxx-	Mental Health Diagnosis - three	C-19		xxx.xxxx; xxx.xxx-; xxx.xx--; xxx.x- --; xxx.xx--; xxx.xxx-
				999.9996=no third MH diagnosis				999.9996=no third diagnosis
				999.9997=unknown				999.9997=unknown
				999.998=not collected				999.9998=not collected
Diagnostic Code - ICD 10 Compatible, Substance use	SuDS 19			optional reporting for MH	Substance use Diagnosis	C-20		
				xxx.xxxx; xxx-----; xxx.----; xxx.x- --; xxx.xx--; xxx.xxx-				xxx.xxxx; xxx.xxx-; xxx.xx--; xxx.x- --; xxx.xx--; xxx.xxx-
								999.9996 no substance use diagnosis
				999.9997=unknown				999.9997=unknown
				999.9998=not collected				999.9998=not collected
Marital Status	SuDS 14			optional reporting for MH	Marital Status	O-1		optional reporting
				01=never married				01=never married
				02=now married/living as couple or cohabiting				02=married/living as couple
				03=separated				03=separated
				04=divorced				04=divorced
				05=widowed				05=widowed
				97=unknown				97=unknown
				98=not collected				98=not collected
GAF/CGAS Score – Optional	MHA 5		MHD 6	optional reporting	GAF/CGAS Score	O-02		optional reporting
				0-100 GAF/CGAS Score				0-100 GAF/CGAS score
				997 = Unknown				997=unknown
				998 = Not collected				
Prior Treatment Episodes	MDS 6			optional reporting for MH	Not reported			
Referral Source	MDS 7			optional reporting for MH	Not reported			
Substance Use (Primary, Secondary, Tertiary)	MDS 14a MDS 14b MDS 14c		DIS 21a DIS 21b DIS 21c	optional reporting for MH	Not reported			

Appendix A

MH-TEDS/MH-CLD Crosswalk Tables

		MH-TEDS				MH-CLD		
Variable Name	Data Field Reference No.			Categories	Variable Name	Data Field Reference No.		Categories
	Admission Record	Discharge/Update Record				Data Files		
		Value taken from admission record	Value taken at discharge or data update			BCI	SHR	
Route of Administration (Primary, Secondary, Tertiary Substances)	MDS 15a MDS 15b MDS 15c			optional reporting for MH	Not reported			
Frequency of Use (Primary, Secondary, Tertiary Substances)	MDS 16a MDS 16b MDS 16c		DIS 22a DIS 22b DIS 22c	optional reporting for MH	Not reported			
Age at First Use (Primary, Secondary, Tertiary, Substances)	MDS 17a MDS 17b MDS 17c			optional reporting for MH	Not reported			
Medication-Assisted Opioid Therapy	MDS 19			optional reporting for MH	Not reported			
Detailed Drug Code (Primary, Secondary, Tertiary)	SuDS 1 SuDS 2 SuDS 3			optional reporting for MH	Not reported			
Diagnostic Code (DSM/ICD)	SuDS 4			optional reporting	Not reported			
Pregnant at Admission	SuDS 6			optional reporting	Not reported			
Veteran Status	SuDS 7			optional reporting for MH	Veteran Status	O-05		optional reporting
				1= Veteran				1= Veteran
				2 = Not a Veteran				2 = Not a Veteran
				7 = Unknown				7 = Unknown
				8 = Not Collected				8 = Not Collected
Source of Income/Support	SuDS 9			optional reporting for MH	Not reported			
Health Insurance	SuDS 10			optional reporting for MH	Not reported			
Payment Source, Primary (Expected or Actual)	SuDS 11			optional reporting for MH	Not reported			
Detailed Criminal Justice Referral	SuDS 13			optional reporting	Not reported			
Days Waiting to Enter Treatment	SuDS 15			optional reporting	Not reported			
Attendance at Self-Help Groups in Past 30 Days	SuDS 17		DIS 27	optional reporting for MH	Not reported			
Not reported					Type of Funding Support	O-03		optional reporting
								01 = Medicaid Only
								02 = Medicaid and Non-Medicaid Sources
								03 = Non-Medicaid Only
								97 = Unknown
Not reported					Mental Health Block Grant funded services	O-04		optional reporting
								01 = Yes, MHBG Used to pay for services and supports
								02 = No, MHBG funds were not used
								97 = Unknown

Appendix A

MH-TEDS/MH-CLD Crosswalk Tables

		MH-TEDS				MH-CLD			
Variable Name				Categories		Variable Name	Data Field Reference No.	Categories	
	Admission Record	Discharge/Update Record					Data Files		
		Value taken from admission record	Value taken at discharge or data update				BCI		
									98 = Not Collected
LEGEND									
Key field = must not be blank. Key fields, taken together, are used to uniquely identify and link records.									
Variable highlighted in green = variable not reported in MH-CLD but required in MH-TEDS.									
Variable highlighted in orange = variable reported in MH-CLD and incorporated in MH-TEDS.									

Appendix B
Table B1. MH-TEDS and MH-CLD 2011–2020 reporting methods and 2020 reporting period, by state or jurisdiction

State or jurisdiction	Data reporting method (C: MH-CLD, T: MH-TEDS)										2020 reporting period
	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	
Alabama	-	-	C	C	C	C	C	C	C	C	10/1/2019 to 9/30/2020
Alaska	-	-	-	-	-	-	-	-	T	T	7/1/2019 to 6/30/2020
American Samoa	-	-	-	-	-	-	-	-	-	-	-
Arizona	C	C	C	C	C	C	C	C	-	C	7/1/2019 to 6/30/2020
Arkansas	C	C	C	C	C	C	C	C	C	C	7/1/2019 to 6/30/2020
California	-	-	C	C	C	C	C	C	C	C	7/1/2018 to 6/30/2019
Colorado	C	-	C	C	C	C	C	C	C	C	7/1/2019 to 6/30/2020
Connecticut	-	C	C	T, C ¹	T, C ¹	T, C ¹	T, C ¹	T, C ¹	T, C ¹	T, C ¹	7/1/2019 to 6/30/2020
Delaware	-	C	C	C	C	T	T	T	T	T	7/1/2019 to 6/30/2020
District of Columbia	-	-	-	T*	T*	T	T	T	T	T	10/1/2019 to 9/30/2020
Federated States of Micronesia	-	-	-	-	-	-	-	-	-	-	-
Florida	C	C	C	C	C	C	C	C	T	T	7/1/2019 to 6/30/2020
Georgia	-	-	-	-	-	-	-	-	-	T	7/1/2019 to 6/30/2020-
Guam	-	-	-	-	-	-	-	-	-	-	-
Hawaii	-	C	C	C	C	C	C	C	C	C	7/1/2019 to 6/30/2020
Idaho	-	C	C	C	C	C	C	C	C	C	7/1/2019 to 6/30/2020
Illinois	-	-	C	C	C	C	C	C	C	C	7/1/2019 to 6/30/2020
Indiana	C	C	C	C	C	C	C	C	C	C	7/1/2019 to 6/30/2020
Iowa	-	-	C	C	C	C	C	C	-	C	7/1/2019 to 6/30/2020
Kansas	C	C	C	C	*	*	*	*	*	T	7/1/2019 to 6/30/2020
Kentucky	C	C	C	C	C	C	C	C	C	C	7/1/2019 to 6/30/2020
Louisiana	C	C	C	T	T	T	T	T	T	T	7/1/2019 to 6/30/2020
Maine	-	-	C	*	*	*	C	**	**	**	7/1/2019 to 6/30/2020
Marshall Islands	-	-	-	-	-	-	-	-	-	-	-
Maryland	-	C	C	C	C	C	C	C	C	***	<i>n.a</i>
Massachusetts	-	-	C	C	C	C	C	C	C	C	7/1/2019 to 6/30/2020
Michigan	-	-	-	-	-	-	T	T	T	T	10/1/2018 to 9/30/2019
Minnesota	-	-	-	C	C	C	C	C	C	C	1/1/2019 to 12/30/2019
Mississippi	-	-	C	T	T	T	T	T	T	T	7/1/2019 to 6/30/2020
Missouri	C	C	C	C	C	C	C	C	C	C	7/1/2019 to 6/30/2020
Montana	C	-	C	C	C	C	C	C	C	C	7/1/2019 to 6/30/2020
Nebraska	-	C	C	C	C	C	T	T	T	T	7/1/2019 to 6/30/2020

Continued. See notes at end of table.

Table B1. MH-TEDS and MH-CLD 2011–2020 reporting methods and 2020 reporting period , by state or jurisdiction (continued)

State or jurisdiction	Data reporting method (C: MH-CLD, T: MH-TEDS)										2020 reporting period
	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	
Nevada	C	C	C	C	C	C	C	C	C	C	7/1/2019 to 6/30/2020
New Hampshire	-	-	-	-	-	-	-	-	-	-	-
New Jersey ²	-	C	C	C	C	C	C	C	C	C	7/1/2019 to 6/30/2020
New Mexico	-	C	C	C	C	C	C	C	C	C	7/1/2019 to 6/30/2020
New York ³	-	C	C	C	C	C	C	C	C	C	4/1/2019 to 3/30/2019
North Carolina	-	-	C	C	C	T	T	T	T	T	7/1/2019 to 6/30/2020
North Dakota	-	C	C	C	C	C	C	C	C	C	7/1/2019 to 6/30/2020
Commonwealth of the Northern Mariana Islands	-	C	C	C	C	C	C	C	C	C	10/1/2019 to 9/30/2020
Ohio	-	C	C	C	C	C	C	C	C	C	7/1/2019 to 6/30/2020
Oklahoma	C	C	C	T	T	T	T	T	T	T	7/1/2019 to 6/30/2020
Oregon	-	-	C	C	C	C	C	C	C	C	1/1/2019 to 12/30/2019
Republic of Palau	-	-	-	-	-	C	C	C	C	C	10/1/2018 to 9/30/2019
Pennsylvania	-	-	C	C	*	T	T	T	T	T	7/1/2018 to 6/30/2019
Puerto Rico	-	-	-	T	T	T	T	T	T	T	7/1/2019 to 6/30/2020
Rhode Island	C	C	C	C	C	C	C	C	C	C	7/1/2019 to 6/30/2020
South Carolina	C	C	C	C	C	C	C	C	C	C	7/1/2019 to 6/30/2020
South Dakota	-	-	C	C	C	C	C	C	C	C	7/1/2019 to 6/30/2020
Tennessee ³	-	C	C	C	C	C	C	C	C	C	7/1/2019 to 6/30/2020
Texas	C	C	C	C	C	C	C	C	C	C	9/1/2019 to 8/30/2020
Utah	C	C	C	C	C	C	C	C	C	C	7/1/2019 to 6/30/2020
Vermont	-	C	C	C	C	T	T	T	T	T	7/1/2019 to 6/30/2020
Virginia	-	C	C	C	C	C	C	C	C	C	7/1/2019 to 6/30/2020
U.S. Virgin Islands	-	-	-	-	-	-	-	-	-	-	-
Washington	C	C	C	C	C	C	C	C	C	C	7/1/2019 to 6/30/2020
West Virginia	-	-	-	-	-	-	-	T	-	-	n.a
Wisconsin	C	C	C	C	C	C	C	C	C	C	1/1/2019 to 12/30/2019
Wyoming	C	C	C	C	C	C	C	C	C	C	7/1/2019 to 6/30/2020

n.a. Not reporting; - State is building capacity for future reporting of client-level data; * State is building capacity to switch to MH-TEDS reporting; ** State switched to building capacity (for MH-CLD reporting) due to staff turnover; *** The state transitioned to a new ASO and the ASO experienced extreme systems functionality issues affecting data collection.

T* District of Columbia: 2014 and 2015 reporting periods: 7/1–6/30. Reporting period changed in FY 2016.

¹ Connecticut, 2015–2020: MH-TEDS for adults, MH-CLD for children/adolescents.

² New Jersey reported 2017 and 2018 data as well as 2020 data in the 2020 reporting year.

³ New York and Tennessee report only a subset of the total individuals served through their SMHAs.

Appendix C

MH-TEDS and MH-CLD Data Dictionary of Included Variables

Variable Name: Age

Description: This field is calculated from the client's date of birth at midpoint of the state's elected reporting period.

Guidelines: Do not round up age. For example, if the calculated age is 13 and 8 months, report only 13. When the client's calculated age is less than 12 months, report age as zero.

If the reported date of birth of a client changes during the reporting period, use the most recent available information in calculating age.

Variable Name: Competitive Employment Status (Aged 16 + Years)

Description: This data element is reported for all individuals (16 years old and over) who are receiving services in non-institutional settings. Institutional settings include correctional facilities like prison, jail, detention centers, and mental health care facilities like state hospitals, other psychiatric inpatient facilities, nursing homes or other institutions that keep a person, otherwise able, from entering the labor force.

Full Time – use state definition for full time employment; includes members of the Armed Forces, and individuals in full-time Supported Employment.

Part Time – use state definition for part time; includes individuals in part-time Supported Employment.

Unemployed – defined as actively looking for work or laid off from job (and waiting to be recalled) in the past 30 days.

Not in labor force – defined as not employed and not actively looking for work during the past 30 days, includes Homemaker, Student, Retired, Disabled, Other Reported Classification (e.g., volunteers), Sheltered/Not-Competitive Employment, an inmate of an institution.

Guidelines: **Employment Status** is defined by SAMHSA as an outcome measure and is collected at admission and discharge/update to assess change.

Coding of individuals with overlapping employment statuses: When individuals are engaged in two or more activities (have overlapping status) during the period when their status is collected by the SMHA, use DOL's system of priorities to determine the appropriate employment status. The prioritization rule is, labor force activities (such as working or looking for work) take precedence over non-labor force activities (such as student and homemaker); and working or having a job takes precedence over looking for work.

Reporting of a person in an internship program: The Individual with Disabilities Education Act (IDEA) ceases to apply for young individuals (18–21 years old) once they received their high school diploma. The following rules should be observed in making determination whether the adult client in an internship program should be reported as ‘Not in Labor Force’ or ‘Employed’.

1. If the internship is a school requirement, whether paid or not, the person should be considered a “student” (Not In Labor Force).
2. If the internship is not a school requirement, an unpaid position, does not displace regular employees or does not entitle the person for the job at the end of the internship period, then report the person’s status as Not In Labor Force - (Other).
3. If the internship is not a school requirement, paid minimum wage and overtime provision, the employer benefits from the internship through the client’s engagement in actual operations of the business and performing productive work, then the person is employed (Full time or Part time).

Variable Name: Detailed Not In Labor Force

Description: This field provides more detailed information about those individuals who are coded as “Not in labor force” in Employment Status.

Homemaker, Student, Retired, Disabled

Resident of Institution – Individuals receiving services from institutional facilities such as hospitals, jails, prisons, long-term residential care, etc.

Other – includes volunteer, seasonal worker, other categories used by the state not specified.

Guidelines: **Detailed Not in Labor Force** is defined by SAMHSA as an outcome measure, and is collected at admission and at discharge/update to assess change.

This field is reported for individuals aged 16 years and older whose **Employment Status** is coded “Not in labor force.”

Variable Name: Co-occurring mental and substance use disorders

Description: This field indicates whether the client has co-occurring substance use and mental health disorders.

Yes – Client has co-occurring substance use and mental health disorders.

No – Client does not have co-occurring substance use and mental health disorders.

Guidelines: The assessment of co-occurring mental and substance use disorders may be based on clinical diagnoses, screening results, claims information, or self-report.

Variable Name: Ethnicity

Description: This field identifies whether or not the client is of Hispanic origin.

Guidelines: If the ethnicity of a client changes during the reporting period, report the most recent available information at the end of the reporting period.

Variable Name: Gender

Description: This field identifies the client's most recent reported sex at the end of the reporting period.

Guidelines: States that collect transgender as an option must report the data as follows: Transgender man (female sex at birth but identifies as male) – code as female; transgender woman (male sex at birth but identifies as female) – code as male.

If the gender of a client changes during the reporting period, report the most recent available information.

Variable Name: Living Arrangements (Residential Status)

Description: This field specifies client's residential status at time of admission (new individuals) or start of the reporting period (continuing individuals).

Homeless – person has no fixed address; includes homeless shelters.

Foster Home/Foster Care – individual resides in foster home. A foster home is a home that is licensed by a county or state department to provide foster care to children, adolescents, and/or individuals. This includes therapeutic foster care facilities. Therapeutic foster care is a service that provides treatment for troubled children within private homes of trained families.

Residential Care – individual resides in a residential care facility. This level of care may include a group home, therapeutic group home, board and care residential treatment, rehabilitation center, or agency-operated residential care facilities.

Crisis Residence – a time-limited residential (24 hours/day) stabilization program that delivers services for acute symptom reduction and restores clients to a precrisis level of functioning.

Institutional Setting – individual resides in an institutional care facility with care provided on a 24 hour, 7 days a week basis. This level of care may include skilled nursing/intermediate care facility, nursing homes, institute of mental disease (IMD), inpatient psychiatric hospital, psychiatric health facility, veterans' affairs hospital, state hospital, or intermediate care facility/MR.

Jail/Correctional Facility – individual resides in a jail and/or correctional facility with care provided on a 24 hour, 7 days a week basis. This includes a jail, correctional facility, detention centers, and prison.

Dependent Living: Private Residence – adult individuals living in a house, apartment, or other similar dwelling who are heavily dependent on others for daily living assistance.

Independent Living: Private Residence – individuals living alone or with others in a private residence and capable of self-care. Includes adult children (age 18 years and over) living with parents and adolescents living independently. Also includes individuals who live independently with case management and supported housing support.

Private Residence – for adult with unknown dependency status, or for all children living in a private residence regardless of living arrangement.

Other – Other residential status

Guidelines: **Living Arrangements** is defined by SAMHSA as an outcome measure and is collected at admission and at discharge/update to assess change.

Variable Name: Marital Status (Optional Reporting)

Description: This field identifies the client’s marital status. The following categories are compatible with categories used in the U.S. Census.

Never married – includes individuals who are single or whose only marriage was annulled.

Currently married – includes married couples, those living together as married, living with partners, or cohabitating.

Separated – includes those legally separated or otherwise absent from spouse because of marital discord.

Divorced – legally divorced.

Widowed

Guidelines: Report marital status at time of discharge or the most recent available status at the end of the reporting period.

Variable Name: Mental Health Diagnosis

Description: This field specifies the client’s current mental health diagnoses during the reporting period.

Guidelines: This measure can be reported by using either the *Diagnostic and Statistical Manual of Mental Disorders (DSM)* from the American Psychiatric Association or the *International Classification of Diseases (ICD)* from the World Health Organization.

States should report the most recent mental health diagnoses that are current during the reporting period. Most recent is defined by the date when the diagnosis is reported.

States can report up to three mental health diagnoses by completing the Mental Health Diagnosis 1, 2, and 3 data elements in sequential order.

If the client has more than three most recent mental health diagnoses, use the algorithm below. Use this if the diagnoses are collected through administrative method (i.e., based on the clinician's evaluation of the person and reported in the client's case record):

1. Report the primary and secondary diagnoses (if available). If the state does not classify diagnosis into primary/secondary; then
2. Conduct your search for diagnostic codes using both Axis I and II. Report all diagnoses in Axis I (clinical disorders) first followed by diagnosis in Axis II (personality disorders and mental retardation) unless a personality disorder in Axis II was labeled as primary diagnosis, then it should be reported first (note: this method is applicable if using DSM-IV and other earlier editions or if the state continues to use the DSM axial structure;
If primary/secondary labels and Axis classifications (I and II) are not used by the state, then
3. Report in chronological order starting from the diagnosis that appears on top of the list or first cited in the clinician's report. Report only mental health diagnosis.

If the state is using claims/encounter data to collect a client's diagnosis, use the three most frequently reported mental health diagnoses in the client's service claims/encounters data throughout the reporting period.

Variable Name: Race

Description: This field specifies the client's most recent reported race at the end of the reporting period.

Alaska Native – used in MH-TEDS only

American Indian or Alaska Native – A person having origins in any of the original people of North and South America (including Central America and the original people of Alaska) and who maintain cultural identification through tribal affiliation or community attachment.

Asian – A person having origins in any of the following people of the Far East, the Indian Subcontinent, or Southeast Asia, including Cambodia, China, India, Japan, Korea, Malaysia, Philippine Islands, Thailand, and Vietnam.

Asian or Pacific Islander (temporary code) – older racial category used only by some states that have not yet adopted the OMB recommended racial categories.

Black or African American – A person having origins in any of the Black racial groups of Africa.

Hispanic (temporary code) – used in MH-CLD only

Native Hawaiian or Other Pacific Islander – A person having origins in any of the original peoples of Hawaii, Guam, American Samoa, or Other Pacific Islander.

White – A person having origins in any of the original people of Europe, North Africa or the Middle East.

Some Other Race Alone – use this category for instances in which the client is not identified in any category above or whose origin group, because of area custom, is regarded as racial class distinct from the above categories (do not use this category for individuals indicating multiple or mixed races).

Two or more races – use this code when the state data system allows multiple race selection and more than one race is indicated (see guidelines).

Guidelines: For states that collect “Other Race” or allow individuals to specify a single race different from the OMB racial categories, use “Some Other Race Alone.” Note that this category should not be used if the client is indicating multiple or mixed race.

[MH-TEDS only] States that collect “Alaska Native” as a separate category should use “American Indian/Alaska Native” for other American Indians.

In cases where the method of collecting race information differs between community programs and state hospitals, or where the race reported differs between data systems, the state should resolve the issue in accordance to the state business rule in resolving data quality issues. States may use the following guidelines in the absence of a state business rule or to supplement an existing one:

1. Use the most recent race reported if variation in a person’s race was reported by the same provider across time
2. Use the most frequently reported race if variation in a person’s race was reported by different providers across time

Variable Name: Treatment Status at the End of the Reporting Period

Description: This field indicates client status at the end of the reporting period—continuing to receive services or reason for discontinuance of treatment.

Continuing Client

Discharged with Treatment Completed – All parts of the treatment plan or program were completed.

Discharged due to Lost Contact/Administrative Discharge

Discharged to Corrections, Jail – Individuals whose course of treatment is terminated because the client has been subject to jail, prison, or house confinement, or has been released by or to the courts.

Discharged due Death Discharged due to Aged Out

Discharged due to Other Specified Reasons Discharged, Reason Unknown

Discharged, Reason not Collected

Variable Name: School Grade Level – Education

Description: This field specifies the school grade level of three sub-populations of individuals, as follows:

Current grade level for school-age children (3–17 years old and 18–21 years of age if the client is in special education) who attended school at any time in the past three months.

Highest grade level completed for school-age children who have not attended school at any time within the past three months.

Highest educational attainment for all individuals, whether currently in school or not.

Guidelines: Information for this data element should be collected and reported for all individuals.

“School” includes, but is not limited to, any one or combination of home-schooling, online education, alternative school, vocational school, or regular school (public, private, charter, traditional, military, magnet, independent, parochial, etc.), at which the child is enrolled in any of the following school grade levels: nursery/preschool (including Head Start), kindergarten, elementary/middle school (Grades 1–8), high school (Grades 9–12, including General Equivalency Degree or GED), vocational school (including business, technical, secretarial, trade, or correspondence courses which are not counted as regular school enrollment and are not for recreation or adult education classes), or college/professional degree.

This measure is to be reported only once during the reporting period, either at time of client discharge or at the end of the reporting period if the client is continuing to receive services.

If the information is collected at the time when the school year just ended, report the recent school grade level completed (not the grade level the child is advancing to in the next school year).

For children who are homeschooled or children in Special Education but have been mainstreamed in regular school grades, please report the equivalent grade level.

“A nursery school is defined as a group or class that is organized to provide educational experiences for children during the year or years preceding kindergarten. It includes instruction as an important and integral phase of its program of child care. Private homes in which essentially custodial care is provided are not considered nursery schools. Children attending nursery school are classified as attending during either part of the day or the full day. Part-day attendance refers to those who attend either in the morning or in the afternoon, but not both. Full-day attendance refers to those who attend in both the morning and the afternoon. Children enrolled in Head Start programs or similar programs sponsored by local agencies to provide preschool education to young children are counted under nursery school” (The American Community Survey [ACS]).

A person who had reached the age of 18 years prior to collecting this information should report the last grade completed, unless they meet the exception criteria for young individuals.

Report the highest educational attainment completed for all individuals 18 years and older (except the young individuals aged 18–21 years in Special Education and

receiving mental health services from the Children Mental Health System) whether they are currently in school or not.

Variable Name: Service Setting Status – Mental Health Service Setting

Description: This field indicates the type of mental health treatment setting(s) in which the client received services throughout the reporting period.

State Psychiatric Hospital – all SMHA-funded and SMHA-operated organizations operated as hospitals that provide primary inpatient care to individuals with mental illness from a specific geographical area and/or statewide.

SMHA-Funded/Operated Community-Based Program – include community mental health centers (CMHCs), outpatient clinics, partial care organizations, partial hospitalization programs, PACT programs, consumer run programs (including Club Houses and drop-in centers), and all community support programs (CSP).

Residential Treatment Center – an organization, not licensed as a psychiatric hospital, whose primary purpose is the provision of individually planned programs of mental health treatment services in conjunction with residential care for children and youth, and in some cases, adult care.

Other Psychiatric Inpatient – a private provider or medical provider licensed and/or contracted through the SMHA. This includes public/private hospitals with separate psychiatric wards.

Institutions under the Justice System – mental health services provided in a jail, prison, juvenile detention center, etc.

Guidelines: This data element refers to the place where services were provided and not particularly the residence of the client.

Variable Name: SMI/SED Status

Description: This field indicates whether the client has serious mental illness (SMI) or serious emotional disturbance (SED) using the state's definition.

Guidelines: Report the most recent available SMI/SED status at the end of the reporting period.

Variable Name: Treatment Status at the Start of the Reporting Period

Description: This field indicates whether the client is already receiving services at the start of the reporting period (continuing) or is a new admission.

New admission – Admission that happened on the first day of the reporting period or thereafter.

Continuing – Already receiving services at the start of the reporting period. The admission date should have occurred at least a day prior to the start of the reporting period.

Variable Name: Veteran Status

Description: This field specifies whether the client has served in the uniform services (Army, Navy, Air Force, Marine Corps, Coast Guard, Public Health Service Commissioned Corps, Coast and Geodetic Survey, etc.).

Guidelines: A veteran is a person 16 years or over who has served (even for a short time), but is not now serving, on active duty in the U.S. Army, Navy, Air Force, Marine Corps, Coast Guard, or Commissioner Corps of the U.S. Public Health Service or the National Oceanic and Atmospheric Administration, or who served as a Merchant Marine seaman during World War II. Individuals who served in the National Guard or Military Reserves are classified as veterans only if they have ever been called or ordered to active duty (excluding the 4-6 months of initial training and yearly summer camps)

Appendix C Table 1. Number of states reporting data, MH-CLD and MH-TEDS 2015–2020

<i>Data item</i>	<i>Number of states</i>					
	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>
Any data	45	47	49	49	48	50
Service settings	45	47	49	49	48	50
Timing of admission	45	47	49	49	48	50
First mental health diagnosis	45	47	49	49	48	50
Gender	45	47	49	49	48	50
Age	45	47	49	49	48	50
Race	45	47	49	49	48	50
Ethnicity	43	45	48	47	47	49
Detailed Hispanic	29	28	28	27	28	27
Marital status	37	40	42	42	43	44
SED/SMI status	45	47	48	49	48	50
Co-occurring substance use	44	44	46	46	46	49
Employment	45	47	49	49	48	50
Detailed not in labor force	44	46	48	48	48	49
Living arrangement	45	47	48	49	48	50
Education	34	39	42	42	43	44
Veteran status	n/c	n/c	23	35	36	38

n/c Not collected.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015–2020 reporting periods. Data as of 03.03.2022.

Appendix C Table 2. Item percentage response rate, by state or jurisdiction: MH-CLD and MH-TEDS 2020

[Based on administrative data reported to MH-CLD or MH-TEDS by all reporting states and jurisdictions. Comparisons among states should be made with caution.]

State or jurisdiction	Service setting	Timing of admission	First-mentioned mental health diagnosis ¹	Gender	Age	Race	Ethnicity	Marital status	SED/SMI	Co-occurring substance use	Employment status	Residence	Education	Veteran status
Total	100	100	98	100	100	91	90	58	95	87	61	60	47	38
Alabama	100	100	100	100	100	100	100	100	93	100	85	77	99	85
Alaska	100	100	100	100	100	98	96	98	100	100	98	95	94	89
Arizona	100	100	99	100	100	65	66	57	100	100	50	†	†	22
Arkansas	100	100	100	100	100	79	87	†	100	67	87	93	†	†
California	100	100	100	100	100	84	81	1	100	100	51	50	†	†
Colorado	100	100	100	100	100	100	90	100	100	100	100	100	99	99
Commonwealth of the Northern Mariana Islands	100	100	85	100	100	100	†	80	85	97	91	81	†	†
Connecticut	100	100	100	100	100	94	94	55	99	63	76	72	60	56
Delaware	100	100	100	99	100	92	85	87	95	100	80	81	65	70
District of Columbia	100	100	86	100	100	93	44	59	100	10	42	5	8	†
Florida	100	100	100	100	100	100	99	96	100	100	87	91	92	96
Georgia	100	100	100	100	100	97	94	93	94	100	57	96	93	84
Hawaii	100	100	100	98	100	69	65	57	99	41	30	25	49	†
Idaho	100	100	100	100	100	89	86	46	35	44	99	33	4	81
Illinois	100	100	100	100	100	90	90	93	93	84	91	88	79	89
Indiana	100	100	100	100	100	100	99	99	97	99	90	90	89	59
Iowa	100	100	100	100	100	75	78	†	100	11	36	93	†	†
Kansas	100	100	100	100	100	99	98	22	88	†	78	82	88	66
Kentucky	100	100	100	99	100	99	97	99	96	98	97	97	93	98
Louisiana	100	100	100	100	100	93	94	56	52	100	57	46	48	83
Massachusetts	100	100	100	100	100	83	80	90	100	100	6	64	50	86
Michigan	100	100	100	100	100	99	92	97	96	98	99	100	94	94
Minnesota	100	100	100	100	100	82	92	†	100	100	99	99	95	†
Mississippi	100	100	100	100	100	99	95	99	81	49	96	98	91	97
Missouri	100	100	100	100	100	97	100	65	92	100	72	82	54	†
Montana	100	100	100	100	100	92	100	†	100	100	37	19	7	24
Nebraska	100	100	100	97	100	94	73	85	67	100	88	87	77	79
Nevada	100	100	100	100	100	79	78	66	68	90	56	49	12	8
New Jersey	100	100	99	100	100	79	91	79	90	86	17	74	72	75
New Mexico	100	100	100	100	100	88	88	33	100	100	36	100	34	98

Continued. See notes at end of table.

Appendix C Table 2. Item percentage response rate, by state or jurisdiction: MH-CLD and MH-TEDS 2020 (continued)

[Based on administrative data reported to MH-CLD or MH-TEDS by all reporting states and jurisdictions. Comparisons among states should be made with caution.]

<i>State or jurisdiction</i>	<i>Service setting</i>	<i>Timing of admission</i>	<i>First-mentioned mental health diagnosis¹</i>	<i>Gender</i>	<i>Age</i>	<i>Race</i>	<i>Ethnicity</i>	<i>Marital status</i>	<i>SED/SMI</i>	<i>Co-occurring substance use</i>	<i>Employment status</i>	<i>Residence</i>	<i>Education</i>	<i>Veteran status</i>
Total	100	100	98	100	100	91	90	58	95	87	61	60	47	38
New York	100	100	100	100	100	93	89	92	100	100	94	81	88	20
North Carolina	100	100	99	100	100	100	100	99	100	†	99	97	98	99
North Dakota	100	100	100	100	100	99	100	98	99	100	93	79	92	90
Ohio	100	100	100	98	100	90	93	37	100	100	25	2	†	†
Oklahoma	100	100	100	100	100	100	100	86	84	100	86	84	83	97
Oregon	100	100	100	100	100	91	91	73	100	99	73	85	76	11
Palau	100	100	100	100	100	100	100	59	100	18	42	58	10	42
Pennsylvania	100	100	100	100	100	98	100	99	100	100	34	†	†	†
Puerto Rico	100	100	100	100	100	95	100	97	96	98	84	98	95	79
Rhode Island	100	100	100	100	100	85	90	86	94	84	80	79	†	88
South Carolina	100	100	100	100	100	98	77	97	100	100	78	93	89	100
South Dakota	100	100	100	100	100	100	100	94	90	51	54	35	6	†
Tennessee	100	100	64	100	100	93	57	51	62	69	52	60	34	47
Texas	100	100	72	100	100	100	100	80	95	72	73	72	72	†
Utah	100	100	100	100	100	92	90	90	91	100	93	94	85	90
Vermont	100	100	100	99	100	75	67	80	†	100	53	60	2	64
Virginia	100	100	100	100	100	99	97	†	94	100	97	93	90	†
Washington	100	100	100	99	99	97	95	41	86	100	43	27	25	23
Wisconsin	100	100	100	100	100	94	88	†	100	100	44	39	†	29
Wyoming	100	100	100	100	100	94	95	97	99	100	100	99	97	45

† Less than 1 percent of individuals.

¹ The first-mentioned mental health diagnosis is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD). Data received 03.03.2022 as part of 2020 reporting period.

APPENDIX D

SMI/SED Federal Definitions

Note that identification of individuals with a serious mental illness (SMI) or serious emotional disturbance (SED) is based on each state or jurisdiction's definitions for SMI and SED; some states and jurisdictions may use definitions that vary from the federal definitions that appear in this appendix.

Serious Mental Illness (SMI)

Pursuant to section 1912 (c) of the Public Health Service Act, as amended by Public Law 102-321 "adult[s] with a serious mental illness" are individuals:

1. age 18 years and older;
2. who currently or at any time during the past year, have had a diagnosable mental, behavioral, or emotional disorder of sufficient duration to meet diagnostic criteria specified within DSM-III-R; and
3. this disorder has resulted in functional impairment which substantially interferes with or limits one or more major life activities.

"These disorders include any mental disorders (including those of biological etiology) listed in DSM-III-R or their ICD-9-CM equivalent (and subsequent revisions), with the exception of DSM-III-R "V" codes, substance use disorders, and developmental disorders, which are excluded, unless they co-occur with another diagnosable serious mental illness..." (p. 29425).

Serious Emotional Disturbance (SED)

Pursuant to section 1912 (c) of the Public Health Service Act, as amended by Public Law 102-321 "children with a serious emotional disturbance" are individuals:

4. from birth up to age 18;
5. who currently or at any time during the past year have had a diagnosable mental, behavioral, or emotional disorder of sufficient duration to meet diagnostic criteria specified within DSM-III-R; and
6. this disorder resulted in functional impairment, which substantially interferes with or limits the child's role or functioning in family, school, or community activities (p.29425).

"These disorders include any mental disorder (including those of biological etiology) listed in DSM-III-R or their ICD-9-CM equivalent (and subsequent revisions) with the exception of DSM-III-R 'V' codes, substance use, and developmental disorders, which are excluded, unless they co-occur with another diagnosable serious emotional disturbance..." (p. 29425).

"Functional impairment is defined as difficulties that substantially interfere with or limit a child or adolescent from achieving or maintaining one or more developmentally appropriate social,

behavioral, cognitive, communicative, or adaptive skills. Functional impairments of episodic, recurrent, and continuous duration are included unless they are temporary and expected responses to stressful events in their environment. Children who would have met functional impairment criteria during the referenced year without the benefit of treatment or other support services are included in this definition....” (p. 29425) [Source: Federal Register 58(96): 29422–29425. May 20, 1993; <https://www.samhsa.gov/sites/default/files/federal-register-notice-58-96-definitions.pdf>].

APPENDIX E

Mental Health and Substance Use Diagnosis Codes

Mental Health Diagnostic Groups With Corresponding ICD-9 Diagnostic Codes

The listings below group ICD-9 codes into a modified form of Clinical Classification Software (CCS) mental health “multi-level” diagnostic category codes (5.1, 5.2, and so on) and corresponding labels (trauma- and stressor-related disorders, anxiety disorders, and so on). Notes:

- **Bold** diagnostic groups are displayed in one or more tables in this report; records with ICD-9 codes in other diagnostic groups are grouped as “other” mental health diagnoses.
- *Italicized codes* in parentheses are nonstandard ICD-9 codes in the data that were recognized to belong to the same CCS mental health diagnostic group.
- Entries in brackets ([650], [651], and so on) are single-level diagnosis categories that are otherwise ignored in this report but are noted here for convenience.
- The ICD-9 diagnostic codes shown below have omitted the dot separator after the first three digits.
- The groupings of certain DSM-IV/ICD-9 codes have been reorganized according to DSM-5 nomenclature.

5.1 **Trauma- and stressor-related disorders** [650]^{16, 17}

3083	3090	3091	30922	30923	30924	30928	30929	3093
3094	30981	30982	30983	30989	3099			
(30830	309	30900	30910	30920	30930	30940	30980	3098
30990)								

5.2 **Anxiety disorders** [651]¹⁸

29384	30000	30001	30002	30009	30010	30020	30021	30022
30023	30029	3003	3005	30089	3009	3080	3081	3082
3084	3089	30921	3130	3131	31321	31322	3133	31382
31383								
(3000	300	3001	3002	30030	30050	30090	30800	308
30810	30820	30840	30890	313	31300	3138	31380)	

¹⁶ For these and other bolded (i.e., tabulated) CCS diagnostic groups, table column and row headers drop the word “disorders” from the description.

¹⁷ Codes for posttraumatic stress disorder and acute stress disorder diagnoses have been grouped under *trauma- and stressor-related disorders*, consistent with DSM-5. Codes for adjustment disorders have also been subsumed into this diagnostic group.

¹⁸ The code for separation anxiety disorder has been grouped with anxiety disorders, consistent with DSM-5.

5.3 Attention deficit, conduct, and disruptive behavior disorders [652]

5.3.1 Conduct disorder [6521]

31200	31201	31202	31203	31210	31211	31212	31213	31220
31221	31222	31223	3124	3128	31281	31282	31289	3129
(312	3120	3121	3122	31240	31280	31290)		

5.3.2 Oppositional defiant disorder [6522]

31381

5.3.3 Attention deficit/hyperactivity disorder [6523]

31400	31401	3141	3142	3148	3149
(314	3140	31410	31480	31490)	

5.4 Delirium, dementia, and amnestic and other cognitive disorders [653]¹⁹

2900	29010	29011	29012	29013	29020	29021	2903	29040
29041	29042	29043	2908	2909	2930	2931	2940	2941
29410	29411	29420	29421	2948	2949	3100	3102	3108
31081	31089	3109	3310	3311	33111	33119	3312	33182
797								
(29000	290	2901	2902	29030	2904	29080	29090	293
29300	29400	294	2942	29480	29490	310	31000	31020
31090	331	33100	33110)					

5.5 Developmental disorders [654]

5.5.1 Communication disorders [6541]

3070	3079	31531	31534	31535	31539	V401
(30700	307	30790)				

5.5.2 Developmental disabilities [6542]

31501	31502	31509	31532	3155	3158
(31550	31580)				

5.5.3 Intellectual disabilities [6543]

317	3180	3181	3182	319		
(3170	31700	318	31800	31810	31820	31900)

5.5.4 Learning disorders [6544]

31500	3151	3152	3159	V400	
(3150	315	31510	31520	31540	V40)

5.5.5 Motor skill disorders [6545]

3154
(31540)

¹⁹ Referred to as “delirium/dementia” in tables and text.

5.6 Disorders usually diagnosed in infancy, childhood, or adolescence [655]

5.6.1 Elimination disorders [6551]

3076 3077
(30760 30770)

5.6.2 Other disorders of infancy, childhood, or adolescence [6552]

3073 31323 31389 3139
(31390)

5.6.3 Pervasive developmental disorders [6553]

29900 29901 29910 29911 29980 29981 29990 29991
(299 2991 2998 2999)

5.6.4 Tic disorders [6554]

30720 30721 30722 30723

5.7 Impulse control disorders, not elsewhere classified [656]

31230 31231 31232 31233 31234 31235 31239
(3123)

5.8 Mood disorders [657]

5.8.1 Bipolar disorders [6571]

29600 29601 29602 29603 29604 29605 29606 29610 29611
29612 29613 29614 29615 29616 29640 29641 29642 29643
29644 29645 29646 29650 29651 29652 29653 29654 29655
29656 29660 29661 29662 29663 29664 29665 29666 2967
29680 29681 29682 29689 29690 29699
(296 2960 2961 2964 2965 2966 29670 2968 2969)

5.8.2 Depressive disorders [6572]

29383 29620 29621 29622 29623 29624 29625 29626 29630
29631 29632 29633 29634 29635 29636 3004 311
(2962 2963 30040 31100 3110)

5.9 Personality disorders [658]

3010 30110 30111 30112 30113 30120 30121 30122 3013
3014 30150 30151 30159 3016 3017 30181 30182 30183
30184 30189 3019
(30100 301 3011 3012 30130 30140 3015 30160 30170
30190)

5.10 Schizophrenia and other psychotic disorders [659]²⁰

29381	29382	29500	29501	29502	29503	29504	29505	29510
29511	29512	29513	29514	29515	29520	29521	29522	29523
29524	29525	29530	29531	29532	29533	29534	29535	29540
29541	29542	29543	29544	29545	29550	29551	29552	29553
29554	29555	29560	29561	29562	29563	29564	29565	29570
29571	29572	29573	29574	29575	29580	29581	29582	29583
29584	29585	29590	29591	29592	29593	29594	29595	2970
2971	2972	2973	2978	2979	2980	2981	2982	2983
2984	2988	2989						
(295	2950	2951	2952	2953	2954	2955	2956	2957
2958	2959	29700	297	29710	29720	20730	20780	20790
29800	298	29810	29820	29830	29840	29880	29890)	

5.11 Alcohol-related disorders [660]²¹

2910	2911	2912	2913	2914	2915	2918	29181	29182
29189	2919	30300	30301	30302	30303	30390	30391	30392
30393	30500	30501	30502	30503	3575	4255	5353	53530
53531	5710	5711	5712	5713	76071	9800		
(29100	291	29110	29120	29130	29140	29150	29180	20190
303	3030	3039	305	3050)				

5.12 Substance-related disorders [661]²²

2920	29211	29212	2922	29281	29282	29283	29284	29285
29289	2929	30400	30401	30402	30403	30410	30411	30412
30413	30420	30421	30422	30423	30430	30431	30432	30433
30440	30441	30442	30443	30450	30451	30452	30453	30460
30461	30462	30463	30470	30471	30472	30473	30480	30481
30482	30483	30490	30491	30492	30493	30520	30521	30522
30523	30530	30531	30532	30533	30540	30541	30542	30543
30550	30551	30552	30553	30560	30561	30562	30563	30570
30571	30572	30573	30580	30581	30582	30583	30590	30591
30592	30593	64830	64831	64832	64833	64834	65550	65551
65553	76072	76073	76075	7795	96500	96501	96502	96509
V6542								
(292	29200	20220	20290	304	3040	3041	3042	3043
3044	3045	3046	3047	3048	3049	305	3052	3053
3054	3055	3056	3057	3058	3059	6555	965)	

²⁰ Referred to as “Schizophrenia, other psychotic” in tables.

²¹ Regardless of their “co-occurring” flag in the data, individuals reported to be diagnosed with only alcohol-related disorders and/or substance-related disorders are not counted as having co-occurring mental health and substance use disorders for the purpose of Chapter 6; some other, non-alcohol and non-substance-related disorder diagnosis is necessary.

²² See footnote 21.

5.13 Suicide and intentional self-inflicted injury [662]

E9500	E9501	E9502	E9503	E9504	E9505	E9506	E9507	E9508
E9509	E9510	E9511	E9518	E9520	E9521	E9528	E9529	E9530
E9531	E9538	E9539	E954	E9550	E9551	E9552	E9553	E9554
E9555	E9556	E9557	E9559	E956	E9570	E9571	E9572	E9579
E9580	E9581	E9582	E9583	E9584	E9585	E9586	E9587	E9588
E9589	E959	V6284						

5.14 Screening and history of mental health and substance use codes [663]

33392	V110	V111	V112	V114	V118	V119	V154	V1541
V1542	V1549	V1582	V6285	V663	V701	V702	V7101	V7102
V7109	V790	V792	V793	7903	V113	V798	V791	V799
3051	30510	30511	30512	30513				

5.15 Miscellaneous mental disorders [670]

29389	2939	30011	30012	30013	30014	30015	30016	30019
3006	3007	30081	30082	3021	3022	3023	3024	30250
30251	30252	30253	3026	30270	30271	30272	30273	30274
30275	30276	30279	30281	30282	30283	30284	30285	30289
3029	3060	3061	3062	3063	3064	30650	30651	30652
30653	30659	3066	3067	3068	3069	3071	30740	30741
30742	30743	30744	30745	30746	30747	30748	30749	30750
30751	30752	30753	30754	30759	30780	30781	30789	3101
316	64840	64841	64842	64843	64844	V402	V403	V4031
V4039	V409	V673						
(306	316	3025	3027	3075	3078	3160	6484	29390
30060	30070	30210	30220	30230	30240	30260	30290	30600
30610	30630	30640	30680	30690	30710	3074	31010	31600)

Mental Health Diagnostic Groups with Corresponding ICD-10 Diagnostic Codes

The listings below group ICD-10 codes into a modified form of Clinical Classification Software (CCS) mental health “multi-level” diagnostic category codes (5.1, 5.2, and so on) and corresponding labels (trauma- and stressor-related disorders, anxiety disorders, and so on). Notes:

- **Bold** diagnostic groups are displayed in one or more tables in this report; records with ICD-10 codes in other diagnostic groups are grouped as “other” mental health diagnoses.
- *Italicized* codes in parentheses are nonstandard ICD-10 codes in the data that were recognized to belong to the same CCS mental health diagnostic group.
- Entries in brackets ([650], [651], and so on) are single-level diagnosis categories that are otherwise ignored in this report but are noted here for convenience.
- The ICD-10 diagnostic codes shown below have omitted the dot separator after the first three digits.
- Some diagnostic groups implemented by CCS for ICD-9 codes were not implemented for ICD-10 codes (conduct disorders, oppositional defiant disorders, attention deficit and attention deficit hyperactivity disorders, pervasive developmental disorders, bipolar disorders, and depressive disorders). ICD-10 codes from broader CCS diagnostic categories were assigned to these more specific diagnostic groups, based on the ICD-10 code’s detailed meaning.
- The groupings of certain ICD-10 codes have been reorganized according to DSM-5 nomenclature.

5.1 Trauma- and stressor-related disorders [650]^{23, 24}

F4310	F4320	F4321	F4322	F4323	F4324	F4325	F4329	F438
F439								
(F431	F432)							

5.2 Anxiety disorders [651]²⁵

F064	F4000	F4001	F4002	F4010	F4011	F40210	F40218	F40220
F40228	F40230	F40231	F40232	F40233	F40240	F40241	F40242	F40243
F40248	F40290	F40291	F40298	F408	F409	F410	F411	F413
F418	F419	F42	F422	F423	F424	F428	F429	F430
F4311	F4312	F488	F489	F930	R452	R453	R454	R455
R456	R457	R4581	R4582	R4583	R4584			
(F40	F401	F4021	F4022	F4023	F4024	F4029	F41	F93)

5.3.1 Conduct disorder [6521]²⁶

F910	F911	F912	F918	F919

²³ For these and other bolded (i.e., tabulated) CCS diagnostic groups, table column and row headers drop the word “disorders” from the description.

²⁴ Codes for posttraumatic stress disorder and acute stress disorder diagnoses have been grouped under *trauma- and stressor-related disorders*, consistent with DSM-5. Codes for adjustment disorders have also been subsumed into this diagnostic group.

²⁵ The code for separation anxiety disorder has been grouped with anxiety disorders, consistent with DSM-5.

²⁶ Codes in this category were assigned from a more general CCS diagnostic group to this one based on the code’s detailed meaning.

5.3.2 **Oppositional defiant disorder [6522]**

F913

5.3.3 **Attention deficit/hyperactivity disorder [6523]²⁵**

F900	F901	F902	F908	F909	R460	R461	R462	R463
R464	R465	R466	R467	R4681	R4689			
(F90	R46)							

5.4 **Delirium, dementia, and amnestic and other cognitive disorders [653]²⁷**

F0150	F0151	F0280	F0281	F0390	F0391	F04	F05	F070
F0781	F0789	F079	F09	F482	G300	G301	G308	G309
G3101	G3109	G311	G3183	R4181	R54			
(F01	F07	G30)						

5.5 **Developmental disorders [654]**

F70	F71	F72	F73	F78	F79	F800	F801	F802
F804	F8081	F8089	F809	F810	F812	F8181	F8189	F819
F82	F88	F89	F985	R4183	R480			
(F80	F81	R48)						

5.6 **Disorders usually diagnosed in infancy, childhood, or adolescence [655]**

F642	F842	F938	F939	F940	F941	F942	F948	F949
F950	F951	F952	F958	F959	F980	F981	F9821	F9829
F983	F984	F988	F989					
(F94	F95	F98)						

5.6.3 **Pervasive developmental disorders [6553]**

F840	F843	F845	F848	F849
(F84)				

5.7 **Impulse control disorders not elsewhere classified [656]**

F630	F631	F632	F633	F6381	F6389	F639	R45850
(F63	R4585)						

5.8 **Mood disorders [657]²⁸**

F0630	F0631	F0632	F0633	F0634	F3010	F3011	F3012	F3013
F302	F303	F304	F308	F309	F340	F341	F348	F3481
F3489	F349	F39	R4586					
(F063	F301	F34)						

²⁷ Referred to as “delirium/dementia” in tables and text.

²⁸ Mood disorder codes that could not be assigned to bipolar or depressive disorders.

5.8.1 Bipolar disorders [6571]

F310	F3110	F3111	F3112	F3113	F312	F3130	F3131	F3132
F314	F315	F3160	F3161	F3162	F3163	F3164	F3170	F3171
F3172	F3173	F3174	F3175	F3176	F3177	F3178	F3181	F3189
F319								
<i>(F31</i>	<i>F311</i>	<i>F313</i>	<i>F316</i>	<i>F317</i>	<i>F318)</i>			

5.8.2 Depressive disorders [6572]

F320	F321	F322	F323	F324	F325	F328	F3281	F3289
F329	F330	F331	F332	F333	F3340	F3341	F3342	F338
F339								
<i>(F32</i>	<i>F33</i>	<i>F334)</i>						

5.9 Personality disorders [658]

F600	F601	F602	F603	F604	F605	F606	F607	F6081
F6089	F609	F69						
<i>(F60)</i>								

5.10 Schizophrenia and other psychotic disorders [659]²⁹

F060	F062	F200	F201	F202	F203	F205	F2081	F2089
F209	F21	F22	F23	F24	F250	F251	F258	F259
F28	F29							
<i>(F06</i>	<i>F20</i>	<i>F25)</i>						

5.11 Alcohol-related disorders [660]³⁰

F1010	F1011	F10120	F10121	F10129	F1014	F10150	F10151	F10159
F10180	F10181	F10182	F10188	F1019	F1020	F1021	F10220	F10221
F10229	F10230	F10231	F10232	F10239	F1024	F10250	F10251	F10259
F1026	F1027	F10280	F10281	F10282	F10288	F1029	F10920	F10921
F10929	F1094	F10950	F10951	F10959	F1096	F1097	F10980	F10981
F10982	F10988	F1099	G621	I426	K2920	K2921	K700	K7010
K7011	K702	K7030	K7031	K7040	K709	O99310	O99311	O99312
O99313	O99314	O99315	P043	Q860				
<i>(F10</i>	<i>F101</i>	<i>F1012</i>	<i>F1015</i>	<i>F1018</i>	<i>F102</i>	<i>F1022</i>	<i>F1023</i>	<i>F1025</i>
<i>F1028</i>	<i>F109</i>	<i>F1092</i>	<i>F1095</i>	<i>F1098</i>	<i>K70</i>	<i>K703</i>	<i>K704</i>	<i>O9931</i>
<i>Q86)</i>								

5.12 Substance-related disorders [661]³¹

F1110	F1111	F11120	F11121	F11122	F11129	F1113	F1114	F11150
F11151	F11159	F11181	F11182	F11188	F1119	F1120	F1121	F11220
F11221	F11222	F11229	F1123	F1124	F11250	F11251	F11259	F11281
F11282	F11288	F1129	F1190	F11920	F11921	F11922	F11929	F1193
F1194	F11950	F11951	F11959	F11981	F11982	F11988	F1199	F1210

²⁹ Referred to as “Schizophrenia, other psychotic” in tables.

³⁰ Regardless of their “co-occurring” flag in the data, individuals reported to be diagnosed with only alcohol-related disorders and/or substance-related disorders are not counted as having co-occurring mental health and substance use disorders for the purpose of Chapter 6; some other, non-alcohol and non-substance-related disorder diagnosis is necessary.

³¹ See footnote 26.

F12120	F12121	F12122	F12129	F1213	F12150	F12151	F12159	F12180
F12188	F1219	F1220	F1221	F12220	F12221	F12222	F12229	F1223
F12250	F12251	F12259	F12280	F12288	F1229	F1290	F12920	F12921
F12922	F12929	F12950	F12951	F12959	F12980	F12988	F1299	F1310
F1311	F13120	F13121	F13129	F1314	F13150	F13151	F13159	F13180
F13181	F13182	F13188	F1319	F1320	F1321	F13220	F13221	F13229
F13230	F13231	F13232	F13239	F1324	F13250	F13251	F13259	F1326
F1327	F13280	F13281	F13282	F13288	F1329	F1390	F13920	F13921
F13929	F13930	F13931	F13932	F13939	F1394	F13950	F13951	F13959
F1396	F1397	F13980	F13981	F13982	F13988	F1399	F1410	F14120
F14121	F14122	F14129	F1414	F14150	F14151	F14159	F14180	F14181
F14182	F14188	F1419	F1420	F1421	F14220	F14221	F14222	F14229
F1423	F1424	F14250	F14251	F14259	F14280	F14281	F14282	F14288
F1429	F1490	F14920	F14921	F14922	F14929	F1494	F14950	F14951
F14959	F14980	F14981	F14982	F14988	F1499	F1510	F15120	F15121
F15122	F15129	F1514	F15150	F15151	F15159	F15180	F15181	F15182
F15188	F1519	F1520	F1521	F15220	F15221	F15222	F15229	F1523
F1524	F15250	F15251	F15259	F15280	F15281	F15282	F15288	F1529
F1590	F15920	F15921	F15922	F15929	F1593	F1594	F15950	F15951
F15959	F15980	F15981	F15982	F15988	F1599	F1610	F1611	F16120
F16121	F16122	F16129	F1614	F16150	F16151	F16159	F16180	F16183
F16188	F1619	F1620	F1621	F16220	F16221	F16229	F1624	F16250
F16251	F16259	F16280	F16283	F16288	F1629	F1690	F16920	F16921
F16929	F1694	F16950	F16951	F16959	F16980	F16983	F16988	F1699
F17200	F17201	F17203	F17208	F17209	F17210	F17211	F17213	F17218
F17219	F17220	F17221	F17223	F17228	F17229	F17290	F17291	F17293
F17298	F17299	F1810	F1811	F18120	F18121	F18129	F1814	F18150
F18151	F18159	F1817	F18180	F18188	F1819	F1820	F1821	F18220
F18221	F18229	F1824	F18250	F18251	F18259	F1827	F18280	F18288
F1829	F1890	F18920	F18921	F18929	F1894	F18950	F18951	F18959
F1897	F18980	F18988	F1899	F1910	F19120	F19121	F19122	F19129
F1914	F19150	F19151	F19159	F1916	F1917	F19180	F19181	F19182
F19188	F1919	F1920	F1921	F19220	F19221	F19222	F19229	F19230
F19231	F19232	F19239	F1924	F19250	F19251	F19259	F1926	F1927
F19280	F19281	F19282	F19288	F1929	F1990	F19920	F19921	F19922
F19929	F19930	F19931	F19932	F19939	F1994	F19950	F19951	F19959
F1996	F1997	F19980	F19981	F19982	F19988	F1999	F550	F551
F552	F553	F554	F558	O355XX0	O355XX1	O355XX2	O355XX3	O355XX4
O355XX5	O355XX9	O99320	O99321	O99322	O99323	O99324	O99325	P0441
P0449	P961	P962	T400X1A	T400X1D	T400X1S	T400X3A	T400X3D	T400X3S
T400X4A	T400X4D	T400X4S	T400X5A	T400X5D	T400X5S	T400X6A	T400X6D	T400X6S
T401X1A	T401X1D	T401X1S	T401X3A	T401X3D	T401X3S	T401X4A	T401X4D	T401X4S
T401X5A	T401X5D	T401X5S	T405X1A	T405X1D	T405X1S	T405X3A	T405X3D	T405X3S
T405X4A	T405X4D	T405X4S	T405X5A	T405X5D	T405X5S	T405X6A	T405X6D	T405X6S
T407X1A	T407X1D	T407X1S	T407X3A	T407X3D	T407X3S	T407X4A	T407X4D	T407X4S
T407X5A	T407X5D	T407X5S	T407X6A	T407X6D	T407X6S	T408X1A	T408X1D	T408X1S
T408X3A	T408X3D	T408X3S	T408X4A	T408X4D	T408X4S	T408X5A	T408X5D	T408X5S
T40901A	T40901D	T40901S	T40903A	T40903D	T40903S	T40904A	T40904D	T40904S
T40905A	T40905D	T40905S	T40906A	T40906D	T40906S	T40991A	T40991D	T40991S
T40993A	T40993D	T40993S	T40994A	T40994D	T40994S	T40995A	T40995D	T40995S
T40996A	T40996D	T40996S						

5.13 Suicide and intentional self-inflicted injury [662]

R45851	T1491	T360X2A	T360X2D	T360X2S	T361X2A	T361X2D	T361X2S	T362X2A
T362X2D	T362X2S	T363X2A	T363X2D	T363X2S	T364X2A	T364X2D	T364X2S	T365X2A
T365X2D	T365X2S	T366X2A	T366X2D	T366X2S	T367X2A	T367X2D	T367X2S	T368X2A
T368X2D	T368X2S	T3692XA	T3692XD	T3692XS	T370X2A	T370X2D	T370X2S	T371X2A
T371X2D	T371X2S	T372X2A	T372X2D	T372X2S	T373X2A	T373X2D	T373X2S	T374X2A
T374X2D	T374X2S	T375X2A	T375X2D	T375X2S	T378X2A	T378X2D	T378X2S	T3792XA
T3792XD	T3792XS	T380X2A	T380X2D	T380X2S	T381X2A	T381X2D	T381X2S	T382X2A
T382X2D	T382X2S	T383X2A	T383X2D	T383X2S	T384X2A	T384X2D	T384X2S	T385X2A
T385X2D	T385X2S	T386X2A	T386X2D	T386X2S	T387X2A	T387X2D	T387X2S	T38802A
T38802D	T38802S	T38812A	T38812D	T38812S	T38892A	T38892D	T38892S	T38902A
T38902D	T38902S	T38992A	T38992D	T38992S	T39012A	T39012D	T39012S	T39092A
T39092D	T39092S	T391X2A	T391X2D	T391X2S	T392X2A	T392X2D	T392X2S	T39312A
T39312D	T39312S	T39392A	T39392D	T39392S	T394X2A	T394X2D	T394X2S	T398X2A
T398X2D	T398X2S	T3992XA	T3992XD	T3992XS	T400X2A	T400X2D	T400X2S	T401X2A
T401X2D	T401X2S	T402X2A	T402X2D	T402X2S	T403X2A	T403X2D	T403X2S	T404X2A
T404X2D	T404X2S	T405X2A	T405X2D	T405X2S	T40602A	T40602D	T40602S	T40692A
T40692D	T40692S	T407X2A	T407X2D	T407X2S	T408X2A	T408X2D	T408X2S	T40902A
T40902D	T40902S	T40992A	T40992D	T40992S	T410X2A	T410X2D	T410X2S	T411X2A
T411X2D	T411X2S	T41202A	T41202D	T41202S	T41292A	T41292D	T41292S	T413X2A
T413X2D	T413X2S	T4142XA	T4142XD	T4142XS	T415X2A	T415X2D	T415X2S	T420X2A
T420X2D	T420X2S	T421X2A	T421X2D	T421X2S	T422X2A	T422X2D	T422X2S	T423X2A
T423X2D	T423X2S	T424X2A	T424X2D	T424X2S	T425X2A	T425X2D	T425X2S	T426X2A
T426X2D	T426X2S	T4272XA	T4272XD	T4272XS	T428X2A	T428X2D	T428X2S	T43012A
T43012D	T43012S	T43022A	T43022D	T43022S	T431X2A	T431X2D	T431X2S	T43202A
T43202D	T43202S	T43212A	T43212D	T43212S	T43222A	T43222D	T43222S	T43292A
T43292D	T43292S	T433X2A	T433X2D	T433X2S	T434X2A	T434X2D	T434X2S	T43502A
T43502D	T43502S	T43592A	T43592D	T43592S	T43602A	T43602D	T43602S	T43612A
T43612D	T43612S	T43622A	T43622D	T43622S	T43632A	T43632D	T43632S	T43692A
T43692D	T43692S	T438X2A	T438X2D	T438X2S	T4392XA	T4392XD	T4392XS	T440X2A
T440X2D	T440X2S	T441X2A	T441X2D	T441X2S	T442X2A	T442X2D	T442X2S	T443X2A
T443X2D	T443X2S	T444X2A	T444X2D	T444X2S	T445X2A	T445X2D	T445X2S	T446X2A
T446X2D	T446X2S	T447X2A	T447X2D	T447X2S	T448X2A	T448X2D	T448X2S	T44902A
T44902D	T44902S	T44992A	T44992D	T44992S	T450X2A	T450X2D	T450X2S	T451X2A
T451X2D	T451X2S	T452X2A	T452X2D	T452X2S	T453X2A	T453X2D	T453X2S	T454X2A
T454X2D	T454X2S	T45512A	T45512D	T45512S	T45522A	T45522D	T45522S	T45602A
T45602D	T45602S	T45612A	T45612D	T45612S	T45622A	T45622D	T45622S	T45692A
T45692D	T45692S	T457X2A	T457X2D	T457X2S	T458X2A	T458X2D	T458X2S	T4592XA
T4592XD	T4592XS	T460X2A	T460X2D	T460X2S	T461X2A	T461X2D	T461X2S	T462X2A
T462X2D	T462X2S	T463X2A	T463X2D	T463X2S	T464X2A	T464X2D	T464X2S	T465X2A
T465X2D	T465X2S	T466X2A	T466X2D	T466X2S	T467X2A	T467X2D	T467X2S	T468X2A
T468X2D	T468X2S	T46902A	T46902D	T46902S	T46992A	T46992D	T46992S	T470X2A
T470X2D	T470X2S	T471X2A	T471X2D	T471X2S	T472X2A	T472X2D	T472X2S	T473X2A
T473X2D	T473X2S	T474X2A	T474X2D	T474X2S	T475X2A	T475X2D	T475X2S	T476X2A
T476X2D	T476X2S	T477X2A	T477X2D	T477X2S	T478X2A	T478X2D	T478X2S	T4792XA
T4792XD	T4792XS	T480X2A	T480X2D	T480X2S	T481X2A	T481X2D	T481X2S	T48202A
T48202D	T48202S	T48292A	T48292D	T48292S	T483X2A	T483X2D	T483X2S	T484X2A
T484X2D	T484X2S	T485X2A	T485X2D	T485X2S	T486X2A	T486X2D	T486X2S	T48902A
T48902D	T48902S	T48992A	T48992D	T48992S	T490X2A	T490X2D	T490X2S	T491X2A
T491X2D	T491X2S	T492X2A	T492X2D	T492X2S	T493X2A	T493X2D	T493X2S	T494X2A
T494X2D	T494X2S	T495X2A	T495X2D	T495X2S	T496X2A	T496X2D	T496X2S	T497X2A
T497X2D	T497X2S	T498X2A	T498X2D	T498X2S	T4992XA	T4992XD	T4992XS	T500X2A
T500X2D	T500X2S	T501X2A	T501X2D	T501X2S	T502X2A	T502X2D	T502X2S	T503X2A

T503X2D	T503X2S	T504X2A	T504X2D	T504X2S	T505X2A	T505X2D	T505X2S	T506X2A
T506X2D	T506X2S	T507X2A	T507X2D	T507X2S	T508X2A	T508X2D	T508X2S	T50902A
T50902D	T50902S	T50992A	T50992D	T50992S	T50A12A	T50A12D	T50A12S	T50A22A
T50A22D	T50A22S	T50A92A	T50A92D	T50A92S	T50B12A	T50B12D	T50B12S	T50B92A
T50B92D	T50B92S	T50Z12A	T50Z12D	T50Z12S	T50Z92A	T50Z92D	T50Z92S	T510X2A
T510X2D	T510X2S	T511X2A	T511X2D	T511X2S	T512X2A	T512X2D	T512X2S	T513X2A
T513X2D	T513X2S	T518X2A	T518X2D	T518X2S	T5192XA	T5192XD	T5192XS	T520X2A
T520X2D	T520X2S	T521X2A	T521X2D	T521X2S	T522X2A	T522X2D	T522X2S	T523X2A
T523X2D	T523X2S	T524X2A	T524X2D	T524X2S	T528X2A	T528X2D	T528X2S	T5292XA
T5292XD	T5292XS	T530X2A	T530X2D	T530X2S	T531X2A	T531X2D	T531X2S	T532X2A
T532X2D	T532X2S	T533X2A	T533X2D	T533X2S	T534X2A	T534X2D	T534X2S	T535X2A
T535X2D	T535X2S	T536X2A	T536X2D	T536X2S	T537X2A	T537X2D	T537X2S	T5392XA
T5392XD	T5392XS	T540X2A	T540X2D	T540X2S	T541X2A	T541X2D	T541X2S	T542X2A
T542X2D	T542X2S	T543X2A	T543X2D	T543X2S	T5492XA	T5492XD	T5492XS	T550X2A
T550X2D	T550X2S	T551X2A	T551X2D	T551X2S	T560X2A	T560X2D	T560X2S	T561X2A
T561X2D	T561X2S	T562X2A	T562X2D	T562X2S	T563X2A	T563X2D	T563X2S	T564X2A
T564X2D	T564X2S	T565X2A	T565X2D	T565X2S	T566X2A	T566X2D	T566X2S	T567X2A
T567X2D	T567X2S	T56812A	T56812D	T56812S	T56892A	T56892D	T56892S	T5692XA
T5692XD	T5692XS	T570X2A	T570X2D	T570X2S	T571X2A	T571X2D	T571X2S	T572X2A
T572X2D	T572X2S	T573X2A	T573X2D	T573X2S	T578X2A	T578X2D	T578X2S	T5792XA
T5792XD	T5792XS	T5802XA	T5802XD	T5802XS	T5812XA	T5812XD	T5812XS	T582X2A
T582X2D	T582X2S	T588X2A	T588X2D	T588X2S	T5892XA	T5892XD	T5892XS	T590X2A
T590X2D	T590X2S	T591X2A	T591X2D	T591X2S	T592X2A	T592X2D	T592X2S	T593X2A
T593X2D	T593X2S	T594X2A	T594X2D	T594X2S	T595X2A	T595X2D	T595X2S	T596X2A
T596X2D	T596X2S	T597X2A	T597X2D	T597X2S	T59812A	T59812D	T59812S	T59892A
T59892D	T59892S	T5992XA	T5992XD	T5992XS	T600X2A	T600X2D	T600X2S	T601X2A
T601X2D	T601X2S	T602X2A	T602X2D	T602X2S	T603X2A	T603X2D	T603X2S	T604X2A
T604X2D	T604X2S	T608X2A	T608X2D	T608X2S	T6092XA	T6092XD	T6092XS	T6102XA
T6102XD	T6102XS	T6112XA	T6112XD	T6112XS	T61772A	T61772D	T61772S	T61782A
T61782D	T61782S	T618X2A	T618X2D	T618X2S	T6192XA	T6192XD	T6192XS	T620X2A
T620X2D	T620X2S	T621X2A	T621X2D	T621X2S	T622X2A	T622X2D	T622X2S	T628X2A
T628X2D	T628X2S	T6292XA	T6292XD	T6292XS	T63002A	T63002D	T63002S	T63012A
T63012D	T63012S	T63022A	T63022D	T63022S	T63032A	T63032D	T63032S	T63042A
T63042D	T63042S	T63062A	T63062D	T63062S	T63072A	T63072D	T63072S	T63082A
T63082D	T63082S	T63092A	T63092D	T63092S	T63112A	T63112D	T63112S	T63122A
T63122D	T63122S	T63192A	T63192D	T63192S	T632X2A	T632X2D	T632X2S	T63302A
T63302D	T63302S	T63312A	T63312D	T63312S	T63322A	T63322D	T63322S	T63332A
T63332D	T63332S	T63392A	T63392D	T63392S	T63412A	T63412D	T63412S	T63422A
T63422D	T63422S	T63432A	T63432D	T63432S	T63442A	T63442D	T63442S	T63452A
T63452D	T63452S	T63462A	T63462D	T63462S	T63482A	T63482D	T63482S	T63512A
T63512D	T63512S	T63592A	T63592D	T63592S	T63612A	T63612D	T63612S	T63622A
T63622D	T63622S	T63632A	T63632D	T63632S	T63692A	T63692D	T63692S	T63712A
T63712D	T63712S	T63792A	T63792D	T63792S	T63812A	T63812D	T63812S	T63822A
T63822D	T63822S	T63832A	T63832D	T63832S	T63892A	T63892D	T63892S	T6392XA
T6392XD	T6392XS	T6402XA	T6402XD	T6402XS	T6482XA	T6482XD	T6482XS	T650X2A
T650X2D	T650X2S	T651X2A	T651X2D	T651X2S	T65212A	T65212D	T65212S	T65222A
T65222D	T65222S	T65292A	T65292D	T65292S	T653X2A	T653X2D	T653X2S	T654X2A
T654X2D	T654X2S	T655X2A	T655X2D	T655X2S	T656X2A	T656X2D	T656X2S	T65812A
T65812D	T65812S	T65822A	T65822D	T65822S	T65832A	T65832D	T65832S	T65892A
T65892D	T65892S	T6592XA	T6592XD	T6592XS	T71112A	T71112D	T71112S	T71122A
T71122D	T71122S	T71132A	T71132D	T71132S	T71152A	T71152D	T71152S	T71162A
T71162D	T71162S	T71192A	T71192D	T71192S	T71222A	T71222D	T71222S	T71232A
T71232D	T71232S	X710XXA	X710XXD	X710XXS	X711XXA	X711XXD	X711XXS	X712XXA
X712XXD	X712XXS	X713XXA	X713XXD	X713XXS	X718XXA	X718XXD	X718XXS	X719XXA

X719XXD	X719XXS	X72XXXA	X72XXXD	X72XXXS	X730XXA	X730XXD	X730XXS	X731XXA
X731XXD	X731XXS	X732XXA	X732XXD	X732XXS	X738XXA	X738XXD	X738XXS	X739XXA
X739XXD	X739XXS	X7401XA	X7401XD	X7401XS	X7402XA	X7402XD	X7402XS	X7409XA
X7409XD	X7409XS	X748XXA	X748XXD	X748XXS	X749XXA	X749XXD	X749XXS	X75XXXXA
X75XXXXD	X75XXXXS	X76XXXXA	X76XXXXD	X76XXXXS	X770XXA	X770XXD	X770XXS	X771XXA
X771XXD	X771XXS	X772XXA	X772XXD	X772XXS	X773XXA	X773XXD	X773XXS	X778XXA
X778XXD	X778XXS	X779XXA	X779XXD	X779XXS	X780XXA	X780XXD	X780XXS	X781XXA
X781XXD	X781XXS	X782XXA	X782XXD	X782XXS	X788XXA	X788XXD	X788XXS	X789XXA
X789XXD	X789XXS	X79XXXXA	X79XXXXD	X79XXXXS	X80XXXXA	X80XXXXD	X80XXXXS	X810XXA
X810XXD	X810XXS	X811XXA	X811XXD	X811XXS	X818XXA	X818XXD	X818XXS	X820XXA
X820XXD	X820XXS	X821XXA	X821XXD	X821XXS	X822XXA	X822XXD	X822XXS	X828XXA
X828XXD	X828XXS	X830XXA	X830XXD	X830XXS	X831XXA	X831XXD	X831XXS	X832XXA
X832XXD	X832XXS	X838XXA	X838XXD	X838XXS	Z915			
(F11	F111	F1112	F1115	F112	F1122	F1125	F1128	F119
F1192	F1195	F1198	F12	F121	F1212	F1215	F1218	F122
F1222	F1225	F1228	F129	F1292	F1295	F1298	F13	F131
F1312	F1315	F1318	F132	F1322	F1323	F1325	F1328	F139
F1392	F1393	F1395	F1398	F14	F141	F1412	F1415	F1418
F142	F1422	F1425	F1428	F149	F1492	F1495	F1498	F15
F151	F1512	F1515	F1518	F152	F1522	F1525	F1528	F159
F1592	F1595	F1598	F16	F161	F1612	F1615	F1618	F162
F1622	F1625	F1628	F169	F1692	F1695	F1698	F17	F172
F1720	F1721	F1722	F1729	F18	F181	F1812	F1815	F1818
F182	F1822	F1825	F1828	F189	F1892	F1895	F1898	F19
F191	F1912	F1915	F1918	F192	F1922	F1923	F1925	F1928
F199	F1992	F1993	F1995	F1998	F55	O35XX)		

5.14 Screening and history of mental health and substance use codes [663]

R780	R781	R782	R783	R784	R785	R786	Z046	Z1330
Z1331	Z1332	Z1339	Z134	Z1340	Z1341	Z1342	Z1349	
Z72810Z72811		Z8651	Z8659	Z87891	Z91410	Z91411	Z91412	Z91419
Z9142	Z9149							
(R78	Z133	Z7281	Z9141)					

5.15 Miscellaneous mental disorders [670]

F061	F068	F440	F441	F442	F444	F445	F446	F447
F4481	F4489	F449	F450	F451	F4520	F4521	F4522	
F4529F4541		F4542	F458	F459	F481	F5000	F5001	F5002
F502F508		F5081	F5089	F509	F5101	F5102	F5103	F5104
F5105	F5109	F5111	F5112	F5113	F5119	F513	F514	F515
F518	F519	F520	F521	F5221	F5222	F5231	F5232	F524
F525	F526	F528	F529	F53	F54	F59	F640	F641
F648F649		F650	F651	F652	F653	F654	F6550	F6551
F6552	F6581	F6589	F659	F66	F6810	F6811	F6812	F6813
F688	F99	O906	R37	R4589	Z87890	Z9183		
(F44	F45	F50	F52	F64	F65	F681	Z8789)	

TEDS Substance Use Diagnosis Codes

291.00–291.99: Alcoholic Psychoses

292.00–292.99: Drug Psychoses

303.00–303.89: Alcohol Intoxication

303.90–303.99: Alcohol Dependence

304.00–304.09: Opioid Dependence

304.20–304.29: Cocaine Dependence

304.30–304.39: Cannabis Dependence

304.10–304.19,

304.40–304.99,

305.10–305.19: Other Substance Dependence 305.00–

305.09: Alcohol Use

305.60–305.69: Cocaine Use

305.20–305.29: Cannabis Use

305.50–305.59: Opioid Use

305.30–305.49,

305.70–305.99: Other Substance Use

APPENDIX F

Terms and Acronyms

Admission

Admission indicates the beginning of mental health service provision to a person with mental illness through programs under the auspices of the state mental health agency (SMHA). This includes new admissions (someone who has never received any services from the SMHA) and readmissions (someone who had previously received services from the SMHA, had been discharged, and started receiving services during the reporting period).

ADHD – Attention deficit/hyperactivity disorder

ADHD is a neurodevelopmental disorder defined by impairing levels of inattention, disorganization, and/or hyperactivity-impulsivity.

CBHSQ – Center for Behavioral Health Statistics and Quality

A center within the Substance Use and Mental Health Services Administration (SAMHSA) under the U.S. Department of Health and Human Services (HHS), CBHSQ is the lead government agency for behavioral health statistics, as designed by the Office of Management and Budget (OMB).

CD – Conduct disorder

CD is a serious behavioral and emotional disorder that can occur in children and teens. A child with this disorder may display a pattern of disruptive and violent behavior and have problems following rules.

DSM-5 – Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition

DSM-5 is the standard classification of mental disorders used by mental health professionals in the United States. DSM consists of three major components: the diagnostic classification, the diagnostic criteria sets, and the descriptive text [Source: <https://www.psychiatry.org/psychiatrists/practice/dsm>].

GED[®] – General Educational Development

GED is designed for people who, for various reasons, did not graduate from high school but want a certificate equivalent to the traditional high school diploma.

ICD-10-CM/ICD-9-CM – International Classification of Diseases, Tenth Revision, Clinical Modification/International Classification of Diseases, Ninth Revision, Clinical Modification

ICD-10-CM/ICD-9-CM is based on the World Health Organization's International Classification of Diseases (ICD-10/ICD-9). ICD-10-CM and ICD-9-CM are used as the official system of assigning codes to diagnoses and procedures associated with hospital utilization in the United States.

ODD – Oppositional defiant disorder

ODD is a childhood disorder that is characterized by negative, defiant, disobedient, and often hostile behavior toward individuals and authority figures primarily.

PDD – Pervasive developmental disorders

PDD refers to a group of conditions that involve delays in the development of many basic skills. Most notable among them are the ability to socialize with others, to communicate, and to use imagination.

Sources: Mental Health Client-Level Data (MH-CLD State Instruction Manual ([version 2.9](#)) and Combined Substance Use and Mental Health Treatment Episode Data Set (TEDS) State Instruction Manual With Data Submission System (DSS) Guide.

APPENDIX G

List of Contributors

This report was prepared for the Center for Behavioral Health Statistics and Quality (CBHSQ), Substance Abuse and Mental Health Services Administration, U.S. Department of Health and Human Services (HHS) under Contract No. 75S20320C00001.

Contributors and reviewers at SAMHSA, listed alphabetically by last name, include Herman Alvarado, Suparna Das, Hamid Ferdosi, Heydy Juarez (COR), Kasimu Muhetaer, and Karuna Nerurkar.